Standord Formulary

# Boston Medical Library 8 The Fenway

DR. W. S. POARDMAN: 57 HANCOCK ST., BOSTON, MASS.

DR. W. S. FORFTMAN; 67 HANGOCK ST., 8051CN, MASS. Digitized by the Internet Archive in 2011 with funding from Open Knowledge Commons and Harvard Medical School DR. VV. S. BOARDMAN: 57 HANCOCK ST., 80SIGN, MASS.



# THE

# STANDARD FORMULARY

A COLLECTION OF

NEARLY FIVE THOUSAND FORMULAS

FOR

PHARMACEUTICAL PREPARATIONS, FAMILY REMEDIES, TOILET
ARTICLES, VETERINARY REMEDIES, SODA FOUNTAIN
REQUISITES, AND MISCELLANEOUS PREPARATIONS ESPECIALLY ADAPTED
TO THE REQUIREMENTS
OF RETAIL DRUGGISTS.

TENTH EDITION, REVISED.

ALBERT E. EBERT, PH. M., PH. D.

AND

A. EMIL HISS, PH. G.

DR. W. S. BOARDMAN: 57 HANCOCK ST., 90510N, MASS.

CHICAGO:

G. P. ENGELHARD & CO.

1900.

22.5.319.

COPYRIGHT, 1897.
By G. P. Engelhard & Co.



# INTRODUCTORY.

No book is of such profit-making value in a drug store as a good formulary—a truly practical work which tells not only what to make, but how to make it. Such a book must be more than a mere compilation of formulas drawn from miscellaneous sources, many of them untried and untrustworthy. A score of carefully selected and thoroughly tested formulas are worth a thousand of the kind embodied in most formularies, which usually consist in great part of mere clippings from current journals or of random and untested selections from prior publications. Pharmacists who possess formulas of merit cling to them with the greatest tenacity; this being one feature which has assisted in fostering the creation of monopoly preparations—the bane of pharmacy of to-day.

This formulary contains a vast collection of formulas, covering everything the pharmacist may desire to make, and every one, it is believed, thoroughly trustworthy. A large proportion have been collected in years past, as the outgrowth of practical experience, and are now offered to American pharmacists for the first time. Dieterich's and Hager's celebrated manuals—the great German authorities—are, by discriminating translations and adaptations, for the first time made accessible to the pharmacists of this country. The British, German, French, Swedish, Norwegian, Belgian, Austrian, and Italian pharmacopæias, Parrish's Pharmacy, the Eclectic Dispensatory, and the National Formulary have contributed their best and most useful formulas, while completeness has been assured by selections of the best from other authorities. While the policy has been not to burden the book with references to the sources of all formulas, due credit has been given in each case as opportunity permitted or equity required.

Attention is called here to the various divisions of the work. Part I embraces what are commonly known as pharmaceutical preparations, containing selections from the various pharmacopæias, the Eclectic Dispensatory, the National Formulary, and other authoritative works. Part III is designed to give the ingredients and quantities of preparations similar to the leading proprietary medicines of the market in order that druggists may know what they are called upon to dispense or sell. The introductory note to the chapter more fully explains its purpose. Part VII embraces such formulas as could not properly be included in the remaining division.

It is believed that the quantities of all formulas have been adapted to the usual needs of pharmacists. Preparations which are usually required in amounts of one pint or one pound bear formulas indicating this amount. If they are commonly prepared in smaller or in larger amounts, the quantities have been duly decreased or increased. Quantities of a formula are usually even amounts, such as 4 ounces, 8 ounces, 16 ounces, one-half gallon, etc. Due consideration has been given the fact that foreign preparations frequently differ in strength and specific gravity from similar preparations as made in this country;

#### INTRODUCTORY.

for example, solution of iron chloride of the German pharmacopæia represents 10 per cent. of metallic iron, and has a specific gravity of 1.28, whereas the similar preparation of the United States pharmacopæia represents 13 per cent. of metallic iron, and has a specific gravity of 1.387. This fact has necessitated almost constant recalculation of formulas taken from foreign sources, and sometimes even the entire recasting of formulas, in order that the product made by the use of American preparations shall be the same as that designed by the original formula.

It is the design of this Formulary that all liquids be measured and all solids be weighed. Hence, foreign formulas usually required the conversion of parts of liquids into volume, and this again necessitated consideration of specific gravities.

Owing to the greater familiarity of American pharmacists with the old system of grains, ounces, etc., quantities are all expressed in these in preference to the metric system. Troy ounces and pounds being now practically obsolete, these weights are replaced by equivalent quantities in the avoirdupois system. Liquid quantities are expressed in the apothecaries' system. Frequently it was considered advisable to alter working directions for making foreign preparations; for example, by the substitution of percolation for maceration.

It is especially advised that pharmacists carefully peruse the introductories to the various chapters or divisions. Failure to produce satisfactory results may be caused by neglect of this precaution. The introductories in Part V are especially complete.

Taken all in all, this volume can not fail, it is believed, to be of incalculable service to every pharmacist who sees in a properly utilized laboratory a means of asserting his personal independence of other's preparations and for extending his business and professional reputation while adding in an important measure to his profits.

THE EDITORS.

#### NOTE TO LATER EDITIONS.

The editors desire to express their sincere appreciation of the very flattering reception accorded this work by pharmacists and others in all parts of the country. It is believed that with the additions and other changes embodied in the successive revised Editions, the book will be found in even closer harmony with the demands, not only of scientific pharmacy, but of the retail drug trade in its relations to the ever vital problem of profit and loss.

The Index to the volume has been adjudged by several critics not sufficiently extensive to facilitate ready reference. This possible objection has, it is believed, been answered in the rigid alphabetical arrangement adopted for the chief classes of preparations including the Elixirs, Extracts, Fluid Extracts, Liniments, Ointments, Powders, Solutions, Syrups Tinctures, Wines, etc. With this fact in mind, reference to any desired preparation will be attended with little difficulty.

The work is undergoing constant revision, in order that the successive editions may be kept fully up to date in every department. Suggestions or contributions to this end will be highly appreciated.

THE EDITORS.

CHICAGO, January 2, 1897.

# TABLE OF CONTENTS.

TABLE OF ABBREVIATIONS		
PART I.		
PHARMACEUTICA	L PREPARATIONS.	
Abstracts 11	Juices 107	
Acids	Kneipp's remedies 108	
Arquebusade 12	Lards	
Balsam 12	Liniments	
Bougies	Lotions 111	
Camphor preparations	Mixtures 112	
Catguts 17	Mucilages	
Caustics	Oils	
Cerates	Ointments	
Chlorodyne	Oleates	
Chloroforms	Pastes	
Collodions	Pills	
Concentrations	Plasters 121	
Confections	Powders 121	
Copper preparations 24	Rademacher's preparations 123	
Cordials	Salts	
Cottons	Silks	
Decoctions	Snuffs	
Discs	Solutions 129	
Drops	Species 135	
Electuaries 32	Spirits	
Elixirs 32	Sponges	
Emulsions	Suets	
Essences 85	Sugars	
Extracts 86	Syrups	
Fluid extracts 86	Thompsonian remedies 150	
Glycerites 104	Tinctures 150	
Honeys	Transfusion fluids	
Infusions	Vinegars	
Inhalations	Waters 159	
Injections	Wines 161	
PART II.		
HOUSEHOLI	REMEDIES.	
Ague cures 164	Burns and scalds, applications for. 171	
Asthma remedies 164	Catarrh and cold in the head, reme-	
Barber's itch remedies 166	dies for	
Bitters 166	Cathartics	
Blood purifiers 168	Chilblain cures	
Bunion cures	Cholera remedies	

# TABLE OF CONTENTS.

Corn eradicators	Liniments 190 Liver remedies 192 Nervous debility remedies 192 Neuralgia remedies 194 Nipples, fissured, cures for 194 Ointments or salves 194 Pile remedies 195 Rheumatism and gout remedies 196 Ringworms, applications for 197 Soothing and teething remedies 197 Syphilis remedies 198 Throat affections, remedies for 198 Tonics 199 Toothache remedies 200 Vermifuges 202 Wart eradicators 204 Miscellaneous 205	
PART III.		
PROPRIETARY PREPARATIONS	207	
PAR'	r IV.	
VETERINARY 1	PREPARATIONS.	
	Swine medicines	
	Dog and cat medicines	
	Poultry medicines	
Dird incdicines	·················· ~**	
DAD	71. 37	
PAK	т v.	
TOILET PRE	EPARATIONS.	
SECTION IDISTINCTIVELY ODOR	Bay rum 306	
PREPARATIONS.	Toilet vinegars	
Perfumes	Smelling salts	
Handkerchief extracts	SECTION II.—SKIN PREPARATIONS.	
Essences	Ointments and creams 309	
Spirits	Toilet milks         316           Tollet lotions         317	
Tinctures	Toilet jellies	
Sachet powders 294	Camphor ice	
Pot-pourris	Toilet lanolins	
Fumigating pastilles, powders, paper,	Almond paste	
essences, vinegar, etc 300	Almond meal 322	
Cologne waters 302	Toilet powders	
Toilet waters 304	Liquid cosmetics 325	

# TABLE OF CONTENTS.

Face bleach 326	Hair tonics 344
Remedies for blackheads, freckles,	Hair promoters 345
pimples, tan and sunburn326-329	Hair dyes 347
Rouges 330	Brilliantines 349
Grease paints 331	Mustache pomades349, 350
Depilatories	SECTION IV MOUTH PREPARATIONS.
Liquid soaps 333	Tooth powders351, 352, 356, 357, 358
Shaving creams and powders 333	Tooth powders351, 352, 357, 358
Manicure cosmetics	Tooth washes
SECTION III. —PREPARATIONS FOR THE	Tooth washes 354
HAIR, SCALP, AND MUSTACHE.	Mouth washes
Shampoos and sea foams 334	Lip salves or pomades
Hair oils	Cachous
Hair pomatums	SECTION V.—BATH PREPARATIONS.
Bandolines	
Hair-curling liquid 341	Bath powders
Remedies for dandruff, baldness, and	Bath tablets 362
loss of hair	Sulphur baths 362
PAR	T VI.
SODA WATER	PREPARATIONS.
Colors	Root beer
	Soda foam
	Syrups
	Tonics
	Vinegars 371
Mineral waters	371–373
, and the second	
PART	r VII.
MISCELLANEOUS PREPARATIONS	5 374–468

DR. VV. S. BOANDMAN: 57 HANGOCK ST.: 80510N, MASS.

Marine Commence of the Commenc

# TABLE OF ABBREVIATIONS.

Dieterich's ManualD.	French PharmacopœiaCodex.	
Hager's ManualH.	Norwegian PharmacopœiaNorw. Pharm.	
American DispensatoryEclectic.	Swedish PharmacopæiaSwed. Pharm.	
German PharmacopæiaGerm. Pharm.	Austrian PharmacopæiaAustr. Pharm.	
German Formulary	Belgian Pharmacopæia Belg. Pharm.	
British Pharmacopœia Brit. Pharm. Italian Pharmacopœia Ital. Pharm.		
British FormularyBrit. Form.	United States PharmacopœiaU. S. P.	
National FormularyN. F.		

#### PART I.

# PHARMACEUTICAL PREPARATIONS

#### Abstracts.

These preparations were first introduced into the United States Pharmacopœia of 1880, and were dropped from the edition of 1890. They are used to some extent; the following is a general process for their manufacture:

Drug, No. 60 powder.....av.oz. 8 Menstruum.

Powdered sugar of milk of each, sufficient

Moisten the drug with menstruum, and pack, macerate, and extract in the usual way for making fluid extracts, reserving the first 61/4 fluidounces of percolate obtained and continuing percolation until the drug is exhausted. Evaporate the second percolate, at a temperature not exceeding 50 degrees C., to 11/4 fluidounces; mix this with the reserved percolate, place the whole in a broad evaporating dish or other suitable vessel; add 4 av. ounces of milk sugar, mix well, cover the vessel with a piece of thin muslin gauze to exclude dust, and set the whole aside in a warm place, where the temperature will not rise above 50 degrees C., until the mixture is dry. Then add enough milk sugar to make the whole weigh 4 av. ounces, reduce the whole to a uniform fine powder, and keep in a well-stopped bottle.

Essentially, therefore, the process of manufacture consists in making a fluid extract, evaporating this to a dry solid extract, and adding enough milk sugar to make up a quantity of one-half of the weight of the original drug.

Eleven abstracts were official, viz., aconițe, belladonna (root), conium (fruit), digitalis, henbane (leaves), ignatia, jalap, nux vomica, podophyllum, senega and valerian. The menstruum usually employed in extracting the drugs was alcohol, the exceptions being nux vomica and ignatia, in which the menstruum was a mixture of alcohol and water in the proportion of eight of the former to one of the latter, and conium, in which the first four fluidounces of alcohol used as men-

struum is mixed with three fluidrams of diluted hydrochloric acid.

Acid, Carbolic, Camphorated. (Phenol Camphor.—Carbolized Camphor.)

Camphor, coarse powder...av.oz. 10
Carbolic acid, crystal...av.oz. 3½
Alcohol....fl.oz. ½

Triturate together until an oily liquid is obtained, or mix in a bottle and agitate frequently until solution occurs.

# Acid, Carbolic, No. 33.

This is a dilution of carbolic acid recommended as more convenient and safe to use than liquefied carbolic acid, i.e., crystal carbolic acid melted and maintained in a liquid state by the addition of 5 per cent of water. It is prepared as follows:

Carbolic acid, crystal.av.oz. 2 or fl.oz. 2 Glycerin.....av.oz. 5 or fl.oz. 4

Melt the acid and add the glycerin.

This No. 33 acid mixes readily with water in all proportions, and, not being as caustic as the ordinary liquefied acid, cannot result in as much mischief or fatality if used improperly, or if taken accidentally or purposely.

# Acid, Hydrocyanic, Scheele's.

This is to be prepared from potassium ferrocyanide and sulphuric acid according to the process of the U. S. Pharmacopœia. It should contain 4 per cent. of absolute hydrocyanic acid.—Brit. Form.

# Acid, Sulphocarbolic, Crude.

Carbolic acid, crude......fl.oz. 5 Sulphuric acid, commercial...fl.oz. 1½ Water......fl.oz. 9

Pour the carbolic acid into an earthenware jar surrounded by cold water; add the sulphuric acid in a fine stream, stirring constantly; then dilute this mixture with the water, also added gradually. Any marked rise in temperature should be avoided. It may be necessary to keep the temperature down by the use of ice added to the cold water.

This forms an economical and effective dis-

infectant for cesspools, urinals, sewers, etc.

—D.

#### Alcohol, Deodorized.

Many methods for deodorizing alcohol have been recommended, but the following will be found satisfactory:

A convenient amount of alcohol is shaken with powdered potassium permanganate until it assumes a decided color. Then allow to stand for several hours until the permanganate has become decomposed, and brown manganese dioxide has deposited. A pinch of pulverized calcium carbonate should then be added, and the whole subjected to distillation, using a well-cooled receiver. Distil very slowly at first, testing the distillate frequently, until a mixture of the distillate and a strong (syrupy) solution of pure caustic soda or potassa, in the proportion of 10 of the former to 1 of the latter, gives no perceptible yellow coloration, on standing for 20 minutes or half an hour. The first portion of distillate that yields this coloration should be rejected; the last one-eighth of liquid should not be distilled, and should also be rejected. The remaining portion only is adapted for use.

This alcohol is adapted to all chemical purposes and for use in the manufacture of perfumes.

#### Alcoolats.

These are a class of French preparations produced by distillation of drugs with alcohol. Balsamam (or baume de) floravanti is an example.

#### Alcoolatures.

A class of French preparations produced by the action of alcohol upon fresh plant parts. They correspond to the tinctures of fresh herbs of our pharmacopœia.

#### Alcooles.

A class of French preparations which consist of alcoholic solutions of volatile oils, and therefore correspond to the spirits of our pharmacopæia.

# **Arquebusade.** (Brown Arquebusade. Wund-Wasser.)

Acetic acid, dilutedfl.oz. 81/4
Diluted alcoholfl.oz. 41/2
Sulphuric acid, dilutedfl.oz. 14
Clarified honeyfl.oz. 2
Mix and filter.—H. and D.

# Arquebusade, White.

Oil of sage	drops	7
Oil of wormwood	drops	7
Oil of rue		7
Oil of peppermint	drops	7
Oil of rosemary	drops	7
Oil of marjoram	drops	7
Oil of lavender flowers	drops	7
Alcohol		
Water		

Dissolve the oils in the alcohol and then add the water.—H.

# Balsam, Blackberry.

•	
Fluid extract of blackberry root.fl.oz.	2
Fluid extract of geraniumfl.oz.	1
Tincture of gingerfl.oz.	1
Syrup of rhubarb, aromaticfl.oz.	4
Oil of cinnamondrops	5
Oil of nutmegsdrops	5
Oil of cloves, oil of pimento,	
eachdrops	10
Cimple distance of the second	

Simple elixir, enough to make.fl. oz. 16 Mix the oils with the tinctures; add the syrup and elixir, and filter.

Refer also to page 180.

# Balsam, Friar's. (Traumatic Balsam.— Turlington's Balsam.)

8 ,		
Benzoin	.av.oz.	11/2
Storax	.av.oz.	1/2
Balsam of tolu	.av.oz.	1/2
Balsam of Peru	.av.oz.	1/4
Myrrh		60
Aloes		
Angelica root		
Alcohol		

Macerate for ten days, and filter. The compound tincture of benzoin may be substituted for the above.

# Balsam Fioravanti. (Baume de Fioravanti.—Spiritus Balsamicus,)

Peru balsam	drops 15
Rectified oil of turpenti	
Oil of cassia	drops 15
Oil of cloves	drops 15
Oil of juniper berries	drops 15
Oil of mace	drops 15
Oil of thyme (white)	drops 15
Alcoholenough to	make fl.oz. 16.
	—Н.

This is a simpler and more rational formula than that of the Codex, which requires distillation.

# Balm of Gilead, Factitious. (Artificial Balsam of Mecca.)

The original is an oleoresin derived from a tree growing on the shores of the Red Sea.

Imitations are much more common and are prepared as stated below:

| Benzoin, coarsely powdered av.oz. 2 | Liquid storax | av.oz. 1½ | Tolu balsam | av.oz. 1 | Balsam of fir | av.oz. 12

Place in a glass flask or bottle, and subject to the heat of a water bath for several hours; agitate frequently until liquefied; allow to cool, and decant the clear portion, to which add sufficient of the oils of lemon, cassia, rosemary, and nutmeg and vanilla extract to give it a strong aromatic odor.

II.

Benzoin, coarsely powderedav.oz.	1
Peru balsamav.oz.	1
Vanilla, cut smallgr.	60
Nutmeg, brokengr.	
Balsam of firav.oz.	

Digest the whole as above, decant, and to decanted liquid add same essential oils.

#### Balsam of Guaiac.

Guaiac resinav.oz. 8	
Peru balsamfl.dr. 1½	
Alcoholfl.oz. 10	
Macerate for 7 days and strain.	

An old remedy for rheumatism, ague, etc. Dose, ¼ to 1 teaspoonful.

# Balsam of Honey. (Pectoral Balsam.)

Opium, powderedgr. 1	5
Tolu balsamgr. 1	5
Storaxgr. 4	5
Honeyav.oz.	
Diluted alcoholfl.oz. 1	6

#### Balsam de Maltha. (Balsam di Malta.)

Benzoin, powderedav.oz.	134
Peru balsamav.oz.	1
Aloesgr.	120
Alcoholfl.oz.	16

Macerate for 7 days and filter.

#### Balsam, Metz's.

Linseed oil, olive oil, eachav.oz.	6
Oil of laurel berries, expressed av.oz.	1
Oil of turpentinefl.oz.	2
Aloes. powdered dr.	2
Verdigris, powdereddr.	3
Sulphate of zinc, powdereddr.	11/2
Oil of juniperfl.oz.	1/2
Oil of clovesfl.dr.	1

Melt the oils by gentle heat, and apply the powders as a dressing to wounds and ulcers.

Balsam, Nutmeg. (Balsamum Nucistæ.

--Muskat Balsam.--Magen Balsam.-Nutmeg Cerate.)

 Yellow wax
 av.oz.
 1¾

 Olive oil
 av.oz.
 3½

 Expressed oil of nutmeg
 av.oz.
 10½

Melt the wax and add the oils.—Germ. Pharm.

A cheaper preparation can be prepared according to the following formula:

Olive oilfl.oz.	6
Yellow waxav.oz.	2
Spermacetiav.oz.	
Expressed oil of nutmegav.oz.	
Alkanetgr.	3
Annattoigr.	15
Alcohol fl.dr.	2

Melt the wax and spermaceti, add the olive, divide it into two parts; in one portion, digest the alkanet for 5 minutes, add the nutmeg oil and strain; triturate the annatto with the alcohol, digest the mixture with the second portion of oily liquid for 5 minutes, strain, add this to the first colature, mix well, and pour into molds if desired. Of course, a cheaper preparation may be produced by using a cheaper oil than olive oil, such as cottonseed or benne oil.—D. modified.

#### Balsam, Riga, Factitious.

The genuine, which is derived from a tree grown in northern Europe and Asia, is scarcely ever seen in this country, and the following mixtures are used as imitations:

I.

Oleo-balsam mixture, N. F. fl.oz. 14
Spirit of sage (1 of oil to 49 of alcohol)......fl.oz. 1½
Tincture of Spanish saffron...fl.dr. 3
—H.

#### Balsam of Soap.

White castile soap, powdered av. oz.	2
Camphorav.oz.	2
White oil of thymefl.dr.	2
Acetic etherfl.oz.	

Mix and digest in a closed vessel at a gentle heat until dissolved, and decant the clear portion.

# 

Boil together in an iron vessel, stirring constantly, until a uniform liquid is obtained, being cautious in regulating the heat so that the liquid will not boil over.

#### Balsam of Turpentine.

Olive oilfl.oz.	6
Oil of turpentinefl.oz.	2
Yellow waxav.oz.	1
Peru balsamfl.dr.	
Camphorgr.	
Essential oil of nutmegfl.dr.	

Melt the wax, add the olive oil, and then incorporate the other ingredients.

#### Balsam, Universal.

Liniment of camphorfl.oz.	2
Infused oil of henbanefl.oz.	
Cottonseed oilfl.oz.	2
Yellow waxav.oz.	2
Solution of lead subacetatefl.oz.	11/2

Melt the wax, add the oils, allow to cool, and when fairly cool, thoroughly incorporate the lead solution with the mixture.

Under the above title, many kinds of preparations are offered, but the formula given is believed to be the most sensible and will furnish as good a product as any.—D.

#### Bandages, Plaster of Paris.

These are made by taking gauze of suitable kind and of the width desired, and rolling it up just as in making roller bandages, and while doing so, sprinkling over it freshly burned plaster of Paris sufficient to fill the pores. The bandage is then to be wrapped in waxed paper and put into a tin box, or it is put directly into the box, which latter should then be well closed.—D.

#### Benzin, Deodorized.

Mix together 8 fluidounces sulphuric acid and 56 fluidounces of water and when cold pour it into a two-gallon bottle; add 1 av. ounce potassium permanganate and agitate until dissolved; then add 1 gallon of benzin and thoroughly agitate, and allow the mixture to remain in contact for twenty-four hours, frequently agitating. Separate the benzin and wash in a similar bottle with a mixture of 120 grains of potassium permanganate, 240 grains of caustic soda, and

32 fluidounces of water, agitating frequently during several hours. Then separate the benzin and wash it thoroughly with water.

On agitating the benzin with the acid permanganate solution, an emulsion-like mixture is produced which separates in a few seconds, the permanganate solution slowly subsiding and showing considerable reduction.

The quantity of permanganate necessary is in direct proportion to the impurities existing in the benzin. The quantity ordered in the formula is sufficient for a very crude article and may be reduced when manipulating with a purer distillate.

# Bitters, Thompsonian. ("Number Four")

Barberry bark, Balmony,

Poplar bark......of each, equal parts.

Other Bitters will be mentioned in Parts II and III.

# Blood, Dried, Defibrinated. (Sanguis Bovinus Inspissatus or Exsiccatus.)

This may be prepared by evaporating fresh defibrinated bullock's blood on a water bath, stirring constantly, until it assumes a granular condition; then spread on glass plates and keep at a temperature of 35 to 40 degrees C. (in a drying oven) until perfectly dry, after which it is powdered and put into well-stopped bottles.—D.

The defibrinated blood may be prepared by vigorously beating fresh blood in a broad dish with a stick or twig until there is no further separation of fibrin.

# Bougies. (Urethral Suppositories.)

These are prepared from three different kinds of materials, the first kind being prepared with cacao butter, and are known as "cacao butter" bougies; the second with gelatin and glycerin, and are known as "gelatin" bougies, and the last with mucilage, and are known as "gum bougies."

# Bougies, Cacao Butter.

The mass for these is prepared by mixing the medicating substance with grated cacao butter, adding a little petrolatum or bland fixed oil. To form the bougies from this, the mass is put into a bougie syringe made of metal, having an opening below like an ordinary syringe, and having a piston which screws down instead of simply pushing down. When the mass is prepared, it is packed into the syringe after removing the piston; the latter is replaced and is slowly screwed downward. As the piston moves forward, the mass begins to make its exit at the opening below in the form of a slender cylinder. This cylinder may subsequently be cut into suitable lengths. In the absence of a syringe, the bougies may be fashioned by rolling them out on a board just as a pill mass is rolled into a "pipe."

Providing the medicating substance is not in powder form, and cannot conveniently or advantageously be reduced to this form, the method of mixing is not so simple as outlined above. If something like an extract is to be incorporated, this must first be softened with water, glycerin, diluted alcohol, or alcohol, after which it may be mixed with the cacao butter as before; or it may be that some fixed oil or other substance will be a more suitable softening agent. Possibly the substance is soluble in the cacao butter in a melted condition; it should, of course, be so dissolved, and after cooling and hardening, the mass should be reduced to powder by grating or otherwise, mixed possibly with a little petrolatum or fixed oil, and be fashioned into bougies as before. Large quantities of liquids cannot, of course, be incorporated with bougie masses; in many instances, however, it will be possible to concentrate the liquid by evaporation without injury to its medicinal principles; then, subsequently, the concentrated residue may be mixed with the cacao butter as before.

These bougies have the disadvantage of being brittle, and hence are used but little.

#### Bougies, Gelatin.

The best kind of gelatin to use in this process is the purest white of the kind known as "French." The mass employed is made from gelatin, glycerin and water, but the proportion cannot be the same in all cases, as the nature of the medicating substance to be combined may require a modification; also the gelatin may vary to some extent, and modification of the proportions may be required on this account.

dered, it is best to keep a supply of several suitable gelatin masses. These should preferably be kept in glass vessels, and be covered with a thin layer of alcohol to prevent moulding, the whole, of course, being well covered. When wanted for use, the alcohol should be poured from the surface of the mass and the adhering alcohol removed by wiping with a tuft of absorbent cotton, after which the mass may be removed either by cutting out a piece with a knife or by melting in a water bath and pouring out the liquefied mixture.

To prepare the mass the following plan should be adopted:

Soak the requisite quantity of gelatin in a portion of the distilled water contained in a porcelain vessel for several hours, or until it is thoroughly softened; add enough water to make 3, 4 or 5 times the weight of original gelatin; add the prescribed quantity of glycerin; heat on a water bath with frequent stirring until dissolved, and then continue heating to remove excess of water.

Medicaments added to gelatin bougie masses may be (1) without any influence upon the latter, or they may (2) cause it to become tenacious, or they may (3) render it thin or unctuous. Examples of the different kinds are appended, also of the different mixtures of glycerin, gelatin and water for use in various cases.

#### Hard Glycerin-Gelatin Mass.

Gelatinav.o.	z. 3
Waterfl.os	z. 9
Glycerinfl.o	z. 5

This mass is to be prepared as directed, and is then to be evaporated until it weighs 12 av.ounces.

#### Soft Glycerin-Gelatin Mass.

Gelatin Water		 			 		 	.av.oz	11/2
Water					 			.fl.oz.	41/2
Glycerin.		 		_		Ĺ		fl.oz.	4

Evaporate this also until it weighs 10 av.ounces.

In preparing the bougies, the medicating substance must be added either in the form. of fine powder, or in the form of a concentrated solution. If it be an extract or similar substance, it is to be softened with water, glycerin, etc., just as in the case of the cacao Where gelatin bougies are regularly or- butter bougies. As in making the latter, no

large volume of liquid can be incorporated into the mass. A larger volume, however, can be incorporated with the gelatin mass than with the cacao butter mass. If necessary, after adding liquid, the consistence of the mass may be restored by a little powdered tragacanth.

Having melted the gelatin mass by aid of a water bath and added the medicaments, the whole may be poured into molds. fore doing so, it may frequently be of advantage to heat the mixture for a few moments carefully over a naked flame so as to render it still more fluid. During heating, the mixture is stirred carefully to liberate all air bub-This prepared mixture is now poured quickly into molds.

The molds used are of the hinged kind, so that they may be opened and the bougies taken out; they are constructed of brass, block fin, or nickel-plated iron. Before using, the interior of the molds must be rubbed with petrolatum or oil to prevent adhesion of the mass; then they must be made quite warm or, in the case of rather hard masses, even quite hot. This warming is necessary to permit escape of air bubbles. The hot mass may then be poured into the molds. The only time an exception is made to pouring the mass in a hot condition into the molds, is when the medicament is an entirely insoluble solid; then the latter must be triturated to an exceedingly fine powder and added to the mass. Before pouring the latter into molds it should be allowed to cool sufficiently, so that when dropped upon a cold stone it will congeal almost immediately. This mass should be poured into the cold molds.

After pouring the mixture into molds, the latter are placed upon ice. After cooling, the mold should be opened and the bougies taken out and exposed to the air for several hours, that they may harden on the surface, after which they should be wrapped in waxed paper and laid horizontally in boxes. The bougies may also be kept in a box rolled in lycopodium.

In the absence of a mold, gelatin bougies suitable size. Oil this tube by sucking into flame, pour into the hot molds, allow to

it a small quantity of olive oil or liquid petrolatum and allowing this to run out again. Now place the tube into the gelatin mass, suck up the latter to the desired height, cover the upper end quickly with the finger, and place the lower end on ice until the lower portion of the mass has solidified; now remove the finger and lay the tube down on the ice in an inclined position. When the bougie has hardened it may be removed by pushing it out with a smaller glass tube or a rod of some kind.

Examples of the three kinds of mixture in which the consistency of the bougie mass is either unaltered or altered:

I.—The consistency is unaltered:

# Bougies, Silver Nitrate.

Silver nitrate.....gr. 7 Distilled water.....sufficient. Hard glycerin-gelatin mass....av.oz. 8

Dissolve the silver nitrate in a few drops of water, add this solution to the melted gelatin mass, and form into bougies in the prescribed manner.

These bougies contain one-half per cent of the silver salt. They soon become discolored and must be made fresh .- D.

## Bougies, Chloral Hydrate.

Chloral hydrate.....gr. 23 Hard glycerin-gelatin mass....av.oz. 1

Pulverize the chloral hydrate, add to the melted gelatin mass, stir until dissolved, and pour into bougies. The product contains 5 per cent of chloral hydrate.-D.

#### Bougies, Iodoform.

These are to be prepared like the chloral hydrate bougies.-D.

#### Bougies, Potassium Iodide.

Prepare these like the chloral hydrate bougies.-D.

2. The mass becomes tenacious:

## Bougies, Alum.

Alum, powdered.....av.oz. Glycerite of tragacanth.....av.oz. 21/2 Soft glycerin-gelatin mass....av.oz. 7

Triturate the alum with the glycerite to a smooth paste, add to the melted gelatin mass, may be formed by means of a glass tube of heat the whole for a moment over a naked

stand for a moment, and then cool rapidly by placing upon ice.

The product contains 5 per cent. of alum.

#### Bougies, Ferric Chloride.

Solution of ferric chloride....fl.dr. 51/2 Glycerite of tragacanth.....av.oz. 11/4 Soft glycerin-gelatin mass...av.oz. 31/2

Melt the glycerite and gelatin mass, add the iron solution, and then proceed as with the alum bougies.—D.

The product contains 10 per cent. of the iron solution.

3. The mass becomes thin or unctuous:

#### Bougies, Tannin.

Tanningr.	100
Alcohol fl.oz.	
Tragacanth, powderedgr.	30
Hard glycerin-gelatin massav.oz.	4

Dissolve the tannin in the alcohol, triturate this with the tragacanth, incorporate with the melted gelatin mass, expose to the heat of a water bath until the alcohol has evaporated, pour into molds, and cool as rapidly as possible. The product contains 5 per cent of tannin.-D.

#### Bougies, Gum.

These may be prepared from mixtures of powdered tragacanth, starch, dextrin, sugar and medicament, rubbed to a paste with water and glycerin, or they may be prepared by massing the medicating ingredient with mucilage of acacia, glycerin and water, and rolling into the proper form. If the bougies contain too large a proportion of medicating ingredient, the latter may be diluted with some inert or harmless body; powdered boric acid will serve acceptably.

#### Camphor Cream.

White castile soapgr. 19	05
Boiling watersufficien	
Ammonium carbonategr. 12	0.
Camphor, powdered gr. 12	
Tincture of opiumfl.dr.	2
Oil of origanumfl.dr.	
Ammonium carbonategr. Camphor, powderedgr. Tincture of opiumfl.dr.	

Dissolve the soap in the boiling water, allow the solution to cool, add the remaining ingredients, and mix well.

Sometimes this is made with double the amount of soap and is also made to contain 1 fluidounce of oil of turpentine to the pint.

A formula for a toilet preparation by the

name of "cream of camphor," may also be found in Part III.

# Camphor, Chloral. (Camphorated Chloral.)

Chloral.....av.oz, 4 Camphor . . . . . . . . . . . . . . av.oz. 4

Mix by agitation in a bottle or trituration in a warm mortar until liquefied and combined.—N. F.

# Camphor Julep, Thompsonian.

Camph	or										.gr.	60
Myrrh		 									gr.	210
Sugar.		 									.gr.	120
Water.		 								fl	.oz.	4

# Camphor Salicylate.

Camphor								.av.oz. 9	
Salicylic acid.								.av.oz. 7	

Reduce the camphor to powder in the usual way and mix well with the acid.

#### Catgut.

This is prepared from the intestines of different animals, but usually the sheep. To prepare the gut, the intestines are cleaned, freed from fat, and steeped for some time in water, after which the external membrane is scraped off with a blunt tool like the back of a knife. The membrane is then cut into strips, bleached possibly and washed in water, dried, twisted or rolled, and then smoothed, the two last-named operations being done by machinery only.

In the absence of catgut ligatures, one may prepare suitable sizes of violin or banjo strings for surgical purposes.

The catgut after being prepared as above, must be freed from contained fat, a convenient method being by maceration in ether or chloroform. The gut may be preserved in alcohol, oil of juniper or other suitable liquid, using a well-closed, wide-mouth bottle as receptacle.

In medicating catgut, which is in skeins, the cord binding the gut should be cut so as to impregnate uniformly.

After impregnation, it is generally directed to wind the gut upon glass spools.

#### Catgut, Carbolated.

#### I. Lister's process:

Carbolic acid, crystal gr.	540
Distilled waterfl.dr.	
Olive oilfl.oz.	71/4

Mix in a wide-mouth glass bottle or other suitable vessel. Place in the mixture as much catgut as is to be impregnated, adding more liquid if necessary, to completely immerse the gut.

Allow the gut to remain in the turbid fluid until the latter becomes clear, agitating from time to time. When the liquid becomes transparent, the gut will have become soft and have absorbed water and acid. It is now wound upon glass spools, the whole then being immersed in a mixture of carbolic acid and olive oil in the proportion of 1 of the former to 4 of the latter. - D.

#### II. Block's process:

Roll the cleansed catgut upon glass spools, macerate in a 5-per-cent aqueous solution of carbolic acid for 48 hours, then unwind in a dish containing a freshly prepared 5-per-cent aqueous carbolic acid solution, and now rewind (tightly) the gut upon the spool. serve in a 5-per-cent solution of carbolic acid in alcohol.-D.

III. Block's carbolic alcohol process:

Prepare like the preceding, using an alcoholic solution of carbolic acid instead of an aqueous one as above. - D.

#### Catgut, Chromated.

Chromic	acidgr.	2
Carbolic	acid, crystalgr.	400
Distilled	waterfl.oz.	1734

Dissolve and add an amount of catgut equal in weight to the carbolic acid used; allow to remain in the solution for 48 hours, then remove, dry, wind upon glass spools, and place in carbolized oil.

This is Lister's process for hardening gut so that it will not be readily absorbed.

#### Catgut, Juniper, Kocher.

Macerate catgut for 24 hours in oil of juniper berries, then preserve either in this oil or in the following solution:

Mercuric chloridegr.	5
Glycerinfl.oz.	2
Alcoholfl.oz.	

#### Catgut, Mercuric Chloride. (Sublimated Catgut.)

#### 1. Bergman's process:

Catgut rolled on glass spools is to be

mercuric chloride, renewing the solution every 2 days, until it remains clear, then preserve the gut in this solution.

## 2. Schede and Kuemmel's process:

Catgut rolled upon glass spools is to be placed for 12 hours in a 5-per-cent aqueous solution of mercuric chloride; then preserve in 1/2 per-cent alcoholic solution of the same salt to which has previously been added 10 per cent of glycerin.-D.

#### Caustic, Arsenical, Ratier's.

Arsenious	acid			 			.gr.	20
Kino								
Cinnabar.				 			.ør.	320

All should be in fine powder.

#### Caustic, Black, Velpeau's.

Triturate powdered licorice root in a mortar, adding sulphuric acid until a suitable mass is formed.

## Caustic, Vienna. (Potassa cum Calce.)

Triturate equal parts by weight of caustic potassa and lime together in a warm iron mortar so as to form a powder, and keep in a well-stoppered bottle.—U. S. P.

Cerate, Brown. (Brown Ointment. -Mother salve. - Unguentum fuscum. -Emplastrum fuscum molle.)

I. Lead plas	ster	 .av.oz. 8

Melt the lead plaster and stir constantly while liquid, until it assumes a dark brown tint; add the wax and lard; allow to cool somewhat, and pour into molds.-Austr. Pharm.

II. A more rational and easy method of preparation is the following:

Black mother plasterav.oz.	8
Lardav.oz.	61/2
Yellow waxav.oz.	11/2

Melt the plaster and wax, add the lard, allow to cool somewhat, and pour into molds as before. - D.

#### Cerate of Calamine. (Turner's Cerate.

-Cerate of Zinc Carbonate.) Zinc carbonate.....av.oz. 1 Simple cerate.....av.oz. 5 —Eclectic.

Under the name "Turner's cerate," the placed in 5-per-cent alcoholic solution of National Formulary gives a preparation of with simple ointment.

# Cerate, Calendula.

Lard, fresh		av.oz. 8
Fluid extract	of calendula	fl.oz. 1

Heat on a water bath until the alcohol has evaporated, stirring frequently meanwhile.

Another method consists in digesting the flowers with melted lard for about 10 minutes, stirring occasionally; then strain, and stir frequently until cooled. It is advisable to add about 2 av.ounces of yellow wax.

#### Cerate of Copaiba.

Yellow	wax,	filtered	 	av.oz.	23/
Balsam	of c	opàiba	 	av.oz.	51/2

Melt the wax and, when it begins to cool, add the balsam; the two will mix more readily if the balsam be warmed to 30 or 40 degrees C. before adding to the wax. -D.

# Cerate, Green. (Ceratum Aeruginis.)

Yellow waxav.oz.	8
Resinav.oz.	434
Gum turpentineav.oz.	
Verdigris, powderedgr.	350
Benzoinated lardgr.	180
Olive oilfl.dr.	3

Melt the wax, resin, and turpentine together, add the verdigris, which has previously been triturated to a smooth paste with the lard and oil, mix well, and pour into molds.-D.

# Cerate of Nutmeg.

Refer to "Balsams."

#### Cerate of Soap.

Soap plasterav.oz.	31/2
Yellow waxav.oz.	
Olive oilfl.oz. 8	

Melt the wax and plaster, add the oil, and stir until cool,-U. S. P. 1870.

#### Chlorodyne.

Under the name "J. Collis Browne's Chlorodyne," an English nostrum was at one time used largely, not only in England, but in this country as well. A number of substitutes have been and are still in use, the formulas for which differ from one another more or less, sometimes quite materially. The National Formulary recognizes one of these preparations under the name of "Mis-

the same strength as the above, but made tura Chloroformi et Cannabis Indicæ Composita," or chloroform anodyne, made as follows:

I.		
	Chloroformfl.oz.	2
	Etherfl.dr.	4
	Tincture of cannabis indicafl.oz.	2
	Tincture of capsicumfl.oz.	1
	Morphine sulphategr.	18
	Oil of peppermint m.	15
	Glycerinfl.oz.	2
	Waterfl.oz.	1
	Alcohol, enough to makefl.oz.	16

Dissolve the oil in 8 fluidounces of alcohol. add the chloroform, ether and tinctures, mix well, add the morphine sulphate previously dissolved in the water and glycerin; finally add the remainder of the alcohol.

#### II.

Chloroformfl.dr.	2
Morphine	
Etherfl.dr.	1
	8
Diluted hydrocyanic acidfl.dr.	2
Tincture of capsicumfl.dr.	2
Molasses	21/2
Extract of licoricegr.	30

This is said to resemble Browne's very closely.

#### III. Chandler's formula:

Morphine hydrochlorate gr. 16	3
Oil of peppermintdrops 20	
Tincture of capsicumdrops_30	)
Fluid extract of cannabis indica.fl.dr	l
Alcoholfl.oz.	9
Glycerinfl.oz.	3

#### IV. Modified Smith's formula:

Chloroformfl.dr. 4
Morphine hydrochlorategr. 20
Oil of peppermintdrops 8
Tincture of capsicumdrops 50
Diluted hydrocyanic acidfl.dr. 2
Mucilage of acaciafl.oz. 1
Fluid extract of cannabis indica. fl.dr. 2
Simple syrup, enough to makefl.oz. 4
Solution of caramel, sufficient to color
properly.

## V. Squires' formula:

Chloroform . ! fl.oz.	1
Stronger etherfl.dr.	2
Alcoholfl.oz.	1
Molassesfl.oz.	1
Extract of licorice, powdered.gr.	300
Morphine hydrochlorategr.	2
Oil of peppermintdrops	4
Simple syrupfl.oz.	41/2
Diluted hydrocyanic acidfl.dr.	4

Dissolve the morphine and oil of peppermint in the alcohol, mix the chloroform and ether with this solution, mix the licorice with the syrup, add the molasses, shake these two mixtures well together, and, lastly, add the hydrocyanic acid and again shake well.

# VI. Gilman's formula:

Chloroform	fl.dr.	2
Glycerin	fl.oz.	2
Alcohol	.fl.oz.	2
Spirit of peppermint	fl.dr.	2
Diluted hydrocyanic acid	.fl.dr.	2
Tincture of capsicum		
Morphine hydrochlorate	gr.	8
Molasses	. H. oz.	3

#### VII. Fenner's formula No. 1:

Oleoresin of capsicumdrops 3	Chloroform	1/2
------------------------------	------------	-----

Dissolve the oils in the chloroform, add the compound spirit of ether, and mix.

#### VIII. Fenner's formula No. 2:

Morphine sulphategr.	15
Chloroform fl.oz.	1
Fluid extract of cannabis indica.fl.oz.	1
Glycerinfl.oz.	1
Alcohol fl.oz.	
Diluted hydrocyanic aciddrops	15
Spirit of peppermintdrops	15

Mix the liquids and dissolve the morphine in the mixture.

# Chloroform of Aconite.—(Aconite Chloroform.)

Aconite root	av. oz 11
Water of ammonia	
Distilled water	fl.oz. 10
Chloroform	sufficient

Bruise the aconite, moisten thoroughly with the distilled water and ammonia previously mixed, macerate for 4 hours, dry carefully, reduce to No. 40 powder, pack tightly in a percolator (such as would be used for highly volatile liquids), macerate for 24 hours with 11 fluidounces of chloroform, and then percolate slowly, adding more choloroform until 16 fluidounces of product are obtained.

—Brit. Form.

"Chloroforms" of other alkaloidal drugs, such as belladonna, hyoscyamus, etc., may be produced in the same manner.

# Chloroform, Camphorated.

Chloroform	.fl.oz.	4
Camphor	av.oz.	8
Mix and dissolve.—Brit. Form.		

## Collodion, Aconite.

Aconite root, in fine powder.av.oz.	81/2
Stronger etherfl.oz.	
Alcoholsufficie	ent.
Balsam of firgr.	240
Pyroxylingr.	60

Pack the powder very tightly in a percolator intended for volatile liquids, mix the ether with 4 fluidounces of alcohol, saturate the drug with this liquid, macerate for about 16 hours, then percolate slowly, adding enough alcohol through the percolator so as to make 16 fluidounces of percolate. In this dissolve the balsam and pyroxylin.

#### Collodion, Belladonna.

This may be prepared in the same way as the preceding, using belladonna leaves instead of aconite root.

# Collodion, Cantharidin.

I.		,	
Cantharidin		gr.	3
Gum turpentine		ıv.oz.	1
Acetone		fl.dr.	3
Collodion, enough to make	e	fl.oz.	8

Triturate the cantharidin with the turpentine to as fine a powder as possible, then add the acetone and heat the mixture very cautiously until solution occurs; then add to the collodion. If a green color be desired, add a small amount of English extract of cannabis indica.—D. modified.

Triturate the cantharidin with the oil, add to the collodion, and agitate until dissolved. If desired of a green tint, color as before.—
D. modified.

#### Collodion, Carbolated.

Carbolic acid, crystal	gr.	150
Collodion	fl.oz.	8
Oil of rose	drops	2
	_	

The product contains 5 per cent of acid.

#### Collodion, Chrysarobin.

Chrysarobin Collodion	 	gr. 320 fl.oz. 8
		_

The chrysarobin should be in very fine

powder and be dissolved in the collodion by agitation.—D.

The product contains 10 per cent of chrysarobin.

# Collodion, Carbolated Salicylic, Unna.

Carbolic acid, crystalav.oz.	1
Salicylic acidav.oz.	1
Collodionfl.oz.	
Mix and dissolve by agitation.—D.	

The product contains one-third by weight of the combined acids.

# **Collodion, Diachylon.** (Collodion with Lead Plaster.)

Lead plastergr.	280
Alcohol fl.dr.	6
Stronger etherfl.dr.	
Collodion, enough to makefl.oz.	8

Melt the plaster by warming, add the alcohol and ether, stir quickly until dissolved, and add at once to the collodion.—D.

The product contains 5 per cent of lead plaster.

#### Collodion, Iodized.

Iodine, reduced to powder gr.	160
Flexible collodionfl.oz.	8

Introduce the iodine into a bottle, add the flexible collodion and agitate until the iodine is dissolved.—N. F.

#### Collodion, Iodoform.

Iodoformgr.	160
Flexible collodionfl.oz.	8

Mix and dissolve the iodoform in the flexible collodion by agitation.—N. F.

#### Collodion, Iodol.

Iodol gr.	300
Alcoholfl.oz.	11/4
Etherfl.oz.	51/6
Pyroxylin gr.	120
Castor oil.fl.dr.3 (or gr. 160 if wei	

Dissolve the iodol in the mixture of alcohol and ether, add the pyroxylin in small portions, agitate until dissolved, and finally add the oil. The product contains 10 per cent, of iodol.

#### Collodion, Iron.

Solid chloride of irongr.	300
Flexible collodionfl.oz.	
Oil of sagedrops	10
Dissolve by agitation.—D.	

The product contains 10 per cent of ferric chloride.

# Collodion, Mercuric Chloride. (Corrosive Sublimate Collodion.—Sublimated Collodion.)

Corrosive sublimate, powdered..gr. 150 Flexible collodion..........fl.oz. 8

Dissolve the salt in the collodion by agitation. If the salt be on hand only in the crystalline form, it should be triturated dry to powder.—D.

The product contains 5 per cent of corrosive sublimate.

# Collodion, Photographer's.

Pyroxylingr	. 275
Alcohol fl.oz	. 21/4
Alcohol	41/2
Stronger etherfl.oz. 10	to 12

Add the pyroxylin to the alcohol, shake well, and add the absolute alcohol and ether.

—H.

This contains almost 5 per cent of gun cotton, and is therefore stronger than the official collodion.

#### Collodion, Salol.

Salolgr.	280
Stronger etherfl.oz.	1
Collodionfl.oz.	7

Dissolve the salol in the ether and add the collodion.—D.

The product contains 10 per cent of salol.

#### Collodion, Thymol.

Thymolgr.	
Collodionfl.oz.	8
Dissolve by agitation.—D.	

#### Collodion, Thiol, Jacobsen.

Thiol, powdergr	. 75
Flexible collodionfl.oz	
	D

#### Concentrations.

This class of preparations was originally introduced by physicians of the Eclectic school of medicine and subsequently was employed by physicians of other schools. They are now employed in Europe; also one, at least, has found recognition in the United States Pharmacopæia, viz., podophyllin.

The general plan for their manufacture originally consisted in extracting the drug with strong alcohol, evaporating this tincture to small bulk, adding to cold water, stirring constantly meanwhile, allowing the precipi-

tate formed to subside, collecting the latter, and drying and powdering it.

If the drug contain an oleoresin, the precipitate cannot be dried sufficiently to powder, but remains a soft, sticky mass. In this case, the precipitate should be dried sufficiently to remove the water, then add enough of the original drug, in powdered form, to reduce to a rather tough mass, break the latter into small pieces, dry in warm air, and pulverize as before.

In many instances, there would be, by the use of water alone as a precipitating agent, an inappreciable amount of precipitate. In such cases, alum is added to the water and some ammonium carbonate to the alcoholic liquid. The precipitate contains aluminium hydrate, and the product will very often be green from precipitated chlorophyll.

Other substances are added to the water to facilitate precipitation, such as acids and alkalies, depending, of course, upon the character of the drug.

While the above methods of preparing concentrations were advised by Eclectic practitioners, other methods are in vogue among manufacturers. In many instances, the concentration is simply a powdered extract, the so-called "euonymin," for example, the drug often being exhausted with dilute alcohol or water. Instead of using the powdered drug as a drying agent, as is stated above, most manufacturers use milk sugar, magnesia or other absorbent powder.

Most of the concentrations are supposed to be resinous in character, and are termed "resinoids," while others are alkaloidal in character. The latter are believed to be prepared by exhausting the drug with a very dilute acid (from ½ to 3 per cent), usually hydrochloric or sulphuric, evaporating the liquid obtained to moderate bulk, adding ammonia to neutralize the acid, collecting the precipitate, and washing and drying it. Hydrastis and sanguinaria are drugs treated in this manner.

As may be surmised from what has been stated, concentrations from different manufacturers differ from each other greatly in quality and strength and color, as well as in other properties.

The nomenclature of the concentrations is very confusing. The resinoids have names ending in "in"-cornin, helenin, etc.-while those of alkaloidal character are known by names ending either in "in" or "ia," and are coupled with names indicating the acid employed in extraction, e.g., hydrastia sulphate, sanguinarin nitrate, etc. Manufacturers of pharmaceuticals have not only modified the processes of preparation, but have introduced innovations in the titles. When it is borne in mind that glucosides have names terminating in "in," and that alkaloidal names end in "ine" ("ia" is also used), and that almost all drugs contain glucosides or alkaloids or even both, one can readily see that almost inextricable confusion must result, often to the serious detriment of sick persons. This is the case particularly with the derivatives of hydrastis and sanguinaria.

The following table is presented because of its convenience for reference. It will not apply to the products of all manufacturers, nor does it mention all concentrations, but it may prove useful nevertheless.

Class I. Concentrations consisting largely of resin and nearly or entirely soluble in alcohol:

Aletridin,\* Asclepidin,\* Cimicifugin (Macrotin), Eryngin,\* Helonin,\* Iridin \* Liatrin,\* Podophyllin (U. S. P.), Ptelein.\*

The pulverulent resinoids of aletris and iris versicolor are known as above, while the soft, oleoresinous articles are called aletrin and irisin.

Class II. Concentrations which generally contain inorganic constituents from the process of manufacture, and are frequently of a green character:

Barosmin, Euonymin (green variety), Lobelin, Lycopin, Podophyllin (yellow variety), Scutellarin, Senecin.

Class III. Concentrations which contain a number (probably most) of the soluble drug constituents:

Aletrin,\* Alnuin, Ampelopsin, Apocynin, Asclepidin,\* Baptisin, Betulin, Caulophyllin, Cerasin, Chelonin, Chimaphilin, Chionanthin, Collinsonin, Colocynthin, Cornin, Corydalin, Cypripedin, Dioscorin, Euonymin (brown variety), Eupatorin, Euphorbin, Eupurpurin, Fraserin, Gelsemin, Geranin, Gossypin, Hamamelin, Humulin, Inulin, Irisin,\* Jalapin, Juglandin, Leontodin, Leptandrin, Liatrin,\* Liriodendrin, Menispermin, Myricin, Phytolaccin, Prunin, Rhusin, Rumicin, Smilacin, Stillingin, Taraxin, Trillin, Viburnin, Xanthoxylin.

Class IV. Concentrations which consist of alkaloids or alkaloidal salts in more or less impure form:

Hydrastin, Hydrastia Sulphate, Muriate, etc.; Sanguinarin, Sanguinarina Nitrate, Sulphate, etc.

Those concentrations whose names have the stars appear in the market in both pulverulent and oleoresinous forms.

# Confection, Aromatic. (Electuaries.)

Mix aromatic powder with an equal weight of honey or a sufficient quantity to form a stiff paste.—U. S. P. 1870.

# Confection of Calamus. (Sugared Calamus.—Candied Sweet Flag.)

Macerate the root in the water for twelve hours, add the sugar, and heat, stirring constantly until perfectly dry. At first the evaporation may be conducted over a direct flame or fire, but toward the latter end of the operation, water-bath temperature only must be employed.

# Confection of Copaiva. (Electuaire de Copahu.)

Balsam of copaibaav.oz. 4	
Cubeb, powderedav.oz. 6	
Catechu, powderedav.oz. 2	
Oil of peppermintfl.dr. 1	

Mix the whole well together.—Codex.

# Confection of Figs. (Medicated Figs.)

Figsav.oz	. 8
Waterfl.oz	
Cinnamon, fine powdergr	. 120
Sugarav.oz	
Senna, fine powderav.or	

Pour hot water on the figs, macerate until softened, strain with expression, add the sugar, heat until a soft pulpy mass is ob-

tained, add the senna and cinnamon, and mix well.

See also "Confection of Senna," U. S. P...

# Confection of Hollyhock, Thompsonian.

Poplar barkav.oz.	1/2
Bayberry barkav.oz.	1/2
Golden sealav.oz.	1/2
Clovesav.oz.	1/2
Cinnamon av.oz.	1/2
Cypripediumav.oz.	1/2
Capsicumgr.	110
Oil of pennyroyalfl.dr.	2
Hollyhock flowersav.oz.	8
•	

Mix the first six ingredients in powder form, add the oil, and incorporate the whole with the flowers, mixing as well as possible, and forming balls the size of small marbles.

# **Confection of Opium.** (Electuary of Theriac.—Theriac.)

Compound powder of opium.av.oz. 5½
Simple syrup......fl.oz. 11½
—Brit. Pharm.

II.	
Powdered opiumgr.	260
Aromatic powderav.oz.	61/4
Clarified honevav.oz.	15
II S P	1870

#### III.

Opiumgr.	55
Angelica rootgr.	275
	220
Valerian rootgr.	110
Squillgr.	110
Zedoarygr.	110
Cassia barkgr.	550
Cardamom gr.	55
Cloves gr.	55
Myrrh gr.	55
Iron sulphate, crystalgr.	55
Glycerin,	
Simple syrup,	
Honey, equal parts of each,	
enough to makeav.oz.	$12\frac{1}{2}$

Mix the ingredients above in powder form, or preferably mix them whole, and then reduce to powder; then add the glycerin, syrup and honey.—H.

The cardamom should be used without the capsule or enveloping membrane.

These preparations differ in many respects, but the most noteworthy difference is in the proportion of opium, the first two containing about  $2\frac{1}{2}$  per cent of this drug, the last about 1 per cent, and the kind of confection

to be dispensed will depend on the nationality of the consumer or the kind he has been in the habit of using.

# Confection of Pepper.

Black pepper, powderedgr. 350
Caraway, powderedgr. 525
Clarified honeyav.oz. 6

# Confection of Prunes. (Medicated Prunes.)

Prunes, stonedav.oz.	8
Waterfl.oz.	16
Citric acid, powderedgr.	60
Tincture of gingerfl.oz.	1
Sugarav.oz.	
Senna, fine powderav.oz.	21/2

Pour hot water on the prunes, macerate until softened, strain with expression, add the sugar and acid; heat until a pulpy mass is obtained, add the senna and tincture, and mix well.

See also "Confection of Senna," U. S. P.

# Confection of Rhubarb, Compound.

Rhubarb, powderedgr.	350
Fennel, powderedgr.	350
Licorice root, powdered av. oz.	134
Senna, powderedav.oz.	13/
Sugar, powderedav.oz.	31/2
Tamarind pulp, purifiedav.oz.	31/2
Syrup of mannafl.oz.	4
, 1	—D.

#### Confection of Sulphur.

Suphurav.oz. 63/4
Cream of tartarav.oz. 13/4
Syrup of orangefl.oz. 5
Tragacanth, powderedav.oz. 3/4
-Brit. Pharm.

#### Confection of Tamarind.

Tamarind pulpav.oz. Sugar, powderedav.oz.	$\frac{6\frac{1}{2}}{9\frac{1}{2}}$
II.	—Н.
Tamarind pulpav.oz.	21/2
Sugar, powderedav.oz.	21/2
Senna, powderedgr.	
Cream of tartargr.	70
Manna av.oz.	4
Warm waterav.oz.	8

Dissolve the manna in the water, strain, add the other ingredients, mix well, and evaporate the whole at a low temperature to proper consistency.—H.

#### Conserves.

In conformity to the custom of the United States pharmacopæia, preparations formerly

under this title are here classed with the confections.

# Copper, Aluminated. (Sapis divinus. Augen Stein.)

Copper sulphate, pure	av.oz.	2
Potassium nitrate		
Potassa alum	av.oz.	2
Camphor	gr.	55

Triturate the three salts separately to fine powder, then mix, melt carefully in a porcelain evaporating dish over a hot fire, then quickly add the camphor in a powdered state and previously mixed with an equal weight of powdered alum, and pour the whole out on a porcelain slab.—Germ. Pharm.

When cold, it may be broken into pieces or rubbed to powder, and then preserved in well-stoppered bottles.

# Copper, Ammoniated.

Ammonium carbonatea	v.oz.	3
Copper sulphate	av oz. 4	£

Triturate together until effervescence ceases, then lay between folds of bibulous paper and dry. Keep in well-stoppered bottles.

## Cordials.

The above title is applied to many preparations which vary greatly in character, and could not properly be placed under one heading. Some occur under other more approprize te titles in this part; others may be found in Parts II and IV. Consult index.

# Cotton, Absorbent. (Purified Cotton, U. S. P.)

Ordinary cotton contains, in addition to the dirt and other matter that accompany it, some fatty and coloring matter. The fatty matter does not permit the cotton to absorb water or aqueous fluids to any appreciable extent, and the unpurified is therefore of no value for surgical and pharmaceutical purposes.

There are several methods by which ordinary cotton may be rendered absorbent. An easy process is to wash it repeatedly with ether, which, of course, extracts or dissolves out the fatty matter. The first portions of ether may be economically replaced by gasolin or petroleum ether, finishing the washing with ether so as to avoid the odor of gasolin

in the product. By means of this process cotton can be rendered absorbent and be dried in a very few minutes.

The process adopted by manufacturers on the large scale is usually about as follows:

Boil any desired quantity of the best corded cotton with a 5-per-cent, solution of caustic potassa or soda for one-half hour, or until the cotton is entirely saturated with the solution, and the alkali has saponified all the fatty matter; wash thoroughly in clear water to remove all the soap and nearly all the alkali; press out the excess of water; place in a 5-per-cent solution of chlorinated lime, allowing to remain for 15 or 20 minutes; again wash, first in some clear water, then dip in water acidulated with hydrochloric acid, and wash again thoroughly in clear water; press out the excess of water and again boil for 15 or 20 minutes in 5-per-cent alkali solution; now wash well in clear water, dip in water acidulated with hydrochloric acid, and again wash thoroughly in clear water; now press out the water and dry.

Owing to the cellular character of cotton, it is very likely to absorb a liquid and not readily give it up again; hence when cotton is to be well washed, it should be kneaded with the hands or otherwise. In removing excess of water or other liquid, an ordinary clotheswringer will be found to serve the purpose admirably. Cotton batting, as purchased, ordinarily occurs in rolled sheets; if care be taken in the above manipulation, the product can be made to retain this "sheet" form.

Medicated Cottons.—These are prepared by impregnating good absorbent cotton by means of immersion in liquid containing the medicating substance. Sometimes a large excess of liquid is employed for impregnation, the excess being removed by subsequent expression; or else only so much is used as that when the whole is taken up by the cotton, the latter will contain the requisite or prescribed amount of medicament. In the former case, the expression may be by means of an instrument like an ordinary clotheswringer, conducting the expression so that the liquid which the cotton is allowed to retain will yield a proper strength of finished

product. In either case, therefore, the result is identical, and while preference is usually given, in the formulas which follow, to the use of a large amount of liquid, the other process may be substituted, the result always depending more on the care and skill of the operator than on any other circumstance. In impregnating cotton, it may be necessary to knead the latter with the liquid, and sometimes even to macerate for one or two hours; if the small quantity of liquid be used, then the cotton must be weighted down in some convenient manner. In expressing cotton after impregnating in a large quantity of liquid, it may be advisable to wrap the cotton in parchment paper to protect it from the press; if salicylic acid be present, the solution and cotton must not be allowed to come in contact with any iron parts. Cotton must always be passed through the press evenly to medicate uniformly.

Drying of impregnated cotton may be done on screens either in a drying closet or in a room which is dry and perfectly aseptic or clean.

Good absorbent cotton, when dipped in water, will take up or retain, after expression, twice its weight of liquid, each pound of dry cotton yielding therefore three pounds of moist cotton.

All prepared cottons should be kept in suitable receptacles or wrappers such as glass, parchment paper, paraffined paper, paraffined or resin-coated pasteboard boxes, etc., to prevent loss by evaporation or accession of septic matter.

#### Cotton, Aluminium Acetate, Burow.

Solution of aluminium acetate. fl.oz. 16
Distilled water... fl.oz. 32
Absorbent cotton......av.oz. 16
Proceed as with borated cotton.—D.

# Cotton, Antirheumatic. (Gicht Watte.)

Oil of birch tar, rectified...drops 12
Oil of turpentine, rectified...drops 12
Oil of juniper wood...drops 12
Oil of cloves...drops 12
Oil of rosemary...drops 12
Camphor...gr. 20
Alcohol...fl.dr. 5½
Absorbent cotton...av.oz. 16

the liquid which the cotton is allowed to retain will yield a proper strength of finished hol, filter, and moisten the cotton with the filtrate in any convenient way, as, for example, by means of an atomizer. During this moistening the cotton should be picked into thin layers, and be turned about frequently so as to impregnate evenly: Dry by exposure to atmosphere for one hour, and wrap in waxed paper or other suitable container.—D. II.

Red saundersgr.	96
Benzoingr.	20
Peru balsam,gr.	
Alcoholfl.dr.	8

Macerate for several days, filter and impregnate cotton with filtrate as in the preceding.—H. modified.

#### Cotton, Arnicated.

Tincture of arnica	.fl.oz.	21/2
Glycerin	.fl.oz.	21/2
Alcohol	.fl.oz.	34
Distilled water	.fl.oz.	12
Absorbent cotton	av.oz.	16

Immerse the cotton in the liquid, press out to the weight of 48 av. ounces, dry carefully, and pack into containers.—D. modified.

The finished product represents 10 per cent of tincture of arnica of the weight of the cotton used. Inasmuch as tincture of arnica U. S. P. is double the strength of the tincture of the German Pharmacopæia, this 10 per cent corresponds to 20 per cent of the tincture of arnica of the latter work.

#### Cotton, Benzoated, Bruns, Jr.

	3	р. с.	4 p. c
Benzoic acid	gr.	315	420
Castor oilfl.	oz.	1	1
Alcohol fl.	oz.	54	54
Absorbent cottonav.	oz.	16	16
	5	р. с.	10 p. c
Benzoic acid	gr.	525	1050
Castor oilfl.	oz.	1	2
Alcoholfl.	oz.	54	51
Absorbent cottonav.	oz.	16	16

Dissolve the acid in the alcohol, add the oil, saturate the cotton with this liquid, prepared after any of the given proportions, then press it until it weighs 48 av. ounces, and dry it at a temperature not exceeding 30 degrees C.—D.

## Cotton, Borated.

-		
	ner	cent

1	
Boric acidgr.	525
Distilled water, hotfl.oz.	
Absorbent cotton,,,,,av.oz.	16

#### 10 per cent.

Boric acidgr.	1050
Distilled water, hotfl.oz.	40 1/4
Distilled water, hotfl.oz. Absorbent cottonav.oz.	16

#### 20 per cent.

Boric acidav.oz.	43/4
Distilled water, hotfl.oz.	40
Absorbent cottonav.oz.	16

Dissolve the acid in the water, immerse the cotton in the solution, press out to the weight of 48 av. ounces, and proceed as before.—D. modified.

# Cotton, Carbolated, Bruns, Jr.

#### 5 per cent.

Carbolic acid, crystal gr.	525
Castor oilfl.dr	4
Resin av.oz.	43/
Alcoholfl.oz.	481/
Absorbent cottonav.oz.	
10	

#### 10 per cent

To per cent.	
Carbolic- acid, crystalgr.	1050
Castor oilfl.oz.	
Resinav.oz.	$7\frac{1}{4}$
Alcoholfl.oz.	7 ½ 42 ½
Absorbent cottonav.oz.	16

Dissolve the resin in the mixed oil and alcohol by agitation, filter, impregnate the cotton as before, pressing out to the weight of 48 av ounces, and drying without heat. Pack immediately in air-tight containers.—D.

## Cotton, Cocaine.

Cocaine hydrochlorate	gr.	210
Distilled water	fl.oz.	91/4
Absorbent cottona	v.oz.	16

Dissolve the cocaine salt in the water, add the alcohol, saturate the cotton with the liquid (the cotton being kept below it by means of weights, but on removal, it is to retain the whole of the liquid), and dry at 30 degrees C.—D.

The product contains 3 per cent of the alkaloidal salt.

#### Cotton, Cocaine-Borated.

Cocaine hydrochlorategr.	140
Boric acidgr.	
Carbolic acid, crystalgr.	
Glycerinfl.dr.	
Alcohol fl.oz.	
Distilled water, hotfl.oz.	113/
Absorbent cottonav.oz.	16

Dissolve the boric acid in the distilled water and glycerin, add the cocaine salt, car-

bolic acid and alcohol, saturate the cotton as in the making of cocaine cotton, and dry by exposure to the air. This is considered useful in dressing burns and scalds.—D.

#### Cotton, Cocaine-Morphine.

Cocaine hydrochlorategr.	210
Morphine hydrochlorategr.	
Alcohol fl.oz.	
Distilled waterfl.oz.	
Absorbent cottonav.oz.	16

Prepare this like the two preceding. This is used for tamponing carious teeth, to allay toothache.—D.

# Cotton, Ichthyol.

20 per cent

No per centi	
Ichthyol-ammoniumav.oz.	434
Alcohol fl.oz.	
Distilled waterfl.oz.	291/2
Absorbent cottonav.oz.	

#### 50 per cent.

Ichthyol-ammoniumav.oz.	12
Alcoholfl.oz.	
Distilled waterfl.oz.	22
Absorbent cottonav.oz.	16

Dissolve the ichthyol in the alcohol and water, saturate the cotton with the solution, press out to the weight of 48 av.ounces, and dry at a temperature not to exceed 25 degrees C.—D.

#### Sotton, Iodized.

Todine													۶.	ŗr.		70	0	
Lotton	L											а	v.	.02	z.,	1	6	

Place the iodine at the bottom of a widemouth glass vial, insert the cotton, tie over the mouth with parchment paper wetted with glycerin, place the vial in a water-bath of from 50 to 60 degrees C., and continue the heat until all of the iodine has been vaporized and the cotton is evenly impregnated with it. Pack in well-closed glass containers.—D.

The product is called a 10-per-cent cotton; practically it contains but 9 per cent of iodine.

#### Cotton, Iodol.

Iodolgr.	1050
Glycerinfl.dr.	10
Alcohol fl.oz.	51
Absorbent cottonav.oz	16

Dissolve the iodol in the alcohol with the aid of a little heat (50 degrees C.), gradually add the glycerin, saturate the cotton with the solution, kneading thoroughly, and proceed

as described under iodoform cotton, pressing out to 48 av. ounces.—D.

The product is a 10-per-cent medicated cotton.

#### Cotton, Iodoform, Mosetig.

#### 5 per cent.

Iodoformgr.	525
Etherfl.oz.	163/
Ether	401/2
Absorbent cottonav.oz	16

#### 10 per cent.

I	
Iodoformgr.	1050
Castor oilfl.dr.	
Resingr.	350
Ether fl.oz.	251/
Alcoholfl.oz.	271/2
Absorbent cottonav.oz.	16
00	

#### 20 per cent.

Iodoformav.oz.	43/
Castor oilfl.dr.	$12^{'}$
Resin av.oz.	11/2
Etherfl.oz.	421/2
Alcohol fl.oz.	91/
Absorbent cottonav.oz.	16

Dissolve the iodoform in the ether and alcohol, add the resin and castor oil if they be used, agitate until dissolved, saturate the cotton with the solution, wrap in thin parchment paper, puncture a number of holes along the edge, and press out to the weight of 48 av. ounces. This work must be performed with a certain amount of celerity. Dry in the open air, excluding daylight during the entire operation.—D.

A better mode of preparation for iodoform cotton is to use only so much solution that, when all is absorbed, the fabric will contain the proper proportion.

# Cotton, Mercuric Chloride. (Sublimated Cotton.)

#### 1. Schede's process:

½ per cent.	
Mercuric chloridegr.	26
Glycerinfl oz.	33/4
Alcoholfl.oz.	13
Distilled waterfl.oz.	32
Absorbent cottonav.oz.	16
½ per cent.	

Mercuric chloridegr.	53
Glycerinfl.oz.	33/
Alcohol	
Distilled waterfl.oz.	32
Absorbent cottonav.oz.	

Dissolve the corrosive sublimate in the

mixed liquids, filter, and impregnate the cotton as before.

#### 2. Link and Voswinkel's process:

#### ¼ per cent.

Mercuric chloridegr.	26
Lithium chloridegr.	
Alcohol fl.oz.	
Distilled waterfl.oz.	
Absorbent cottonav.oz.	16
. /	

#### ½ per cent.

/2 per cent.	
Mercuric chloridegr.	52
Lithium chloridegr.	
Alcohol fl.oz.	$18\frac{1}{2}$
Distilled waterfl.oz.	32
Absorbent cottonav.oz.	16

Proceed as before and press out to the weight of 48 av. ounces; dry at a temperature of 25 to 30 degrees C.

Ordinarily sublimated cotton suffers diminution in strength due to reduction of the mercuric chloride. This is ascribed to lack of care in preparing the absorbent cotton used, the presence of glycerin and of stearic acid to impart a brilliant whiteness, and creaking sound when pressed between the fingers, etc. The second process is supposed to yield a permanent product. The lithium chloride is intended to replace the glycerin of other processes.

3. With tartaric acid, ¼ per cent.:	
Mercuric chloridegr.	26
Tartaric acidgr.	
Alcohol fl.oz.	273/
Distilled.waterfl.oz.	24
Absorbent cottona.v.oz.	16

Proceed according to the usual mode, press out to the weight of 48 av. ounces, and dry under exclusion of daylight.—D.

4. Lister's sero-sublimate cotton, ½ per cent.:

Mercuric chlorideg	r. 52
Horseblood-serumav.d	z. 12
Distilled waterfl.c	z. 34½
Absorbent cottonav.c	z. 16

Dissolve the corrosive sublimate by trituration in the blood-serum, add the water, and saturate the cotton with the liquid; press out to 48 av. ounces.

If horseblood-serum cannot be had, dissolve 52 gr. of corrosive sublimate and 210 gr. of sodium chloride, by trituration in 770 gr. of egg-albumen diluted with 46 fluid-ounces of distilled water, and in this soak the cotton.

Dry at a temperature not exceeding 30 degrees C., and keep from the light.—D.

This cotton really contains mercury in the form of albuminate.

#### 5. Sal alembroth cotton:

Mercuric chloridegr. 2	6
Ammonium chloridegr. 1	
Alcohol fl.oz.	91/4
Distilled waterfl.oz. 3	$8\frac{1}{2}$
Absorbent cottonav.oz. 1	6

Immerse the cotton in the solution and press out to the weight of 48 av. ounces; dry in the dark.—D.

## Cotton, Naphthalin.

Naphthalingr.	1050
Resingr.	
Castor oilfl.dr.	
Alcoholfl.oz.	$52^{\circ}$
Absorbent cotton av.oz.	16

Dissolve the solids in the liquids with the aid of heat, soak the cotton in the hot solution, and quickly express to 48 av. ounces; dry by exposure to air.—D.

The product is a 10-per-cent medicated cotton.

#### Cotton, Resorcin.

# 3 per cent.

Resorcin gr.	315
Glycerinfl.dr.	54
Alcohol fl.oz.	
Distilled waterfl.oz.	
Absorbent cottonav.oz.	

#### 5 per cent.

Resorcing	r. 525
Glycerinfl.os	
Alcohol fl.o.	
Distilled waterfl.oz	
Absorbent cottonav.o	z. 16

Proceed in the usual manner, pressing out to the weight of 48 av. ounces; dry at a temperature between 25 and 30 degrees C.

# Cotton, Salicylated.

# 1. Process of Bruns, Jr.:

#### 5 per cent.

Salicylic acidgr.	525
Castor oilfl.dr.	
Alcohol fl.oz.	53
Absorbent cottonav.oz.	16
1.0	

#### 10 per cent.

Salicylic acidgr.	1050
Castor oilfl.dr.	
Alcoholfl.oz.	51
Absorbent cottonav.cz.	16

Dissolve the acid in the alcohol, add the oil, and proceed in the usual way, pressing

out to the weight of 48 av. ounces; dry at a in, then press it until the product weighs 48 temperature between 25 and 30 degrees C.

# 2. Thiersch's process:

#### 4 per cent.

Salicylic acid gr.	420
Glycerinfl.dr.	1
Alcohol flo.z.	8
Distilled water, hot:fl.oz!	39
Absorbent cottonav.oz.	16
10 per cent.	

Salicylic acidgr.	1050
Glycerinfl.dr.	
Alcohol fl.oz.	$15\frac{1}{2}$
Distilled water, hotfl.oz.	31
Absorbent cottonav.oz.	16

Dissolve the acid in alcohol, add the glycerin and water, and proceed as before.

# Cotton, Styptic. (Hemostatic Cotton.— Ferrated Cotton.)

Absorbent cotton, Solution of chloride of iron, Glycerin,

Water . . . . . . . . of each sufficient.

Mix the liquids in the proportion of 5 parts of the iron solution, 1 part of glycerin, and 4 parts of water, in such quantities that the cotton shall be completely immersed in the liquid when gently pressed. Allow the cotton to remain in the liquid one hour, then remove it, press it until it has been brought to twice its original weight, spread it out in thin layers, in a warm place, protected from dust and light, and when it is sufficiently dry, transfer it to well-closed receptacles.-N. F.

II.

Solution of ferric chloride	fl.oz.	$6\frac{1}{2}$
Glycerin	.fl.oz.	1
Distilled water		
Alcohol	fl.oz.	18‡
Absorbent cotton		

Impregnate and press the cotton in the usual manner; dry with exclusion of daylight, and keep the product in amber-colored bottles.-D. modified.

III.

Solution of chloride of fron
(Germ. Pharm.sp.gr. 1.28).fl.oz. 9
Glycerinfl.oz. 1
Waterfl.oz. 191/4
Alcohol
Purified cottonav.oz. 16

av. ounces, and dry it at a gentle heat, with exclusion of light,

One hundred parts contain about 25 parts of anhydrous ferric chloride.

Keep the product protected against light. --Germ. Form.

The 9 fluidounces of solution of ferric chloride of the German pharmacopæia used in the last formula corresponds to 7 fluidounces of the solution of the United States Pharmacopœia.

## Cotton, Tannin-Carbolated.

(Tannin, 10 per cent; carbolic acid, 8 per cent)

Tannic acidgr.	1050
Carbolic acid, crystalgr.	
Castor oilfl.oz.	
Alcohol fl.oz.	$46\frac{1}{2}$
Absorbent cottonav.oz.	16

Dissolve the acids in the alcohol and oil, impregnate and press the cotton in the usual manner, and dry in the open air without heat.—D.

# Cotton, Thymolated, Ranke.

#### 2 per cent.

•	
Thymolgr.	210
Resingr.	420
Spermacetiav.oz.	$6\frac{1}{2}$
Alcohol fl.oz.	$46\frac{1}{2}$
Absorbent cottonav.oz.	16
5 per cent.	
Thymolgr.	525
Resingr.	1050
Spermacetiav.oz.	71
Alcohol fl.oz.	43
Absorbent cottonav.oz.	16
Absorbent cotton	10

Effect solution and saturation at an elevated temperature, pressing out, while still warm, to the weight of 48 av. ounces; dry in the open air. - D.

#### Cotton, Zinc Chloride, Bardeleben.

Zinc chloridegr.	1050
Distilled water, hotfl.oz.	$43\frac{1}{2}$
Absorbent cottonav.oz.	16

Proceed as in making borated cotton. - D. The product is a 10-per-cent medicated cotton.

# **Decoction of Barley.** (Barley Water.)

Pearl ba	rley				 		av.oz.	11
Distilled								

Wash the barley with cold water, and re-Mix the liquids, immerse the cotton there- ject the washings; boil the washed barley with the distilled water for 20 minutes in a covered vessel, and strain. The product is about 16 fluidounces.—Brit, Pharm.

#### Decoction of Broom.

Broom to	ops (scopari	as)	gr.	350
Distilled	water		fl.oz.	16

Boil in a covered vessel for 10 minutes, strain and pour water over the strainer until the colature measures 16 fluidounces.—Brit. Pharm.

#### Decoction of Buckthorn, Compound.

Buckthorn bark, cutgr. 720
Rhubarbgr. 144
Hopsgr. 36
Carduus mariana seedsgr. 36
Distilled watersufficient

Heat the buckthorn with 18 fluidounces of distilled water for 30 minutes, then add the other ingredients, heat again for 10 minutes, strain, and add enough distilled water through the strainer to make the colature measure 16 fluidounces.

It is advisable to macerate the buckthorn with the water for at least 2 hours before heating.—D.

#### Decoction of Dandelion.

Dandelion, sliced and bruised...gr. 350 Water, enough to make.....fl.oz. 16

Boil the dandelion with 16 fluidounces of water for 10 minutes, strain and add enough water through the strainer to make the colature measure the required amount.—Brit. Pharm.

#### Decoction of Granatum.

Pomegranate bark,	cutav.oz 13/4
Distilled water	sufficient

Add 32 fluidounces of water to the bark, boil down to 16 fluidounces, strain and add, if necessary, enough water through the strainer to make the colature measure 16 fluidounces.—Brit. Pharm.

# Decoction of Guaiac, Compound. (Decoctum Lignorum.)

Guaiac woodgr. 360
Sarsaparilla, cutgr. 360
Licorice root, cutgr. 72
Sassafras wood, cutgr. 72
Watersufficient

Macerate the sarsaparilla with 20 fluidounces of water for 24 hours, add the guaiac,

heat for 1 hour, then add the licorice and sassafras, and strain in 15 minutes, adding enough water through the colature to make up 16 fluidounces.—Belg. Pharm.

#### Decoction of Pareira.

Pareira root, in No. 20 powder.av.oz. 1 Distilled water.....sufficient.

Boil the drug with 16 fluidounces of water for 15 minutes in a covered vessel, strain and add enough water through the strainer to make up 16 fluidounces.—Brit. Pharm.

# Decoction of Sarsaparilla, Stronger Compound.

(Stronger Zittmann's Decoction.)

Sarsaparilla, cut finegr.	600
Waterfl.oz.	64
Red sulphide of mercurygr.	5
Calomelgr.	24
Alum, powderedgr.	40
Sugar, powderedgr.	40
Anise, bruisedgr.	24
Fennel, bruisedgr.	24
Licorice, cutgr.	72
Senna, cutgr.	120

Macerate the sarsaparilla with the water for 24 hours, and strain; triturate the two mercury salts, alum, and sugar together, tie the mixed powders into a folded piece of muslin, suspend this into the infusion of sarsaparilla previously put into an earthen vessel, and evaporate the liquid down to 32 fluidounces. While yet hot, add the remaining drugs, and when cold, strain without pressure, set the decoction aside to settle, and decant the clear liquid.

The formula given above is the one formerly in vogue. The formula recognized by the German pharmacopæia contains no mercurials whatever; the proportions of the remaining ingredients is somewhat different, although the difference is a very immaterial one, and the water added to the drugs is 34 fluidounces, the colature; after heating for some hours, being made up to 32 fluidounces.

The formula given is an unscientific one, but contains mercury in some form, unless the evaporation has been conducted in metallic vessels, and hence is often preferred on this account.

The second edition of the German pharmacopeaia recognized this preparation under the title here given, but the present (third) edition calls it simply "compound decoction of sarsaparilla" and does not recognize the weaker decoction at all.

# Decoction of Sarsaparilla, Weaker Compound.

(Weaker Zittmann's Decoction.)

Residue from preceding decoction.	
Sarsaparilla, cut finegr.	384
Waterfl.oz.	96
Cardamom, bruised gr.	20
Cinnamom, bruisedgr.	20
Lemon peel, cutgr.	20
Licorice root, cutgr.	20

Mix the residue above specified with the sarsaparilla, and boil the whole with the water until the whole is reduced to 32 fluid-ounces, and while still hot, add the remaining drugs; allow to cool, strain, set aside to settle, and decant the clear liquid.

As stated in the preceding article, this preparation is not recognized at all by the present German pharmacopæia, but the second edition did consider it, and gave for it the following formula:

Sarsaparilla, cut	gr. 288
Watersuf	ficient
Lemon peel, cut	gr. 30
Cinnamon, bruised	gr. 30
Cardamom, bruised	gr. 30
Licorice root, cut	

Macerate the sarsaparilla with 30 fluidounces of water for 24 hours, then heat in a covered vessel on a water bath for 3 hours, stirring occasionally; add the other ingredients, macerate for fifteen minutes, strain the liquid with expression, allow the decoction to settle, pour off all the clear liquid, and add enough water to it to make 32 fluidounces.

#### Discs. (Lamellæ.)

These are prepared by adding to a concentrated solution of gelatin some glycerin and a solution of a medicating ingredient. While hot this is poured on a perfectly level and polished surface, and after drying the discs are cut out from the sheet, each disc being 1-25th inch in thickness and weighing about 1-50th grain. The discs of the British pharmacopæia are used only in ophthalmic practice. They are, as follows:

Discs of Atropine, each containing 1-5000th grain of atropine sulphate.

Discs of Cocaine, each containing 1-200th grain of cocaine hydrochlorate.

Discs of Physostigmine, each containing 1-1000th grain of physostigmine.

#### Drops, Cholera.

Various preparations commonly known by this title will be found under the "Mixtures."

#### Drops, Cordial Warner's.

Sennagr.	12
Coriandergr.	36
Fennel gr.	36
Cochineal gr.	
Extract of licorice gr.	15
Spanish saffron gr.	15
Raisinsav. oz.	
Water fl.oz.	8
Alcohol fl.oz.	91/4
	, .

Mix, macerate for 7 days, agitating occasionally, and filter.—H. modified.

#### Drops, Cramp. (Krampf Tropfen.)

The "red," also known as Herzstaerkungs Tropfen and Tinctura Apoplectica rubra, is made after one of the following formulæ:

1. Aromatic tincturefl	. dr.	4
Tincture of catechu	l.dr.	4
Tincture of cinnamonf	l.dr.	4
Red saunders, rasped	.gr.	64
Alcohol	l.oz.	63/
Spirit of ether	l.oz.	714

Macerate for several days, and strain through cotton.—H.

2. (	Oil of peppermintdrops	3
	Oil of anisedrops	
	Oil of cinnamondrops	
-	Tincture of catechu fl.oz.	1
1	Aromatic tincture fl.oz.	2
(	Chloroform fl.dr.	1/2
5	Spirit of etherfl.oz.	5
	Alcohol fl.oz.	
		−Ĥ.

# **Drops, Gold.** (Gold Tropfen.—Tinctura Dulcis.)

Potassium acetategr.	112
Caramelgr.	
Hydrochloric ether (ethyl chlo-	
ride)fl.oz.	1.
A	- 4

modified.

In the absence of the above the aromatic

tincture of the National Formulary may be dispensed.—H.

The ethereal tincture of chloride of iron of the National Formulary is also known as Lamotte's Gold Drops, and this is what may be desired when "gold drops" are requested.

#### Drops, Dysmenorrhœa,

Rademacher. (Guttæ ad Menstrua Tormentosa Rademacheri.)

Tincture of nux vomica.

Tincture of castor, equal parts of each by measure.—H.

# Drops, Jesuit. (Guttæ Jesuitarum.)

Guaiac resin, powderedav.oz.	$3^{\frac{1}{2}}$
Peru balsamgr.	96
Sassafras bark, cutav.oz.	21/4
Alcoholfl.oz.	16

Mix, macerate for 7 days, agitating occasionally, and filter.

#### Drops, Mother.

Valerianav.oz.	21/
Galanga gr.	
Red saundersgr.	10
Compound spirit of ether fl.dr.	2
Spirit of cinnamonfl.dr.	
Diluted alcoholsuffici	

Reduce the three drugs to fine powder and extract by percolation with diluted alcohol; the percolate obtained should be of such amount that when added to the two spirits the product will measure just 16 fluidounces.

Very frequently simply tincture of valerian is dispensed as "mother drops."

Other "drops" may be found scattered throughout this work (consult index).

#### Eau Sedative de Raspail.

Water of ammonia	.fl.oz.	2
Spirit of camphor	.fl.dr.	11/
Chloride of sodium	av.oz.	1
Water, sufficient to make	.fl.oz.	16

## ELIXIRS.

The presentation of this multifarious array of elixirs perhaps requires an explanation. The catalogues of the large manufacturers list all of the elixirs herein mentioned, and this fact indicates a demand for them. In order, therefore, that retail pharmacists may supply this demand without recourse to the manufacturer or the wholesaler, we have given reliable formulæ for these preparations.

# Elixir of Acetanilid. (Elixir of Antifebrin.)

Acetanilid.....gr. 128
Simple elixir .....fl.oz. 16

Dissolve by agitation. Each fluidram contains one grain of acetanilid.

# Elixir, Adjuvant.

Sweet orange peel	.gr.	120
Wild cherry		
Licorice root, Russian		
Coriander	gr.	60
Caraway	gr.	60
Simple syrupfl		5
Alcohol.		
Water of each, s	uffic	ient

Reduce the wild cherry to a moderately coarse powder, moisten it with 4 fluidrams of water, and set aside for 12 hours. Reduce the other solids also to a moderately coarse powder, mix this intimately with the wild cherry, and having mixed 1 volume of alcohol with 2 volumes of water, moisten the powder with 4 fluidrams of the mixture, and pack tightly in a percolator. Then gradually pour menstruum on top until 11 fluidounces of percolate are obtained. Mix this with the syrup and filter.—N. F.

#### Elixir of Aletris.

Fluid extract of aletris farinosa. fl.oz. 2 Simple elixir......fl.oz. 14

Mix, let stand for several days, and filter. Each fluidram represents 7½ grains of aletris farinosa.

## Elixir of Aloin, Strychnine and Belladonna.

Fluid extract of belladonna root..m. 64 Elixir of aloin and strychnine, enough to make.....fl.oz.16

Each fluidram contains ¼ gr. of aloin, and 1-100 gr. of strychnine, and represents ½ gr. of belladona root.

# Elixir of Aloin and Strychnine.

Aloin	-	gr.	32
Strychnine sulpha			
Simple elixir		fl.oz	16

Dissolve the alkaloidal salt in the elixir by trituration in a mortar, or agitation in a bottle; add the aloin, agitate until dissolved, and filter if necessary.

Each fluidram contains ¼ gr. aloin, and 1-100 gr. of strychnine sulphate.

#### Elixir of Ammonium Bromide.

Ammonium bromide gr.	640
Citric acid gr.	30
Aromatic elixir enough to make fl.oz.	16

Dissolve the ammonium bromide and the citric acid in about 8 fluidounces of aromatic elixir, by agitation; then add the remainder of the elixir, and filter, if necessary.

Each fluidram contains 5 gr. of ammonium bromide.—N. F.

#### Elixir of Ammonium Chloride.

Ammonium chloride......gr. 1280 Simple elixir, enough to make fl. oz. 16 Dissolve by agitation, and filter, if necessary.

Each fluidram contains 10 gr. of ammonium chloride.

## Elixir of Ammonium Chloride and Licorice, Compound.

Ammonium chloride......gr. 640
Compound elixir of licorice,
enough to make.....fl.oz. 16
Dissolve by agitation, and filter, if necesary.

Each fluidram contains 5 gr. of ammonium chloride.

#### Elixir of Ammonium Valerianate.

Elixii of Ammonium valerianat	e.
1. Ammonium valerianategr.	256
Chloroformm.	61
Tincture of vanilla fl.dr.	2
Compound tincture of cudbear fl.dr.	2
Water of ammonia,	
Aromatic elixirof each suffici	ient

Dissolve the ammonium valerianate in about 10 fluidrams of aromatic elixir, in a graduated vessel, and add enough water of ammonia, in drops, until a faint excess of it is perceptible in the liquid; then add the chloroform, tincture of vanilla, and compound tincture of cudbear, and finally enough aromatic elixir to make 16 fluidounces. Filter if necessary.

Each fluidram contains 2 gr. of ammonium valerianate.—N. F.

#### 2 Goddard's formula modified:

Goddard's formula, modified:
Valerianic acid, from the root . fl.dr. 3
Ammonium carbonatesufficient
Distilled waterfl.oz. 4
Elixir of curacao fl.oz. 10
Orange flower waterfl.oz. 4
Mucilage of gum arabicfl.oz. 1
Tincture of cudbearfl.dr. 2

Add the acid to the water and neutralize well as  $\frac{1}{100}$  gr. of strychnine sulphate.

with ammonium carbonate; mix with the other ingredients, and filter.

## Elixir of Ammonium Valerianate and Chloral Hydrate.

Chloral hydrate......gr. 640
Elixir of ammonium valerianate,
N. F., enough to make....fl.oz. 16

Dissolve by agitation and filter if necessary. Each fluidram contains 5 gr. of chloral hydrate and 2 gr. of ammonium valerianate.

## Elixir of Ammonium Valerianate with Cinchonidine.

Cinchonidine sulphate......gr 64
Elixir of ammonium valerianate,
N. F., enough to make....fl.oz. 16

Dissolve by agitation. The elixir of ammonium valerianate employed in making this preparation should be exactly neutral.

Each fluidram contains  $\frac{1}{2}$  gr. of cinchonidine sulphate and 2 gr. of ammonium valerianate.

#### Elixir of Ammonium Valerianate, Cinchonidine and Iron Pyrophosphate.

Iron pyrophosphate, soluble....gr. 64
Distilled water, hot.....fl.dr. 4
Elixir of ammonium valerianate
with cinchonidine, enough to
make.....fl.oz. 16

Dissolve the iron salt in the water and add the elixir of ammonium valerianate with cinchonidine.

Each fluidram contains nearly 2 gr. of ammonium and  $\frac{1}{2}$  gr. of cinchonidine sulphate, as well as  $\frac{1}{2}$  gr. of iron pyrophosphate.

#### Elixir of Ammonium Valerianate with Cinchonidine, Iron Pyrophosphate and Strychnine.

Strychnine sulphate.....gr. 11/4
Distilled water.....fl.dr. 2
Elixir of ammonium valerianate, cinchonidine and iron
pyrophosphate, enough to
make......fl.oz. 16

Dissolve the strychnine sulphate in the water, and add the elixir.

Each fluidram contains nearly 2 gr. of ammonium valerianate,  $\frac{1}{2}$  gr. of cinchonidine sulphate and  $\frac{1}{2}$  gr. of iron pyrophosphate, as well as  $\frac{1}{100}$  gr. of strychnine sulphate.

#### Elixir of Ammonium Valerianate with Cinchonidine, Iron Pyrophosphate and Quinine.

Iron pyrophosphate, soluble....gr. 64
Distilled water, hot.....fl.dr. 4
Elixir of ammonium valerianate,
with cinchonidine and quinine,
enough to make....fl.oz. 16

Dissolve the iron salt in the water and add the elixir.

Each fluidram contains nearly 2 gr. of ammonium valerianate,  $\frac{1}{2}$  gr. of cinchonidine sulphate, and  $\frac{1}{4}$  gr. of quinine hydrochlorate, as well as  $\frac{1}{2}$  gr. of iron pyrophosphate.

#### Elixir of Ammonium Valerianate with Cinchonidine, Iron Pyrophosphate, Quinine and Strychnine.

Strychnine sulphate......gr. 1½
Distilled water.....fl.dr. 2
Elixir of ammonium valerianate, cinchonidine, iron pyrophosphate, and quinine, enough to make.....fl.oz. 16

Dissolve the strychnine salt in the water and add the elixir.

## Elixir of Ammonium Valerianate with Cinchonidine and Quinine.

Quinine hydrochlorate......gr. 32 Cinchonidine sulphate......gr. 64 Elixir of ammonium valerianate, N. F., enough to make.....fl.oz. 16

Mix, dissolve by agitation and filter.

Each fluidram contains 2 gr. of ammonium valerianate,  $\frac{1}{2}$  gr. of cinchonidine sulphate and  $\frac{1}{4}$  gr. of quinine hydrochlorate.

#### Elixir of Ammonium Valerianate with Cinchonidine, Quinine and Strychnine.

Strychnine sulphate......gr. 1½
Distilled water.....fl.dr. 2
Elixir of ammonium valerianate
with cinchonidine and quinine, enough to make....fl.oz. 16

Dissolve the strychnine in the water and add the elixir.

## Elixir of Ammonium Valerianate with Cinchonidine and Strychnine.

Strychnine sulphate .......gr. 1½
Distilled water .......fl.dr. 2
Elixir of ammonium valerianate
with cinchonidine, enough to
make ........fl.oz. 16
Dissolve the strychnine sulphate in the

water, add the elixir and filter if necessary.

Each fluidram contains  $\frac{1}{100}$  gr. of strychnine sulphate,  $\frac{1}{2}$  gr. of cinchonidine sulphate and 2 gr. of ammonium valerianate.

## Elixir of Ammonium Valerianate and Iron.

Iron pyrophosphate, soluble...gr. 128
Distilled water, hot.....fl.oz. 1
Elixir of ammonium valerianate,
N. F. . . . . . fl.oz. 15

Dissolve the iron salt in the water and add the elixir.

Each fluidram contains 1 gr. of iron pyrophosphate and 2 gr. of ammonium valerianate.

## Elixir of Ammonium Valerianate with Iron and Quinine.

Quinine hydrochlorate......gr. 32
Iron pyrophosphate, soluble....gr. 64
Distilled water, hot......fl.dr. 4
Elixir of ammonium valerianate,
N. F., enough to make....fl.oz. 16

Add the quinine salt to 15 fluidounces of elixir, dissolve by agitation, dissolve the iron salt in the water, mix the two solutions, and add the remainder of the water.

Each fluidram contains ¼ gr, of quinine hydrochlorate, ½ gr. of iron pyrophosphate and 2 gr. of ammonium valerianate.

## Elixir of Ammonium Valerianate with . Iron, Quinine and Strychnine.

Dissolve the strychnine sulphate in the water and add the elixir.

## Elixir of Ammonium Valerianate with Quinine.

Quinine hydrochlorate.....gr. 32 Elixir of ammonium valerianate.fl.oz. 16

Dissolve the quinine hydrochlorate in the elixir by agitation, and, if necessary, by occasionally immersing the bottle containing the ingredients in hot water, until solution has been effected; filter if necessary.

Each fluidram contains ¼ gr. of quinine hydrochlorate. and 2 gr. of ammonium valerianate.—N. F.

## Elixir of Ammonium and Morphine Valerianates.

Dissolve by agitation. The elixir of ammonium valerianate used in making the above should not be alkaline.

Each fluidram contains  $\frac{1}{16}$  gr. of morphine valerianate and 2 gr. of ammonium valerianate.

## Elixir of Ammonium, Quinine and Strychnine Valerianates.

Strychnine (alkaloid)......gr. 1¼
Valerianic acid........sufficient
Quinine valerianate.....gr. 64
Elixir of ammonium valerianate,
N. F., enough to make...fl.oz. 16

Dissolve the strychnine in 2 fluidrams of the elixir of ammonium valerianate by the aid of a slight excess of valerianic acid. Triturate the quinine salt with this solution and add the remainder of the elixir of ammonium valerianate, agitate occasionally until dissolved, then filter.

In case the valerianic acid is in such excess that its odor is perceptible, the liquid must be cautiously neutralized by stirring it with a glass rod which is repeatedly moistened with very dilute ammonia water. Any excess of the latter must be avoided, as otherwise alkaloidal strychnine will be precipitated.

Each fluidram contains  $_{10}^{1}$  gr. of strychnine valerianate,  $_{2}^{1}$  gr. of quinine valerianate and  $_{2}^{2}$  gr. of ammonium valerianate.

## Elixir of Ammonium Valerianate with Sumbul. (Elixir of Ammonium

Valerianate with Musk Root.)

Fluid extract of sumbul......fl.oz. 2 Elixir of ammonium valerianate.fl.oz. 14 Mix, let stand for several hours and filter through purified talcum.

Each fluidram contains nearly 2 gr. of ammonium valerianate and represents 7½ gr. of sumbul root.

## Elixir of Ammonium Valerianate with Strychnine.

Strych	nnine sulp	hate		.gr.	11/4
			ı valeria- f	l.oz.	16
Mix,	dissolve	by	agitation,	and	filter.

The elixir of ammonium valerianate must not be alkaline.

#### Elixir of Anise. (Aniseed Cordial.)

Anethol m.	25
Oil of fenneldrops	4
Spirit of bitter almondfl.dr.	14
Deodorized alcoholfl.oz.	4
Simple syrupfl.oz.	10
Distilled waterfl.oz.	2
Magnesium carbonate, powder.gr.	120

Mix the anethol, oil and spirit with the alcohol, add the syrup and water, and set the mixture aside for 12 hours. Then mix it intimately with the magnesium carbonate, and filter it through a wetted filter, returning the first portions of the filtrate until it runs through clear.—N. F.

#### Elixir of Antipyrin.

Dissolve by agitation.

Each fluidram contains 5 gr. of antipyrin.

## Elixir of Arbor Vitæ. (Elixir of Thuja Occidentalis.)

Fluid extract of arbor vitæ....fl.dr. 10} Simple elixir, enough to make..fl.oz. 16

Mix, allow to stand for several hours and filter.

Each fluidram represents 5 gr. of arbor vitæ.

## Elixir of Arsenic. (Elixir of Potassium Arsenite.)

Fowler's solution......fl.dr. 10<sup>1</sup>/<sub>1</sub> Simple elixir, enough to make.fl.oz. 16

Each fluidram contains 5 minims of Fowler's solution, which represents  $\frac{1}{20}$  gr. of arsenious acid.

#### Elixir, Aromatic.

Compound spirit of orange...fl.dr. 6½ Simple syrup......fl.oz. 24 Calcium phosphate, precipitated.av.oz. 1 Alcohol,

Distilled water, of each.....sufficient.

To the spirit add enough alcohol to make 16 fluidounces. To this solution add the syrup in several portions, agitating after each addition, and afterwards add, in the same manner, 24 fluidounces of water. Mix the calcium phosphate intimately with this liquid, and filter through a well-wetted filter, returning the first portions until a clear

liquid is obtained. Lastly wash the filter with a mixture of 1 volume of water and 3 of alcohol, until the filtrate measures 64 fluidounces.—U. S. P.

#### Elixir of Arsenic and Iron Chloride.

Refer to "Elixir of Chlorides of Arsenic and Iron."

#### Elixir of Arsenic, Iron and Mercury Chlorides.

Refer to Elixir of Chlorides of Arsenic, Iron and Mercury.

#### Elixir of Arsenic, Iron and Quinine.

Refer to Elixir of Iron, Quinine and Arsenic.

## Elixir of Arsenic and Mercury Iodides.

Donovan's solution.....fl.dr.  $10\frac{1}{2}$  Simple elixir, enough to make fl.oz. 16

Each fluidram contains  $\frac{1}{20}$  gr. each of red iodide of mercury and iodide of arsenic.

#### Elixir of Arsenic and Quinine.

Solution of arsenious acid...fl.dr. 10½ Quinine sulphate.....gr. 128 Simple elixir, enough to make.fl.oz. 16

Dissolve by agitation and filter if necessary.

Each fluidram contains & gr. of arsenious acid, and 2 gr. of quinine sulphate.

#### Elixir of Arsenic and Strychnine.

Solution of arsenious acid...fl.dr. 10½ Strychnine sulphate.....gr. 1½ Simple elixir, enough to make.fl.oz. 16

Dissolve by agitation and filter.

Each fluidram contains  $\frac{1}{20}$  gr. of arsenious acid and  $\frac{1}{100}$  gr. strychnine sulphate.

#### Elixir of Beef.

Extract of beefgr.	256
Distilled waterfl.oz.	1
Simple elixir, enough to make.fl.oz.	16

Dissolve the extract in the water, add the elixir, let stand for several days if possible, and filter.

Each fluidram contains 2 gr. of extract of beef.

The extract of beef suitable for this and similar preparations is that which is prepared by Liebig's method.

## Elixir of Beef, Bismuth, Cinchona and Iron.

Extract of beefg	r. 256
Distilled waterfl.o	
Elixir of bismuth, cinchona and	
iron, enough to makefl.o	z. 16

Dissolve the extract in the water add the elixir, let stand for several days if possible, and filter.

#### Elixir of Beef and Cinchona.

Extract of beefgr. Distilled waterfl.oz.	
Detannated elixir of cinchona,	
N. F., enough to make, fl.oz.	- 16

Dissolve the extract in the water, add the elixir, let stand for several days if possible, and filter.

Each fluidram contains 2 gr. of extract of beef and represents about  $1\frac{1}{2}$  gr. of cinchona.

#### Elixir of Beef, Cinchona and Iron.

Extract of beefgr.	256
Distilled waterfl.oz.	
Elixir of cinchona and iron,	
N. F., enough to make flor	16

Dissolve the extract in the water, add the elixir, allow to stand for several days if possible, and filter.

Each fluidram contains 2 gr. of extract of beef and nearly 2 gr. of iron phosphate and represents about  $1\frac{1}{2}$  gr. of cinchona.

## Elixir of Beef, Cinchona, Iron and Strychnine.

Strychnine sulphategr.	11/
Distilled waterfl.oz.	
Elixir of beef, iron and cin-	
chona, enough to makefl.oz.	16

Dissolve the alkaloidal salt in the water, add to the elixir, and filter.

#### Elixir of Beef, Coca and Iron.

Iron phosphate, solubleg	r. 256
Extract of beef	
Distilled water, hotfl.d	
Elixir of coca, N. F., enough	
to makefl.c	z. 16

Dissolve the iron salt and extract each in one ounce of water, mix with the elixir, allow to stand for several days, and filter.

Each fluidram contains 2 gr. each of iron phosphate and extract of beef and represents about 7 gr. of coca.

#### Elixir of Beef and Iron.

Citrate of iron and ammonium ..gr. 128
Distilled water, warm.....fl.oz. 1
Elixir of beef, enough to make.fl.oz. 16
Dissolve the iron salt in the water and add

the elixir.

Each fluidram contains 1 gr. of iron salt and 13/4 gr. of extract of beef.

#### Elixir of Beef, Iron and Malt.

Extract of beefgr.	256
Extract of malt (thick) av.oz.	4
Citrate of iron and ammoniumgr.	128
Spirit of orangefl.dr.	1
Alcoholfl.oz.	2
Sherry winefl.oz.	9
Water,	
Ferric hydrate, of eachsuffic	ient

Dissolve the extract of beef in one fluidounce of hot water, and add the alcohol containing the spirit of orange, then the wine with which the malt extract has previously been mixed; shake frequently during 2 or 3 days, filter, and wash the filter with a mixture of alcohol and water in the proportion of 1 of the former to 4 of the latter by measure, so as to obtain a filtrate of 15 fluid ounces. Dissolve the iron salt in 6 fluidrams of water, add to the filtrate, and then add enough water to make 16 fluidounces.

The ferric hydrate may be prepared as described under the heading of elixir of gentian; the amount to be used must be sufficient to detannate the mixture, and if an insufficient amount has been used, more must be added, allowing to stand for several days more. The test to be applied is the usual one—filtering a small amount of liquid and testing the filtrate with solution of iron chloride to note if any discoloration occur.

#### Elixir of Berberine.

Berberine phosphategr. &	32
Distilled water, hotfl.oz.	1
Simple elixirfl.oz. 1	

Dissolve the berberine in the water and add the elixir.

Each fluidram contains 1/2 gr. of berberine

Each fluidram contains ¼ gr. of berberine phosphate.

#### Elixir of Berberine and Iron.

Iron pyrophosphate, soluble	gr.	128
Distilled water, hot	fl.oz.	1
Elixir of berberine, enough	to	
make		16

Dissolve the iron salt in the water, add the elixir, and filter if necessary.

Each fluidram contains 1 gr. of iron pyrophosphate and nearly 1/4 gr. of berberine phosphate.

#### Elixir of Bismuth.

Ι.	
	Bismuth and ammonium citrategr. 256
	Water, hotfl.oz. 1
	Water of ammonia,
	Aromatic elixir, of eachsufficient

Dissolve the bismuth and ammonium citrate in the hot water, allow the solution to stand until any undissolved matter has subsided; then decant the clear liquid, and add to the residue just enough water of ammonia to dissolve it. Then mix this with the decanted portion and add enough aromatic elixir to make 16 fluidounces; filter, if necessary.

Each fluidram represents 2 gr. of bismuth and ammonium citrate.—N. F.

II.

Bismuth subnitrate......gr. 180
Nitric acid, C. P.....fl.dr. 3
Tartaric acid,
Sodium bicarbonate, C. P.,

Distilled water, of each.....sufficient Simple elixir, enough to make fl.oz. 16

Mix 3 fluidrams of nitric acid with an equal measure of distilled water, and to this add the bismuth subnitrate, stirring until solution is effected; add enough distilled water to make 3 fluidounces. Now dissolve 135 grains of tartaric acid in 12 fluidrams of distilled water, and pour this into the bismuth solution, stirring constantly during mixing. To this mixture gradually add 150 grains of sodium bicarbonate, stirring constantly during mixing. Dilute the magma thus obtained with 5 fluidounces of distilled water, and after the lapse of several hours, pour the whole upon a plain filter; allow the liquid to drain, and wash the precipitate with distilled water until the washings pass taste-

Now mix 150 grains of sodium bicarbonate with 1 fluidounce of distilled water, and add 135 grains of tartaric acid gradually, with constant stirring. When reaction has ceased, and a clear solution has formed, add the washed precipitate of bismuth tartrate and stir until it is dissolved. Now add enough

distilled water to make 3 fluidounces and then the elixir. Allow to stand for 24 hours and filter.

The 3 fluidounces of solution to which elixir is added is a solution of tartrate of bismuth and sodium, and may be used to advantage, instead of citrate of bismuth and ammonium, in all preparations containing a soluble compound of bismuth.

Each of the elixirs of bismuth contains about 2 gr. of the respective bismuth salt.

#### Elixir of Bismuth and Cinchona.

Citrate of bismuth and ammo-
niumgr. 128
Distilled water, hotfl.dr. 4
Ammonia watersufficient
Detannated elixir of cinchona,
N. F., enough to makefl.oz. 16

Mix the bismuth salt with the hot water, allow the solution to stand to permit any undissolved matter to subside; decant the clear liquid and add to the residue just enough ammonia water to dissolve. Mix this solution with the decanted liquid, and if alkaline, neutralize the mixture with dilute solution of citric acid gradually added. To the whole add the elixir of cinchona, let stand 24 hours, and filter if necessary.

Each fluidram contains 1 gr. of bismuth salt and represents 1½ gr. of calisaya.

## Elixir of Bismuth, Cinchona, Iron and Pepsin.

Citrate of bismuth and ammo-
nium gr. 128
Detannated tincture of cin-
chonafl.oz. 21
Iron pyrophosphate, solublegr. 256
Pepsin, puregr. 128
Distilled water, hot fl.oz. $1\frac{1}{2}$
Aromatic spiritfl.oz. 1
Simple syrupfl.oz. 5
Simple elixir, enough to make. fl. oz. 16
Water of ammoniasufficient

Add the bismuth salt to 4 fluidrams of the water, triturate well for a moment, allow to stand until the insoluble portion subsides, decant the clear liquid, carefully add to the residue just sufficient ammonia water to dissolve it, carefully avoiding any excess, and mix this solution with the decanted portion.

Add the pepsin to 5 fluidounces of simple elixir and agitate occasionally until dis-

solved; also dissolve the iron salt in the remainder of the water.

Now mix the three liquids, add the tincture, the spirit, syrup, and the remainder of the elixir, allow to stand for 24 hours, and filter

Each fluidram represents 1 gr. each of citrate of bismuth and ammonium and of pepsin, nearly 2 gr. of cinchona, and 2 gr. of iron pyrophosphate.

## Elixir of Bismuth, Cinchona and Iron.

Bismuth and ammonium citrategr. 1	128
Water, hotfl.dr.	4
Water of ammonia sufficient	ent
Elixir of cinchona and iron.	
enough to makefl.oz.	16

Dissolve the bismuth and ammonium citrate in the hot water, allow the solution to stand until any undissolved matter has subsided; then decant the clear liquid, and add to the residue enough water of ammonia to dissolve it, carefully avoiding an excess. Then mix the solution with the elixir of cinchona and iron; let the mixture stand 24 hours, if convenient, and filter.

Each fluidram contains 1 gr. of bismuth and ammonium citrate, and nearly 2 gr. of iron phosphate.—N. F.

#### Elixir of Bismuth, Cinchona, Iron, Pepsin and Strychnine.

Strychnine sulphategr	11/4
Distilled waterfl.dr.	4
Elixir of bismuth, cinchona,	
iron and pepsinfl.oz.	$15\frac{1}{2}$

Dissolve the strychnine sulphate in the water, add the elixir, and filter.

## Elixir of Bismuth, Cinchona and Pepsin.

Detannated tincture of cin- chonafl.oz.	$2\frac{1}{2}$
Citrate of bismuth and ammo-	100
niumgr.	120
Pepsin, puregr.	128
Distilled water, hotfl.dr.	
Water of ammoniasuffic	ient
Aromatic spiritfl.oz.	1
Simple syrupfl.oz.	4
Simple elixir, enough to make.fl.oz.	16

Triturate the citrate of bismuth and ammonium with the water, allow to stand until the insoluble matter subsides, to the residue add ammonia water until solution takes

place, carefully avoiding any excess, and mix the two liquids. Add the pepsin to 7 fluidounces of elixir, agitate occasionally until dissolved; mix this with the preceding liquid, add the tincture, spirit, syrup and remainder of the elixir, allow to stand for 24 hours, and filter.

Each fluidram represents nearly 2 gr. of cinchona and contains 1 gr. each of pepsin, and of citrate of bismuth and ammonium.

## Elixir of Bismuth, Cinchona, Iron and Strychnine.

Strychnine sulphategr. 1	1/4
Water, hot	
Elixir of cinchona, iron and	
bismuth; enough to makefl.oz. 16	

Dissolve the strychnine sulphate in the hot water, add the elixir of cinchona, iron and bismuth, and filter, if necessary.—N. F.

#### Elixir of Bismuth and Gentian.

Citrate of bismuth and ammo-	
niumgr.	128
Distilled water, hotfl.dr.	4
Ammonia watersuffic	ient
Elixir of gentian, N. F., enough	
to makefl.oz.	16

Treat the bismuth salt as described under Elixir of Bismuth and Cinchona, add the elixir of gentian, allow to stand for 24 hours, and filter if necessary.

Each fluidram contains 1 gr. of bismuth salt and represents about 2 gr. of gentian.

#### Elixir of Bismuth, Gentian and Iron.

Citrate of bismuth and ammo-	
nium	128
Distilled water, hotfl.dr.	
Ammonia watersuffic	ient
Elixir of gentian and iron phos-	
phate, enough to makefl.oz.	16

Treat the bismuth salt as in the preceding elixir, add the elixir of gentian and iron, let stand 24 hours, and filter if necessary.

Each fluidram contains 1 gr. of bismuth salt and nearly 1 gr. of iron phosphate, and represents about  $1\frac{1}{2}$  gr. of gentian.

## Elixir of Bismuth, Gentian, Iron and Strychnine.

Strychnine sulphategr.	
Distilled water fl.dr.	4
Elixir of bismuth, gentian and	
iron, enough to makefl.oz.	16

Dissolve the strychnine in the water, add pyrophosphate and bismuth salt.

the elixir, let stand a few hours and filter.

Each fluidram contains  $i\delta\sigma$  gr. of strychnine sulphate, nearly 1 gr. bismuth salt and nearly 1 gr. of iron phosphate, and represents about  $1\frac{1}{2}$  gr. of gentian.

## Elixir of Bismuth, Gentian and Strychnine.

Strychnine sulphategr. 11/4
Distilled waterfl.dr. 4
Elixir of bismuth and gentian,
enough to makefl.oz. 16

Dissolve the strychnine in the water, add the elixir, let stand a few hours, and filter. Each fluidram contains 180 gr. of strychnine sulphate, and nearly 1 gr. of bismuth salt and represents about 18 gr. of gentian.

## Elixir of Bismuth, Golden Seal and Iron.

Glycerite of hydrastisfl.oz.	1
Iron pyrophosphate, solublegr.	128
Distilled water, hotfl.dr.	4
Elixir of bismuth, enough to	
makefl.oz.	16

Dissolve the iron salt in the water, and the glycerite and elixir, let stand for a day or two, and filter. The elixir must be perfectly neutral.

Each fluidram contains 1 gr. of iron pyrophosphate, and nearly 2 gr. of bismuth salt and represents 33/2 gr. of hydrastis:

#### Elixir of Bismuth and Golden Seal.

(Elixir of Bismuth and Hydrastis.)

Glycerite of hydrastis......fl.oz. 1

Elixir of bismuth........fl.oz. 15

Mix, let stand for several days if possible, and filter. The elixir of bismuth must be exactly neutral before adding the glycerite.

Each fluidram represents 33/4 gr. of hydrastis and contains nearly 2 gr. of citrate of bismuth and ammonium.

#### Elixir Bismuth and Iron.

Iron pyrophosphate, solublegr.	128
Distilled water, hotfl.dr.	4
Elixir of bismuthfl.oz.	8
Simple elixir, enough to make.fl.oz.	16

Dissolve the iron salt in water and add the elixirs.

Each fluidram contains 1 gr. each of iron pyrophosphate and bismuth salt.

#### Elixir of Bismuth, Iron and Pepsin.

Citrate of bismuth and ammo-	
niumgr.	128
Iron pyrophosphategr.	128
Pepsin, puregr.	
Distilled water, hotfl.oz.	1
Water of ammoniasuffic	
Simple elixir, enough to make.fl.oz.	16

Triturate the bismuth with 4 fluidrams of water, allow the insoluble matter to subside, decant the clear liquid, to the residue add gradually ammonia water until solution occurs, carefully avoiding any excess, and mix this with the decanted portion. Dissolve the iron pyrophosphate in the remainder of the water; also dissolve the pepsin in 12 fluidounces of elixir by frequent agitation. Mix the three liquids, add the remainder of the elixir, and filter.

Each fluidram contains 1 gr. each of iron pyrophosphate, pepsin, and citrate of bismuth and ammonium.

## Elixir of Bismuth, Iron, Pepsin and Quinine.

	hydrochlorategr. &	32
Elixir of	bismuth, iron and pep-	
sin	fl.oz. 1	16

Mix, dissolve by agitation, and filter, if necessary.

Each fluidram contains 1 gr. each of pepsin, iron pyrophosphate and citrate of bismuth and ammonium and ¼ gr. of quinine hydrochlorate.

## Elixir of Bismuth, Iron and Strychnine.

Iron pyrophosphate, solublegr.	128
Strychnine sulphategr.	11/4
Distilled waterfl.oz.	1
Elixir of bismuthfl.oz.	8
Simple elixir, enough to make fl. oz.	16

Dissolve the iron salt and strychnine salt separately in 4 fluidrams of the water; add the two elixirs, and filter if necessary. The elixir of bismuth must be perfectly neutral.

Each fluidram contains The gr. of strychnine sulphate and 1 gr. each of iron pyrophosphate and bismuth salt.

## Elixir of Bismuth, Nux Vomica and Pepsin.

Tincture of nux vomica....fl.dr. 5½ Elixir of pepsin and bismuth, N. F., enough to make...fl.oz. 16 Each fluidram contains nearly 1 gr. of pepsin and 2 gr. of citrate of bismuth and ammonium and represents about ½ gr. of nux vomica.

#### Elixir of Bismuth and Pancreatin.

Citrate of bismuth and ammo-	
niumgr.	128
Pancreatin, puregr.	
Distilled waterfl.oz.	1
Water of ammoniasuffici	ent
Tincture of cudbearfl.dr.	2
Simple elixir, enough to make.fl.oz.	16

Triturate the bismuth salt with the water, allow the insoluble portion to subside, decant the clear liquid, add sufficient ammonia water to dissolve the residue, add this solution and the decanted portion to 12 fluidounces of elixir mixed with the tincture, then add the pancreatin, agitate occasionally until the latter is apparently dissolved, filter in a well-covered funnel, and add enough elixir through the filter to make the filtrate measure 16 fluidounces.

Each fluidram contains 1 gr. each of pancreatin and citrate of bismuth and ammonium.

## Elixir of Bismuth, Pancreatin and Pepsin.

Refer to "Elixir of Pancreatin" and its combinations.

#### Elixir of Bismuth and Pepsin.

Pepsin, puregr.	128
Bismuth and ammonium citrategr.	256
Water of ammoniasuffic	ient
Glycerinfl.oz.	2
Alcoholfl.oz.	
Simple syrupfl.oz.	4
Compound elixir of taraxacum.fl.oz.	1
Purified talcumgr.	120
Water, enough to makefl.oz	

Dissolve the pepsin in 4 fluidounces of water. Dissolve the bismuth and ammonium citrate in 1 fluidounce of warm water, allow the solution to stand until clear, if necessary; then decant the clear liquid, and add to the residue just enough water of ammonia to dissolve it, carefully avoiding an excess. Then mix the two solutions, and add the glycerin, compound elixir of taraxacum and alcohol. Thoroughly incorporate the talcum with the mixture, filter it through a wetted filter, and pass enough water through the

b

filter to make the filtrate measure 12 fluidounces. To this add the syrup.

Each fluidram represents 1 gr. of pepsin and 2 gr. of bismuth and ammonium citrate.

-N. F.

## Elixir of Bismuth, Pepsin and Quinine.

Quinine hydrochlorategr.		32
	pepsin and bismuth,	10
N. F.,		16

Mix and dissolve by agitation.

Each fluidram contains ¼ gr. of quinine hydrochlorate, of pepsin, and 2 gr. of citrate of bismuth and ammonium.

## Elixir of Bismuth, Pepsin and Strychnine.

Ctural min a autobata	11/
Strychnine sulphate.	gr. 1%
Elixir of pepsin and	bismuth, fl.oz. 16

Dissolve the strychnine sulphate in the elixir by agitation.

Each fluidram represents 180 gr. of strychnine sulphate, 1 gr. of pepsin, and 2 gr. of bismuth and ammonium citrate.—N. F.

## Elixir of Bismuth, Pepsin and Wafer Ash.

Fluid extract of	wafer ashfl.oz.	2
Elixir of pepsin	and bismuth,	
	fl.oz.	14

Mix, allow to stand 24 hours, and filter through purified talcum.

Each fluidram represents  $7\frac{1}{2}$  gr. of wafer ash, nearly 1 gr. of pepsin and nearly 2 gr. of citrate of bismuth and ammonium.

#### Elixir of Bismuth and Quinine.

Quinine hydrochlorategr. 32	
Elixir of bismuth, enough to	
makefl.oz. 16	

Dissolve the quinine salt in the elixir (which should be neutral) by agitation and filter, if necessary.

Each fluidram contains 1 gr. of quinine hydrochlorate and 2 gr. of bismuth salt.

#### Elixir of Bismuth and Strychnine.

· Strychnine sulphategr.	1 1/2
Distilled waterfl.dr.	$4^{\prime}$
Elixir of bismuthfl.oz. 18	$5\frac{1}{2}$

Dissolve the alkaloidal salt in the water and add to the elixir, which latter should be neutral.

Each fluidram contains 180 gr. of strychnine sulphate and nearly 2 gr. of bismuth salt. fuga.

#### Elixir, Bitter. (Elixir Amarum.)

Extract of wormwoodav. oz.	31/4
Oleosaccharate of peppermint.av.oz.	13/4
Aromatic tincture, N. Ffl.oz.	13/4
Bitter tincture, N. Ffl.oz.	13/4
Waterfl.oz.	81

Triturate the extract and oleosaccharate with the water to a smooth condition and add the other ingredients. This preparation should be cloudy and of a dark brown color.—Germ. Pharm.

#### Elixir of Blackberry.

Fluid extract of rubus	fl. oz.	2
Tincture of vanilla	fl.dr.	4
Compound elixir of taraxacur	n fl. oz.	4
Simple elixir, enough to make	fl.oz.	16
Each fluidram represents 7½	gr. of b	lack-
erry root bark.		

#### Elixir of Blackberry, Compound.

Blackberry rootgr.	480
Gallsgr.	
Cinnamon, Saigongr.	480
Clovesgr.	
Mace	60
Gingergr.	60
Blackberry juicefl.oz.	24
Simple syrupfl.oz.	12
Glycerinfl.oz.	12
Diluted alcoholsuffic	

Reduce the solids to a moderately coarse powder, moisten it with diluted alcohol, and percolate it with this menstruum in the usual manner, until 16 fluidounces of percolate are obtained. To this add the blackberry juice, syrup and glycerin, and mix thoroughly.

## Elixir of Black Haw. (Elixir of Viburnum Prunifolium.)

Fluid extract of black haw...fl.oz. 2
Compound tincture of cardamom...........fl.dr 9½
Aromatic elixir, enough to
make............fl.oz. 16

Mix, allow the mixture to stand a few days, if convenient, and filter.

## Elixir of Black Cohosh. (Elixir of Cimicifuga.)

Fluid extract of black cohoshfl.oz.	4
Alcohol	1
Simple elixirfl.oz.	11

Mix, let stand 24 hours, and filter through purified talcum.

Each fluidram represents 7½ gr. of cimicifuga.

#### Elixir of Black Cohosh, Compound.

(Compound Elixir of Cimicifuga.)

Fluid extract of cimicifugafl.dr.	$9\frac{1}{2}$
Fluid extract of wild cherryfl.dr.	8
Fluid extract of sanguinariafl.dr.	31/4
Fluid extract of licoricefl.dr.	31/4
Simple elixirfl.oz.	13

Mix, allow to stand for 24 hours, and filter through purified talcum.

Each fluidram represents about  $4\frac{1}{2}$  gr. of cimicifuga, 4 gr. of wild cherry, and  $1\frac{1}{2}$  gr. each of sanguinaria and licorice.

Each fluidram represents about 7½ gr. of black haw.—N. F.

#### Elixir of Black Haw, Compound.

(Compound Elixir of Viburnum Prunifolium.)

Fluid extract of black hawfl.oz.	2
Fluid extract of hydrastisfl.oz.	2
Fluid extract of Jamaica dog-	
woodfl.oz.	1
Simple elixirfl.oz.	
Simple clixii	TT

Mix, allow to stand for 24 hours, and filter.

Each fluidram represents  $7\frac{1}{2}$  gr. each of black haw and golden seal and  $3\frac{3}{4}$  gr. of Jamaica dogwood.

This is listed in manufacturers' catalogues under the names "Sedative Cordial" and "Sedative Elixir."

#### Elixir of Blue Flag.

Fluid extract of blue flagfl.oz.	4
Alcoholfl.oz.	1
Simple elixirfl.oz.	
Simple enxir	11

Mix, allow to stand for 24 hours, and filter. Each fluidram represents 15 gr. of blue flag.

#### Elixir of Blue Flag and Wahoo.

Fluid extract of blue flagfl.oz.	23/
Fluid extract of wahoofl.oz.	
Alcoholfl.oz.	
Simple elixirfl.oz. 1	.0

Mix, allow to stand for 24 hours and filter through talcum.

Each fluidram represents about 10 gr. each of blue flag and wahoo.

#### Elixir of Three (or Triple) Bromides.

Potassium Sodium bro				
Elixir of	caffeine,	enough	to	

Mix, dissolve by agitation, and filter, if necessary.

Each fluidram contains 8 gr. of each of the bromides of potassium, sodium, and caffeine.

#### Elixir of Six Bromides.

Potassium bromidegr.	640
Sodium bromidegr.	640
Ammonium bromidegr.	384
Calcium bromidegr.	192
Lithium bromidegr.	64
Iron bromidegr.	64
Compound tincture of cudbear.fl.dr.	2
Simple elixir, enough to make.fl.oz.	16

Dissolve by agitation and filter, if necesary.

Each fluidram contains 5 gr. each of potassium and sodium bromides, 3 gr. of ammonium bromide,  $1\frac{1}{2}$  gr. of calcium bromide, and 1 gr. each of lithium and iron bromides.

#### Elixir of Bromide of Ammonium.

Refer to Elixir of Ammonium Bromide.

#### Elixir of Bromide of Caffeine.

Refer to Elixir of Caffeine.

#### Elixir of Bromide of Calcium.

Refer to Elixir of Calcium Bromide.

#### Elixir of Bromide of Lithium.

Refer to Elixir of Lithium Bromide.

#### Elixir of Bromide of Potassium.

Potassium Citric acid	bromidegr.	1280
Aromatic	elixir, enough to	
make	fl.oz.	16

Dissolve the potassium bromide and citric acid in the elixir, by agitation, and filter.

Each fluidram contains 10 gr. of potassium bromide.—N. F.

#### Elixir of Bromide of Sodium.

Sodium bromidegr. 1	280
Citric acidgr.	30
Aromatic elixir, enough to	
make fl.oz.	16

Dissolve the sodium bromide and citric acid in the elixir, by agitation, and filter, if necessary.

Each fluidram contains 10 gr. of sodium bromide.—N. F.

## Elixir of Bromide of Sodium and Lupulin.

Refer to Elixir of Lupulin and its combinations.

#### Elixir of Bromide of Zinc.

Zinc bromide.								gr.	128
Citric acid									
Simple elixir						•		fl.oz.	16

Dissolve by agitation and filter, if necessary.

Each fluidram contains 1 gr. of zinc bromide.

#### Elixir of Buchu.

Fluid extract of buchufl.oz.	2
Alcohol	1
Simple syrupfl.oz.	
Magnesium carbonategr.	
Aromatic elixir, enough to make fl.oz.	

Mix the fluid extract of buchu with the alcohol, then add 12 fluidounces of aromatic elixir, and the syrup; incorporate with it the magnesium carbonate, and filter. Finally, pass enough aromatic elixir through the filter to make 16 fluidounces.

Each fluidram represents about  $7\frac{1}{2}$  gr. of buchu.—N. F.

#### Elixir of Buchu, Compound.

Compound fluid extract of bu-	
chufl.oz.	4
Alcoholfl.oz.	1
Simple syrupfl.oz.	1
Magnesium carbonategr.	120
Aromatic elixir, enough to make fl. oz.	16

Mix the compound fluid extract of buchu with the alcohol, then add 8 fluidounces of aromatic elixir, and the syrup; incorporate with it the magnesium carbonate, and filter. Finally, pass enough aromatic elixir through the filter to make 16 fluidounces.

Each fluidram represents 15 m. of compound fluid extract of buchu.—N. F.

#### Elixir of Buchu and Juniper, Compound. (Rheumatic Elixir.)

Fluid extract of buchufl.dr.	61
Fluid extract of barberry bark fl.dr.	31/4
Fluid extract of juniper berries fl.dr.	31/4
Sodium salicylategr.	160
Simple syrupfl.oz.	1
Alcohol fl.oz.	1
Simple elixir enough to makefl.oz.	16

Mix all, let stand for 24 hours, and filter through purified talcum.

Each fluidram contains  $1\frac{1}{4}$  gr. of sodium salicylate, and represents 3 gr. of buchu, and  $1\frac{1}{2}$  gr. each of barberry bark and juniper berries.

## Elixir of Buchu, Juniper and Potassium Acetate.

Fluid extract of buchufl.dr.	12
Fluid extract of juniper berries fl.dr.	4
Potassium acetategr.	192
Alcohol fl. oz.	1
Simple syrup fl.oz.	1
Simple elixirfl.oz.	12

Mix, allow to stand for 24 hours and filter through talcum.

Each fluidram contains  $1\frac{1}{2}$  gr. of potassium acetate, and represents about  $5\frac{1}{2}$  gr. of buchu, and 2 gr. of juniper berries.

## Elixir of Buchu, Juniper, Uva Ursi and Potassium Acetate.

Fluid extract of buchufl.oz.	2
Fluid extract of uva ursi fl.dr.	11
Fluid extract of juniper berries fl.dr.	$5\frac{1}{2}$
Potassium acetateav.oz.	$1\frac{7}{2}$
Alcoholfl.oz.	1
Simple syrupfl.oz.	1
Simple elixir enough to make fl.oz.	16

Mix, allow to stand for 24 hours, and filter through purified talcum.

Each fluidram contains 5 gr. of potassium acetate, and represents  $7\frac{1}{2}$  gr. of buchu, 5 gr. of uva ursi, and  $2\frac{1}{2}$  of juniper berries.

#### Elixir of Buchu and Pareira.

Fluid extract of buchufl.oz.	2
Fluid extract of pareira bravafl.oz.	2
Alcohol fl.oz.	1
Simple syrup fl.oz.	1
Simple elixirfl.oz.	

Mix, allow to stand for 24 hours, and filter through purified talcum.

Each fluidram represents  $7\frac{1}{2}$  gr. each of buchu and pareira brava.

#### Elixir of Buchu and Pareira, Compound.

Fluid extract of buchufl.dr.	8
Fluid extract of juniper berries fl.dr.	4
Fluid extract of pareira bravafl.dr.	2
Fluid extract of stone-rootfl.dr.	2
Alcoholfl.oz.	1
Simple syrupfl.oz.	1
Simple élixir fl.oz.	12

Mix, allow to stand for 24 hours, and filter through purified talcum.

Each fluidram represents about 4 gr. of buchu, 2 gr. of juniper berries, and 1 gr. each of pareira brava and collinsonia.

## Elixir of Buchu and Potassium Acetate.

Potassium acetate.....gr. 640 Elixir of buchu, enough to make fl. oz. 16

Dissolve the potassium acetate in about 12 fluidounces of elixir of buchu, filter, if necessary, and add the remainder of the elixir of buchu.

Each fluidram represents 5 gr. of potassium acetate, and about 7 gr. of buchu.— N. F.

## Elixir of Buckthorn. (Elixir of Frangula.)

Fluid extract of frangulafl.oz.	4
Alcoholfl.oz.	
Compound elixir of taraxacumfl.oz.	
Aromatic elixirfl.oz.	

Mix them, allow the mixture to stand 48 hours, if convenient, and filter.

Each fluidram represents 15 gr. of frangula.—N. F.

#### Elixir of Buckthorn and Senna.

Fluid extract of frangulafl.oz.	2
Elixir of sennafl.oz.	14

Each fluidram represents  $7\frac{1}{2}$  gr. of buckthorn bark, and 26 gr. of senna.

#### Elixir of Butyl Chloral Hydrate.

Refer to "Elixir of Croton Chloral Hydrate" for above and its combinations.

#### Elixir of Caffeine.

Caffeine	128
Diluted hydrobromic acid fl.dr.	
Syrup of coffeefl.oz.	4
Aromatic elixir enough to make fl.oz.	16

Rub the caffeine in a mortar, with the diluted hydrobromic acid and about 2 fluidounces of aromatic elixir, until solution is effected; then add the syrup, and lastly the remainder of the aromatic elixir. Filter, if necessary.

Each fluidram contains 1 gr. of caffeine.

-N. F.

#### Elixir of Calcium Bromide.

Calcium bromide gr.	640
Citric acidgr.	
Aromatic elixir, enough to make fl.oz.	16

Dissolve the calcium bromide and the citric acid in about 12 fluidounces of aromatic elixir by agitation, then add the remainder of the aromatic elixir, and filter, if necessary.

Each fluidram contains 5 gr. of calcium bromide.—N. F.

#### Elixir of Calcium Hypophosphite.

Calcium hypophosphitegr.	256
Citric acidgr.	
Aromatic elixir, enough to make fl. oz.	

Dissolve the calcium hypophosphite in 14 fluidounces of aromatic elixir, and filter; dissolve the citric acid in the filtrate, and pass the remainder of the aromatic elixir through the filter.

Each fluidram contains 2 gr. of calcium hypophosphite.—N. F.

## Elixir of Calcium and Sodium Hypophosphites and Cherries.

Refer to "Elixir of Cherries" for this and its combinations.

#### Elixir of Calcium and Sodium Hypophosphites with Cinchona.

Refer to "Elixir of Cinchona and Hypophosphites."

#### Elixir of Calcium and Sodium Hypophosphites with Malt.

Calcium hypophosphitegr.	128
Sodium hypophosphitegr.	128
Adjuvant elixirfl.oz.	8
Fluid extract of malt, N. Ffl.oz.	8

Dissolve the salts in the elixir by trituration, filter, and add the malt extract.

Each fluidram contains 1 gr. each of the hypophosphites of calcium and sodium.

#### Elixir of Calcium and Sodium Hypophosphites with Tar.

Calcium hypophosphitegr.	128
Sodium hypophosphitegr.	
Distilled water, hot fl.oz.	2
Elixir of tarenough to make fl.oz.	16

Dissolve the salts in the water, add the elixir, and filter.

Each fluidram contains 1 gr. each of the hypophosphites.

#### Elixir of Calcium Iodide.

I. Calcium iodide......av.oź. 1½
Simple elixir..enough to make fl.oz. 16
Dissolve by agitation, and filter.

II. Inasmuch as calcium iodide is an unstable compound, it should be prepared as needed, and the following formula should therefore receive preference:

Hydrate the calcium oxide with 6 fluidounces of water, add the solution of iron oxide, heat to boiling, allow to stand a few minutes, decant the clear liquid, add to the residue a fresh portion of distilled water, heat again to boiling, decant as before, and repeat the process again until the mixed decantates measure 10 fluidounces; add the alcohol containing the spirit, let stand for an hour or more, filter, in the filtrate dissolve the sugar by agitation, and strain if necessary.

Each fluidram contains 5 gr. of calcium iodide.

#### Elixir of Iodo-Bromide of Calcium, Compound. (Compound Elixir of Calcium Bromide with Iodides.)

Calcium bromide gr.	256
Sodium iodidegr.	256
Potassium iodidegr.	256
Magnesium chloridegr.	256
Compound fluid extract of sar-	
saparillafl.oz.	2
Compound fluid extract of stil-	
lingiafl.oz.	2
Aromatic elixirfl.oz.	4
Sugarav.oz.	$4\frac{1}{2}$
Waterenough to make fl.oz.	16

Dissolve the salts in the water, add the sugar, and to this syrup add the fluid extracts previously mixed with the aromatic elixir; after standing for 2 days, filter and add the remainder of the water.

#### Elixir of Calcium Lactophosphate.

Calcium lactategr	. 128
Phosphoric acid (85 per cent.) fl.dr	. 1
Waterfl.oz	. 1
Simple syrup fl.oz	. 1
Aromatic elixir, enough to make fl. or	z. 16
Triturate the calcium lactate w	vith th

Triturate the calcium lactate with the phosphoric acid, water and syrup, until the salt is dissolved, then add the remainder of the aromatic elixir, and filter.

Each fluidram represents 1 gr. of calcium lactate, or about 1½ gr. of so-called calcium lactophosphate.—N. F.

## Elixir of Calcium Lactophosphate and Cinchona.

Detannated elixir of cinchona...fl.oz. 8 Elixir of calcium lactophosphate fl.oz. 8

#### Elixir of Calcium Lactophosphate, Cinchona and Iron.

Calcium lactategr. 6	4
	1/2
Water of ammoniafl.dr.	4
Citric acid gr. 12	0.5
Elixir of cinchona and iron,	
enough to makefl.oz. 1	.6

Dissolve the calcium lactate in 7 fluidounces of elixir of cinchona and iron, with the aid of the phosphoric acid; then add the citric acid, and when this is dissolved, the water of ammonia; finally, add the remainder of the elixir of cinchona and iron, and filter.

Each fluidram contains ½ gr. of calcium lactate (or about ¾ gr. of so-called calcium lactophosphate,) and nearly 2 gr. of iron phosphate.—N. F.

#### Elixir of Calcium Phosphate.

Calcium phosphategr.	640
Hydrochloric acid, concentra-	
tedfl.dr.	5
Water fl.oz.	1
Tincture of cudbear fl.dr.	$^2$
Simple elixir, enough to make fl.oz.	16

Mix the calcium phosphate with the water, add the acid, dissolve, add the clixir, and then the tincture.

Each fluidram contains 5 gr. of calcium phosphate.

#### Elixirs of Calisaya.

Elixir of calisaya, and its various combinations, will be referred to under the head of "Elixir of Cinchona."

#### Elixir of Cascara Sagrada.

Fluid extract of cascara sagrada fl.oz. 4 Compound elixir of taraxacum fl.oz. 12

Mix them, allow the mixture to stand a few days, if convenient, and filter.

Each fluidram represents 15 gr. of cascara sagrada.—N. F.

11.	
Tincture of orange peelfl.oz.	14
Alcohol	6
Cinnamon water fl.oz.	21
Simple syrupfl.oz.	43/4
Fluid extract of cascara sagrada,	
of the Brit. pharmfl.oz.	
-Brit, Form	a.

#### Elixir of Cascara Sagrada, Compound. | Elixir of Celery, Compound.

(Laxative Elixir.)

Fluid extract of cascara sagradafl.oz. Fluid extract of senna fl.dr.	
Fluid extract of butternutfl.oz.	
Compound elixir of taraxacum, enough to makefl.oz	16

Mix them, allow to stand a few days, if convenient, and filter .- N. F.

#### Elixir of Cascara Sagrada with Sodium Salicylate.

Elixir of cascara sagradafl.oz.	5
Sodium salicylateav.oz.	$2\frac{1}{2}$
Simple elixir, enough to make fl.oz.	16

Mix, dissolve by shaking, and filter if necessary.

Each fluidram represents approximately 2 gr. cascara sagrada, and contains very nearly I gr. of sodium salicylate.

#### Elixir, Castillon's.

Cinchona, coarsely powderedgr.	160
Gentian, coarsely powderedgr.	
Ipecac, coarsely powdered gr.	80
Columbo, coarsely powderedgr.	80
Cinnamon, coarsely powdered gr.	20
Aqueous extract of opiumgr.	20
Diluted alcoholsuffic	ient

Macerate the drugs with 16 fluidounces of diluted alcohol for 7 days, and filter, adding enough menstruum through the filter to make up 16 fluidounces of filtrate.

#### Elixir, Cathartic, Compound.

Fluid extract of sennafl.oz.	. 2
Fluid extract of podophyllum.fl.oz.	
Fluid extract of leptandrafl.dr. Fluid extract of jalapfl.dr.	6
Rochelle saltav.oz. Sodium bicarbonategr.	
Compound elixir of taraxacum, fl.oz.	
Elixir of licorice, enough to makefl.oz.	16

Mix the fluid extracts with the compound elixir of taraxacum; in the mixture, dissolve the salts by agitation, and add the elixir of

The product should not be filtered, and should be shaken up whenever any of it is dispensed.-N. F.

Compound elixir of cascara sagrada, N. F., is also known as "laxative elixir," or "elixir purgans," and may be used as a cathartic elixir instead of the above.

Fluid extract of celery seed fl.oz.	1
Fluid extract of cocafl.oz.	1
Fluid extract of kolafl.oz.	1
Fluid extract of black haw fl. oz.	1
Alcohol fl.oz.	2
Aromatic elixir, enough to make fl.oz.	16

Mix the alcohol with 4 fluidounces of aromatic elixir; to this add the fluid extract of celery in several portions, shaking after each addition, and afterwards the other fluid extracts; finally, add the remainder of the elixir, allow the mixture to stand 24 hours, and filter .- N. F.

#### Elixir of Celery and Guarana.

Fluid extract of celery seed fl. oz.	2
Fluid extract of guaranafl.oz.	2
Aromatic elixirfl.oz.	

Mix, allow to stand for 24 hours, and filter through talcum.

Each fluidram represents 7½ gr. each of celery and guarana.

#### Elixir of Cherries. (Elixir Cerasorum.)

Ripe, sour cherries, free from
stems av. oz. 8
Alcoholfl.oz. 2
Glycerin fl.oz. 1
Simple syrupsufficient.

Crush the cherries and stones to a pulp, add the alcohol and glycerin, macerate for 7 days, press and filter, and to the filtrate add simple syrup enough to make 16 fluidounces.

#### Elixir of Cherries with Calcium and Sodium Hypophosphites.

Calcium hypophosphite gr.	
Sodium hypophosphitegr. Elixir of cherries, enough to	128
makefl.oz.	16

Triturate the two salts to fine powder, add to the elixir, dissolve by agitation, and

Each fluidram contains 1 gr. each of sodium and calcium hypophosphites.

#### Elixir of Chloral Hydrate.

Chloral hydrate, crystal.....gr. 640 Simple elixir, enough to make. .fl.oz 16

Mix, dissolve by agitation, and filter, if necessary.

Each fluidram contains 5 gr. of chloral hydrate.

# Elixir of Chirata. fl.oz. 4 Simple elixir fl.oz. 12

Each fluidram represents 1½ gr. of chirata.

## Elixir of Chloral Hydrate and Ammonium Valerianate.

Refer to "Elixir of Ammonium Valerianate" and its combinations.

## Elixir of Chlorides of Arsenic and Iron. (Elixir of Two Chlorides.)

Solution of arsenious acid.....fl.dr. 10½ Tincture of citrochloride of iron fl.dr. 5½ Simple elixir.....fl.oz. 14

Each fluidram contains  $\frac{1}{2}$  gr. of arsenious acid (as so-called "chloride of arsenic") and about  $\frac{1}{2}$  gr. of iron chloride.

#### Elixir of Chlorides of Arsenic, Iron and Mercury. (Elixir of Three Chlorides.)

Each fluidram contains ½ gr. of iron protochloride, 1½ gr. of mercuric chloride and 2½ gr. of arsenious acid (as so-called "chloride of arsenic").

## Elixir of Four Chlorides. (Four Chlorides.)

Mercuric chloridegr.	2
Solution of arsenicfl.dr	$5\frac{1}{2}$
Tincture of ferric chloridefl.oz.	2
Hydrochloric acid, diluted fl.dr.	11
Syrup of gingerfl.oz.	4
Water, enough to makefl.oz.	16
3.51 11 1 1 (1)	

Mix, dissolve and filter, if necessary.

Each fluidram contains about 46 gr. of arsenious acid (as so-called "chloride of arsenic"), 44 gr. of mercuric chloride, about 34 gr. of ferric chloride, and about 5 minims of diluted hydrochloric acid.

#### Elixir of Chloroform.

Chloroformfl.dr	4
Alcohol fl.oz.	2
Simple elixir, enough to makefl.oz.	16

Mix the alcohol and chloroform, and add the elixir.

Each fluidram contains very nearly 2 minims of chloroform.

#### Elixir of Chloroform, Compound.

(Chloroform Paregoric.)

Chloroform. fl.oz. 3 Tincture of opium, fl.oz. 3 Spirit of camphor, fl.oz. 3	
Spirit of camphorfl.oz. 3	fl.oz. 3
Spirit of camphorfl.oz. 3	fl.oz. 3
Aromatic spirit of ammoniafl.oz. 3	
Alcoholfl.oz. 33	
Oil of cassia	

Mix the chloroform with the alcohol, then add the oil, aromatic spirit of ammonia, spirit of camphor and tincture of opium. Allow the mixture to stand a few hours, and filter in a well-covered funnel.

Each fluidram represents about 1 gr. of opium and 11 minims of chloroform.—N. F.

#### Elixir of Cinchona.

Tincture of cinchonafl.oz.	24
Simple syrupfl.oz.	
Glycerin	
Aromatic elixirfl.oz.	$9\frac{1}{2}$

Mix the liquids, allow to stand as long as convenient, and filter through a wetted filter.

Each fluidounce represents about 14 gr. of yellow cinchona.—N. F.

The compound elixir of quinine, N. F., is sometimes dispensed as elixir of cinchona or calisaya.

#### Elixir of Cinchona, Detannated.

Detannated tincture of cinchona. fl. oz.  $2\frac{1}{2}$  Simple syrup. fl. oz. 2 Glycerin fl. oz. 2 Aromatic elixir fl. oz.  $9\frac{1}{2}$  Mix the liquids, and filter, if necessary.

Each fluidounce represents about 14 gr. of yellow cinchona.—N. F.

II. Compound elixir of quinine is often used as a detannated elixir of calisaya and is recommended to be used even by some of the best pharmacists, who see no advantage in employing a tedious process of detannation, when simple solution will yield practically the same results.

ш

Yellow cinchonagr. 240
Curacoa orange peelgr. 160
Coriandergr. 40
Cardamomgr. 15
Cinnamongr. 30
Anisegr. 10
Cacao (Baker's)gr. 80
Purified talcumgr. 120
Alcohol,
Waterof each, sufficient
Spirit of orangefl.dr. 1½
Sugar av oz 5

Reduce the cinchona, orange peel, spices, and cacao together to a moderately fine powder, extract by slow percolation with a menstruum, consisting of 1 part, by measure, of alcohol, and 3 parts of water, so as to obtain 8 fluidounces of percolate.

Prepare-hydrate of iron as directed under "Elixir of Gentian," and detannate the above percolate as there directed, washing the residue, with a mixture similar to the menstruum used, until the liquid measures 12 fluidounces. To the latter add the spirit of orange and the talcum, shake well, filter, washing the filter with the same liquid that was used before until the filtrate measures 12½ fluidounces. To the filtrate add the sugar, and dissolve by agitation; strain, or filter, if this be necessary.

IV.

Yellow cinchona, in fine powder.gr. 480 Quicklime....gr. 360

Slake the lime in the least quantity of water, mix intimately with the bark, moisten with alcohol, pack in a glass percolator, and percolate slowly with alcohol as a menstruum in the usual manner until 8 fluidounces are obtained. The first portions of the percolate must be tested with solution of ferric chloride; so long as this liquid affords a discoloration it must be returned to the percolator.

Prepare flavoring as follows:

Oil of orange peelfl.dr.	2
Oil of carawayfl.dr.	
Oil of cassiafl.dr.	
Oil of anise	
Alcohol fl.oz.	1

Triturate one fluidounce of this mixture with 60 gr. of purified talcum, 3 fluidrams of alcohol and 1 fluidounce of water, and filter.

To 12 fluidounces of simple syrup add 5 gr. of citric acid and mix this with the tincture first obtained, subsequently adding 11½ fluidounces of water. Then to this add the flavoring mixture, mix well, and filter the whole.

· Control of the cont	
Yellow cinchonagr. 4	480
Saigon cassiagr.	80
Coriandergr.	80
Nutmeggr.	20
Star anisegr.	20
Sugarav.oz.	10
Alcohol,	
Water, of eachsuffici	ent
Spirit of orangefl.dr.	2
Purified talcum or.	$120^{\circ}$

Reduce the cinchona, cassia, coriander, nutmeg and anise to a moderately fine powder, and extract by percolation with a mixture of alcohol and water, in the proportion of 1 by measure of the former to 3 of the latter, until 22 fluidounces of percolate are obtained. Now beat the white of 1 egg with a portion of the percolate, add the remainder of the percolate, and set aside for 24 hours, agitating occasionally. Test at the end of the specified period of time with solution of ferric chloride and if discoloration occurs, the white of another egg may be added as before, allowing to stand 24 hours, then filtering. Wash the filter with a liquid similar to the menstruum used until 25 fluidounces of filtrate are obtained. To this add the spirit of orange and purified talcum, filter; to the filtrate add the sugar, dissolve by agitation and strain, or filter, if necessary.

Each fluidram represents about 13/4 gr. of cinchona.

## Elixir of Cinchona, Compound. (Elixir of Cinchona and Coca.)

Fluid extract of cinchona	.fl.dr.	10
Fluid extract of coca	.fl.dr.	10
Tincture of cacao	.fl.oz.	$2\frac{1}{2}$
Simple elixir	.fl.oz.	11
Mix, and filter, if necessary.		

Each fluidram represents about  $4\frac{1}{2}$  gr. each of cinchona and coca.

Elixir of Cinchona and Beef.
Elixir of Cinchona, Beef and Iron.
Elixir of Cinchona, Beef, Iron and
Strychnine.

Refer for above to Elixir of Beef and its combinations.

Elixir of Cinchona and Bismuth.

Elixir of Cinchona, Bismuth and Iron.

Elixir of Cinchona, Bismuth, Iron
and Pepsin.

Elixir of Cinchona, Bismuth, Iron, Pepsin and Strychnine.

Elixir of Cinchona, Bismuth, Iron and Strychnine.

Elixir of Cinchona, Bismuth and Pepsin.

Refer for above to Elixir of Bismuth and its combinations.

## Elixir of Cinchona and Calcium Lactophosphate.

#### Elixir of Cinchona, Calcium Lactophosphate and Iron.

Refer for above Elixir of Calcium Lacto phosphate and its combinations.

#### Elixir of Cinchona and Gentian.

Extract of gentiangr.	70
Tincture of vanillafl.dr.	2
Simple syrupfl.oz.	1
Aromatic spiritfl.dr.	
Elixir of cinchona, enough to	
makefl.oz.	16

Dissolve the extract in about 8 fluidounces of elixir, add the tincture, spirit and syrup, and the remainder of the elixir of cinchona, and filter the whole.

Each fluidram represents about 2 gr. of gentian and about  $1\frac{1}{2}$  gr. of cinchona.

## Elixir of Cinchona, Gentian and Iron Malate.

Malate of iron ("scales")gr.	128
Extract of gentiangr.	35
Simple syrupfl.oz.	4
Elixir of cinchonafl.oz.	6
Tincture of vanillafl.dr.	2
Oil of cinnamondrop.	1
Water, hotfl.oz.	1
Aromatic elixir, enough to make fl.oz.	16

Dissolve the iron salt and extract in the water, add the other ingredients and filter.

Each fluidram represents about 1 gr. of gentian and  $\frac{1}{2}$  gr. of cinchona, and contains 1 gr. of iron malate.

The malate of iron to be used should not be the ferrated extract of apples, but the pure malate of iron which appears in the scale form.

## Elixir of Cinchona and Hypophosphites.

Calcium hypophosphitegr. Sodium hypophosphitegr.	
Citric acidgr. Waterfl.oz.	30
Elixir of cinchona, enough to make	

Dissolve the hypophosphites and citric acid in the water, add the elixir of cinchona, and filter.

Each fluidram contains 1 gr. each of the hypophosphites of calcium and sodium.— N. F.

## Elixir of Cinchona and Iron. (Ferrated Elixir of Cinchona.)

Iron phosphate, solublegr. \$	256
Water, boilingfl.oz.	1
Compound elixir of quininefl.oz.	15

Dissolve the iron phosphate in the water, add the compound elixir of quinine, and filter.

Each fluidram contains 2 gr. of iron phosphate.—N. F.

## Elixir of Cinchona and "Protoxide" of Iron.

	" protoxide		
- iron		fl.oz.	$1\frac{1}{2}$
Glycerin		fl.oz.	$1\frac{1}{2}$
Elixir of cinc	hona, N. F	fl.oz.	13

Mix the solution and the glycerin, and add the elixir.

#### Elixir of Cinchona, Iron and Pepsin.

Pepsin, puregr.	128
Hydrochloric acidfl.dr.	1
Water	
Elixir of cinchona and iron,	
enough to makefl:oz.	16

Dissolve the pepsin in the water mixed with the hydrochloric acid, add the elixir of cinchona and iron; let the mixture stand a few days, if convenient, and filter.

Each fluidram represents 1 gr. of pepsin and about  $1\frac{1}{2}$  gr. of iron phosphate.—N. F.

#### Elixir of Cinchona, Iron and, Phosphorus.

Spirit of phosphorusfl.dr.	74
Elixir of cinchona and iron,	-
enough to makefl.oz.	16

Each fluidram contains 200 gr. of phosphorus, 2 gr. of iron phosphate, and represents nearly 2 gr. of cinchona.

## Elixir of Cinchona, Iron and Strychnine.

Strychnine sulphategr.	
Distilled water	z
enough to makefl.oz.	16

Dissolve the strychnine sulphate in the water and add the elixir.

Each fluidram contains 185 gr. of strychnine sulphate, and about 2 gr. of iron phosphate.—N. F.

#### Elixir of Cinchona and Pepsin.

I,	7					
-	Quinine	sulphate.	 	gr.	16	
		ne sulph				
7	Elixir of	pepsin	 	.fl.oz.	16	
	- //					ı

Dissolve the alkaloidal salts in the elixir and filter if necessary.

II. Pure pepsin gr.	128
Hydrochloric acidfl.dr.	
Detannated elixir of cinchona, enough to makefl.oz.	16

Dissolve by agitation and filter, using purified talcum, if necessary.

## Elixir of Cinchona, Pepsin and Strychnine.

Quinine sulphategr.	16
Cinchonine sulphategr.	8
Strychnine sulphategr.	
Elixir of pepsin, enough to make fl.oz.	

Dissolve the alkaloidal salts in the elixir, and filter, if necessary.

Each fluidram represents small quantities of cinchona alkaloids, 180 gr. of strychnine sulphate, and 1 gr. of pepsin.—N. F.

## Elixir of Cinchona with Phosphates.

Syrup of calcium lactophos-	
phate, U. S. Pfl oz.	4
Syrup of iron lactophosphatefl.oz.	2
Diluted phosphoric acidfl.oz.	1
Quinine sulphategr.	
Alcoholfl.oz.	
Spirit of orangefl.dr.	4
Waterfl.oz.	
	_

Dissolve the quinine salt in the alcohol previously mixed with the acid and spirit, pour this solution into the syrups previously mixed with the water, allow to stand for 2 days, and filter.

II.

Elixir of cinchona......fl.oz. 8 Compound syrup of phosphate..fl.oz. 8

III. The Elixir of Cinchona, Iron and Calcium Lactophosphate, N. F., may be dispensed under the above title.

#### Elixir of Cinchona and Strychnine.

Strychnine sulphategr. Detannated elixir of cinchonafl.oz.	1¼ 16
Dissolve by agitation.	

Each fluidram contains 180 gr. of strychnine sulphate, and represents nearly 2 gr. of cinchona.

#### Elixir of Cinchonidine.

Cinchoni	$_{ m dine}$	e sulphate.		gr.	128
Simple e	lixir			fl.oz.	16
Dissolve	by	agitation,	and	filter, if	neces-
arv.					

Each fluidram contains 1 gr. of cinchonidine sulphate.

#### Elixir of Cinchonidine and Ammonium Valerianate.

Elixir of Cinchonidine, Ammonium Valerianate and Iron Pyrophosphate.

Elixir of Cinchonidine, Ammonium Valerianate, Iron Pyrophosphate and Quinine.

Elixir of Cinchonidine, Ammonium Valerianate, Iron Pyrophosphate and Strychnine.

Elixir of Cinchonidine, Ammonium Valerianate and Quinine.

Elixir of Cinchonidine, Ammonium Valerianate, Quinine and Strychnine.

## Elixir of Cinchonidine, Ammonium Valerianate and Strychnine.

Refer for above to Elixir of Ammonium Valerianate and its combinations.

#### Elixir of Cinchonidine and Iron.

Iron pyrophosphate, soluble.	gr.	256
Cinchonidine sulphate	gr.	128
Distilled water, hot	.fl.dr.	6
Simple elixir		

Dissolve the iron pyrophosphate in the water, and the cinchonidine in the elixir; mix the two solutions and filter if necessary.

Each fluidram contains  $1~{\rm gr.}$  of cinchonidine sulphate and  $2~{\rm gr.}$  of iron pyrophosphate.

## Elixir of Cinchonidine, Iron and Strychnine.

Make this either by adding 1¼ gr. of strychnine sulphate to the preceding, or the elixir of iron phosphate, cinchonidine and strychnine may be employed.

#### Elixir of Coca.

Fluid extract of coca	.fl.oz.	2
Alcohol	.fl.oz.	1
Simple syrup	.fl.oz.	2
Tincture of vanilla	.fl.dr.	2
Purified talcum		
Aromatic elixir, enough to mak	e.fl.oz.	16

Mix the fluid extract with the alcohol, syrup and 10 fluidounces of aromatic elixir,

add the purified talcum and incorporate the latter thoroughly. Let the mixture stand during 48 hours, if convenient, shaking occasionally; then filter, add the tincture of vanilla to the filtrate, and pass the remainder of the elixir through the filter.

Each fluidram represents 71 gr. of coca .-N. F.

#### Elixir of Coca, Beef and Iron.

Refer for above to Elixir of Beef and its combinations.

#### Elixir of Coca and Cinchona.

Refer to Compound Elixir of Cinchona.

#### Elixir of Coca and Guarana. (Compound Elixir of Coca.)

Fluid extract of coca.....fl.oz. Fluid extract of guarana.....fl.oz. Purified talcum.....gr. 120 Compound elixir of taraxacum.fl.oz. 12

Mix the liquids, and thoroughly incorporate the purified talcum with the mixture; let it stand during 48 hours, if convenient, occasionally agitating, then filter.

Each fluidram represents 74 gr. each of coca and guarana.-N. F.

#### Elixir of Coca and Phosphorus.

Spirit of phosphorus......fl.dr. 15 Elixir of coca, enough to make.fl.oz. 16

Mix and filter if necessary.

Each fluidram contains 100 gr. of phosphorus and represents 61 gr. of coca.

#### Elixir of Codeine.

Codeine sulphategr.	16
Simple elixirfl.oz.	

Dissolve by agitation.

Each fluidram contains 1/8 gr. of codeine sulphate.

#### Elixir of Corydalis, Compound. (Alterative Elixir.)

attie Zimiii,	
Fluid extract of corydalisfl.oz.	1
Fluid extract of stillingiafl.oz.	1
Fluid extract of prickly ash bark fl.dr.	4
Fluid extract of blue flag fl.oz.	11
Alcoholfl.oz.	2
Potassium iodidegr.	
Aromatic elixir, enough to make fl. oz.	16

Mix the alcohol with the fluid extracts, dissolve the potassium iodide in the mixture, and add the aromatic elixir. Let the mixture stand a few days, if convenient, and filter.

iodide, and small quantities of the several fluid extracts.

#### Elixir of Codeine and Terpin Hydrate.

Codeine sulphategr.	16
Terpin hydrategr.	
Simple elixir, enough to make.fl.oz.	16

Dissolve by agitation and filter if necessary.

Each fluidram contains 1/8 gr. of codeine sulphate, and 2 gr. of terpin hydrate.

#### Elixir of Crampbark, Compound. (Compound Elixir of Viburnum Opulus.)

Fluid extract of crampbarkfl.dr.	10
Fluid extract of trilliumfl.oz.	$2\frac{1}{2}$
Fluid extract of aletrisfl.dr.	10
Compound elixir of taraxacumfl.oz.	11

Mix, allow to stand a few days, if convenient, and filter.-N. F.

#### Elixir of Croton Chloral Hydrate. (Elixir of Butyl Chloral Hydrate.)

Croton chloral hydrategr.	256
Alcoholfl.oz.	1
Tincture of cacaofl.oz.	2
Simple elixir, enough to make.fl.oz.	16

Dissolve the croton chloral in the alcohol, add the tincture and elixir, and filter, if necessary.

Each fluidram contains 2 gr. of croton chloral hydrate.

#### Elixir of Croton Chloral Hydrate and Quinine.

Ouining	aulphoto es	100
Quimine	sulphategr.	120
Till:	croton chloral hydrate fl.oz.	16

Reduce the quinine salt to fine powder, add the elixir, dissolve by agitation, and filter, if necessary.

Each fluidram contains 1 gr. of quinine sulphate and 2 gr. of croton chloral hydrate.

#### Elixir of Curacao. (Curacao Cordial.)

Spirit of curacaofl.dr.	2
Orris root, fine powdergr.	30
Deodorized alcoholfl.oz.	4
Citric acidgr.	50
Simple syrupfl.oz.	8
Magnesium carbonategr.	120
Distilled water, enough to make fl. oz.	

Mix the spirit of curacao with the alcohol, add the orris root, magnesium carbonate, and 3 fluidounces of water. Allow the mixture to stand 12 hours, occasionally agitating; Each fluidram contains 3 gr. of potassium then pour it on a wetted filter, returning the first portions of the filtrate until it runs through clear, and pass enough water through the filter to make the filtrate measure 8 fluidounces. In this dissolve the citric acid, and finally add the syrup.-N. F.

#### Elixir of Damiana. (Elixir of Turnera.) Fluid extract of damiana.....fl.oz. Magnesium carbonate.....gr. 240 Alcohol.....fl.oz. Glycerin.....fl.oz. Aromatic elixir, enough to make fl. oz. 16

Mix the fluid extract with the alcohol, glycerin and 8 fluidounces of aromatic elixir. Incorporate the magnesium carbonate thoroughly with the mixture by trituration, then filter through a wetted filter, and pass the remainder of the aromatic elixir through the

Each fluidram represents nearly 10 gr. of damiana.-N. F.

#### Elixir of Damiana, Iron, Nux Vomica and Phosphorus.

Fluid extract of damianafl.oz.	2
Tineture of nux vomicafl.dr.	101
Iron pyrophosphate, solublegr.	128
Elixir of phosphorusfl.oz.	4
Alcoholfl.oz.	2
Distilled water, hotfl.dr.	4
Simple elixir, enough to make.fl.oz.	16
201 11 0 11 1 11 11 11	

Mix the fluid extract, tincture, elixir of phosphorus, alcohol, and 6 fluidounces of simple elixir, also dissolve the iron salt in the water, mix the two liquids, add the remainder of the simple elixir, and filter, if necessary, in a well-covered funnel.

Each fluidram represents 7½ gr. of damiana and about 1 gr. of nux vomica, and contains 200 gr. of phosphorus and 1 gr. of iron pyrophosphate.

#### Elixir of Damiana, Iron and Phosphorus.

P	
Fluid extract of damianafl.oz.	2
Elixir of phosphorusfl.oz.	4
Iron pyrophosphate, solublegr.	128
Alcoholfl.oz.	1
Distilled water, hotfl.dr.	4
Simple elixir, enough to make.fl.oz.	16

Mix the fluid extract, elixir of phosphorus, alcohol, and 8 fluidounces of simple elixir, dissolve the iron pyrophosphate in the water, mix the two liquids, add the remainder of the elixir, and filter, if necessary, in a well-covered funnel.

phorus and 1 gr. of iron pyrophosphate and represents 7½ gr. of damiana.

#### Elixir of Damiana, Nux Vomica and Phosphorus.

Fluid extract of damianafl.oz.	2
Tincture of nux vomicafl.dr.	104
Elixir of phosphorusfl.oz.	2
Alcoholfl.oz.	2
Simple elixir, enough to makefl.oz.	16

Mix the above ingredients in the order given and filter, if necessary, in a well-covered funnel.

Each fluidram represents 200 gr. of phosphorus, about 1 gr. of nux vomica, and 71 gr. of damiana.

#### Elixir of Damiana and Phosphorus.

Elixir of phosphorus	fl.oz.	8
Fluid extract of damiana	fl.oz.	2
Alcohol	fl.oz.	2
Simple elixir	fl.oz.	4

Mix the elixir of phosphorus, alcohol, and fluid extract and add the simple elixir.

Each fluidram represents 100 gr. of phosphorus and 7½ gr. of damiana.

#### Elixir of Damiana, Phosphorus and Strychnine.

Elixir of phosphorus	,fl.oz. 8
Fluid extract of damiana	
Alcohol	
Strychnine sulphate	gr. 11
Simple elixir	.fl.oz. 4

Mix the elixir of phosphorus, alcohol, and fluid extract and add the simple elixir, having first dissolved the alkaloidal salt in the latter

Each fluidram represents 71 gr. of damiana and contains 180 gr. each of phosphorus and strychnine sulphate.

#### Elixir of Dewberry Root, Compound.

Dewberry root, in coarse pow-	
der av.oz.	21
Galls, powderedgr.	120
Kino, powderedgr.	120
Cinnamon, powderedgr.	60
Cloves, powderedgr.	30
Capsicum, powderedgr.	5
Tincture of opiumfl.dr.	4
Spirit of peppermintm.	45
Brandyfl.oz.	16
Sugarav.oz.	$7\frac{1}{2}$

Macerate all of the above, sugar excepted, for 14 days, shaking occasionally; express, Each fluidram contains 200 gr. of phos- filter, and in the filtrate dissolve the sugar.

#### Elixir of Dandelion.

Fluid extract of dandelion....fl.oz. 6 Simple elixir, enough to make..fl.oz. 16

Each luidram represents 22½ gr. of dandelion.

## Elixir of Dandelion, Compound. (Compound Elixir of Taraxacum.)

Fluid extract of dandelion....fl.dr. 4
Fluid extract of sweet orange peel fl.dr. 2
Fluid extract of wild cherry...fl.dr. 2
Fluid extract of licorice root...fl.oz. 1
Tincture of cinnamon......fl.dr. 4
Compound tincture of cardamom fl.dr. 4
Aromatic elixir, enough to make.fl.oz. 16

Mix, let stand a few days, and filter.—N. F. (last edition).

# | II. | Dandelion | gr. 320 | Wild cherry | gr. 320 | Sweet orange peel, recently dried.gr. 320 | Licorice, Russian, peeled | av. oz. 21 | Cinnamon, Saigon | gr. 80 | Cardamom | gr. 80 | Cardamom | gr. 80 | Caraway | gr. 80 | Cloves | gr. 80 | Cloves | gr. 27

Mix the drugs, reduce them to a moderately coarse powder, and extract by percolation with a mixture of 1 volume of alcohol and 2 of water, so as to obtain 11 fluidounces of percolate; to this add the syrup; let stand a few days, if possible, and filter.—N. F. (1st edition).

III. A formula for a preparation of the same name, which is also much in use and which is much different in some respects from either of the preceding, 's the following.

Fluid extract of dandelion....fl.dr. 5
Fluid extract of wild cherry...fl.dr. 3
Fluid extract of gentian.....fl.dr. 1
Fluid extract of licorice root...fl.dr. 1
Simple elixir, enough to make..fl.oz. 16
Mix and filter.

#### Elixir, Digestive, Compound.

See Compound Elixir of Pepsin.

#### Elixir, Emmenagogue.

, 00	
Ruegr. 96	5
Spanish saffrongr. 96	
Savin	
Socotrine aloes gr. 192	
Adjuvant elixir, enough to make fl. oz. 16	
Reduce the drugs to moderately fine po	w

der, mix with 12 fluidounces of adjuvant elixir, macerate for 7 days, agitating occasionally, filter, and through the filter add the remainder of the elixir.

Each fluidram represents  $\frac{3}{4}$  gr. each of rue, saffron and savin, and  $1\frac{1}{2}$  gr. of aloes.

#### Elixir of Eucalyptus.

Fluid extract of eucalyptusfl.oz. Alcoholfl.oz.	
Magnesium carbonategr.	120
Syrup of coffee fl.oz.	
Compound elixir of taraxacum fl.oz.	6

Mix the fluid extract with the alcohol, then add the other ingredients, shake the mixture occasionally during 48 hours, and filter.

Each fluidram represents 7½ gr. of eucalyptus.—N. F.

This preparation is also the same as what is generally sold or dispensed as aromatic or compound elixir of eucalyptus.

#### Elixir Flavoring No. 29.

Oil of sweet orange peelfl.oz.	31
Oil of caraway seedm.	100
Oil of coriander seedm.	100
Oil of cassia m.	100
Oil of anise (or oil of nutmeg)m.	50
Alcoholfl.oz.	

The oils used must be perfectly fresh. One fluidounce of this flavor is used for 1 gallon of elixir.

#### Elixir of Galls, Aromatic.

Gallsav.oz. 1
Nutmegsav. oz. ½
Cinnamonav.oz.
Brandysufficient.
Elixir of orangefl.oz. 10

Reduce the drugs to moderately coarse powder, moisten with brandy, pack in a percolate and percolate until 6 fluidounces of liquid are obtained, to which add the elixir.

#### Elixir de Garus. (Elixir Gari.)

٠	Cinnamongr.	30	
	Canellagr.	30	
	Clovesgr.	30	
	Nutmeggr.	30	
	Myrrhgr. 1	10	
	Aloesgr. 2		
	Spanish saffrongr.	8	
	Orange flower waterfl.oz.	1	
	Waterfl.oz.	8	
	thingle sprap	16	
	Alcohol fl.oz.	16	
	Reduce the drugs except the saffron	to	2

moderately coarse powder, macerate for 24 hours in a small still with 8 fluidounces of alcohol and the water, then distil off 8 fluidounces; to this distillate add the saffron, the remainder of the alcohol and the orange flower water, macerate for 2 days, agitating occasionally; add the syrup, and filter. II.

Oil of cassiadrops	8
Oil of clovesdrops	
Oil of macedrops	8
Saffrongr.	20
Tincture of vanillafl.dr	$\frac{1}{2}$
Alcohol	5
Orange flower waterfl.oz.	$6\frac{1}{2}$
Sugarav.oz.	7

Mix the oil, saffron, tincture and alcohol, macerate for 2 days, agitating occasionally; strain to remove the saffron, add the orange flower water and sugar, agitate until the latter is dissolved, and filter.—H. modified.

#### Elixir of Gentian.

Distilled water,

Aromatic elixir, of each.....sufficient. Dilute the solution of tersulphate of iron with 4 fluidounces of cold water, and add it, constantly stirring, to the water of ammonia, previously diluted with an equal volume of cold water. Collect the precipitate on a wellwetted muslin-strainer, allow it to drain completely, return it to the vessel, mix it intimately with 4 fluidounces of water, and again drain. Repeat this operation once more with the same quantity of water. When the precipitate has been completely drained for the third time, fold the strainer, and press it gently so as to remove the water as completely as possible without loss of magma; then remove the magma into a tared bottle, and ascertain its weight. Now add to the magma one-fifth of its weight of alcohol, the fluid extract, compound tincture and 12 fluidounces of aromatic elixir, and shake the mixture occasionally during 24 hours. Filter through paper, and pass enough aromatic elixir through the filter to make the product measure 16 fluidounces.

Each fluidram represents about 2 gr. of gentian.—N. F. (last edition).

II.

Extract of gentiangr.	70
Aromatic spiritfl.dr.	3
Tincture of vanillafl.dr.	
Simple syrupfl.oz.	1
Aromatic elixir, enough to make fl.oz.	16

Dissolve the extract in about 2 fluidounces of aromatic elixir, add the syrup, spirit, tine-ture and remainder of the elixir. Filter, if necessary. This is of about the same strength as the preceding.—N. F. (1st edition).

III. This preparation is also made according to one of the formulas for compound elixir of gentian which follow.

#### Elixir of Gentian, Compound.

I.

Stronger compound infusion of	
gentian, N. Ffl.oz.	41
Aromatic elixirfl.oz.	113
II.	
Gentiangr.	256
Coriander	64
Bitter orange peelgr.	64
Alcohol,	
Water, of eachsuffic	eient
Sugar av.oz.	5
Aromatic spiritfl.oz.	1
Egg albumengr.	120
Citric acidgr.	

Mix alcohol and water in the proportion of 1 of the former to 2 of the latter by measure, and with this mixture percolate the drugs, previously ground to moderately fine powder, until 12 fluidounces of percolate are obtained. To this percolate add the albumen and citric acid, agitate until the latter is dissolved, add the aromatic spirit and filter. In the absence of dried egg albumen, the white of 1 egg may be employed.

III.

Compound tincture of gentian. fl.oz. 5¼ Simple elixir.....fl.oz. 10¾

Each fluidram represents 2 gr. of gentian.

Elixir of Gentian and Bismuth.

Elixir of Gentian, Bismuth and Iron. Elixir of Gentian, Bismuth, Iron and Strychnine.

Elixir of Gentian, Bismuth and Strych-

Refer for above under Elixir of Bismuth and its combinations.

#### Elixir of Gentian and Cinchona. Elixir of Gentian, Cinchona and Iron

#### Malate.

Refer for above under Elixir of Cinchona and its combinations.

#### Elixir of Gentian with Iron Chloride Tincture.

Tincture of citro-chloride of iron fl.dr. 12¾
Elixir of gentian, enough to
make.....fl.oz. 16
Mix and filter, if necessary.

Each fluidram represents about ¾ gr. of ferric chloride, and nearly 2 gr. of gentian.—N. F.

#### Elixir of Gentian and Iron Phosphate.

(Ferrated Elixir of Gentian.)	
Iron phosphate, solubleg	r. 128
Distilled water, hotfl.d	r. 4
Elixir of gentian, enough to	
makefl.o	z. 16
Dissolve the iron phosphate in th	e wate

Dissolve the iron phosphate in the water, add the elixir of gentian, and filter, if necessary.

Each fluidram represents 1 gr. of iron phosphate and nearly 2 gr. of gentian.—N. F.

## Elixir of Gentian and Iron Pyrophosphate.

Iron pyrophosphate, solublegr. 128 Distilled water, hot fl.dr. 4
Elixir of gentian, enough to makefl.oz. 16
Dissolve the iron salt in the water, add the

elixir, and filter, if necessary.

Each fluidram contains 1 gr. of iron salt

and represents nearly 2 gr. of gentian.

#### Elixir of Gentian, Iron Phosphate, Nux Vomica and Quassia.

Trun Common with Quantities	
Tincture of nux vomicam.	256
Iron phosphategr.	128
Distilled water, hotfl.dr.	4
Fluid extract of quassiafl.dr.	4
Compound fluid extract of gen-	
tianfl.dr.	4
Simple elixirenough to make fl.oz.	16
Dissolve the iron phosphate in the	water
add the other ingredients and filter.	

Each fluidram contains 1 gr. of iron phosphate, and represents  $\frac{2}{5}$  gr. of nux vomica, nearly 2 gr. of quassia, and 1 $\frac{1}{4}$  gr. of gentian.

#### Elixir of Golden Seal and Bismuth. Elixir of Golden Seal, Bismuth and Iron.

Refer for above to Elixir of Bismuth and its combinations.

#### Elixir of Gentian and Phosphorus.

Fluid extract of gentianfl.dr.	
Elixir of phosphorusfl.oz.	8
Compound elixir of taraxacumfl.oz.	5
Aromatic clixir, enough to make fl.oz.	16

Each fluidram represents 100 gr. of phosphorus and 5 gr. of gentian.

## Elixir of Golden Seal. (Elixir of Hydrastis.)

Glycerite of hydrastis......fl.dr. 10½ Simple elixir, enough to make..fl.oz. 16

Each fluidram represents 5 gr. of golden seal.

#### Elixir of Grindelia.

Fluid extract of grindeliafl.oz.	1
Compound spirit of orangefl.dr.	11
Alcohol	
Compound elixir of taraxacumfl.oz.	13

Mix them, allow the mixture to stand a few days, if convenient, then filter.

Each fluidounce represents 30 gr. of grindelia.—N. F.

#### Elixir of Guaiac.

Tincture of guaiacfl.oz.	4
Potassium carbonategr.	20
Waterfl.dr.	
Glycerinfl.oz.	4
Compound elixir of taraxacumfl.oz.	4
Simple syrupfl.oz.	4

Dissolve the potassium carbonate in the water, add to the tincture of guaiac and to this mixture add the remaining ingredients in the order given above.

#### Elixir of Guarana.

	I.	
ı	Fluid extract of guaranafl.oz.	31
	Aromatic elixirfl.oz.	31
	Compound elixir of taraxacumfl.oz.	$9\frac{1}{2}$

Mix them; allow the mixture to stand during 48 hours, if convenient, and filter.

Each fluidram represents about 12 gr. of guarana.—N. F.

Mix the guarana and magnesia, moisten with  $2\frac{1}{2}$  fluidounces of diluted alcohol, set aside for 24 hours, then mix with the sand,

pack in a percolator, percolate until 13 fluidounces of liquid are obtained, then remove the mass from the percolator, inclose it in a cloth and express in a tincture press; to the percolate add the oil and syrup, and make up to 16 fluidounces by addition of the expressed liquid, previously concentrating the latter, if necessary, by evaporation.—H. modified and Brit. Form.

Each fluidram represents about 11 gr. of guarana.

## Elixir of Guarana, Compound. (Elixir of Guarana and Celery.)

Refer to Elixir of Celery and Guarana.

#### Elixir of Guarana and Coca.

Refer to Elixir of Coca and Guarana.

#### Elixir of Helonias.

Fluid extract of heloniasfl.oz.	4
Simple elixirfl.oz.	

Mix, allow to stand for 24 hours and filter. Each fluidram represents 15 gr. of helonias.

# Elixir of Helonias, Compound. (Compound Elixir of Squaw-vine.--Compound Elixir of Mitchella.)

Fluid extract of false unicorn	
(helonias dioica)fl.oz.	2
Fluid extract of mitchellafl.oz.	4
Fluid extract of blue cohoshfl.oz.	2
Fluid extract of crampbarkfl.oz.	2
Purified talcumav.oz.	ļ
Aromatic elixir, enough to make fl.oz. 1	6
Mix and filter.	

Each fluidram represents nearly 14 gr. of mitchella, and 7 gr. each of helonias, blue cohosh and crampbark.

#### Elixir of Hops.

Fluid extract of hopsfl.oz.	2
Magnesium carbonategr.	120
Tincture of vanillafl.dr.	4
Compound elixir of taraxacum.fl.oz.	2
Aromatic elixir, enough to	
makefl.oz.	16

Triturate the fluid extract with the magnesium carbonate, then gradually add the compound elixir of taraxacum, tincture of vanilla, and the aromatic elixir. Allow the mixture to stand several days, if convenient, occasionally agitating and then filter.

Each fluidram represents  $7\frac{1}{2}$  gr. of hops.—N. F.

#### Elixir of Hypophosphites.

Calcium hypophosphitegr.	
Sodium hypophosphitegr.	128
Potassium hypophosphitegr.	128
Citric acidgr.	30
Waterfl.oz.	4
Glycerinfl.dr.	4
Compound spirit of cardamom, fl.dr.	4
Aromatic elixir, enough to	
makefl.oz.	16

Dissolve the hypophosphites and the citric acid in the water; then add the glycerin, compound spirit and the aromatic elixir. Filter, if necessary.

Each fluidram contains 3 gr. of calcium hypophosphite and 1 gr. each of sodium and potassium hypophosphite.—N. F.

#### Elixir of Hypophosphites, Compound.

The elixir of hypophosphites with iron is sold and dispensed under this name.

#### Elixir of Hypophosphite of Calcium.

Refer to Elixir of Calcium Hypophosphite.

#### Elixir of Hypophosphites and Cherries.

Refer for this to Elixir of Cherries and its combinations.

#### Elixir of Hypophosphites and Cinchona.

Refer to Elixir of Cinchona and Hypophosphites.

#### Elixir of Hypophosphites with Iron.

Calcium hypophosphitegr.	188
Sodium hypophosphitegr.	128
Potassium hypophosphite gr.	64
Sulphate of iron, clear crystalsgr.	96
Citric acidgr.	30
Distilled waterfl.oz.	4
Simple syrup fl.oz.	4
Aromatic elixir, enough to	
make fl.oz.	16

Dissolve the hypophosphites in 3 fluidounces of water, and add the syrup. Dissolve the sulphate of iron in the remainder of the water, and mix this with the other solution. Then add 6 fluidounces of aromatic elixir, set the mixture aside, in a cold place, for 12 hours, and filter from the deposited calcium sulphate. Finally, dissolve the citric acid in the filtrate, and pass enough aromatic elixir through the filter to make 16 fluidounces.

Each fluidram contains about ½ gr. of hypophosphite of iron (ferrous), about 1 gr.

each of the hypophosphites of calcium and sodium, and ½ gr. of potassium hypophosphite.

—N. F.

## Elixir of Hypophosphites of Iron and Quinine.

Iron hypophosphitegr. 128
Potassium citrate gr. 128
Quinine sulphategr. 128
Calcium hypophosphitegr. 30
Spirit of orangefl.dr. 2
Orange flower waterfl.oz. 1
Sugarav.oz. 5
Alcohol,
Distilled water of each, sufficient.

Dissolve the iron hypophosphite with the aid of the potassium citrate in the orange flower water, and enough water to make the solution measure 6½ fluidounces, and in this dissolve the sugar. Triturate the quinine sulphate with 5 fluidounces ot alcohol, add a solution of the calcium hypophosphite in 4 fluidrams of water, and shake the mixture occasionally during 1 hour; filter, and wash the filter with enough alcohol to make 6½ fluidounces. Add this solution to the spirit of orange, mix this with the iron solution and sugar solution previously prepared, and filter the whole.

Each fluidram contains 1 gr. each of the hypophosphites of iron and quinine.

11. Solution of iron hypophosphite,

Mix the quinine hypophosphite with 8 fluidounces of elixir, add enough of the acid to dissolve the quinine, add the solution of iron hypophosphite, and then enough elixir to make 16 fluidounces, and filter.

This is of the same strength as the preceding.

## Elixir of Hypophosphites of Iron, Quinine and Strychnine.

This may be prepared by dissolving 1½ gr. of strychnine sulphate in 4 fluidrams of distilled water, and adding enough of the preceding elixir to make 16 fluidounces.

#### Elixir of Hypophosphites with Malt.

Refer for above to Elixir of Calcium Hypophosphite and its combinations.

#### Elixir of Hypophosphite of Iron.

Refer to Elixir of Iron Hypophosphite.

#### Elixir of Hypophosphite of Sodium.

Sodium hypophosphitegr.	256
Citric acidgr.	30
Aromatic elixir, enough to	
make fl.oz.	16

Dissolve the sodium hypophosphite and the citric acid in the elixir by agitation, and filter, if necessary.

Each fluidram contains 2 gr. of sodium hypophosphite.—N. F.

#### Elixir of Hypophosphites with Tar.

Refer to Elixir of Calcium Hypophosphite and its combinations.

#### Elixir of Six Iodides.

Arsenic iodidegr.	1
Mercuric iodidegr.	1
Manganese iodidegr.	13
Sodium iodidegr.	128
Potassium iodidegr.	128
Solution of iron iodide, N. Fm.	15
Sodium hypophosphite suffici	ent
Simple elixir, enough to make fl.oz.	16

Add the six iodides to the elixir, dissolve by agitation, add a few grains of sodium hypophosphite, or sufficient to decolorize the liquid, and filter.

Each fluidram contains 1-128 gr. each of arsenic and mercury iodides, 1-12 gr. of ferrous iodide, 1-10 gr. of manganese iodide, and 1 gr. each of sodium and potassium iodides.

## Elixir of Iodides of Arsenic and Mercury.

Refer for this to Elixir of Arsenic and its combinations.

#### Elixir of Iodide of Calcium.

Refer for the above to Elixir of Calcium Iodide.

## Elixir of Iodide of Potassium Compound. (Alterative Elixir.)

Potassium iodidegr.	640
Tincture of citrochloride of	
iron fl.dr.	101
Spirit of orangefl.dr.	4
Fluid etrxact of saxifragefl.dr.	12
Fluid extract of stillingiafl.dr.	12
Fluid extract of menispermum fl.dr.	12
Fluid extract of helonias fl.dr.	12
Sugarav.oz.	41
Water, enough to makefl.oz.	16

Dissolve the potassium iodide in the water,

add the tincture of iron, and in this mixture dissolve the sugar by agitation. Mix the fluid extracts, add the spirit, then the syrup, allow the whole to stand for two days, and filter.

Each fluidram contains 5 gr. of potassium iodide, and represents about  $\frac{1}{2}$  gr. of ferric chloride, and about  $5\frac{1}{2}$  gr. each of saxifraga, stillingia, menispermum and helonias.

#### Elixir of Iodide of Potassium.

Potassium iodide .......gr. 640
Aromatic elixir of licorice,
enough to make ......fl.oz. 16
Dissolve by agitation.

Each fluidram contains 5 gr. of potassium iodide

## Elixir of Iron and Ammonium Valerianate.

Refer for this to Elixir of Ammonium Valerianate and its combinations.

Elixir of Iron, Beef and Cinchona.

Elixir of Iron, Beef, Cinchona and Strychnine.

Elixir of Iron, Beef and Coca. Elixir of Iron, Beef and Malt.

Refer for above under Elixir of Beef and its combinations.

# Elixir of Iron and Arsenic Chlorides. Elixir of Iron, Arsenic and Mercury Chlorides.

Refer for above to Elixir of Arsenic and its combinations.

#### Elixir of Iron and Berberine.

Refer for above to Elixir of Berberine and its combinations.

Elixir of Iron and Bismuth.

Elixir of Iron, Bismuth and Cinchona. Elixir of Iron, Bismuth, Cinchona and Pepsin.

Elixir of Iron, Bismuth, Cinchona, Pepsin and Strychnine.

Elixir of Iron, Bismuth, Cinchona and Strychnine.

Elixir of Iron, Bismuth and Gentian. Elixir of Iron, Bismuth, Gentian and Strychnine.

Elixir of Iron, Bismuth and Golden Seal. Elixir of Iron, Bismuth and Pepsin.

Elixir of Iron, Bismuth, Pepsin and Quinine.

Elixir of Iron, Bismuth and Strychnine.

Refer for the above to Elixir of Bismuth and its combinations.

Elixir of Iron, Damiana, Nux Vomica and Phosphorus.

Elixir of Iron, Damiana and Phosphorus.

Refer for the above to Elixir of Damiana and its combinations.

Elixir of Iron, Calcium Lactophosphate and Cinchona.

Elixir of Iron and Cinchona.

Elixir of Iron, Cinchona and Pepsin.

Elixir of Iron, Cinchona and Strychnine.

Refer for the above to the National Formulary.

## Elixir of Iron, Cinchona and Phosphorus.

Refer for above to Elixir of Cinchona and its combinations.

Elixir of Iron and Cinchonidine.

## Elixir of Iron, Cinchonidine and Stry-chnine.

Refer for above to Elixir of Cinchonidine and its combinations.

#### Elixir of Iron and Malt.

Refer to the National Formulary.

#### Elixir of Iron, Malt and Beef.

Refer for this to Elixir of Beef and its combinations.

Elixir of Iron and Pepsin. (Ferrated Elixir of Pepsin.)

Refer to Elixir of Pepsin and Iron.

#### Elixir of Iron, Pepsin and Quinine.

Iron pyrophosphate, soluble...gr. 256
Quinine hydrochlorate....gr. 32
Distilled water, hot....fl.oz. 1
Elixir of pepsin, N. F., enough
to make....fl.oz. 16

Dissolve the iron salt in the water, add the elixir and the quinine salt, agitate occasionally until dissolved, and filter. Each fluidram contains 2 gr. of iron pyrophosphate,  $\frac{1}{2}$  gr. of quinine hydrochlorate, and nearly 1 gr. of pepsin.

#### Elixir of Iron, Quinine and Arsenic.

Iron pyrophosphategr.	128
Quinine hydrochlorategr.	
Solution of arsenious acid m.	
Distilled water, hot fl.dr.	4
Simple elixir, enough to make fl.oz.	

Dissolve the iron pyrophosphate in the water, dissolve the quinine in about 12 fluid-ounces of clixir, by agitation, mix the solutions, add the acid solution and the remainder of the clixir, then neutralize exactly with ammonia water, carefully added, and filter.

Each fluidram contains 1 gr. of iron pyrophosphate, ½ gr. of quinine hydrochlorate, and 1-32 gr. of arsenious acid.

#### Elixir of Iron, Quinine and Strychnine

Tincture of citrochloride of iron fl.oz.	2
Quinine hydrochlorategr.	64
Strychnine sulphategr.	
Alcohol fl.dr.	
Aromatic elixir, enough to make fl.oz.	

Dissolve the alkaloidal salts in 12 fluidounces of elixir, then add the tincture and the alcohol, and finally, the remainder of the elixir; filter, if necessary.

Each fluidram represents about 1 gr. of ferric chloride, ½ gr. of quinine hydrochlorate, and 1-100 gr. of strychnine sulphate.— N. F.

## Elixir of Iron and Wild Cherry. (Ferrated Elixir of Wild Cherry.)

Iron pyrophosphate gr.	128
Distilled water, hotfl.dr.	4
Fluid extract of wild cherryfl.oz.	2
Alcoholfl.oz.	2
Simple elixir, enough to make fl.oz.	16

Mix the alcohol and fluid extract, add the elixir, and then iron salt previously dissolved in the water, and filter through purified talcum.

Each fluidram contains 1 gr. of iron pyrophosphate, and represents 7½ gr. of wild cherry.

#### Elixir of Iron Chloride Tincture and Gentian.

Refer to Elixir of Gentian and Iron Chloride Tincture.

#### Elixir of Iron and Quinine Citrate.

(Elixir of Iron and Quinine.)

Citrate of iron and quinine....gr. 256
Water ...........fl.oz. 1
Aromatic elixir, enough to make fl.oz 16

Dissolve the citrate in the water, add the elixir, and filter. Each fluidram contains 2 gr. of iron and quinine citrate.

#### Elixir of Iron and Hypophosphites.

Refer for this to the Elixirs of the Hypophosphites.

#### Elixir of Iron Hypophosphite.

Solution of iron hypophosphite fl.dr. 1234 Aromatic elixir, enough to make fl.oz. 16

Mix, allow the mixture to stand a few days in a cool place, and filter, if necessary.

Each fluidram contains 1 gr. of ferric hypophosphite.—N. F.

#### Elixir of Iron and Quinine Hypophosphites.

#### Elixir of Iron, Quinine and Strychnine Hypophosphites.

Refer for the above to the Elixirs of Hypophosphites.

#### Elixir of Iron Lactate.

Iron lactate, in crusts gr.	128
Potassium citrategr.	384
Aromatic elixir, enough to	
make fl.oz.	16

Triturate the iron lactate with the potassium citrate and about 4 fluidounces of aromatic elixir, gradually added, until solution has been effected; then add the remainder of the aromatic elixir, and filter.

Each fluidram contains 1 gr. of iron lactate.—N. F.

## Elixir of Iron Malate, Cinchona and Gentian.

Refer to Elixir of Cinchona and its combinations.

#### Elixir of Iron Peptonate.

٠.			
	Dried egg albumeng	r.	75
	(Or fresh egg albumen gr.		560)
	Distilled watersu	ffic	ient
	Hydrochloric acidfl.	dr.	2
	Pepsin, pureg		
	Solution of iron oxychloride fl. d	lr.	15
	Solution of sodasu		
	Brandyfl.d		
	,		

Dissolve the albumen in 16 fluidounces of distilled water, add the hydrochloric acid and

pepsin, digest the mixture at a temperature of 40 degrees C., until it produces only a faint turbidity with nitric acid; allow to cool, neutralize with solution of soda, strain, mix the colature with the solution of iron oxychloride to which has been added 16 fluidounces of distilled water. The mixture is again neutralized with solution of soda, the precipitate is washed by decantation with distilled water, until the washings are no longer affected by silver nitrate. The precipitate is now drained on a well-wetted muslin strainer, transferred to a porcelain capsule, 10 m. of hydrochloric acid are added, and the mixture heated on a water bath and stirred until solution occurs. To this solution is now added distilled water to make 141 fluidounces, and lastly, the brandy is added.—D.

Iron peptonate may be obtained by spreading the solution in the porcelain capsule upon glass plates and allowing to dry.

TT.

Pepsin, puregr.	4
Dried egg albumengr.	30
Simple syrupfl.dr.	4
Solution of dialized iron or iron-	
oxychloride fl.dr.	$12\frac{1}{2}$
Aromatic elixirfl. dr.	121/2
Distilled water, enough to make fl.oz.	16

Dissolve the albumen in 3½ fluidounces of water, add the pepsin and digest for four hours at 50 degrees C. Mix the syrup and solution of iron with 9 fluidounces of the water, then add to the pepsin solution and heat to 90 degrees C. Cool, add the elixir and the remainder of the water. Set aside for 8 days and then decant the clear solution.

If to the above be added 32 grains of crystmanganese chloride, previously dissolved in 1 fluidram of water, it will constitute the "Solution of Iron and Manganese Peptonate."

#### Elixir of Iron Phosphate.

Iron phosphate, solublegr.	256
Water, hot	1
Aromatic elixir, enough to make fl. oz.	16

Dissolve the iron phosphate in the water, mix this solution with the aromatic elixir, and filter, if necessary.

Each fluidram contains 2 gr. of iron phosphate.—N. F.

## Elixir of Iron Phosphate, Cinchonidine and Strychnine.

Iron phosphate, solublegr.	256
Potassium citrategr.	32
Cinchonidine sulphategr.	64
Strychnine sulphategr.	11
Alcoholfl.oz.	1
Water, hotfl.dr.	6
Aromatic elixir, enough to make fl. oz.	16

Dissolve the iron phosphate and potassium citrate in the water. To 12 fluidounces of aromatic elixir, contained in a bottle, add the alcohol, and afterwards the alkaloidal salts, and agitate until the latter are dissolved, or nearly so. Then mix the two solutions, and, having shaken the mixture, add the remainder of the aromatic elixir. Finally, filter.

Each fluidram contains 2 gr. of iron phosphate, ½ gr. of cinchonidine sulphate, and rbo gr. of strychnine sulphate.—N. F.

#### Elixir of Iron Phosphate, Gentian, Nux Vomica and Quassia.

Refer to Elixir of Gentian and its combinations.

## Elixir of Iron, Phosphate, Quinine and Strychnine.

1.	<b>\</b>	
	Iron phosphate, solublegr.	128
	Quinine (alkaloid)gr.	64
	Strychnine (alkaloid)gr.	2
	Alcoholfl.oz.	2
	Distilled water, hotfl.dr.	6
	Aromatic elixir, enough to make fl.oz.	16

Dissolve the alkaloids in the alcohol and add 12 fluidounces of aromatic elixir, then dissolve the iron phosphate in the water, and add to the previous mixture. Finally, add the remainder of the aromatic elixir.

Each fluidram contains 1 gr. of iron phosphate, ½ gr. of quinine, and ¼ gr. of strychnine.—N. F.

II.	
	1 2
Quinine sulphategr. 64	1
Citric acidgr.	ŏ
Iron phosphate, solublegr. 256	3
Alcoholfl.oz.	
Simple syrupfl.oz.	ì
Distilled water, hotfl.oz.	ĺ
Orange flower waterfl.oz.	3
Sodium bicarbonatesufficient	t

Triturate the strychnine and quinine sulphate with the acid until well mixed, and rub this mixture with the alcohol gradually added. Heat the syrup to about 65 degrees C., add to it the alcoholic liquid, and stir until clear. Dissolve the iron salt in the water, add the orange flower water, mix this with the preceding liquid, and allow to cool. Then add sodium bicarbonate in very small amounts, stirring thoroughly after each addition, until the elixir remains but slightly acid. Allow to stand for a few hours, then filter through white filter paper. Any excess of soda must be avoided.

#### III.

Strychnine sulphategr.	1‡
Quinine hydrochlorategr.	128
Iron phosphate, solublegr.	256
Potassium citrategr.	32
Alcoholfl.oz.	11
Distilled water, hotfl.oz.	1
Glycerinfl.dr.	18
Aromatic elixir, enough to make fl.oz.	16
The state of the s	

Dissolve the quinine salt in 10 fluidounces of elixir, mixed with the alcohol, by agitation, and mix this solution with the strychnine sulphate previously dissolved in 2 fluidrams of the water.

Dissolve the iron phosphate in 6 fluidrams of the water, add 2 fluidounces of glycerin and mix this solution with the preceding liquid. Now to this mixture add the potassium citrate dissolved in 1½ fluidounces of aromatic elixir mixed with 2 fluidrams of glycerin. Allow the whole to stand for several hours, then filter.

#### IV.

Quinine sulphategr.	
Iron phosphate, solublegr.	256
Strychnine sulphategr.	11
Alcoholfl.oz.	2
Glycerinfl.oz.	2
Simple syrupfi.oz.	2
Distilled water, hotfl.oz.	1
Aromatic elixir, enough to make fl.oz.	16

Dissolve the strychnine salt in the alcohol, and add the quinine; mix the glycerin and syrup, and heat, and when warm add to the alkaloidal solution; continue heating carefully, until quinine is dissolved, and add enough elixir to make 15 fluidounces. Dissolve the iron salt in the water, add this to previous liquid, let stand three or four hours, and filter.

V.

Iron phosphate, solublegr.	256
Quinine sulphategr.	128
Strychnine sulphategr.	1 1
Alcoholfl.oz.	1
Simple syrupfl.oz.	8
Aromatic elixir, enough to make fl.oz.	16

Dissolve the iron phosphate in the syrup by the aid of heat, and raise the temperature to near the boiling point. Dissolve the alkaloidal salts in 6 fluidounces of aromatic elixir, contained in a flask, by the aid of heat, and while still hot add this solution all at once to the iron solution, shaking immediately. Allow to stand 24 hours, then filter.

# Elixir of Iron, Quinine and Strychnine Phosphates. (Elixir of Three Phosphates.)

Nearly all of the preparations dispensed under this name contain the iron as phosphate or pyrophosphate, and the quinine and strychnine in some other form than as phosphate. If it be desired to dispense such a preparation as "elixir of three phosphates," then any of the preparations made according to formulas given in this formulary under elixir of iron phosphate, or pyrophosphate, quinine and strychnine may be dispensed.

The following formula does actually contain the three bases in the form of phosphates, which are maintained in solution by the excess of hydrochloric acid:

Solution of iron chloride, U.S. P.fl.dr. Quinine (alkaloid)gr.	$\frac{7\frac{1}{2}}{110}$
Strychnine (alkaloid)gr.	1
Phosphoric acid, U. S. Pfl.dr.	
Distilled waterfl.dr.	2
Alcoholfl.oz.	1
Simple elixirfl.oz.	10
Simple syrup, enough to make fl.oz.	16

Mix the iron solution, phosphoric acid and water, and in this mixture dissolve the alkaloids; to this solution add the syrup, and then elixir and alcohol previously mixed.

However, any elixir containing iron in the form of phosphate or pyrophosphate will inevitably darken upon exposure to light, and therefore some manufacturers place upon the market a so-called "permanent elixir of three phosphates," which contains the iron as citrochloride; a preparation of this character would be well represented by the elixir of

iron, quinine and strychnine of the National Formulary.

#### Elixir of Iron "Protoxide".

Solution of "protoxide" of iron fl.oz. 2 Simple elixir.....fl.oz: 14

## Elixir of Iron "Protoxide" and Cinchona.

Refer to Elixir of Cinchona and its combinations.

#### Elixir of Iron Pyrophosphate.

Iron pyrophosphate, soluble...gr. 256 Distilled water, hot.....fl.oz. 1 Aromatic elixir, enough to make fl.oz. 16

Dissolve the iron pyrophosphate in the water, add the elixir, and filter, if necessary. Each fluidram contains 2 gr. of iron pyrophosphate.—N. F.

## Elixir of Iron Pyrophosphate and Ammonium Valerianate.

Elixir of Iron Pyrophosphate, Ammonium Valerianate and Cinchonidine.

Elixir of Iron Pyrophosphate, Ammonium Valerianate, Cinchonidine and Quinine.

Elixir of Iron Pyrophosphate, Am monium Valerianate, Cinchonidine, Quinine and Strychnine.

Elixir of Iron Pyrophosphate, Ammonium Valerianate, Cinchonidine and Strychnine.

Elixir of Iron Pyrophosphate, Ammonium Valerianate and Quinine.

Elixir of Iron Pyrophosphate, Ammonium Valerianate, Quinine and Strychnine.

Refer for the above to Elixir of Ammonium Valerianate and its combinations.

## Elixir of Iron Pyrophosphate and Gentian.

Refer to Elixir of Gentian and its combinations.

## Elixir of Iron Pyrophosphate and Quinine.

This may be prepared like elixir of iron pyrophosphate, quinine and strychnine, the strychnine to be omitted, of course.

## Elixir of Iron Pyrophosphate, Quinine and Strychnine.

٠.			
	Iron pyrophosphate, soluble	.gr.	256
	Quinine sulphate	.gr.	64
	Strychnine	.gr.	11
	Citric acid	.gr.	5
	Alcohol		3
	Spirit of orange	fl dr.	11
	Distilled water		7
	Simple syrup	fl.oz.	6
	Ammonia water		ient

Triturate the quinine sulphate, strychnine and acid together, until minutely divided, and add the alcohol and spirit of orange; warm the syrup to about 65 degrees C., and add to the alcoholic mixture, stirring until clear. To this add the iron salt previously dissolved in the water; to the mixture add ammonia water, drop by drop, until the mixture is clear, and finally filter.

TT.

11.		
Strychnine (alkaloid)	gr.	11
Quinine (alkaloid)	gr.	64
Iron pyrophosphate		128
Alcohol	fl.oz.	2
Distilled water, hot	fl.oz.	3
Simple syrup	fl. oz.	3
Aromatic elixir		8

Dissolve the strychnine and quinine in the alcohol, also the iron salt in the water, mix the two solutions, add the syrup and then the elixir, and filter, if necessary.

H.

1.	
Strychnine (alkaloid)gr.	1
Quinine sulphategr.	64
Citric acidgr.	5
Alcoholfl.oz.	3
Simple syrupfl.oz.	6
Distilled water, hotfl.oz.	
Orange flower waterfl.oz.	3
Iron pyrophosphate, solublegr.	256
Sodium bicarbonatesuffic	

Triturate together the alkaloids and the acids until thoroughly mixed; rub this with the alcohol gradually added. Heat the syrup to about 65 degrees C., add it to the alcoholic mixture, and stir until clear. Dissolve the iron salt in the water, and add the orange flower water; mix the two solutions, and when cold, add carefully bicarbonate of sodium in small portions until the elixir remains but slightly acid. Allow to stand for a few hours, then filter through white filter paper. Excess of soda must be carefully avoided.

## Elixir of Iron Pyrophosphate and Strychnine.

Iron pyrophosphategr.	256
Strychnine sulphategr.	
Distilled water, hotfl.oz.	2
Simple elixir, enough to make.fl.oz.	16

Dissolve the iron salt and strychnine sulphate in the hot water, add the elixir, and filter.

Each fluidram contains 2 gr. of iron pyrophosphate and 180 gr. of strychnine sulphate,

#### Elixir of Iron Salicylate.

Iron salicylategr.	640
Distilled water, hotfl.oz.	
Glycerinfl.oz.	
Simple elixir, enough to make.fl.oz.	

Dissolve the iron salt in the hot water and glycerin, add the elixir, allow to stand for a few days and filter.

Each fluidram contains 5 gr. of iron salicylate.

#### Elixir of Iron Salicylate, Compound.

Iron salicylate	_	
Distilled water, hotfl	oz.	$2\frac{1}{4}$
Glycerinfl	.oz.	21
Fluid extract of colchicum root fl	.dr.	9
Deodorized tincture of opiumfl	.dr.	$4\frac{1}{2}$
Simple elixir, enough to make.fl	.oz.	16

Dissolve the iron salt in the hot water and glycerin, add the other ingredients, allow to stand a few days, and filter.

Each fluidram contains 5 gr. of iron salicylate and represents about 4½ gr. of colchicum root and 2 m. of deodorized tincture of opium.

#### Elixir of Iron Valerianate.

Iron valerianategr.	128
Alcoholfl.oz.	1
Simple elixirfl.oz.	15

Dissolve the iron salt in the alcohol, add the elixir, and filter.

Each fluidram contains 1 gr. of iron valerianate.

## Elixir of Jaborandi. (Elixir of Pilocar-

Pusi		
Fluid extract of jaborandifl.	oz.	1
Syrup of coffeefl.	oz.	-3
Tincture of vanillafl.	dr.	4
Compound elixir of taraxacum,		
enough to makefl.	oz.	16

Mix, allow the mixture to stand during 4 days, if convenient, and filter.

Each fluidram represents 3\% gr. of jaborandi.—N. F.

## and Elixir of Iron, Quinine and Zinc Valerianates.

Refer for this to Elixirs of Valerianates of different bases.

#### Elixir of Kola.

Fluid extract of kolafl.oz. 2
Ammoniated glycyrrhizingr. 60
Saccharingr. 60
Oil of orangedrops 5
Waterfl.oz. 7
Alcoholfl oz. 31
Simple syrupfl.oz. 3
Simple elixir, enough to makefl.oz. 16

Dissolve the ammoniated glycyrrhizin in the water and in this dissolve the saccharin; add the syrup and alcohol, followed by the fluid extract of kola, to which has been added the oil of orange; set aside for 5 or 6 hours agitating occasionally; filter, and add the simple elixir.

Each fluidram represents 7½ gr. of kola.

#### Elixir of Lactophosphate of Calcium. Elixir of Lactophosphate of Calcium and Cinchona.

#### Elixir of Lactophosphate of Calcium, Cinchona and Iron.

Refer for above to Elixir of Calcium Lactophosphate and its combinations.

#### Elixir of Licorice.

ı.		
	Fluid extract of licoricefl.oz.	2
	Aromatic elixirfl.foz.	. 14
	Mix and filter, if necessaryN. F.	

Purified extract of licorice

(U. S. P.)av.oz.	31
Fennel waterfl.oz.	$9\frac{1}{4}$
Anisated solution of ammoniafl.oz.	31

Dissolve the extract in the water and add the solution.—Germ. Pharm.

The mixture is turbid and must be shaken before use.

This second preparation is best known by the names Elixir e Succo Liquiritiæ,. Elixir Pectorale, Pectoral Elixir, Liquor Pectoralis, Brust Tropfen and Brust Elixir.

#### Elixir of Licorice, Aromatic.

[.	Fluid extract of licoricefl.oz.	3	
	Oil of clovesdrops	3	
	Oil of cinnamon (Ceylon)drops	3	
	Oil of nutmegdrops	2	
	Oil of fenneldrops	6	
	Magnesium carbonategr. 1	20	
	Aromatic elixir, enough to make fl.oz.		
	Triturate the oils with the magnesium		۳.

bonate, and gradually add 14 fluidounces of aromatic elixir. Shake occasionally during an hour, filter, and pass enough aromatic elixir through the filter to make 14 fluidounces of filtrate. Add the fluid extract to the filtrate, mix, and filter, if necessary.— N. F.

II.

Cardamom (seed without capsule	gr.	16
Cinnamon	.gr.	16
Staranise		16
Coriander		8
Caraway		8
Canella		4
Nutmeg		4
Cloves		4
Vanilla		24
Ammoniated glycyrrhizin		110
Diluted alcoholf	l.oz.	$6\frac{1}{2}$
Water, hotf		1
Simple syrup, enough to make f		16
. ,		

Reduce the drugs to moderately coarse powder, macerate for 7 days in the diluted alcohol, and filter, adding, if necessary, enough diluted alcohol through the filter to make the filtrate measure  $6\frac{1}{2}$  fluidounces. Dissolve the glycyrrhizin in the water, mix this solution with the filtrate, and add the syrup.

III.

Select licorice root, cut and	
slightly bruisedav.oz.	$2\frac{1}{4}$
Water of ammoniafl.dr.	
Glycerinfl.oz.	
Waterfl.oz.	

Macerate for 24 hours, strain, boil for 10 minutes, filter, and evaporate at gentle heat until reduced to 6 fluidounces.

Now add to this evaporated infusion:

Simple syrupfl.oz. 6
Alcoholfl.oz. 4
Spirit of orangefl.dr. 2
Oil of cinnamon (Cevlon)drops 2

This elixir is employed for disguising the taste of bitter medicines, particularly quinine. No acid should be used because it dissolves the quinine and makes its bitter taste more perceptible, and at the same time liberates the glycyrrhizin from its combination with ammonia and renders it insoluble, and therefore valueless for the purpose of disguising or modifying taste.

## Elixir of Licorice with Ammonium Chloride, Compound.

Refer for this to Elixir of Ammonium Chloride, etc

#### Elixir of Licorice Compound.

Pure extract of licorice, (U.	
S. P.)av.oz.	1/2
Wine of antimonyfl.oz.	1
Paregoric	2
Spirit of nitrous etherfl.dr.	
Elixir of cherries, enough to	
makefl.oz.	16

Dissolve the extract in a portion of the elixir and add the remaining ingredients.

The above replaces "brown mixture" in the form of an elixir.

#### Elixir of Lithium Bromide.

Lithium bromidegr. 6	40
Citric acidgr.	
Aromatic elixir, enough to make fl.oz.	16

Dissolve the solids in about 14 fluidounces of aromatic elixir, by agitation; add the remainder of the aromatic elixir and filter.

Each fluidram contains about 5 gr. of lithium bromide.—N. F.

#### Elixir of Lithium Citrate.

Lithium citrate......gr. 640 Aromatic elixir, enough to make fl.oz. 16 Dissolve by agitation, and filter.

Each fluidram contains 5 gr. of lithium citrate.—N. F.

#### Elixir of Lithium Salicylate.

Lithium salicylate.....gr. 640 Aromatic elixir, enough to make fl.oz. 16

Dissolve by agitation, and filter.

Each fluidram contains 5 gr. of lithium salicylate.—N. F.

Elixir of Long Life. ("Elixir ad Longam Vitam."—" Elixir of Life."—Compound Tincture of Aloes. (Germ. Pharm.) ("Swedish Bitters.")

Aloesgr.	900
Rhubarb gr.	35
	35
Zedoarygr.	35
Spanish saffrongr.	35
Waterfl.oz.	4
Alcoholfl.oz.	12

Mix the drugs in coarse powder with the two liquids, macerate for 3 days, agitating frequently; express and filter. Sometimes 35 gr. of agaric is added to the other drugs, and the menstruum generally employed is diluted alcohol.

The following is a simple formula which

may be used for the preparation of this ancient and complicated remedy:

Tincture of aloes and myrrhfl.oz. 8
Tincture of rhubarbfl.oz. 2
Compound tincture of gentianfl.oz. 1
Waterfl. oz. 1
Alcoholfl.oz. 4

#### Elixir of Lupulin.

Fluid extract of lupulin,fl. oz.	1
Magnesium carbonateav.oz.	1
Simple elixir, enough to makefl.oz.	16

Triturate the fluid extract with the talcum, add the elixir, transfer to a bottle, set aside for several hours, and filter.

The above is of the strength usually furnished by manufacturers; Diehl's formula, which is largely used, directs the use of 2 fluidounces of the fluid extract to the pint of finished elixir.

## Elixir of Lupulin and Sodium Bromide.

Fluid extract of lupulinfl.dr.	101
Purified talcumgr.	120
Sodium bromidegr.	640
Aromatic elixir of licorice,	
enough to makefl.oz.	16

Triturate the fluid extract with the talcum, add some of the elixir, transfer to a bottle, add the sodium salt and the remainder of the elixir, dissolve by agitation, and filter after several hours.

Each fluidram represents 5 gr. of lupulin and contains 5 gr. of sodium bromide.

#### Elixir of Malt.

Extract of maltfl.oz.	4
Simple elixirfl.oz.	12

#### Elixir of Malt, Beef and Iron.

Refer to Elixir of Beef and its combinations.

## Elixir of Malt, and Calcium and Sodium Hypophosphites.

Refer to Elixir of Calcium Hypophosphite and its combinations.

#### Elixir of Malt and Iron.

Extract of maltfl.oz.	4
Iron phosphate, solublegr.	
Water, hotfl.dr.	
Aromatic elixir, enough to make fl. oz.	

Dissolve the iron phosphate in the water by the aid of heat, mix the solution with the valerianate.

extract of malt, and add the elixir. Set the mixture aside for 24 hours, and filter.—N. F.

Each fluidram represents 1 gr. of iron phosphate and 15 m. of extract of malt.

Extract of malt, most suitable for this preparation, should have about the consistence of balsam of Peru, at a temperature of about 15 degrees C.

#### Elixir of Malt and Pepsin.

Elixir of 1	nalt		 .fl.oz. 8
Elixir of p	epsin, N.	F	 .fl.oz. 8

Mix and filter.

Each fluidram represents  $\frac{1}{2}$  gr. of pepsin and 15 m. of extract of malt.

#### Elixir of Manaca and Salicylates.

Fluid extract of manacafl.oz.	$2\frac{1}{2}$
Sodium salicylateav.oz.	13/4
Potassium salicylategr.	384
Lithium salicylategr.	96
Simple elixir, enough to make. fl. oz.	16

Dissolve the salicylates in some of the elixir, add the fluid extract and the remainder of the elixir, allow to stand for a few hours, and filter through talcum.

Each fluidram contains 6 gr. of sodium salicylate, 3 gr. of potassium salicylate, and 34 gr. of lithium salicylate and represents nearly 10 gr. of manaca.

#### Elixir of Matico, Compound.

Fluid extract of matico. fl.oz. 3 Fluid extract of buchu. fl.oz. 1 Fluid extract of cubeb fl.oz. 1 Alcohol. fl.oz. 2 Simple elixir fl.oz. 4 Compound elixir of taraxacum. fl.oz. 4 Mix set aside for 3 days, and filter through		
Fluid extract of cubeb       fl.oz. 1         Alcohol       fl.oz. 2         Simple elixir       fl.oz. 4         Compound elixir of taraxacum       fl.oz. 4	Fluid extract of maticofl.oz.	3
Alcohol	Fluid extract of buchufl.oz.	11
Simple elixir	Fluid extract of cubebfl.oz.	11
Compound elixir of taraxacumfl.oz. 4	Alcoholfl.oz.	2
•	Simple elixirfl.oz.	4
Mix. set aside for 3 days, and filter through	Compound elixir of taraxacumfl.oz.	4
	Mix, set aside for 3 days, and filter thro	ugl

talcum.

Each fluidram represents 11 gr. of matico

## and nearly 4 gr. each of buchu and cubeb. Elixir Mercury and Arsenic Iodides.

Refer to Elixir of Arsenic and its combinations.

#### Elixir of Mercury, Arsenic and Iron Chlorides.

Refer to Elixirs of Chlorides for above.

#### Elixir of Morphine Valerianate.

Morphine valerianategr.	16
Simple elixirfl.oz.	16

Dissolve by agitation, and filter.

Each fluidram contains ½ gr. of morphine valerianate.

#### Elixir of Nux Vomica, Bismuth and Pepsin.

Refer to Elixir of Bismuth and its combi-

#### Elixir of Nux Vomica, Damiana, Iron and Phosphorus.

See Elixir of Damiana and its combinations.

#### Elixir of Nux Vomica, Gentian, Iron and Quassia.

Refer to Elixir of Gentian and its combinations.

## Elixir of Nux Vomica and Phospho-

Tincture of nux vomica.....fl.dr. 41 Elixir of phosphorus, enough to

Mix them. This preparation should be freshly made, when wanted for use.

Each fluidram represents 2 m. of tincture of nux vomica, and nearly 1-50 gr. of phosphorus.-N. F.

#### Elixir of Orange.

Oil of orangefl.dr.	41
Alcoholfl.oz.	
Waterfl.oz.	
Simple syrupfl.oz.	28
Purified talcumav.oz.	

Mix the oil and alcohol, add the talcum, shake well, and then add the other ingredients in small portions at a time, agitating well after each addition.-U. S. P. 1880 modified.

The oil used should be a perfectly fresh sweet oil of orange peel.

#### Elixir of Orange, Compound. (Compound Wine of Orange.-Vinum Amarum, Bitter Wine .- Elixir Stomachicum, Stomachic Elixir.-Elixir Viscerale Hoffmanni.)

1410 2101111411111)	
Bitter orange peel, cutgr.	1600
Cinnamon, bruisedgr.	320
Potassium carbonategr.	80
Extract of gentiangr.	160
Extract of wormwoodgr.	160
Extract of buckbeangr.	160
Extract of cascarillagr.	160
Sherry wine, enough to make.fl.oz.	16

Macerate the orange peel, cinnamon, and potassium carbonate with 16 fluidounces of sherry wine for 8 days, agitating occasion-

dissolve the extracts, filter, and add enough sherry wine through the filter to make the filtrate measure 16 fluidounces. - Germ. Pharm.

The National Formulary also recognizes what is identically the same preparation under the title of "compound wine of orange"; in the latter no extracts are used, but the drugs themselves are mixed with the orange peel, cinnamon, and potassium carbonate. the whole being extracted by percolation.

#### Elixir of Pancreas.

Take 1 pig pancreas, chop into pieces, and macerate in a cool place for 3 days in a mixture of

Water	fl.oz.	32
Glycerin	fl.oz.	$6\frac{1}{2}$
Hydrochloric	acidfl.dr.	5

Strain, add & fluidram of oil of orange and enough glycerin to make 48 fluidounces, and filter.

#### Elixir of Pancreatin.

Pancreatin, puregr.	128
Sodium bicarbonategr.	16
Waterfl.oz.	2
Simple elixir, enough to make.fl.oz.	16

Macerate the pancreatin in the water for 24 hours, add the sodium bicarbonate, triturate until dissolved, gradually add the elixir and filter.

Each fluidram represents 1 gr. of pancreatin.

The elixir of pancreas may be substituted for the above, if deemed desirable.

#### Elixir of Pancreatin and Bismuth.

Refer to Elixir of Bismuth and its combinations.

#### Elixir of Pancreatin, Bismuth and Pepsin.

Citrate of bismuth and ammo-	
niumgr.	128
Pancreatin, puregr.	64
Pepsin, puregr.	64
Distilled water, hot fl.oz.	1
Water of ammoniasuffic	ient
Glycerinfl.oz.	2
Waterfl.oz.	2
Tincture of cudbearfl.dr.	2
Simple elixir, enough to make.fl.oz.	16

Triturate the bismuth salt with the water, allow the insoluble portion to subside, decant the clear portion, to the residue add ammonia ally; express the liquid portion, in the latter water very gradually, until solution occurs,

carefully avoiding any excess, and mix this liquid with the decanted portion.

Macerate the pepsin and pancreatin with the glycerin and water for 24 hours, agitating occasionally; add the tincture, the bismuth solution, and the elixir, and filter through purified talcum.

Each fluidram contains 1 gr. each of pepsin and of citrate of bismuth and ammonium, and  $\frac{1}{2}$  gr. of pancreatin.

#### Elixir of Pancreatin and Pepsin.

Pancreatin, puregr.	-64
Pepsin, puregr.	128
Glycerin fl.oz.	
Waterfl.oz.	
Tincture of cudbearfl.dr.	2
Simple elixir, enough to make.fl.oz.	16

Macerate the pepsin and pancreatin with the glycerin and water for 24 hours, agitating occasionally; add the tincture and elixir, and filter through talcum.

Each fluidram contains 1 gr. of pepsin and  $\frac{1}{2}$  gr. of pancreatin.

## Elixir of Pancreatin, Potassium and Rhubarh.

Refer to Elixir of Rhubarb and its combinations.

#### Elixir of Papain.

sionally, and filter.

Papaingr.	256
Glycerinfl.oz.	~90 91
Sherry wine	04
Saccharingr.	10
Chloroform water	10
Chloroform waterfl.oz.	
Mix, let stand for 7 days, agitating	occa-

Each fluidram contains 2 gr. of papain.

#### Elixir of Paraldehyde.

Paraldehydefl.oz.	4
Glycerin fl.oz.	2
Alcohol fl.oz.	5
Tincture of cardamomfl.dr.	21
Oil of orange	$15^{T}$
Oil of cinnamonm.	15
Compound tincture of cudbear.fl.dr.	2
Aromatic elixir, enough to make fl.oz.	16
Mire the increasion to the suder since	

Mix the ingredients in the order given, and filter, if necessary.—N. F.

Each fluidram contains 15 m. of paraldehyde.

Elixir of paraldehyde varies in strength from 10 to 25 per cent, as prescribed in different localities. The formula here given produces a 25 per cent. elixir, and from this the weaker preparations may readily be made

by the addition of aromatic clixir colored with compound tineture of cudbear in the proportion used in the above formula.

To make a 20 per cent elixir of paraldehyde, for instance, 4 fluidounces of the 25 per cent elixir are mixed with 1 fluidounce of colored aromatic elixir. To make 5 fluidounces of 15 per cent elixir, 3 fluidounces of the 25 per cent elixir are required, and to make the same quantity of 10 per cent elixir, 2 fluidounces of the above elixir are required.

#### Elixir of Pareira Brava.

Fluid extract of pareirafl.oz.	2
Simple elixirfl.oz.	14
Mix allow to stand for 24 hours and	

through talcum.

Each fluidram represents 7½ gr. of pareira brava.

#### Elixir of Pepsin.

Pepsin, puregr.	128
Hydrochloric acidfl.dr.	1.
Glycerinfl.oz.	2
Compound elixir of taraxacum.fl.oz.	1
Alcohol fl.oz.	3
Purified talcumgr.	120
Sugarav.oz.	
Water, enough to make fl.oz.	16

Mix the pepsin with 6 fluidounces of water, add the glycerin and acid, and agitate until solution has been effected. Then add the compound elixir of taraxacum, alcohol, and the talcum, and mix thoroughly. Set the mixture aside for a few hours, occasionally agitating. Then filter it through a wetted filter, dissolve the sugar in the filtrate, and pass the remainder of the water through the filter.

Each fluidram represents 1 gr. of pepsin.— N. F.

# Elixir of Pepsin, Compound. (Elixir of Lactated (or Lactinated) Pepsin.—Compound Digestive Elixir.)

1.,	
Pepsin, soluble scalesgr.	75
Pancreatin, puregr.	8
Diastasegr.	8
Lactic acid	20
Hydrochloric acidm.	40
Glycerinfl.oz.	4
Waterfl.oz.	2
Tincture of cudbear, N. Ffl.dr.	2
Talcum, purifiedgr. 1	120
Aromatic elixir, enough to make fl.oz.	16
Add the acid to the water and glycerin	, and

to this mixture add the pepsin pancreatin, and diastase, and macerate until apparently dissolved; then add the tincture and aromatic elixir; thoroughly incorporate the purified talcum and filter.—N. F.

II.

Pepsin, puregr.	80
Pancreatingr.	40
Diastase or ptyolingr.	10
Cudbear, powderedgr	180
Diluted hydrochloric acidm.	20
Lactic aciddrops	3
Alcohol fl.oz.	5
Waterfl.oz.	7
Simple syrupfl.oz.	4

Mix all the above except the syrup, macerate for 3 days, agitating frequently; filter, to the filtrate add the syrup, and then through the filter add enough of a mixture of alcohol and water, in the proportion of 5 to 7 by measure, to make the liquid measure 16 fluid-ounces.

### Elixir of Pareira and Buchu. Elixir of Pareira and Buchu, Compound.

Refer for these to Elixir of Buchu and its combinations.

Elixir of Pepsin and Bismuth.

Elixir of Pepsin, Bismuth and Cinchona.

Elixir of Pepsin, Bismuth, Cinchona and Iron.

Elixir of Pepsin, Bismuth, Cinchona, Iron and Strychnine.

Elixir of Pepsin, Bismuth and Iron. Elixir of Pepsin, Bismuth, Iron and Quinine.

Elixir of Pepsin, Bismuth and Nux Vomica.

Elixir of Pepsin, Bismuth, and Quinine,

Elixir of Pepsin, Bismuth, and Strychnine.

its combinations.

Elixir of Pepsin, Bismuth and Wafer Ash.

Ash.

Refer for above to Elixir of Bismuth and

Elixir of Pepsin and Cinchona. Elixir of Pepsin, Cinchona and Iron. Elixir of Pepsin, Cinchona and Strychnine.

Refer for above to Elixir of Cinchona and its combinations.

#### Elixir of Pepsin and Iron.

Tincture of citrochloride of iron.fl.dr. 9½ Elixir of pepsin, enough to make fl.oz. 16

Mix and filter, if necessary.

Each fluidram represents about  $\frac{1}{2}$  gr. of chloride of iron (ferric), and nearly 1 gr. of pepsin.—N. F.

#### Elixir of Pepsin, Iron and Quinire.

Refer to Elixirs of Iron and combinations.

#### Elixir of Pepsin and Malt.

Refer to Elixir of Malt and its combinations.

#### Elixir of Pepsin and Quinine.

Quinine sulphategr.	32
Elixir of pepsinfl.oz.	

Agitate until dissolved and filter.

Each fluidram contains 1 gr. of pepsin and ‡ gr. of quinine sulphate.

## Elixir of Pepsin, Quinine and Strychnine.

Strychnine sulphategr.	11
Distilled waterfl.dr.	
Elixir of pensin and quininefl.oz.	

Dissolve the alkaloidal salt in the water and add the elixir.

Each fluidram contains 1-100 gr. of strychnine sulphate, nearly  $\frac{1}{4}$  gr. of quinine sulphate, and nearly 1 gr. of pepsin.

#### Elixir of Pepsin and Strychnine.

Strychnine sulphategr	. 14
Distilled water	. 4
Elixir of pepsinfl.oz	. 15 <del>1</del>

Dissolve the alkaloidal salt in the water and add the elixir.

Each fluidram contains 1-100 gr. of strychnine sulphate and nearly 1 gr of pepsin.

#### Elixir of Pepsin and Wafer Ash.

(Elixir Pepsin and Ptelea.)

Pepsin, puregr.	128
Simple elixirfl.oz.	14
Fluid extract of wafer ashfl.oz.	2
Purified talcumgr.	120

Add the pepsin to the simple elixir, agitate until dissolved, add the remaining ingredients, set aside for 24 hours, and filter.

Each fluidram contains 1 gr. of pepsin and represents  $7\frac{1}{2}$  gr. of wafer ash.

#### Elixir of Phosphorus.

L.		
	Spirit of phosphorusfl.oz.	33/4
	Oil of anise	16
	Glycerinfl.oz.	9
	Aromatic elixir, enough to make fl.oz.	

To the spirit contained in a bottle, add the oil and glycerin, and mix by repeatedly inverting bottle until a clear liquid is obtained. Then add the elixir in several portions, gently agitating after each addition, until all is added.—U. S. P.

#### II.

Phosphorusgr.	21/2
Chloroformfl.dr.	
Alcoholfl.oz.	234
Glycerin, enough to makefl.oz.	16

Dissolve the phosphorus in the chloroform, add the alcohol, and then the glycerin.—Brit. Form

Each fluidram contains 1-50 gr. of phosphorus.

### Elixir of Phosphorus, Compound.

Strychnine sulphategr.	11/4
Quinine sulphategr.	64
Iron pyrophosphategr.	
Distilled water, hotfl.oz.	1
Alcoholfl.oz.	1
Elixir of phosphorusfl oz.	8
Simple elixir, enough to make fl.oz.	16

Dissolve the strychnine salt in 4 fluidrams of the water, and the iron salt in the remainder of the water.

Mix the alcohol and elixir of phosphorus, add the two solutions already prepared, then the quinine salt and the simple elixir, agitate until dissolved, and filter in a well-covered funnel.

Each fluidram contains 1-100 gr. of strychnine sulphate, 1 gr. of iron pyrophosphate, ½ gr. of quinine sulphate and 1-100 gr. of phosphorus.

# Elixir of Phosphorus, Cinchona and Iron.

Refer to Elixir of Cinchona and its combinations.

#### Elixir of Phosphorus and Coca.

Refer to Elixir of Coca and its combinations.

### Elixir of Phosphorus and Damiana.

Elixir of Phosphorus, Damiana and Iron.

Elixir of Phosphorus, Damiana, Iron and Nux Vomica.

Elixir of Phosphorus, Damiana and Nux Vomica.

# Elixir of Phosphorus, Damiana and Strychnine.

Refer for above to Elixir of Damiana and its combinations.

### Elixir of Phosphorus and Gentian.

Refer for this to Elixir of Gentian and its combinations.

# Elixir of Phosphorus and Nux Vomica.

Refer to Elixir of Nux Vomica and its combinations.

# Elixir of Phosphorus, Quinine and Strychnine.

Elixir of phosphorusfl.oz.	8
Quinine hydrochlorategr.	32
Strychnine sulphategr.	11
Distilled waterfl.dr.	4
Tincture of cudbearfl.dr.	
Simple elixir, enough to makefl.oz.	16

Dissolve the quinine salt in 7 fluidounces of simple elixir, and the strychnine salt in the water, mix the two solutions, and then add the other ingredients.

Each fluidram contains 1-100 gr. of strychnine sulphate, ½ gr. of quinine sulphate, and 1-100 gr. of phosphorus.

#### Elixir of Phosphorus and Strychnine.

Strychnine sulphategr.	14
Distilled waterfl.dr.	
Elixir of phosphorusfl.oz.	8
Tincture of cudbearfl.dr.	2
Simple elixir, enough to makefl.oz.	16

Dissolve the quinine salt in the water and add the remaining ingredients.

Each fluidram contains 1-100 gr. each of phosphorus and strychnine sulphate.

#### Elixir of White Pine, Compound.

Fluid extract of white pine bark fl.oz. 1
Fluid extract of balsam gilead
buds
Fluid extract of spikenardm. 64
Fluid extract of wild cherry
bark
Sanguinarine nitrategr. 2
Morphine acetategr. 3
Chloroform m. 64
Alcohol fl.oz. 7
Waterfl.oz. 4
Simple syrupfl.oz. 3
Mix the fluid extracts with the alcohol,

water and syrup previously mixed, and filter through purified talcum until clear; add the chloroform and dissolve the sanguinarine and morphine salts in the mixture.

The above represents the now well-known "white pine cough syrup" in the elixir form.

#### Elixir of Potassium Acetate.

Potassium acetate.....gr. 640 Aromatic elixir, enough to make fl. oz. 16

Dissolve the potassium acetate in the elixir, and filter, if necessary.

Each fluidram contains 5 gr. of potassium acetate.—N F.

# Elixir of Potassium Acetate, Buchu and Juniper.

# Elixir of Potassium Acetate, Buchu, Juniper and Uva Ursi.

Refer for abova to Elixi of Buchu and its combinations.

# Elixir of Potassium Acetate and Juniper.

Potassium acetategr.	640
Fluid extract of juniperfl.oz.	2
Magnesium carbonategr.	
Aromatic elixir, enough to make fl. oz.	16

Triturate the fluid extract with the magnesium carbonate, add 12 fluidounces of aromatic elixir, in which the potassium acetate has previously been dissolved, filter and add the remainder of the aromatic elixir through the filter.

Each fluidram represents 5 gr. of potassium acetate, and 7½ gr. of juniper.—N. F.

#### Elixir of Potassium Bromide.

Refer to the Elixirs of Bromides.

### Elixir of Potassium Iodide. Elixir of Potassium Iodide, Compound.

Refer to the Elixirs of Iodides, etc.

# Elixir Proprietatis. (Tincture Aloes Crocata.)

Aloes, coarse powderav.oz.	1
Myrrh, coarse powderav.oz.	
Spanish saffron, cutgr.	110
Alcoholfl.oz.	16

Mix, macerate for 8 days, and filter.

In the United States, the official tincture of aloes and myrrh is frequently, though improperly, dispensed for the above preparation.

### Elixir Proprietatis Paracelsi.

This is very similar to preceding. It contains double the amount of saffron, and 1½ fluidounces of the alcohol is replaced by diluted sulphuric acid.

### Elixir Pulmonic. (Pectoral Elixir.)

300
128
128
128
64
8
4
1
16

Dissolve the morphine salt in the water add the licorice extract, mix well, add the remaining ingredients, and filter.

Each fluidram contains 1-16 gr. of morphine sulphate.

# Elixir of Quassia, Gentian, Iron and Nux Vomica.

Refer to Elixir of Gentian and its combinations.

# Elixir of Quinine. (Elixir of Quinine Sulphate.)

Quinine sulphate.....gr. 128
Simple elixir.....fl.oz. 16
Mix and dissolve by agitation, warming

gently, if necessary, and filtering.

Each fluidram contains 1 gr. of quinine sulphate.

### Elixir of Quinine Bisulphate.

Quinine	bisulphate.		gr.	128
	lixir			
Dissolve	by agitation	n and	filter, if	neces-
sary.				

Each fluidram contains 1 gr. of quinine bisulphate.

# Elixir of Quinine, Compound. (Elixir of Cinchona Alkaloids.)

Quinine sulphategr.	16
Cinchonidine sulphategr.	
Cinchonine sulphategr.	
Aromatic elixir, enough to make fl. oz.	

Add the alkaloidal salts to the aromatic elixir, dissolve them by agitation, and filter.

—N. F.

Each fluidounce contains 1 gr. of quinine sulphate, and  $\frac{1}{2}$  gr. each of cinchonidine and cinchonine sulphate.

If it is desired to impart a color to this

elixir, this may be effected by the addition of 2 fluidrams of compound tincture of cudbear to the above quantity.

# Elixir of Quinine and Ammonium Valerianate.

Elixir of Quinine, Ammonium, Valerianate and Cinchonidine.

Elixir of Quinine, Ammonium Valerianate, Cinchonidine and Iron.

Elixir of Quinine, Ammonium Valerianate, Cinchonidine, Iron and Strychnine.

Elixir of Quinine, Ammonium Valerianate, Cinchonidine and Strychnine.

# Elixir of Quinine, Ammonium and Strychnine Valerianates.

Refer for above to Elixir of Ammonium Valerianate and its combinations.

Elixir of Quinine and Arsenic.

Elixir of Quinine, Arsenic and Iron.

Refer to Elixir of Arsenic and its combinations.

Elixir of Quinine and Bismuth.

Elixir of Quinine, Bismuth, Iron and Pepsin.

Elixir of Quinine, Bismuth and Pep-

Refer to Elixir of Bismuth and its combinations.

Elixir of Quinine and Croton Chloral. Refer to Elixir of Croton Chloral Hydrate and its combinations.

Elixir of Quinine and Iron Citrate. Elixir of Quinine, Iron and Pepsin. Elixir of Quinine, Iron and Strych-

Elixir of Quinine and Iron Hypophosphites.

Elixir of Quinine, Iron and Strychnine Hypophosphites.

Elixir of Quinine, Iron Phosphate and Strychnine.

Elixir of Quinine, Iron and Strychnine Phosphates.

Elixir of Quinine and Iron Pyrophosphate.

Elixir of Quinine, Iron Pyrophosphate and Strychnine.

Refer for above to Elixir of Iron and its combinations.

#### Elixir of Quinine and Pepsin.

Refer to Elixir of Pepsin and its combinations.

# Elixir of Quinine and Phosphates.

Quinine sulphategr.	32
Iron phosphate, sotuble, hotgr.	128
Potassium citrategr.	
Syrup of calcium lactophos-	
phate	4
Waterfl.oz.	1

Aromatic elixir, enough to make fl. oz. 16

Dissolve the quinine sulphate in 10 fluidounces of aromatic elixir, if necessary, with the aid of a gentle heat. Dissolve the iron phosphate and the potassium citrate in the water, and add the solution to that first prepared. Then add the syrup of calcium lactophosphate and lastly the remainder of the elixir; filter, if necessary.

Each fluidram contains ¼ gr. of quinine sulphate, 1 gr. of iron phosphate, and about ¾ gr. of so-called calcium lactophosphate.—
N. F.

# Elixir of Quinine, Phosphorus and Strychnine.

Refer to Elixir of Phosphorus and its combinations.

#### Elixir of Quinine and Strychnine.

Quinine sulphategr.	64
Strychnine sulphate gr.	
Simple elixirfl.oz.	

Dissolve the alkaloidal salts in the elixir by agitation, and filter.

Each fluidram contains  $\frac{1}{2}$  gr. of quinine sulphate and 1-100 gr. of strychnine sulphate.

#### Elixir of Quinine Valerianate.

Quinine valerianategr.	128
Tincture of cudbear fl.dz.	
Simple elixir, enough to make.fl.oz.	16

Triturate the quinine valerianate with a little of the elixir to a smooth paste. Add about 8 fluidounces more of elixir, triturate until dissolved, add the tincture and the remainder of the elixir.

Each fluidram contains 1 gr. of quinine valerianate.

# Elixir of Quinine, Iron and Zinc Valerianates.

Refer for this to the Elixirs of the Valerianates.

# Elixir of Quinine and Strychnine Valerianates.

Strychnine (alkaloid)gr.	11
Valerianic acidsuffic	ient
Quinine valerianategr.	128
Tincture of cudbearfl.dr.	2
Simple elixir, enough to make.fl.oz.	16

Triturate the strychnine and quinine sulphate with a little elixir to a smooth paste, add 4 fluidounces of elixir and just enough valerianic acid to dissolve the alkaloids; then add the tincture and the remainder of the elixir, neutralize any excess of valerianic acid as described in the formula preceding, and filter.

Each fluidram contains 1 gr. of quinine valerianate and 1-100 gr. of strychnine valerianate.

# Elixir of Quinine Valerianate and Strychnine.

Quinine valerianategr.	128
Strychnine sulphategr.	14
Compound tincture of cudbear.fl.dr.	2
Aromatic elixir, enough to make fl.oz.	16

Triturate the quinine valerianate and the strychnine sulphate with about 8 fluidounces of aromatic elixir, until they are dissolved, then add the compound tincture of cudbear and the remainder of the aromatic elixir, and filter, if necessary.

Each fluidram contains 1 gr. of quinine valerianate, and 1-100 gr. of strychnine sulphate.—N. F.

# Elixir of Raspberry, Compound.

Fluid extract of rhatanyfl.oz.	1
Fluid extract of cinnamonfl.dr.	2
Fluid extract of clovesfl.dr.	2
Fluid extract of nutmegfl.dr.	2
Fluid extract of pimentofl.dr.	2
Raspberry juicefl.oz.	8
Simple elixirfl.oz.	6
Purified talcumav.oz.	1/2

Shake the talcum and fluid extracts in a bottle until well mixed, add the other ingredients and filter.

Perhaps a preferable method of preparation, in some respects at least, would be by extracting the drugs with diluted alcohol, adding the juice to the percolate, flavoring and sweetening the mixture, and filtering.

### and Strychnine Elixir of Rhubarb, Aromatic.

Aromatic	fluid	extract	of rhu-	
barb			fl.oz.	1
Simple eli	xir		fl.oz.	15

This is of the same strength as the aromatic syrup of rhubarb of the United States pharmacopæia.

#### Elixir of Rhubarb.

Sweet tincture of rhubarb (U.		
S. P.)	fl.oz.	8
Alcohol		
Water	.fl.oz.	3
Glycerin	fl.oz.	2
Simple syrup		
Mix and filter.		

Each fluidram represents about 2½ gr. of rhubarb.—N. F.

# Elixir of Rhubarb and Magnesium Acetate. (Elixir of Rhubarb and Magnesia.)

7	
Magnesia, calcinedgr.	150
Acetic acidsuffici	ent
Fluid extract of rhubarbfl.oz.	2
Aromatic elixir, enough to make fl.oz.	16

Dissolve the magnesia in  $2\frac{1}{2}$  fluidounces of acetic acid, with the aid of a gentle heat, adding, if necessary, a little more acid, drop by drop, until the solution is neutral to test-paper. Then add the fluid extract and the elixir, and filter.

Each fluidram represents about 4 gr. of magnesium acetate and  $7\frac{1}{2}$  gr. of rhubarb.

N. F.

# Elixir of Rhubarb and Potassium with Pancreatin.

Rhubarbgr.	320
Golden sealgr.	
Cinnamongr.	
Potassium bicarbonategr.	320
Pancreatingr.	320
Spirit of peppermintfl.dr.	1
Simple syrupfl.oz.	2
Diluted alcohol,	
Simple elixirof each, suffici	ent

Moisten the rhubarb, golden seal and cinnamon (first reduced to a suitable powder), with diluted alcohol, and pack moderately in a percolator; allow to macerate 48 hours and then percolate with diluted alcohol until 6 ounces have been obtained; in the percolate dissolve the potassium bicarbonate and add the pancreatin previously dissolved in the syrup, and about 4 fluidounces of elixir; mix

thoroughly, add the spirit and enough elixir | Elixir of Saccharin. to make the whole measure 16 fluidounces, and filter.

This is similar to the preceding, containing only pancreatin in addition. Like the preceding also, it may be prepared with fluid extracts.

### Elixir of Rhubarb and Potassium.

(Neutralizing Elixir.)

Rhubarb
Golden sealgr. 160
Cinnamongr. 160
Potassium bicarbonategr. 320
Spirit of peppermintfl.dr. 1
Simple syrupfl.oz. 2
Diluted alcohol,
Simple elivir of each sufficient

Reduce the three drugs to moderately coarse powder, extract them in the usual way by percolation with diluted alcohol until 6 fluidounces of percolate are obtained. In this percolate dissolve the potassium bicarbonate, add the spirit of peppermint, syrup, and enough elixir to make 16 fluidounces of product, and filter.

This preparation represents the well-known syrup of rhubarb and potassium in the elixir form.

### Elixir of Rhubarb, Magnesia and Senna.

Magnesia, calcinedgr.	144
Acetic acidsuffic	
Fluid extract of rhubarbfl.dr.	
Fluid extract of senna fl.dr.	
Simple elixir, enough to makefl. oz.	

Dissolve the magnesia in  $2\frac{1}{2}$  fluidounces of acetic acid with the aid of a gentle heat, adding, if necessary, a little more acetic acid, drop by drop, until the solution is neutral to test paper; then add the fluid extracts and elixir, and filter.

Each fluidram contains 4 gr. of magnesium acetate and represents 4 gr. each of rhubarb and senna.

#### Elixir of Rhubarb and Senna.

Fluid extract of rhubarbfl.oz.	2
Fluid extract of sennafl.oz.	2
Tincture of cacaofl.oz.	
Simple elixirfl.oz.	10

Mix and filter, if necessary.

Each fluidram represents 7½ gr. each of senna and rhubarb.

Saccharingr.	384
Sodium bicarbonategr.	
Alcoholfl.oz.	2
Distilled water, enough to make fl.oz.	16

Rub the saccharin and sodium bicarbonate in a mortar, with 8 fluidounces of water gradually added; when dissolved, add the alcohol, filter, and wash the filter with the remainder of the water.

Twenty minims contain 1 gr. of saccharin. -Brit. Form.

This preparation is intended as a sweetening agent in place of sugar or syrup.

### Elixir of Salicylic Acid.

Salicylic acidgr.	640
Potassium citrategr.	960
Glycerinfl.oz.	8
Aromatic elixir, enough to make fl. oz.	16

Dissolve the potassium citrate in the glycerin with the aid of a gentle heat, add the acid, and continue the heat until it is dissolved; then add the elixir. This elixir should be freshly made when wanted for use.

Each fluidram contains 5 gr. of salicylic acid.—N. F.

### Elixir of Salicylic Acid, Compound.

Salicylic acidgr.	640
Sodium bicarbonategr.	
Potassium iodidegr.	192
Fluid extract of black cohoshfl.dr.	. 4
Fluid extract of gelsemiumfl.dr.	. 2
Compound spirit of orange:fl.dr.	
Glycerinfl.oz.	. 4
Waterfl.oz.	
Alcoholfl.oz.	
Simple syrup, enough to make.fl.oz.	16

Mix the acid, sodium bicarbonate and water in a capacious mortar, stir occasionally until reaction is completed, add the potassium iodide, stir until dissolved, then add the alcohol, glycerin, fluid extracts, spirit and syrup, and filter.

#### Elixir of Salicylates and Manaca.

Refer for this to Elixir of Manaca, etc.

# Elixir of Salicylate of Iron.

# Elixir of Salicylate of Iron, Compound.

Refer to combinations of elixirs containing

#### Elixir of Salicylate of Lithium.

Refer to Elixir of Lithium Salicylate.

#### Elixir of Salicylate of Sodium.

Refer to Elixir of Sodium Salicylate.

# Elixir of Salicylate of Sodium and Cascara Sagrada.

Refer to Elixir of Cascara Sagrada and its combinations.

#### Elixir of Senna.

	Deodorized fluid extract of senna. fl. oz	. o
	Compound tincture of cardamom fl.oz.	. 1
	Simple elixirfl.oz	
IJ	•	
	Alexandria sennaav.oz.	11
	Sugarav.oz.	8
	Alcohol,	
	Water,	
	Diluted alcoholof each, sufficie	ent
	Chloroformm.	16
	Oil of corianderdrops	2

Mix  $2\frac{3}{4}$  fluidounces of alcohol with  $8\frac{1}{4}$ fluidounces of water, and with it evenly moisten the senna; pack tightly in a closed vessel, macerate for 3 days, express forcibly, break up the marc, macerate it with enough more of the same kind of menstruum to furnish, in all, 11 fluidounces of liquid, express in 24 hours, mix the two liquids, add the sugar, heat in a closed vessel by means of a waterbath to 94 degrees C., maintain at this temperature 10 minutes, allow to cool, strain, add the chloroform, tincture of capsicum, and oil of coriander, first mixed with 2 fluidrams of alcohol, and finally add, if necessary, enough diluted alcohol to make 16 fluidounces,-Brit, Form.

According to American ideas, a better product could be obtained by macerating the finely cut senna with 12 fluidounces of alcohol for 24 hours, expressing, drying, extracting by percolation with diluted alcohol, dissolving the sugar in the liquid by agitation or percolation, and to the solution adding the chloroform and oil first dissolved in the alcohol.

#### Elixir of Senna, Compound.

Fluid extract of sennafl.oz.	2
Purified tamarind pulpav.oz.	4
Oil of corianderdrops	12
Alcoholfl.dr.	2
Simple elixir, enough to makefl.oz.	16

Dissolve the oil in the alcohol, add to the fluid extract and pulp, then add the elixir.

#### Elixir of Senna and Buckthorn.

Refer to Elixir of Buckthorn and its combinations.

# Elixir of Senna, Magnesia and Rhubarb.

### Elixir of Senna and Rhubarb.

Refer to Elixir of Rhubarb and its combinations.

#### Elixir of Sodium Bromide.

Refer to the Elixirs of Bromides.

# Elixir of Sodium Bromide and Lupulin.

Refer to Elixir of Lupulin and its combinations.

# Elixir of Sodium Hypophosphite.

Refer to Elixir of Hypophosphite of Sodium.

Other elixirs containing sodium hypophosphite may be found with Elixir of Calcium Hypophosphite and its combinations, and with Elixirs of the Hypophosphites.

### Elixir of Sodium Salicylate.

Sodium salicylate......gr. 640 Aromatic elixir, enough to make fl.oz. 16

Dissolve the sodium salicylate in elixir, by agitation, and filter, if necessary.

Each fluidram contains 5 gr. of sodium salicylate.—N. F.

# Elixir of Sodium Salicylate with Cascara Sagrada.

Refer to Elixir of Cascara Sagrada and its combinations.

#### Elixir, Simple.

By a simple elixir is commonly understood a flavored and sweetened dilute alcohol, intended as a vehicle for medicinal remedies, the flavoring agent being a volatile oil or combination of oils. The term may, however, include all such elixirs as are employed as vehicles, although these may be something more than a flavored and sweetened dilute alcohol.

If the second definition holds good, then the following elixirs, which have already been mentioned, will come under the head of simple elixirs: Elixir of orange, elixir of cherries, aromatic elixir, compound elixir of taraxacum, elixir de Garus, elixir of anise and adjuvant elixir. Whenever simple elixir is mentioned, as an ingredient of a preparation in this formulary, some one of the elixirs that are prepared by simply flavoring and sweetening dilute alcohol should be selected, such a one was will best disguise the taste of the medicament; at times it will, of course, be found necessary or advantageous to employ one of the others, like compound elixir of taraxacum, elixir of cherries, etc. In case there be no preference whatever, the aromatic elixir of the pharmacopœia should be used.

It is suggested that a finer product will always be obtained by using deodorized alcohol instead of ordinary alcohol; also that no volatile oil be used which is not only perfectly pure, but also perfectly fresh.

Formulas for making other simple elixirs are the following:

Oil of orangefl.dr.	1/2
Oil of cinnamondrops	5
Oil of anisedrops	2
Oil of bitter almonddrops	1
	5
	163/
Waterfl.oz.	36
Sugarav.oz.	26
	240
W' (1 1 1 - 1 - 1 - 1 - 1 - 1 - 1	
	Oil of cinnamon. drops Oil of anise. drops Oil of bitter almond. drops Tincture of cardamom fl.dr. Alcohol. fl.oz. Water fl.oz. Sugar av.oz. Cacao (Baker's) gr. Magnesium carbonate gr.

Mix the oils, tincture and alcohol, and triturate with the cacao and magnesium carbonate, having first mixed the latter intimately; transfer them ixture to a bottle, add the water gradually, agitate occasionally for several hours, filter, express the filter between muslin, filter the expressed liquid, mix the two filtrates, in the liquid dissolve the sugar by agitation, and filter or strain as may be necessary.

I.	
Oil of orange (fresh)	ı. <b>4</b> 8
Oil of lemondrop	s 12
Oil of corianderdrop	
Tincture of vanillafl.oa	
Tincture cardamomfl.or	
Powdered chocolate (Baker's)g	r. 480
Alcoholfl.os	
Simple syrupfl.oz	
Water (filtered through char-	
coal) enough to make fl.o.	z. 64
D: 1 .1 .1 .1 .4 .1.1	c 1

Dissolve the oils in 4 fluidounces of alcohol, add 12 fluidounces water and the remainder of the alcohol, and shake well. Disand filter.

solve the chocolate in 3 fluidounces of hot water, add the syrup, shake well, add the tinctures of cardamom and vanilla, and enough water to make 64 fluidounces. Filter through paper, on which about  $\frac{1}{2}$  av. ounce of talcum has been distributed; return the first part until the filtrate is clear.

III.	
Oil of orangem.	70
Alcoholfl.oz.	
Purified talcumgr.	$120^{-}$
Orange flower waterfl.oz.	18‡
Simple syrupfl.oz.	181

Mix the oil and alcohol, add the talcum, shake well, add the other ingredients, shake again, and filter.

IV.	
Tincture of fresh orange peelfl.oz	. 12
Tincture of fresh lemon peelfl.oz	. 4
Alcohol	. 8
Orange flower waterfl.oz	
Purified talcumav.oz	. 2
Simple syrup	. 32

Mix the whole well and filter.

This and the preceding have been known as elixir of orange.

V.		
	Oil of sweet orangefl.oz.	11
	Oil of carawaydrops	
	Alcohol fl.oz.	
	Spirit of cinnamondrops	32
	Simple syrupfl.oz.	36
	Glycerinfl.oz.	8
	Distilled waterfl.oz.	4
	Calcium phosphateav.oz.	11

Mix the oils and alcohol, add the calcium phosphate, shake well, add the other ingredients, shake again, and filter.

VI.	
Oil of orangefl.d	lr. 21
Oil of Ceylon cinnamondro	ps 3
Oil of anisedro	ps 3
Oil of carawaydro	ps 6
Tincture of vanillafl.d	r. 9
Simple syrupfl.o	z. 26
Sherry winefl.o	z. 3
Alcoholfl.c	z. 12½
Waterfl.o	z. 23
Purified talcumav.c	z. 1

Mix the oils with the talcum; mix the alcohol, wine and water, add to the mixture of talcum and oils, then add the vanilla and the syrup; let stand one hour, shaking often, and filter.

VII.
Soluble saccharingr. 24
Oil of anise
Alcoholfl.oz. 16
Distilled water, enough to make fl.oz. 64

Dissolve the saccharin in 40 fluidounces of water, add the oil of anise, previously dissolved in 16 fluidounces of alcohol, and the remainder of the water. Add 1 av. ounce of purified talcum; let stand 24 hours, occasionally shaking, and filter.

#### VIII.

Cinnamon waterfl.oz.	24
Simple syrupfl.oz.	24
Alcoholfl.oz.	
Spirit of orangefl.oz.	

This may be clarified by shaking with paper pulp or purified talcum, and filtering. The pulp can be made by beating \frac{1}{2} av.ounce filter paper in a mortar with sufficient water just to moisten it.

#### IX.

Ceylon cinnamongr. 90	
Star anisegr. 60	
Coriandergr. 90	
Nutmeggr. 30	
Caraway	
Oil of sweet orangefl.dr.	2
Diluted alcoholsufficient	
Simple syrupfl.oz. 32	

Percolate the aromatics, previously reduced to coarse powder, with diluted alcohol previously mixed with the oil of orange, continuing the percolation until 32 fluidounces of aromatic tincture are obtained, and mix with the syrup, filtering through talcum, if necessary.

X.Oil of sweet orangefl.dr.	2
Oil of carawayfl.dr.	1
Oil of corianderm.	40
Oil of cassia	40
Alcohol fl.oz.	16
Waterfl.oz.	
Sugarfl.oz.	18
Purified talcumav.oz.	

Mix the oils with the alcohol, add the water and sugar, shake till latter is dissolved, add the talcum, shake again, and filter. XI.

. ± •		
Oil of orange	.fl.dr.	1
Oil of Ceylon cinnamon		
Alcohol		
Simple syrup	.fl.oz. 28	5
Distilled water	.fl.oz. 2	7

Dissolve the oils in the alcohol, add the syrup to this solution until a milkiness or nine valerianate.-N. F.

slight precipitation of oil is produced, then pour the mixture into the remaining syrup, constantly stirring during the whole process and filter, using paper pulp or purified talcum, if necessary, to clarify.

### XII

Orange flower water	fl.oz.	32
Bitter almond water	fl. oz.	8
Simple syrup	fl. oz.	8
Glycerin	fl. oz.	8
Alcohol	fl.oz.	8

Mix all and filter through purified talcum.

### Elixir of Stillingia.

Fluid extract of stillingiafl.oz.	2
Alcoholfl.dr.	4
Simple elixir, enough to makefl.oz.	16

Mix the fluid extract and alcohol, add the elixir, and filter through purified talcum.

Each fluidram represents 7½ gr. of stillin-

# Elixir of Stillingia, Compound.

Compound fluid extract of stil-		
lingia fl.	.oz.	4
Aromatic elixirfl.	oz.	12

Mix them, allow the mixture to stand a few days, or longer, if convenient, and filter.

Each fluidram represents 15 m. of compound fluid extract of stillingia.-N. F.

Compound fluid extract of stillingia		
Alcohol	Compound fluid extract of stil-	
Alcohol	lingia fl.oz.	2
Compound elixir of taraxacumfl.oz. 2 Simple elixirfl.oz. 10		
Simple elixirfl.oz. 10	Compound elixir of taraxacumfl.oz.	2
	Simple elixirfl.oz.	10

Mix the fluid extract and alcohol, add the elixirs, and filter through talcum.

### Elixir of Strychnine Valerianate.

Strychnine valerianate	gr.	11
Acetic acid	.sufficie	nt
Tincture of vanilla		
Compound tincture of cudbea		
Aromatic elixir, enough to make	e.fl.oz.	16

Triturate the strychnine valerianate with about 1 fluidounce of aromatic elixir, gradually added, and effect complete solution by the addition of 1 or more drops of acetic acid, avoiding an excess. Then add the tinctures, and lastly, the remainder of the aromatic elixir. Filter, if necessary.

Each fluidram contains 1-100 gr. of strych-

Elixir o	f	Sumbul.	(Elixir	of	Musk
Root	.)				

Fluid extract of sumbulfl.oz.	$-2rac{1}{2}$
Alcohol fl.oz.	1
Adjuvant elixirfl.oz.	121
Purified talcumav.oz.	

Triturate the fluid extract with the talcum, add the alcohol and elixir, and filter.

Each fluidram represents about 10 gr. of sumbul.

# Elixir of Sumbul, Compound.

Fluid extract of sumbulfl.oz.	2
Fluid extract of scullcapfl.dr.	4
Fluid extract of valerianfl.dr.	1
Alcohol fl.oz.	1
Adjuvant elixir, enough to make.fl.oz.	16
Purified talcumav.oz.	
	_

Mix the fluid extracts and alcohol, add the talcum, shake well, then add the elixir, shake again, and filter.

Each fluidram represents  $7\frac{1}{2}$  gr. of sumbul, about 2 gr. of scullcap, and about  $\frac{1}{2}$  gr. of valerian.

II.

Fluid extract of sumbulfl.oz.	2
Alcohol fl.oz.	1
Elixir of ammonium valerianate.fl.oz.	8
Simple elixirfl.oz.	5

Mix the fluid extract and alcohol, add the two elixirs, and filter through talcum.

Each fluidram contains 1 gr. of ammonium valerianate and represents 7½ gr. of sumbul.

These two mixtures are quite dissimilar, but both are dispensed under the name "compound tincture of sumbul."

### Elixir of Sumbul and Ammonium Valerianate.

Refer to Elixir of Ammonium Valerianate and its combinations.

By way of comparison, refer also to the formula immediately preceding.

#### Elixir of Tar.

Glycerite	of tar	fl.oz.	4
Elixir of	cherries	fl.oz.	12

Each fluidram represents nearly 1 gr. of

#### Elixir of Tar. Compound.

, , , , , , , , , , , , , , , , , , , ,		
Syrup of wild cherry	fl. oz.	31
Syrup of tolu	fl.oz.	31
Morphine sulphate	or.	21
Methylic alcohol	fl.dr.	6
Distilled water, hot	.fl.dr	ĭ
Wine of tar, enough to make	fl.oz	16

Dissolve the morphine sulphate in the water, add the solution to the two syrups previously mixed, then add the methylic alcohol and the wine of tar.

Each fluidram contains 1-50 gr. of morphine sulphate.—N. F.

Most of the methyl alcohol of the market is very impure; the above requires an absolutely pure article.

### Elixir of Tar with Calcium and Sodium Hypophosphites.

Refer to Elixir of Calcium Hypophosphite and its combinations.

### Elixir of Terpin Hydrate.

Terpin hydrategr. 1	28
Glycerinfl.oz.	
Alcohol fl.oz.	
Simple elixir, enough to make.fl.oz.	16

Each fluidram contains 1 gr. of terpin hydrate.

# Elixirs Containing Valerianate of Ammonium.

All elixirs containing valerianate of ammonium may be found under the head of Elixir of Ammonium Valerianate.

### Elixir of Valerianate of Iron.

Refer for this to Elixir of Iron combinations.

### Elixir of Valerianates of Iron, Quinine and Zinc.—(Elixir of Triple Valerianates.)

Iron valerianategr. 64
Quinine valerianategr. 64
Zinc valerianategr. 64
Tincture of cudbearfl.dr. 2
Valerianic acidsufficient
Simple elixir, enough to makefl.oz. 16

Triturate the 3 valerianates with 8 fluidounces of elixir to a smooth paste, add, if necessary, a very small amount of the acid, just enough to dissolve the salts, then add the tincture and the remainder of the elixir, and filter.

If too much valerianic acid has been added so that it is betrayed by its odor, it should be exactly neutralized by stirring with a glass rod repeatedly dipped in dilute ammonia water.

Each fluidram contains ½ gr. each of the valerianates of iron, quinine and zinc.

### Elixir of Valerianate of Morphine.

Refer to the Elixirs of Morphine combinations.

# Elixir of Valerianate of Quinine. Elixir of Valerianate of Quinine and Strychnine.

Refer to Elixir of Quinine and its combinations.

### Elixir of Valerianate of Zinc.

Zinc valerianategr.	128
Stronger solution of ammonium	
citratefl.dr.	$12\frac{1}{2}$
Alcohol fl.oz.	$2^{-}$
Spirit of bitter almondfl.dr.	1↓
Compound tincture of cudbear.fl.dr.	2
Aromatic elixir, enough to make fl.oz.	16

Mix the stronger solution of ammonium citrate with 4 fluidounces of aromatic elixir and the alcohol, and triturate the zinc valerianate with this mixture, added gradually and in portions, until solution has been effected. Then add the spirit tincture, and the remainder of the aromatic elixir. Allow the mixture to stand a few days, and filter.

Each fluidram contains 1 gr. of zinc valerianate.—N. F.

# Elixir of Wafer Ash. (Elixir of Ptelea.) Fluid extract of wafer ash....fl.oz. 234 Simple elixir, enough to make..fl.oz. 16

Mix, and allow to stand for about 24 hours, then filter through purified talcum.

Each fluidram represents 10 gr. of wafer ash.

# Elixir of Wafer Ash, Bismuth and Pepsin.

Refer to Elixir of Bismuth and its combinations.

# Elixir of Wafer Ash and Pepsin.

Refer to Elixir of Pepsin and its combinations.

# Elixir of Wahoo. (Elixir of Euonymus.)

Fluid extract of wahoofl.oz.	21
Waterfl.oz.	
Syrup of coffeefl.oz.	
Compound elixir of taraxacumfl.oz.	$9\frac{1}{2}$

Mix them, let the mixture stand 48 hours, and filter.

Each fluidram represents about  $9\frac{1}{2}$  gr. of wahoo.—N. F.

### Elixir of Wahoo and Blue Flag.

Refer to Elixir of Blue Flag and its combinations.

### Elixir of Wild Cherry.

Fluid extract of wild cherryfl.oz.	4
Alcohol fl.oz.	1
Simple elixirfl.oz.	11

Mix, allow to stand for 24 hours, and filter through purified talcum.

Each fluidram represents 15 gr. of wild cherry.

# Elixir of Wild Cherry and Iron.

Refer to Elixir of Iron and its combinations.

# Elixir of Yerba Santa. (Elixir of Eriodictyon.)

Fluid extract of yerba santafl.oz.	2
Pumice stone, powderedav.oz.	1
Alcoholfl.oz.	1
Simple elixir, enough to makefl. oz.	16

Triturate the fluid extract with the pumice stone until well mixed, add the alcohol, mix again, then add 13 fluidounces of elixir, mix once more, let the whole stand for several hours, stirring occasionally, then filter, returning the first portions of filtrate to the filter until the liquid is clear, and finally adding enough simple elixir through the filter until the filtrate measures the requisite amount.

Each fluidram represents 7½ gr. of yerba santa.

### Elixir of Yerba Santa, Aromatic.

(Elixir Corrigens.)

ι.		
	Fluid extract of yerba santafl.oz.	1
	Simple syrupfl.oz.	8
	Pumice, fine powdergr.	240
	Magnesium carbonategr.	80
	Compound elixir of taraxacum,	
	enough to makefl.oz.	16

Mix 7 fluidounces of compound elixir of taraxacum with the syrup and pumice, then add the fluid extract, and mix the whole thoroughly by agitation. Shake the mixture occasionally during 2 hours, then allow it to settle, and carefully decant the liquid into a funnel, the neck of which contains a small pellet of absorbent cotton. Afterwards add the dregs and allow them to drain. To the filtrate add the magnesium carbonate, and shake occasionally during several hours.

Let the mixture stand at rest during 12 hours, if convenient, then decant the liquid and filter it through paper. To the filtrate add enough compound elixir of taraxacum, if necessary, to make 16 fluidounces.—N. F.

Yerba santa, coarse powdergr. 360
Sweet orange peel, recently
dried and in coarse powdergr. 120
Liquor potassafl.dr. 1
Oil of clovesdrops 4
Oil of cinnamondrops 4
Oil of carawaydrops 2
Oil of corianderdrop 1
Compound tincture of carda-
momfl.dr. 1
Sugarav.oz. 7
Glycerin,
Water,
Alcoholof each, sufficient

Mix the oils and tincture with the drugs and extract by percolation in the usual way, employing as a menstruum a mixture of 1 part of alcohol, 1 of glycerin, and 3 of water, all by measure, with 1 per cent of liquor potassa, 10 fluidounces of percolate are to be obtained, which is to be returned to the percolator if not clear; to this add the remainder of the liquor potassa and 2 fluidounces of alcohol, and in the whole dissolve the sugar by agitation.

HI.

Yerba santa av.oz.	1
Sweet orange peelgr. 1	.44
Cardamom (without capsule)gr.	28
Cloves	28
Cinnamongr.	28
Anisegr.	20
Coriander gr.	20
Caraway gr.	20
Red saundersgr.	10
Sugarav. oz.	7
Alcohol,	•
Glycerin	
Distilled waterof each, sufficient	ent

Mix the drugs, reduce to moderately coarse powder, extract by percolation with a menstruum composed of 1 part of alcohol, 1 of glycerin, and 3 of water, all by measure, until 12 fluidounces of percolate are obtained; in the latter, dissolve the sugar by agitation, and filter.

#### Elixir of Yerba Santa, Compound.

Fluid extract of yerba santafl.oz.	1
Fluid extract of grindeliafl.oz.	1
Alcohol fl.oz.	
Pumice stone, powdered av.oz.	1
Simple elixir, enough to makefl.oz.	

Mix the fluid extracts, triturate with pumice stone, add 13 fluidounces of simple elixir, mix again, allow the whole to stand for several hours, stirring occasionally, and filter.

Each fluidram represents nearly 4 gr. each of yerba santa and grindelia.

#### Elixir of Zinc Bromide.

Refer to the Elixirs of the Bromides.

# Elixir of Zinc Valerianate.

### Elixir of Zinc, Iron and Quinine Valerianate.

Refer for the above to the Elixirs of Valeriantes.

### Emulsion of Almond.

(Emulsion Amygdalæ.—Emulsio Amygdalarum Saccharata.—Almond Milk.—Mistura Amygdalæ—Simple Emulsion.)

Sweet almond	
Sugar	gr. 220
Distilled water, sufficient	to
make	fl.oz. 16

Blanch the almonds with hot water, add the acacia and sugar, beat them in a mortar until thoroughly mixed, rub this mixture with 14 fluidounces of the water, gradually added, until a uniform mixture results, strain, and wash mortar and strainer with the remainder of the water.—U. S. P.

II.	
Sweet almondgr.	
Sugargr.	360
Distilled waterfl.oz.	16

Blanch almonds in the usual way, triturate with a little sugar, then with a little water gradually added, then slowly add the remainder of the water, mix well, strain with expression, and in the colature dissolve the remainder of the sugar.—Codex.

111.								
Sweet	almonds					 	.av.oz.	13/4
Sugar		 	 			 	.av.oz.	13/4
	ed water							

From the almond prepare au emulsion in the usual manner, so as to obtain 15 fluid-ounces of emulsion, and in this dissolve the sugar.—Germ. Form.

IV. Prepare like III, however triturating the almond to fine powder with 1 av. oz. of sugar, the remainder of the sugar to be dissolved in the colature.—Austr. Phar.

V

The following is also called "emulsion of almond," but is also known as look album, white linctus, or white drink:

Sweet almondav.oz.	$3_{\frac{1}{4}}$
Bitter almondgr.	90
Sugar, granulatedav. oz.	
Tragacanth, fine powdergr.	
Orange flower water fl. dr.	
Distilled waterfl.oz.	

Make an emulsion with the almonds, 4 fluidounces of water, and nearly the whole of the sugar, and strain. Triturate the tragacanth with the rest of the sugar, then add small portion of the emulsion, and triturate briskly and for a long time, until well mixed; then add the remainder of the emulsion in small portions, while continuing the trituration, and finally add the orange flower water.

# Emulsion of Almond, Compound.

(Compound Almond Milk.)

Sweet almond, blanchedav.oz	$1\frac{1}{2}$
Henbane seedgr.	144
Calcined magnesiagr.	144
Sugar, moderately fine powd., av. oz.	11/2
Bitter almond waterfl.oz.	11/2
Waterfl.oz.	$13\frac{1}{2}$

From the sweet almond and henbane seeds, prepare an emulsion with the water, strain, add the bitter almond water, mix the magnesia and sugar, mix this emulsion, and shake until the sugar is dissolved.—Germ. Form.

#### Emulsion of Almond Oil.

(Oil Emulsion.—Emulsio Oleosa.)

The following is generally dispensed as simple emulsion:

Sweet almond oilfl.oz	1 3/4
Gum arabic, powdergr.	360
Distilled water fl.oz.	1334

Make an emulsion in the usual manner.— Germ. Pharm.

# Emulsion, Camphorated.

(Emulsio Camphorata.)

I.	
Camphorgr	. 15
Acacia powdergr	. 72
Sugarav.oz	z. 14
Emulsion of almondsfl.oz	z. 16
Alcohol	r. 1

Add the alcohol to the camphor contained in a mortar and reduce to fine powder; now add the acacia, and then the sugar, finally

the emulsion of almonds, and mix the whole well.—H.

II.	
Sweet almondav.oz Sugarav.oz.	13/4
Sugarav.oz.	13/4
Camphor waterfl.oz.	14

Blanch the almond, make into an emulsion with the water, strain, and in the colature dissolve the sugar.—D.

# Emulsion of Monobromated Camphor.

Monobromated Camphorgr.	150
Sweet almond oilfl.oz.	
Gum arabic, powdergr.	225
Distilled watersuffi	

Dissolve the camphor compound in the oil, add the gum, and  $2\frac{1}{2}$  fluidounces of water, triturate until an emulsion is formed, and then add enough water to make 16 fluidounces.—D.

# Emulsion of Castor Oil.

Castor oilav.oz.	$5\frac{1}{2}$
Acacia, fine powder gr.	585
Tincture of vanillafl.dr.	3
Simple syrupfl.oz.	
Watersuffic	

Carefully weigh the castor oil and the acacia into a mortar, triturate until well mixed; then add 2 fluidounces of water all at once to the mixture of oil and acacia, triturating briskly until a thick, creamy emulsion is produced. To this add gradually with stirring, a mixture of the syrup and tincture with a portion of the remaining water, and finally enough water to make 16 fluidounces.

—N. F.

11.		
Castor oil	fl.oz.	11
Egg yolk		
Syrupf	l.oz.	3
Oil of cassia	fl.dr.	1.

Beat the yelk thoroughly in a mortar, add the two oils, triturate until emulsified, and finally add the syrup.

#### Emulsion of Cod Liver Oil.

Emulsions of fixed oils may be prepared by the use of a number of emulsifying agents, such as acacia, tragacanth, extract of malt, yolk of egg, glycerite of yolk of egg, condensed milk, mucilage of Irish moss, tincture of quillaja, and dextrin.

In preparing an emulsion of cod liver oil,

any of these may be employed; The National Formulary recognizes five emulsions of cod liver oil made with mucilage of Irish moss, acacia, glycerite of yolk of egg, tincture of quillaja, tragacanth, and dextrin mucilage. These emulsions may be plain; that is, may consist simply of oil, emulsifier, water, sugar and flavoring, or they may be combined with other agents, such as hypophosphites, phosphates, lactophosphates, phosphorus creosote, etc., or perhaps with two or more such agents. The method adopted in incorporating these medicaments varies according to its character; if it be soluble in the oil, it is added to the latter before emulsifying, and if soluble in water it is dissolved in the latter before it is added to the mixture of gum and Should, however, the substance not be oil. soluble in either the oil or water, then other methods still must be adopted; if it be solid in character, it should be triturated to a very fine powder and be added to the emulsion; if it be an alcoholic liquid, like tincture of benzoin or spirit of nitrous ether, it is generally best to add it to the emulsion after all the water has been added.

Emulsions of cod liver oil usually require flavoring, and this consists most frequently in the addition of a volatile, like cassia, wintergreen, bitter almonds, sassafras, etc. which should be added to the cod liver oil, or it may consist of a flavored syrup like syrup of tolu, which should be incorporated with the emulsion after most of the water has been added. No objection can be made to using two or three flavoring agents.

The emulsions given here are made with the use of different emulsifying agents and different flavoring agents, but no one need be restricted to the use of such of either as may be mentioned.

I.	
Cod liver oil fl.oz.	8
Acaciaav.oz.	$2\frac{1}{4}$
Syrup of tolufl.dr.	12
Flavoring (of any desired kind)'	
Water, of each, suffici	ent

Triturate the oil and acacia together in a mortar, add 3 fluidounces of water all at once to the mixture of oil and acacia, triturating briskly until a thick, creamy emulsion is

the syrup of tolu, and enough water to make 16 fluidounces of finished emulsion.-N.F. (preferred formula.)

11.			
C	od liver oil	fl.oz.	8
Co	ondensed milk, thick	.av.oz.	6
	il of sassafras		
	il of wintergreen		
0	il of bitter almond	drops	2
	lycerin		
	ater, enough to make		

Add the cod liver oil gradually to the condensed milk, then incorporate the other oils, I fluidounce of water, the glycerin, and the remainder of the water.

II.	
Cod liver oil fl.oz.	8
Yolk of two eggs.	
Tragacanth, powderedgr.	16
Elixir of saccharinfl.dr.	
Tincture of benzoinfl.dr.	
Spirit of chloroformfl.dr.	
Oil of bitter almonddrops	8
Distilled water, enough to make fl. oz.	16

Measure 5 fluidounces of distilled water, place the powdered tragcanth in a dry mortar, and triturate with a little of the cod liver oil; then add the yolks of eggs and stir briskly, adding water as the mixture thickens. When of a suitable consistence, add the remainder of the oil and water alternately, with constant stirring, avoiding frothing. Transfer to a pint bottle, add the elixir of saccharin, tincture of benzoin, spirit of chloroform, and oil of almonds, previously mixed, shake well, and add distilled water, if necessary, to make 16 fluidounces.-Brit. Form.

Instead of the elixir of saccharin, the liquor saccharini of the National Formulary may be employed. IV.

The emulsions of cod liver oil with malt extract are considered under the head of Extract of Malt and its combinations.

# Emulsion of Cod Liver Oil, Stronger,

Cod liver oilfl.oz.	8
Acacia, fine powderav.oz.	
Glycerinfl.oz.	1
Sugargr.	480
Water, enough to makefl.oz.	14

Mix the acacia with 2 fluidounces of water, and make an emulsion with this and 6 fluidproduced. To this add the desired flavoring, ounces of oil. Now add the sugar and af terwards incorporate with this emulsion alternate portions of the glycerin and the remainder of the water and oil.

This is a stock emulsion (without flavor) to be used for preparing the usual 50 per cent simple emulsion of cod liver oil, or compound emulsions containing such additions as may be desired.

It is advisable to keep this emulsion in the bottles in which the emulsions are usually dispensed, 7 fluidounces in an 8-ounce bottle, 14 fluidounces in a 16-ounce bottle, etc. Whenever an emulsion of cod liver oil, simple compound, is wanted it can then frequently be prepared by simple addition of the requisite ingredients.

# Emulsion of Cod Liver Oil with Coca.

Stronger emulsion of cod liver oil fl.oz. 14
Elixir of coca......fl.oz. 2
Oil of cloves........drops 24

Each half fluidounce represents nearly 4 gr. of coca.

II.

Prepare an emulsion like any of those mentioned under Emulsion of Cod Liver Oil, replacing 2 fluidounces of water by elixir of coca

# Emulsion of Cod Liver Oil with Creosote and Hypophosphites.

Cod liver oil	fl. oz.	8
Creoste, pure		$1\frac{1}{2}$
Acacia powder	.av.oz	$2^{\frac{1}{4}}$
Glycerin	fl. oz.	1
Syrup of orange	fl. dr.	4
Calcium hypophosphite	gr.	128
Sodium hypophosphite	gr.	128
Oil of wintergreen	fl.dr.	1
Oil of sassafras	.fl.dr.	į
Oil of cinnamon	.fl.dr.	1/2
Distilled water, enough to mak	e fl.oz.	16

Mix the cod liver oil, creosote, and essential oils, with the acacia, in a dry mortar; dissolve the hypophosphites in 3 fluidounces of warm water, pour the solution, all at once, into the mixture of oils, creosote, and acacia, and stir briskly in one direction with the pestle until emulsification takes place; then add the glycerin, syrup, and enough water to make 16 fluidounces, and strain through a cloth.

# Emulsion of Cod Liver Oil with Hypophosphites.

Emulsify the oil with the acacia and 3 fluidounces of water, and add the flavoring. Then dissolve the hypophosphite in sufficient water, mix this solution with the syrup, and add the mixture gradually to the emulsified oil; lastly, add enough water to make 16 fluidounces, and mix the whole thoroughly.

If several hypophosphites are required, equal parts of them may be used, amounting altogether to 128 gr. for the above formula. Varying quantities, larger or smaller than the above, may, of course, be used upon prescription.—N. F.

II.

Calcium hypophosphite......gr. 64
Sodium hypophosphite.....gr. 64
Potassium hypophosphite.....gr. 32
Water, hot.........fl.oz. 2
Stronger emulsion of cod liver oil fl.oz. 14

Dissolve the hypophosphites in the water and add to the emulsion. Flavor as desired.

Instead of the stronger emulsion in this formula, a 50 per cent emulsion may be used. Or the simple emulsion may be prepared in any desired manner, dissolving the hypophosphites in a portion of the water.

# Emulsion of Cod Liver Oil with Hypophosphite of Calcium.

Prepare like any of the preceding, using 128 gr. of calcium hypophosphite to every 16 fluidounces of emulsion.

# Emulsion of Cod Liver Oil with Lactophosphate of Calcium.

Cod liver oilfl.oz. 8
Acacia, fine powderav.oz. 21
Calcium lactate gr. 256
Phosphoric acid (U. S. P., 85
per cent)fl.dr. $1\frac{1}{2}$
Syrup of tolufl.oz. $1\frac{1}{2}$
Flavoring (of any desired kind),
Water of each sufficient

Emulsify the oil with the acacia, and 3

fluidounces of water, and add the flavoring. Then dissolve the calcium lactate in 1 fluidounce of water with the aid of the phosphoric acid, add the solution gradually to the emulsified oil, then the syrup, and lastly enough water to make 16 fluidounces. Mix the whole thoroughly.

This emulsion should be freshly prepared when dispensed.—N. F.

# Emulsion of Cod Liver Oil with Malt Extract.

Refer for this to Extract of Malt and its combinations.

# Emulsion of Cod Liver Oil with Phosphate of Calcium.

Emulsify the oil with the acacia, and 3 fluidounces of water, and add the flavoring. Then triturate the calcium phosphate with the syrup and a portion of the remaining water, add the mixture gradually to the emulsified oil, and lastly, enough water to make 16 fluidounces. Mix the whole thoroughly.—N. F.

# Emulsion of Cod Liver Oil with Phosphates of Calcium and Sodium.

Cod liver oil	. fl.oz.	8
Acacia, fine powder		
Calcium phosphate		
Sodium phosphate		
Syrup of tolu		
Flavoring (of any desired kir	nd)	
Waterof each	h, suffic	ient

Emulsify the oil with the acacia, and 3 fluidounces of water, and add the flavoring. Then triturate the salts to a fine powder, incorporate with the syrup and a portion of the remaining water, and triturate with the emulsified oil. Finally, add enough water to make 16 fluidounces, and mix the whole thoroughly together.—N. F.

# Emulsion of Cod Liver Oil with Phosphate of Sodium.

Every 16 fluidounces should contain 128 gr. of sodium phosphate. This should be dissolved in 2 fluidounces of water which may be added to 14 fluidounces of either simple

or stronger emulsion of cod liver oil, or the solution may be incorporated during the process of emulsification in making an emulsion of cod liver oil.

# Emulsion of Cod Liver Oil, Phosphorated. (Phosphatic Emulsion.—Phosphatic Mixture.)

Cod liver oilfl.oz, 4
Glycerite of yolk of egg (U.
S. P.)av.oz. 3
Diluted phosphoric acidfl.dr. 6
Oil of bitter almonddrops 10
Rum, Jamaicafl.oz. 4
Orange flower water, enough
to makefl.oz. 16

To the glycerite of yolk of egg contained in a suitable bottle, gradually add the cod liver oil, in small portions at a time, shaking after each addition, until the added portion is emulsified. Then gradually add the phosphoric acid, rum and oil of bitter almond, incorporating them thoroughly. Finally, add the orange flower water and mix the whole thoroughly.

# Emulsion of Cod Liver Oil with Wild Cherry.

- 0
Cod liver oilfl.oz. 8
Acacia, fine powderav.oz. 24
Fluid extract of wild cherryfl.oz. 1
Syrup of tolufl.oz. $1\frac{1}{2}$
Flavoring (of any desired kind),
Waterof each, sufficient

Emulsify the oil with the acacia and 3 fluidounces of water, and add the flavoring. Mix the fluid extract and syrup with a portion of the remaining water, and add the mixture gradually to the emulsified oil. Lastly, add enough water to make 16 fluidounces and mix the whole thoroughly.—N. F.

# Emulsion of Linseed Oil, Dr. Thomson's.

Linseed oil	fl.oz.	33/
Oil of wintergreen	fl.dr.	į.
Oil of cinnamon	fl.dr.	1/2
Acacia, powder	av.oz.	2
Water	fl.oz.	6
Glycerin	fl.oz.	11
Simple syrup	fl.oz.	$2\frac{1}{2}$
Hydrocyanic acid, dilute	m.	40

Triturate the mixed oils with the acacia, add 3 fluidounces of water, triturate until emulsified, add the syrup, glycerin, acid, and remainder of the syrup and mix well.

#### Emulsion of Phosphorus. (Phosphorated Emulsion.)

Phosphorated oil	.fl.dr.	7
Acacia, powder		
Distilled water		
Peppermint water		
Simple syrup		
Min the oil and own in a day		

Mix the oil and gum in a dry mortar, emulsify by addition of the water, then add the peppermint water and the syrup, and mix well.—D.

Each fluidram represents about 1-30 gr. of phosphorus.

# Emulsion of Paraffin Oil and Hypophosphites. (Compound Emulsion of Liquid Petrolatum.)

i.e		
	Paraffin oil (liquid petrolatum)fl.oz.	$5\frac{1}{4}$
	Acacia, powderav.oz.	$2\frac{1}{2}$
	Glycerin	14
	Calcium hypophosphitegr.	
	Sodium hypophosphitegr.	96
	Water, enough to makefl.oz.	16

Add the acacia to the oil and mix thoroughly (in a large mortar), then add 5 fluidounces of water (all at once) and rub briskly until the emulsion is formed. Dissolve the hypophosphites in 3 fluidounces of water, to which add the glycerin; then add all the emulsion and rub well together—and any water necessary to make up the measure of 16 fluidounces of finished product.

II.

Liquid petrolatumfl.oz.	4
Oil of sweet almondsfl.oz.	2
Acacia, powderav.oz.	1 3/4
Glycerin fl.oz.	14
Sodium hypophosphitegr.	
Calcium hypophosphitegr.	128
Lime waterfl.oz.	8

Mix all the oil and petrolatum and incorporate well with the gum, then add 3 fluid-ounces of the water all at once, stir briskly until an emulsion is formed; dissolve the hypophosphites in 3 fluidounces of the lime water, mix with the preceding liquid, and then add the glycerin.

These preparations may be flavored in any suitable manner.

#### Emulsion of Resorcin.

Resorcingr. 8	0
Emulsion of almondfl.oz. 1	
Syrup of orangefl.oz.	
Dissolve the massucia in the countries	_

add the syrup.—D.

# Emulsion of Salicylic Acid. (Emulsion Salicylica.)

* /	
Salicylic acidgr	r. 170
Oil of sweet almondfl.	
Acacia, powderav.	oz. $1\frac{1}{2}$
Orange flower waterfl.	oz. $2\frac{1}{4}$
Waterfl.c	
Simple syrupfl.	oz. 13/4

Mix the salicylic acid and oil, add the gum, and to this mixture add all the orange flower water, stirring briskly until emulsification is complete; then add the water and finally the syrup.—D.

This forms about a 2 per cent emulsion of salicylic acid.

### Emulsion of Terpin Hydrate.

Terpin hydrategr.	256
Olive oilfl.oz	
Acacia, powderav.oz	$2\frac{1}{4}$
Syrup of cinnamonfl.oz	
Cinnamon water, enough to	
makefl.oz	16

Reduce the terpin hydrate (with the intervention of a little sugar, if necessary) to a fine powder, triturate first with the acacia and then with the oil, and then add, all at once, 2 fluidounces of cinnamon water. When the emulsion is complete, incorporate the syrup, finally adding the required amount of cinnamon water.

#### Emulsion of Spermaceti.

Spermaceti gr	. 256
Acacia, powdergr	
Waterfl.oz.	
Simple syrupfl.oz	2i

Put the syrup and gum into a warm mortar, add the spermaceti and triturate with a warm pestle until united; then add the warm water gradually, and continue agitation or trituration until the whole is cold.

#### Emulsion of Wax.

This may be prepared like the emulsion of spermaceti, substituting yellow wax for the spermaceti, or by the use of the following method:

Yellow wax, filteredav.oz.	$1\frac{3}{4}$
Mucilage of acaciafl.oz.	5
Distilled water, warmfl.oz.	10

Melt the wax in a capacious mortar on a water bath or by any other suitable method, at the same time warming the pestle; to the wax add the mucilage, mix well until an emulsion is formed, then add the distilled water.—D.

#### Emulsion of Tolu Balsam.

Tolu balsamgr.	144
Alcohol fl.dr.	15
Tincture of quillajafl.dr.	13
Distilled water, hot fl.oz.	$12\frac{1}{2}$

Dissolve the balsam in the alcohol, and add the tincture; then make an emulsion by adding the water gradually without constant agitation.—Codex.

### Essences, Flavoring.

Flavoring essences of various kinds, such as for soda water, other beverages and for culinary purposes may be found in Part VI.

# Essence of Pepsin.

I.	
Pepsin, puregr.	128
Hydrochloric acid, dilutedrops.	5
Aromatic elixirfl.oz.	3
Glycerinfl.oz.	1
Waterfl.oz.	6
Angelica wine, enough to make.fl.oz.	16

Mix all, agitate frequently until the pepsin is dissolved, and filter through purified talcum.

II.

1.	
Pepsin, puregr.	240
Hydrochloric acid, dilutefl.dr.	1
Glycerinfl.oz.	1
Simple syrupfl.oz.	2
Compound elixir of taraxacum.fl.oz.	1
Alcoĥolfl.oz.	$2^{\circ}$
Oil of clovesdrops	1
Water enough to make.fl.oz.	

Mix the pepsin, glycerin, acid and 8 fluidounces of water, agitate frequently until the pepsin is dissolved, then add the syrup, elixir, oil of cloves first dissolved in the alcohol, and the remainder of the water, and filter.

III.

11.	
Pepsin, puregr.	
Simple elixir (U. S. P.)fl.oz.	11
Glycerin	$1\frac{1}{2}$
Catawba winefl.oz.	
Angelica winefl.oz.	5
Waterenough to make.fl.oz.	16
Water	10

Mix, dissolve, and filter through calcium phosphate or purified talcum.

IV.

Fresh calves' rennetav.oz. 3½
Glycerinfl.oz. 3½
Alcoholfl.dr. 14
Tincture of fresh orange peelfl.dr. 1½
Waterfl.oz. 11 <sup>1</sup> / <sub>4</sub>
Purified talcumav.oz. 1

Mix the rennet and glycerin, then add the

alcohol, water and tincture of orange, and macerate for 4 or 5 days, with repeated agitation; add the tale, agitate and allow to stand for an hour, or until the tale has been largely deposited; now decant, on a muslin or flannel filter, the supernatant liquid first, and finally the dregs; then filter through paper. V.

Pepsin, puregr.	128
Glycerinfl.oz.	31
Oil of cinnamondrops	1
Oil of pimentodrops	1
Oil of clovesdrops	2
Hydrochloric acidm.	20
Purified talcumgr.	120
Alcohol fl.dr.	4
Sherry wine (good and light	
color)fl.oz.	31
Distilled water, enough to make fl.oz.	16

Mix the wine with the acid and 6 fluidounces of water; add to it the pepsin and shake until dissolved; the oils are added to the alcohol, triturated with the talc, the pepsin solution gradually added and filtered; return the first portions until the filtrate is perfectly bright, and pass the remainder of water through filter; when every portion has passed, add to the filtrate the glycerin.

#### Essences, Perfume.

These may be found in Part V.

#### Essence of Saccharin.

Saccharingr. 150
Distilled waterfl.oz. 15
Sodium carbonate or bicarbon-
atesufficient
Brandyfl.oz. 1

Mix the saccharin with 3 fluidounces of the water, add the sodium salt gradually in small amounts, stirring thoroughly after each addition, until the saccharin is dissolved, then add the remainder of the water and the brandy, and filter. Any excess of soda must be carefully avoided.

This can be used for sweetening preparations of one kind or another instead of elixir or solution of saccharin. It is especially useful in sweetening the tea, coffee and other beverages of diabetic persons.—D.

#### Essence of Tamarinds.

Tamarind pulp, purifiedav.oz.	83/
Alexandria senna (previously	, ,
extracted with alcohol)av.oz.	14
Water, boilingfl.oz.	40
Infuse for 12 hours, then strain, pres	s the

marc, and evaporate the strained liquor by boiling to 19 av.ounces. Then take 11½ fluidounces of the residue, neutralize with solution of soda, and add:

Alcohol	fl.oz.	34
Simple syrup		
Tincture of vanilla		

#### Etheroles.

These are a class of French preparations similar to the ethereal tinctures used to some extent in this country.

#### Extract of Aloes, Fluid.

What is commonly sold under this name is made by dissolving 8 av.ounces of aloes in 16 fluidounces of diluted alcohol by the aid of a moderate heat, straining, and evaporating the colature to 16 fluidounces.

To make the official tincture of aloes, it is directed to mix  $\mathbf{1}_{\frac{1}{2}}$  fluidounces of the above, and 3 fluidounces of fluid extract of licorice with enough diluted alcohol to make 16 fluidounces.

### Extract of Aloes, Acid.

Aloesav.oz.	4
Water, boilingfl.oz.	20
Sulphuric acid, purefl.dr.	1
Distilled water, coldfl.dr.	3

Pour the boiling water upon the aloes, stir well, allow to cool, add the acid previously mixed with the cold distilled water, set aside for 24 hours, decant the clear liquid, and evaporate the decantate to dry extract.

The yield is about 40 per cent.

This preparation may be made from the dried aqueous extract, but it is more convenient to begin with the crude drug.—D.

#### Extract of Aloes and Myrrh, Fluid.

This, like the fluid extract of aloes, is one of the "convenience" preparations put up by the large manufacturers. It may be prepared as follows:

Socotrine	aloesav.oz.	41
Myrrh	av.oz.	$4\bar{i}$
Alcohol	sufficien	t

Mix the drugs in coarse powder with 12 fluidounces of alcohol, macerate for 7 days in a warm place, occasionally agitating, then heat moderately on a water bath for 2 hours, strain through flannel, add through the strainer enough alcohol to make 16 fluidounces, and allow to cool.

To make the tincture of aloes and myrrh, the manufacturers direct that 5 fluidounces of this "fluid extract" be mixed with 11 fluidounces of alcohol. Inasmuch as the official preparation contains licorice, the tincture is not properly made unless  $1\frac{1}{2}$  fluidounces of the alcohol be replaced by fluid extract of licorice.

# Extract of Alstonia Constricta, Fluid. (Fluid Extract of Australian Fever Bark.)

Extract the drug in fine powder by the use of alcohol as a menstruum, using the process of the pharmacopœia or the National Formulary or any other suitable process of making fluid extracts.

# Extract of Bamboo Brier, Compound

Fluid. (Alterative Compound.—Mc-Dade's Compound.—Creek Indian Remedy.—Mistura Smilacis Compositus.)

Bamboo brier rootav.oz. 4
Stillingiaav.oz. 4
Burdock rootav.oz. 4
Poke rootav.oz. 4
Prickly ash barkav.oz. 1
Diluted alcoholsufficient

Mix the drugs, reduce to fine powder, and extract, using diluted alcohol as a menstruum, by the process of the pharmacopæia, or the National Formulary or any other suitable process for fluid extracts, the product to measure 16 fluidounces.

# Extract of Benzoin, Compound Fluid.

Benzoinav.oz.	434
Storaxav.oz.	3
Tolu balsamav.oz.	$2\frac{1}{2}$
Aloesgr. 3	50
Alcoholsufficie	

Reduce the benzoin and aloes to coarse powder, add these and the tolu and storax to 12 fluidounces of alcohol contained in a closed vessel, digest the mixture, at a temperature not exceeding 65 degs. C., for 2 hours, then strain through flannel, and wash the residue and strainer with enough alcohol to make the colature measure 16 fluidounces.

This is another "convenience" preparation from which the compound tincture of benzoin is directed to be prepared by mixing 6 fluidounces with 10 fluidounces of alcohol.

#### Extract of Benzoin, Fluid.

Benzoin, in coarse powder....av.oz. 8½
Alcohol....sufficient

Mix the benzoin with 12 fluidounces of alcohol, macerate the mixture in a warm place, then heat on a water bath until quite warm, strain through flannel, and wash the dregs and strainer with enough alcohol to make the colature measure 16 fluidounces.

This is similar to the "convenience" preparations put up by manufacturers. The tincture is directed to be prepared from it by mixing 6 fluidounces of it with 10 fluidounces of alcohol. It may also be used for making benzoinated lard and for benzoinating other fats and fatty bodies.

# Extract of Berberis Aquifolium,

Fluid. (Fluid Extract of Oregon Grape.)

Reduce the drug to fine powder and extract by means of a mixture of alcohol and water in the proportion of 3 of the former to 2 of the latter, both by measure, using the process of the pharmacopæia, or the National Formulary, or any other suitable process for fluid extracts.

### Extract of Blackberry Root Bark, Compound Fluid.

Blackberry root barkav.oz	. 12
Cinnamonav.oz	. 11
Nutmegav.oz	. 11
Clovesav.oz	. 3/4
Allspiceav.oz	
Diluted alcoholsuffic	ient´

Mix the drugs, grind to fine powder, and extract, using diluted alcohol as a menstruum, by the process of the pharmacopæia, or National Formulary or any other suitable process for fluid extracts, the product to measure 16 fluidounces.

Other spices may be used instead of those in the above formula.

Such a preparation may be employed for making mild and pleasant carminative elixirs, syrups, "balsams," etc., for summer complaint, dysentery, etc. The above may be converted into the aromatic syrup of blackberry (N. F.) by mixing 1 fluidounce with 5 of diluted alcohol, 11 fluidounces of blackberry juice, and 16 av. ounces of sugar, agitating until dissolved and straining. A

syrup without juice may be prepared by mixing 1 fluidounce of fluid extract, 5 of diluted alcohol and 20 of simple syrup.

# Extract of Bittersweet. (Extract of Dulcamara.)

Dulcamara, coarse powder....av.oz. 4
Distilled water.....fl.oz. 24

Mix the drug with 16 fluidounces of water, macerate for 24 hours, strain with expression, upon the residue pour 8 fluidounces of boiling distilled water, macerate for 1 hour, and strain and express once more. Mix the 2 colatures, add 2 drams or so of filter paper torn into shreds—scraps of filter paper from the cutting of filter papers may be used—and heat the liquid to boiling. Now skim the liquid, filter it through flannel, and evaporate to extract consistency.

A preparation that will keep better may be prepared by evaporating the strained liquid after boiling to rather soft extract, adding to the latter 2 fluidounces of alcohol, mixing well, allowing to stand for 48 hours, filtering and evaporating the filtrate to extract consistency.—D. modified.

# Extract of Black Cohosh, Compound Fluid.

Black cohoshav.oz.	83/
Wild cherry av.oz.	
Licorice	
Ipecacav.oz.	1
Senegaav.oz.	
Diluted alcoholsufficie	nt

Mix the drugs, reduce to fine powder, and extract, using diluted alcohol as a menstruum, by the process of the pharmacopæia or the National Formulary, or any other suitable process for fluid extracts, the product to measure 16 fluidounces.

# Extract of Bryony, Fluid.

Moisten 17½ av.ounces (=16 troy oz.) of bryony in fine powder with 2 fluidounces of water. Allow to stand in a covered vessel for one hour, then mix intimately with 3 fluidounces of alcohol, let stand for another hour, then extract by the process of the pharmacopoeia or the National Formulary, using a menstruum composed of alcohol and water in the proportion of 3 of the former to 2 of the latter, both by measure.

### Extract of Buchu, Compound Fluid.

Buchuav.oz.	11
Cubebav.oz.	21
Juniperav.oz.	21
Uva ursiav.oz.	
Alcohol.	*
Waterof each, suffici	ent

Mix the drugs, reduce to moderately fine powder and extract by the usual process for fluid extracts so as to obtain 16 fluidounces, using a menstruum composed of 1 volume of water and 2 of alcohol.—N. F.

# Extract of Buchu, Juniper and Potassium Acetate, Compound.

Buchuav.oz.	131
Juniperav.oz.	
Potassium acetateav.oz.	1
Alcohol,	

Water......of each, sufficient

Mix the buchu and juniper, reduce them to tolerably fine powder, and extract by the process of the pharmacopæia or the National Formulary. The menstruum should consist of alcohol and water in the proportion of 2 of the former to 1 of the latter, both by measure. The percolate, after dissolving the potassium acetate in it, should measure 16 fluidounces.

# Extract of Buchu and Pareira, Fluid.

Mix buchu leaves and cut pareira in equal parts, reduce to fine powder, and extract by the process of the pharmacopæia or the National Formulary, using as a menstruum a mixture of alcohol and water in the proportion of 2 of the former to 1 of the latter, both by measure.

# Extract of Buchu, Pareira and Uva Ursi, Fluid.

Make like the preceding, using  $8\frac{1}{2}$  av. ounces of buchu and  $4\frac{1}{2}$  av.ounces of each of pareira and uva ursi, the product to measure 16 fluidounces.

# Extract of Buckthorn. (Extract of Frangula.)

1.	The second second	
	Frangula, moderately coarse	
	powderav.oz.	4
	Waterfl.oz.	

Mix the drug with 16 fluidounces of water, macerate for 24 hours, strain with expression, upon residue pour 8 fluidounces of boiling water, macerate again for 1 hour, strain

and express, mix the two liquids, and evaporate to dryness.—D.

II.

Buckthorn bark, coarse powder.av.oz. 8

Diluted alcohol.....fl.oz. 20

Distilled water.....sufficient

Mix the bark and diluted alcohol, macerate for 2 days, pack in a percolator, allow the liquid to drain, and then pass enough water through the drug to make the percolate measure 40 fluidounces. Evaporate this percolate on a water bath to thick extract consistency.—Brit. Pharm.

# Extract of Buckthorn, Bitterless

Fluid. (Extractum Frangulae examaratæ fluidum.)

Bitterless buckthorn bark....av.oz.  $17\frac{1}{2}$  Alcohol,

Water.....of each, sufficient

Extract the bark by the process of the pharmacopæia, or the National Formulary, or any other suitable process for fluid extracts. The menstruum should consist of alcohol and water in the proportion of 3 of the former to 1 of the latter, both by measure, and the product should measure 16 fluidounces.— D.

Bitterless buckthorn bark is prepared by intimately mixing 10 av.ounces of buckthorn bark in very fine powder and 1 av.ounce of calcined magnesia with 20 fluidounces of water, allowing to stand for 12 hours, drying on a water bath with constant stirring, powdering again, and sifting.—D.

# Extract of Buckthorn, Palatable Fluid.

Fluid extract of buckthornfl.oz.	16
Ammoniated glycyrrhizin gr.	120
Saccharingr.	60
Solution of potassafl.dr.	2
Waterfl.oz.	2

Dissolve the saccharin and glycyrrhizin in the water to which has been added the solution of potassa, the fluid extract of buckthorn, and mix thoroughly. The result is an elegant product, free from nauseating or disagreeable taste.

This product is, of course, somewhat weaker than a fluid extract is supposed to be.

# Extract of Cactus Grandiflorus, Fluid.

The preparation sold under this name is usually a concentrated tincture, made from

the freshly gathered fleshy stems, leaves and flowers, which are crushed, covered with alcohol, and macerated for 2 weeks. Three species of plants are now generally employed indiscriminately in making this preparation, viz., Cereus grandiflorus, C. Bonplandi and C. McDonaldi.

#### Extract of Calamus.

Calamus, cut	fine	 	 	 av.oz.	4
Alcohol					
***				00 1	

Water.....of each, sufficient

Mix the drug with 9 fluidounces of alcohol and 12 of water, macerate for 4 days, agitating occasionally, and strain with expression. To the residue, add 4½ fluidounces of alcohol and six of water, macerate for 24 hours, and strain with expression. Mix the two liquids and evaporate on a water bath to thick fluid extract.—Germ. Phar.

# Extract of Calendula, Nonalcoholic, Fluid.

Calendula (flowering herb) in
No 40 powderav.oz. 17½
Glycerinfl.oz 8
Alcohol,
Water of each, sufficient

Moisten the flowers with a menstruum composed of 2 parts of water and 1 of alcohol, by measure, and macerate for 12 hours. Precolate to exhaustion with same strength of menstruum, recover the alcohol by distillation, than evaporate carefully to 8 fluid-ounces, and add the glycerin.

# Extract of Cardamom, Compound Fluid.

What is commercially sold under this title is made as follows:

Cardamom (without capsules) av.oz.	31
Cassia cinnamonav.oz.	$3\hat{1}$
Carawayav.oz.	
Cochinealav.oz.	
Diluted alcoholsufficien	t

Grind the drugs to moderately fine powder, and extract with diluted alcohol by the process of the pharmacœpia or the National Formulary, so as to obtain 16 fluidounces of product.

This product is ten times the strength of the official tincture, which may be prepared by mixing 1 fluidounce of this with  $\frac{1}{2}$  fluidounce of glycerin, and  $8\frac{1}{2}$  fluidounces of diluted alcohol.

#### Extract of Cantharides, Acetic.

Cantharides,			
der		 .av.oz.	4
Alcohol		 fl.oz.	20
Diluted acetic	acid	 . fl.oz.	1

Mix the above, macerate for 8 days, express, allow the colature to stand for several days, then filter, and evaporate the filtrate at a temperature not exceeding 60 degrees C., so that on cooling it will have the consistency of butter.

The yield is about 30 per cent.—D.

# Extract of Carduus Benedictus.

Carduus benedictus, cut ......av.oz. 8 Alcohol, Water,

Diluted alcohol.....of each, sufficient

Macerate the drug with 32 fluidounces of water, macerate for 24 hours, express, macerate the residue with 16 fluidounces of water for 24 hours, express again, mix the two liquids, evaporate to 4 av. ounces, and add to it 4 fluidounces of alcohol. Upon allowing to stand undisturbed in a cool place, crystals of potassium salts will separate; the liquid should be filtered, the substance remaining upon the filter should be macerated, or soaked with 2 fluidounces of diluted alcohol, and filter this in two or three days, mix the 2 filtrates, evaporate to 1½ av. ounces allow the extract to cool, set aside for 8 days, then dissolve it in 5 fluidounces of distilled water, set aside in a cool place for 24 hours, filter and evaporate the filtrate to a thick extract upon a water bath.

The yield is about 16 per cent.—D.

#### Extract of Cascara Sagrada.

Cascara sagrada, No. 20 powder. av. oz. 8 Alcohol,

Water..........of each, sufficient

Mix the drugs with 10½ fluidounces of alcohol and 13½ fluidounces of water, macerate for 6 or 7 days and express. Mix the residue with 7 fluidounces of alcohol, and 8¾ fluidounces of water, macerate for 3 days, and express again. Mix the two liquids, set aside in a cool place for several days, filter, and evaporate the filtrate on a water bath either to thick consistency or to dryness. If made into hard plastic extract, the yield is about 28 per cent; if made into dry mass, it is but about 24 per cent.—D.

A more rational method of preparing this

extract would be by extracting the drug by percolation, using a menstruum composed of alcohol and water in the proportion of 3 of the former to 4 of the latter by measure, and evaporating the percolate as before.

# Extract of Cascara Sagrada, Fluid.

Cascara Sagrada, fine powder..av.oz. 17½
Diluted alcohol........sufficient

Extract the drug by the usual process for fluid extracts so as to obtain 16 fluidounces of product.—U. S. P.

TT

Boil the bark with 3 or 4 successive quantities of water, about 32 fluidounces each time, straining after each boiling. Evaporate the united colatures on a water bath to 12 fluidounces, allow to cool, add the alcohol, let stand for several hours, filter, and to the filtrate add enough distilled water, if necessary, to make 16 fluidounces.—Brit. Pharm.

# Extract of Cascara Sagrada, Bitterless or Tasteless Fluid. (Extractum

Cascaræ Sagradæ Examaratæ Fluidum.)

Bitterless cascara sagrada.....av.oz. 17½ Alcohol, Water.......of each, sufficient

Use as menstruum a mixture of alcohol and water in the proportion of 3 of the former and 1 of the latter by measure. The process of extraction may be that of the pharmacopœia, or the National Formulary, or any other suitable process for fluid extracts. The product should measure 16 fluidounces.—D.

Bitterless cascara sagrada is made by intimately mixing 10 av.ounces of cascara sagrada in fine powder and 1 of calcined magnesia with 20 fluidounces of water, allowing to stand for 12 hours, then drying on a water bath with constant stirring, powdering again, and sifting.—D.

ΤT

Cascara sagrada, No.40 powder. av.oz. 17½
Calcined magnesia.....gr. 80
Alcohol,
Water......of each, sufficient

Mix the alcohol and water in the proportions of 2 parts of alcohol to 3 parts of water

by measure, and moisten the mixed powders thoroughly with the menstruum, then let it stand until the bitterness has disappeared, and pack in a cylindrical percolator, not too firmly, and pour on menstruum. When the liquid begins to drop from the percolator, close the lower orifice, and proceed according to the U. S. P. or N. F. method of preparing fluid extracts, the product to measure 16 fluid-ounces.

III.

Glycerin.....of each, sufficient

Mix the drug and magnesia, moisten with water and allow to macerate for several hours. Then pack the mixture in a percolator and allow to macerate for 48 hours; add 12 fluid-ounces of alcohol, allow to macerate for 12 hours longer, and finally begin percolation, using diluted alcohol as the menstruum. The first 12 fluidounces should be reserved, and percolation continued to exhaustion. Recover the alcohol and evaporate the percolate to soft extract, and dissolve it in the reserved portion, adding sufficient glycerin to make 16 fluidounces.

The bitterless fluid extracts may be used as they are or be made still more pleasant by the addition of licorice and cinnamon or other aromatics. The next preparations are examples of aromatic fluid extract of cascara.

# Extract of Cascara Sagrada, Aromatic Fluid. (Cascara Aromatic.)

Cascara sagrada, fine powder .. av.oz. 17½
Licorice root, fine powder .. av.oz. 134
Calcined magnesia .. av.oz. 2½
Glycerin .. fl.oz. 4
Compound spirit of orange .. fl.dr. 1½
Alcohol .. fl.oz 8
Water,
Diluted alcohol ... of each, sufficient

Mix the powdered drugs and the magnesia with 32 fluidounces of water; macerate for 12 hours and then dry the mixture on a waterbath at a gentle heat. Mix the glycerin and the alcohol with 4 fluidounces of water, and percolate the dried powders with this menstruum, followed by diluted alcohol. Reserve the first 13½ fluidounces that pass, and

set this aside. Continue the percolation with diluted alcohol to practical exhaustion, evaporate this second portion to a soft extract, dissolve it in the reserved portion, and add the compound spirit of orange and sufficient diluted alcohol to make 16 fluidounces of fluid extract.—N. F.

TT

Cascara sagrada, No. 40 powder av.oz.	$17\frac{1}{2}$
Calcined magnesiaav.oz.	13
Waterfl.oz.	18
Alcoholfl.oz.	
Glycerinfl.oz.	4
Extract of licorice, purified, N. F. av. oz.	41
Saccharingr.	
Oil of fenneldrops	5
Diluted alcoholsufficie	ent

Mix the first two ingredients very intimately, moisten with water and macerate for several hours. Then pack the mixture in a percolator and allow to macerate for another period of about 48 hours; then add the alcohol and allow to macerate 12 hours longer. Now start percolation with diluted alcohol and continue until the drug is exhausted. The percolation is allowed to go on very slowly at first, the heavier portion being reserved to the amount of about 13 fluidounces and the remainder in succession to about 16 fluidounces. Recover the alcohol, commencing with the most dilute of the liquors, and evaporate until the whole is reduced to 12 fluidounces, including the glycerin; to this product add the extract of licorice, saccharin and oil of fennel, the whole to measure 16 fluidounces.

III.

Cascara sagrada	, No. 40 powder av. oz.	171
Calcined magne	siagr.	80
Alcohol	fl.oz.	19
Water	suffici	ent

Mix the alcohol with the water in the proportion of 2 of the former to 3 of the latter by measure; moisten the mixed powders thoroughly with the menstruum, and let stand until the bitterness disappears; then pack in a percolator and percolate, adding more water, if necessary, to exhaust the drug. Reserve the first 12 fluidounces of percolate, evaporate the remainder to 2 fluidounces, mix the two and add.

Spirit of cinnamonm.	30
Spirit of nutmegm.	15
Fluid extract of licorice or puri-	
fied extract of licorice, N. F. fl. oz.	1
Glycerin	1

This preparation may also be prepared from the bitterless fluid extract of cascara sagrada by the addition of licorice extracts and aromatics.

### Extract of Cascara Sagrada, Water-Miscible Fluid.

Cascara sagrada (1 year old),	
No. 20 powderav.oz.	$17\frac{1}{2}$
Alcoholfl.oz.	4
Distilled watersufficie	nt

Moisten the bark with a portion of the water; allow to remain a few hours to soften and swell; pack loosely in a percolator, and percolate with more water until exhausted. Evaporate on a water bath to the consistency of a brittle extract, which, when cold, treat with cold water until thoroughly disintegrated. Allow this to stand and settle. Strain through flannel, and evaporate the strained liquor to 12 fluidounces, add the alcohol, when cold, and filter if necessary.

This preparation does not deposit either on keeping or on diluting with water, and, although bitter, is free from nauseous taste and smell.

# Extract of Cascara Sagrada, Compound Fluid.

Fluid extract of cascara sagradafl.oz.	$6\frac{1}{2}$
Fluid extract of licoricefl.oz.	61
Fluid extract of berberis aquifo-	~
liumfl.oz.	31/
—I	

# Extract of Cinchona, Compound Fluid.

The article sold commercially under this title may be prepared as follows:

Alcohol.....of each, sufficient

As a menstruum, use a mixture of the three liquids in the proportion of 1½ fluidounces each of the glycerin and water, and 14 of alcohol. Extract the mixed drugs by any suitable process so as to obtain 16 fluidounces of product.

This product is 4 times the strength of the compound tincture. To make the tincture, mix 4 fluidounces of this extract, 1 of water, and 11 of alcohol.

# Extract of Catechu, Compound Fluid.

The article sold under this name may be made as follows:

Catechu, powder......av.oz. 7 Cassia cinnamon, powder.....av.oz.  $3\frac{1}{2}$ Clean, dry sand,

Diluted alcohol.....of each, sufficient

Mix the two drugs and then add about twice the volume of sand; extract this mixture by the usual method for making fluid extracts, using diluted alcohol as a menstruum, the product to measure 16 fluid-ounces.

This product is 4 times the strength of the official tincture, which latter may be prepared from it by mixing 4 fluidounces with 12 fluidounces of diluted alcohol.

#### Extract of Colchicum Seed.

Colchicum seed, coarse powder..av.oz. 8 Alcohol,

Water.....of each, sufficient

Mix the seed with 10 fluidounces of alcohol and 30 of water, macerate for 5 or 6 days, agitating occasionally, and express. Mix the residue with 14 fluidounces of alcohol and 12 of water; macerate for 3 days, and express again. Mix the two liquids, set aside in a cool place for 2 days, filter, and evaporate the filtrate upon a water bath to thick extract.

The yield is 18 to 20 per cent.—D.

#### Extract of Columbo.

Columbo, coarsely powdered....av.oz. 8 Alcohol,

Water.....of each, sufficient

Mix the drug with 9 fluidounces of alcohol and 11 of water, macerate for 3 days at a temperature of 30 to 40 degs. C., and express. Mix the residue with 6 fluidounces of alcohol and 7 fluidounces of water; macerate for 24 hours, and express again. Mix the 2 liquids, place in a cool place for 2 days, filter, and evaporate the filtrate to dryness. The yield is 9 to 11 per cent.—D.

A preferable method of production would be to extract the fine drug with mixture of 4 parts by means of alcohol and 5 of water, by percolation, and evaporate this percolate to thick extract or drynes.

# Extract of Couch Grass. (Extract of Triticum.—Mellago Graminis.)

Triticum,	cut						.av.oz.	8
Water, bo	iling.						fl.oz.	48

Digest for 6 hours, strain, boil down about 12 fluidounces, allow to cool, add 16 fluidounces of water, filter, and evaporate the filtrate to thick consistency.—Germ. Pharm.

The yield is about 30 per cent.

# Extract of Couch Grass, Fluid. (Fluid Extract of Triticum.)

- I. Refer to the United States pharmacopœia.
- II. The following is a quicker process than that of the pharmacopoeia and yields a product equally good:

Triticum, cutav.oz	. 174
Alcoholfl.oz	. 4
Waterpint	s 10

Boil the drug for one-half hour with 5 pints of water, strain and express; boil the residue again in the same way with the same amount of water, strain again and express. Mix the 2 decoctions, evaporate them to 12 fluid-ounces, add 4 fluidounces of alcohol, filter, and add enough through the filter, if necessary, to make 16 fluidounces.

# Extract of Elecampane. (Extract of Inula.—Extractum Helenii.)

Mix the drug with 19½ fluidounces of alcohol and 24 fluidounces of water, macerate for 24 hours, and express; macerate the residue with 9½ fluidounces of alchol and 12 fluidounces of water for 24 hours and express again. Mix the two liquids and evaporate on a water bath to thick extract.—Germ. Pharm.

A better plan would be to exhaust the ground with a mixture of alcohol and water in the proportion of 4 of the former to 5 of the latter by measure, and then evaporate the liquid obtained, as before.

# Extract of Ergot. (Ergotin.)

The extract of ergot of the United States pharmacopœia is prepared by evaporating the fluid extract which is made by extraction of the drug with diluted alcohol. This extract is now generally dispensed in this country when Bonjean's ergotin is demanded. This latter preparation was originally made,

however, by exhausting powdered ergot with cold water by percolation, heating the percolate to 93 degs. C., filtering, evaporating the filtrate to a syrupy consistence, allowing to cool, adding alcohol in considerable excess to precipitate gummy and albuminous matter, allowing to stand, decanting the clear liquid, or else filtering, and evaporating the decantate or filtrate on a water bath to soft extract.

The ergotin of Wiggers, which is probably no longer in use, was simply a dried alcoholic extract of ergot.

### Extract of Ergot, Fluid.

Mix the acid with 15¾ fluidounces of diluted alcohol; moisten, pack in a percolator, and macerate the drug in the usual way, using this mixture as a menstruum, and then percolate. When all this mixture is exhausted, follow it with diluted alcohol, percolating until no more extractive is obtained. Reserve the first 14 fluidounces of percolate, evaporate the remaining percolate, at a temperature not above 50 degs. C., to soft extract, add this to first percolate, and finally add enough diluted alcohol to make 16 fluidounces.—U. S. P.

II. Another preparation highly desirable for hypodermic use is the following:

"Purified ergot" is digested with twice its weight of water at 65 degs. C., for 24 hours and expressed; the residue is again macerated in warm water for 12 hours. After settling, the expressed liquids are strained and evaporated separately; when both together weigh one-half as much as the ergot employed, they are mixed and sufficient alcohol added to make the liquid of 25 per cent alcoholic strength, or one-third as much as the aqueous solution. After standing for several hours, the liquid is filtered and the gummy residue washed with so much 25 per cent alcohol as to make the filtered liquid measure three-fourths or 75 per cent of the amount of crude drug employed (volume for weight). To this glycerin is added to make the finished preparation represent the amount of

crude drug ergot originally used, pint for pound.

This preparation should be kept in small well-filled bottles in a cool, dark place.

Purified ergot employed in making the latter preparation may be made by percolating coarsely powdered ergot with deodorized benzine, until no more fatty matter is extracted, then passing alcohol through the drug until all the benzine is displaced, and then spreading the extracted drug out in thin layers and allowing it to remain exposed until perfectly dry and free from the odor of alcohol and benzine.

Such a prepared ergot will keep indefinitely.

# Extract of Galanga, Fluid.

Extract the powdered drug by any suitable process, using alcohol as a menstruum, and evaporate the tincture obtained upon a water bath to soft extract.

### Extract of Gentian, Compound Fluid.

The article sold commercially under this title may be prepared as follows:

Water ...... of each, sufficient

Reduce the drugs together to moderately coarse powder, and extract by the usual process for making fluid extracts, so as to obtain 16 fluidounces of product, using as a menstruum a mixture of alcohol and water in the proportions of 6 parts by volume of the former to 4 of the latter.

The product is six times the strength of the official compound tincture, which latter may be prepared from it by mixing 1 fluidounce of it with 3 of alcohol and 2 of water.

# Extract of Golden Seal, Aqueous or Non-Alcoholic Fluid.

The preparation which is sold under this name is generally prepared like the "glycerite of hydrastis" of the pharmacopæia. This is prepared according to the following formulæ:

Moisten the hydrastis with 6 fluidounces

of alcohol, pack it firmly in a percolator, and percolate with alcohol until the hydrastis is practically exhausted. To the percolate add 4 fluidounces of water, and then remove the alcohol by evaporation or distillation. After the alcohol is driven off, add enough water to the residue to make it measure 8 fluidounces, set it aside for 24 hours, then filter, pass enough water through the filter to make the filtrate measure 8 fluidounces, and, lastly, add the glycerin.-U. S. P.

# Extract of Golden Seal. (Extract of Hydrastis.)

Exhaust golden seal in fine powder by any suitable process, using as a menstruum a mixture of alcohol and water in the proportion of 3 of the former to 1 of the latter by volume; then evaporate the tincture on a water bath to soft extract consistency.-D. modified.

The yield is about 30 per cent.

II.

Hydrastis, in fine powderav.oz.	171/
Alcoholfl.oz.	32
Glycerin	
Watersufficie	

Exhaust the hydrastis with the alcohol previously reduced to the strength of 85 per cent by the addition of 4 ounces of water in a suitable percolator. To the percolate thus obtained add the glycerin and water, and recover the alcohol by distillation. Let the residue stand for 2 days to separate resinous matter which deposits, decant the clear liquid, filter it, and add sufficient water to make the preparation measure 16 fluidounces.

### Extract of Golden Seal, Colorless Fluid.

The preparation which is sold under this title is the same as the "colorless solution of hydrastis," which see elsewhere in Part I.

# Extract of Hemlock Spruce, Nonalcoholic Fluid.

Pinus	Canadensis,	in No.	40	
	der		av.oz	. 17½
Water				
	ol			
Glycer	in		fl oz	. 8

Mix the alcohol and water in the proportion of 1 of the former to 2 of the latter, and exhaust the drug by the process of percolation. Distill the alcohol from the percolate fluidounces of alcohol and 12 of water for 2

and evaporate the remaining liquid to 8 fluidounces; to the residue add the glycerin.

# Extract of Granatum. (Extract of Pomegranate Root Bark.)

Exhaust granatum in fine powder by any suitable process, using as a menstruum a mixture of alcohol and water in the proportion of 4 of the former to 5 of the latter by measure. Evaporate the tincture on a water bath to thick extract or to dryness. If evaporated to thick extract the yield is about 20 per cent, and if to dryness, about 16 per cent.—D. modified.

#### Extract of Horsechestnut Bark.

Horsechestnut bark, coarse pow-Distilled water. Alcohol.....of each, sufficient

Mix the bark with 28 fluidounces of distilled water for 12 hours, then heat on a water bath for 2 or 3 hours, and express. Heat the residue for 2 hours with 16 fluidounces of water, and again express. Evaporate the expressed liquids to 4 fluidounces, add 21/2 fluidounces of alcohol, set aside for 24 hours, filter, wash the filter with a mixture of 4 fluidrams of alcohol and 61/2 fluidrams of water, and evaporate the filtrate to dryness. The yield is about 14 per cent.—D. modified.

# Extract of Jamaica Dogwood, Fluid.

(Fluid Extract of Piscidia.)

Jamaica dogwood, fine powder, av.oz. 171/2 Alcohol,

Water ..... of each, sufficient

Moisten the drug with a mixture of the glycerin and 51/2 fluidounces of alcohol, pack in a percolator, and exhaust with a mixture of alcohol and water in the proportion of 3 of the former to 1 of the latter by measure. Reserve the first 13 fluidounces, evaporate the remainder to soft extract, dissolve this in the reserved portion, and add enough of a mixture of alcohol and water like that used for extraction, to make the product measure 16 fluidounces.-D. modified.

### Extract of Kola.

Kola, coarse powder .....av.oz. 8 Alcohol,

Distilled water.....of each, sufficient

Macerate the drug with a mixture of 28

days and express. Macerate the residue with 181/2 fluidounces of alcohol and 8 of water for 2 days and again express. Mix the two liquids, filter, and evaporate the filtrate on a water bath to dryness. The yield is about 80 or 85 per cent.—D.

#### Extract of Kousso, Fluid.

Kousso, coarse powder.....av.oz. 171/2 Alcohol . . . . . . sufficient.

Exhaust the drug by any suitable process of percolation, using alcohol as a menstruum, the product to measure 16 fluidounces.—D.

### Extract of Lavender, Compound Fluid.

The article sold commercially under this title may be prepared as follows:

Cinnamonav.oz.	23
Nutmeggr.	
Red saundersgr.	600
Cloves	300
Oil of lavender flowersfl.oz.	
Oil of rosemaryfl.dr.	2
Alcoholfl.oz.	
Waterfl.oz.	4
Diluted alcoholsuffic	

•Reduce the drugs to coarse powder; add the oils to the alcohol, add the water and with this mixture extract the drugs by any suitable process for fluid extracts, so as to obtain 16 fluidounces of extract, adding diluted alcohol, if necessary, to make up the requisite volume.

The product is eight times the strength of the compound tincture, which latter may be prepared from it by mixing 2 fluidounces of it with 101/2 fluidounces of alcohol and 31/2 fluidounces of water.

#### Extracts, Liquid.

A class of preparations by the name of "liquid extracts" are recognized by the British pharmacopæia and British Formulary. These are in general of the same strength as the fluid extracts of this country, and such of the liquid extracts as are recognized in this formulary are mentioned under the title of "fluid extracts."

#### Extract of Licorice.

T.

Licorice, coarse powder	av.oz. 171/2
Ammonia water	$\dots$ fl.oz. $2\frac{1}{2}$
Distilled water	sufficient

of water, macerate the drug with this mixture, macerate for 24 hours, pack in a percolator, and percolate with the remainder of the mixture, and then with distilled water until the drug is practically exhausted. Evaporate the liquid obtained on a water bath to pilular consistence.

This is the U. S. P. formula for pure extract of licorice.

Extract of licorice in sticks, Water .....of each, sufficient

Put a layer of well-washed rye-straw over the bottom of a keg or other suitable tall vessel. Then put a single layer of sticks of extract, broken into coarse pieces, over it. Continue to put in alternate layers of straw and extract until the vessel is full, or the whole of the extract has been disposed of. Fill the vessel with cold water, and allow it to remain for 3 days. Then draw off the solution, which has formed, by means of a faucet, or siphon, or otherwise, refill the vessel with cold water, and proceed as before. Mix the several solutions obtained, allow any suspended matter to subside, decant the clear solution, and strain the remainder without pressure. Finally evaporate the liquid on a water bath to the consistence of a pilular

This is the N. F. formula for purified extract of licorice.

Licorice root, cut.....av.oz. 8 Distilled water.....sufficient

Dry the licorice, reduce it to coarse powder, macerate with 24 fluidounces of water for 12 hours, and express; mix the residue with 16 fluidounces of hot water, macerate for 1 hour and again express. Mix the two liquids, add some filter paper pulp or scraps of filter paper, boil for not less than 15 minutes, removing the scum, and filtering. This filtrate must be returned to the filter until it is absolutely clear or transparent. Then evaporate to the consistence of thick honey, set it aside in a cool place for 2 days, dissolve in 2 parts of water, filter again, and evaporate on a water bath to thick extract.

Inasmuch as the liquids obtained or produced in the manufacture of this preparation Mix the ammonia water with 48 fluidounces "sour" quite easily, this extract is best made during the cool seasons of the year, the different operations also being performed as rapidly as possible.

The yield from Russian licorice is 35 to 38 per cent; from spanish, 20 to 25 per cent.

—D.

IV.

Russian licorice, coarsely pow-
deredav.oz. 8
Alcoholfl.oz. 9
Watersufficient

Pour 40 fluidounces of cold water over the licorice, allow to stand for 4 hours, stirring frequently, and then express. Extract the press cake with 24 fluidounces of boiling water and again express. Mix the two liquid extracts and at once.

Evaporate to 4 av.ounces, and add to the solution while hot the alcohol, and allow to stand for 24 hours. Then filter through paper and from the filtrate distill off 5 fluidounces. Evaporate the residue to a medium thick extract consistence.

The extract is completely soluble in water. The yield is about 18 to 20 per cent. It is necessary, particularly in summer, to perform the operation rapidly; by beginning at 6 in the morning one can be ready by midday to go on with the evaporation and the alcohol can be added by evening.—D.

# Extract of Lobelia, Compound Fluid.

Lobelia herbav.oz.	83/
Skunk cabbageav.oz.	41/
Blood rootav.oz.	
Diluted alcoholsufficier	

Mix the drugs, reduce to powder, and extract in the usual way by percolation, employing diluted alcohol as a menstruum, the product to measure 16 fluidounces.

# Extract of Logwood, Fluid.

Unfermented	logwood,	No. 2	0	
powder			.av.oz.	16
Distilled water				

Boil the logwood with 32 fluidounces of water in a covered copper or enameled pan for half an hour, and strain. Add 32 fluidounces of water to the residue, boil for another half-hour, and again strain. Repeat the process for a third time, and having mixed the strained liquors, evaporate over a water bath (or preferably in vacuo) until the product measures 16 fluidounces. Set aside

for 7 days, and then decant the clear liquor by means of a syphon from any sediment that may have been deposited.—Brit. Form.

# Extract of Lovage. (Extractum Levistici.)

Exhaust powdered lovage in any suitable manner, using as a menstruum a mixture of 4 parts of alcohol and 5 of water by measure. Evaporate the liquid upon a water bath to the consistence of thick extract. The yield of extract is about 18 per cent.—D.

#### Extract of Malt.

Mix the malt with 16 fluidounces of water, macerate for 6 hours, then add the remainder of the water at a temperature of 30 degs. C. and digest the whole on a water bath at a temperature not exceeding 55 degs. C. for an hour. Strain and press out all the liquid from the residue. Evaporate the colature at once on a water bath, or preferably in a vacuum apparatus, at a temperature not exceeding 55 degs. C., as rapidly as practicable, to the consistence of thick honey.

The extract must be preserved in well-closed vessels in a cool place.—U. S. P. 1880 and N. F.

H.

The following is a process for preparing a malt extract of somewhat different character, the quantities given here being, of course, such as would be used upon a tolerably large scale: these may, of course, be reduced as desired:

Use a tube or churn of a capacity of 5 gallons and having a perforated false bottom. Pour into it 3½ gallons of water of a temperature of about 76 to 78 degs. C., then add ½ peck of barley malt, coarsely ground, gradually stirring it in well. Cover the vessel and set away in a warm place, and allow to remain perfectly at rest for 3 or 4 hours, taking care that the temperature does not fall below 65 degs. C. This is the process of mashing.

mixed the strained liquors, evaporate over a water bath (or preferably in vacuo) until the product measures 16 fluidounces. Set aside now sprinkle over the top of the malt some

water a little above the temperature of the Extract of Malt with Beef. extract, and draw off the fluid below, until the malt is practically exhausted. Evaporate the fluid by means of a water bath to thick consistence.

The tub or churn must be kept perfectly clean. Once a week it should be washed with alkaline water and when not in use it should be kept filled with lime water.

# Extract of Malt, Fluid.

Malt	av.oz. 16
Alcohol,-	
Water	of each, sufficient

Reduce the malt to coarse powder, not finer than No. 20. Moisten it with 8 fluidounces of a mixture of 1 volume of alcohol and 3 volumes of water, and set it aside, well-covered, until it has ceased to swell. Then mix it with as much of the menstruum as it will take up without dripping, pack it uniformly, but without pressure, in a percolator, and add enough of the before-mentioned menstruum to cover it. When the liquid begins to drop from the orifice, close the latter, and allow the contents to macerate during 24 hours, adding from time to time more menstruum, if necessary, to keep the malt just covered. Then remove the cork and allow the percolation to proceed until the percolate weighs 12 av.ounces. Set this aside, well-corked, until any suspended matters have been deposited. Then decant the clear liquid and preserve it for use.

The product thus obtained may be regarded as being practically equivalent to the drug in the proportion of minim for grain, the apparent excess of dissolved matters present in the first portions of the percolate being about offset by the soluble matters still remaining in the drug, when the percolation is interrupted.—N. F.

#### Extract of Malt with Alteratives.

Calcium bromidegr.	128
Sodium bromide gr.	
Potassium iodidegr.	160
Water, hotfl.oz.	
Extract of malt, enough to	
makefl.oz.	16

Dissolve the salts in the water, and add to the extract.

Extract of	f	beef,	Liebig'sav.oz.	1/2
Water			fl.oz.	1
Extract of	οf	malt.	fl.oz.	15

Dissolve the extract of beef in the water and mix with the malt extract.

### Extract of Malt with Beef, Wine and Iron.

Extra	ct of	malt		 	 	fl.oz.	8
Beef,	wine	and	iron.	 	 	fl.oz.	8

### Extract of Malt with Cascara Sagrada.

Fluid extract of ca	scara sagrada fl.oz.	2
	fl.oz. 1	

# Extract of Malt with Cod Liver Oil.

Extract of maltfl.oz.	8
Cod liver oil fl.oz.	
Oil of wintergreendrops	
Oil of bitter almondsdrops	5

Rub the cod liver oil very gradually and thoroughly with the malt extract, then add the flavoring oils. If the mixture becomes too thick at any time, thin by the addition of a little water.

II.

Cod liver oil fl.oz.	8
Tragacanth, powdergr.	
Extract of maltfl.oz.	
Water fl. oz.	

Triturate the gum with the malt extract until well mixed, then add the oil gradually with uninterrupted trituration, and then add the water.

#### III.

Cod liver oil	.fl.oz.	8
Mucilage of dextrin		
Extract of malt.		

To the mucilage of dextrin contained in a suitable bottle, add the extract of malt, and mix them thoroughly by agitation, and then gradually add the cod liver oil, first in small portions, agitating each time until the lastadded portion is perfectly incorporated.

Extract of malt, most suitable for this preparation, should have about the same consistence as balsam of peru, at a temperature of 15 degs. C .- N. F.

Extract of Malt with Cod Liver Oil and Iron. (Extract of Malt with Cod Liver Oil, Ferrated.)
I.  Cod liver oil
Add the oil gradually to the extract by trituration in a mortar until an emulsion is formed, then add the syrup.
Solution of dialyzed ironfl.dr. 5½  Extract of malt with cod liver oil, enough to makefl.oz. 16
Mix well by trituration.
III.  Citrate of iron and ammoniumgr. 64 Water, hot
Dissolve the iron salt in the water and incorporate this solution with malt extract and oil.  IV.
Iron phosphate, solublegr. 64 Water, hotfl.dr. 4 Extract of malt with cod liver oilfl.oz. 15½ Prepare like the preceding.
V. Replace the iron phosphate in IV, with soluble iron pyrophosphate.
Extract of Malt with Cod Liver Oil

### Extract of Malt with Cod Liver Oil, Pancreatin and Pepsin.

Pancreatin, saccharatedgr. Pepsin, saccharatedgr.	64 128
Extract of malt with cod liver oilfl.oz.	

Triturate the pepsin and pancreatin together to fine powder, add a portion of the malt extract with oil, mix well, and add the remainder.

### Extract of Malt with Cod Liver Oil and Pancreatin. (Malt Extract with Pancreatized Cod Liver Oil.)

Cod liver oil fl.oz	. 6
Water fl.oz.	
Extract of maltfl.oz.	
Pancreatin, puregr.	20
Sodium chloridegr.	40
Sodium bicarbonate gr.	60
Oil of pimentosuffici	ent

water, add the oil, and keep at a tempera- to the extract of malt in the usual manner.

ture of about 32 degs. C. for 3 hours, stirring occasionally. Put the malt extract in a mortar, add the pancreatized oil gradually with constant stirring, and flavor the whole with oil of pimento (or other suitable flavoring.)

II. In this preparation, the oil is not first pancreatized as in the foregoing formula:

Pancreatin, saccharated . . . . . . . . gr. 64 Extract of malt with cod liver . . . . . . . fl.oz. 16

Triturate the pancreatin to fine powder, add a small portion of the extract with oil, mix well, and add the remainder.

# Extract of Malt with Cod Liver Oil and Pepsin.

Pepsin, saccharated .....gr. 128 Extract of malt with cod liver

Triturate the pepsin to fine powder, add a portion of the malt extract with oil, mix well, and add the remainder.

# Extract of Malt with Cod Liver Oil and Hypophosphites.

Calcium hypophosphite gr.	64
Sodium hypophosphitegr.	
Potassium hypophosphitegr.	33
Glycerinfl.oz.	1
Water, hotfl.oz.	1
Extract of malt with cod liver	
oil	14

Triturate the hypophosphites to a fine powder, dissolve them as nearly as possible in the water and glycerin, and incorporate with malt extract and oil.

# Extract of Malt with Cod Liver Oil and Phosphorus.

Phosphorus	gr.	1
Cod liver oil	fl.oz.	24
Extract of malt		

Dissolve the phosphorus by the heat of a water bath in 4 fluidounces of the oil in a stoppered bottle, shake thoroughly and when still warm incorporate the extract of malt, then add the remainder of the oil slowly to form an emulsion.

II. Instead of using phosphorus as above, 100 minims of phosphorated oil may be employed, this latter to be added to the cod Dissolve the pancreatin and salines in the liver oil, and this mixture then to be added

III. Instead of using either free phospho-
rus or phosphorated oil, solution of phospho-
rus N. F. may be employed as according to
the following:

Solution of phosphorusfl.oz.	1
Extract of malt with cod liver	
oilfl.oz.	15

Mix well by trituration.

Each half fluidounce contains about 1-100 gr. of phosphorus.

# Extract of Malt with Gentian and Chloride of Iron.

Extract of maltfl.oz.	8
Elixir gentian and chloride of	
iron	8

# Extract of Malt with Hypophosphites.

Calcium hyphophosphitegr.	64
Sodium hypophosphitegr.	
Potassium hypophosphite gr.	
Water, hotfl.oz.	
Glycerinfl.oz.	
Extract of maltfl.oz.	

Rub the salts in a mortar with the water until dissolved or nearly so, add the glycerin and finally the malt extract.

The above is also made to contain, 32 gr. of iron hypophosphite.

# Extract of Malt with Hypophosphite of Calcium.

Calcium hypophosphitegr.	128
Water, hotfl.oz.	
Glycerin	
Extract of malt fl.oz.	14

Triturate the calcium salt to fine powder, dissolve as nearly as possible in the water and glycerin, and add the malt extract.

# Extract of Malt and Iron. (Ferrated Extract of Malt.)

I.

Iron pyrophosphate gr.	64
Water, hotfl.dr.	4
Extract of malt, enough to make fl.oz.	16

Dissolve the iron salt in the water and incorporate the solution with the malt extract.

This is the usual method of preparing ferrated extract of malt.

TT

Solution of dialyzed iron . . . . . fl.dr. 5½ Extract of malt, enough to make fl.oz. 16

#### III.

Citrate	of i	ron	and	an	nno	nium	gr.	64
Water,	ho	ι					fl.dr.	4
Extract	of	mal	t				fl.oz.	151/2

Dissolve the iron salt in the water and add the extract.

IV. Instead of iron citrate in the last formula, soluble iron phosphate may be employed.

v.

Tincture of	citrochloride of iron	fl.dr.	4
Extract of	malt	fl.ož.	151/2

### Extract of Malt with Iron and Quinine Citrate.

Citrate of iron and quininegr.	128
Water, hotfl.oz.	1
Extract of maltfl.oz.	15

Dissolve the iron and quinine salt in the water and incorporate with the malt extract.

# Extract of Malt with Iron, Quinine and Strychnine Citrate.

Strychnine sulphate	2 <sup>5</sup> /8
Extract of malt with iron and	
quinine citrate, enough to makefl.oz. 1	6

Dissolve the strychnine salt in the water and incorporate this solution with the compound malt extract.

#### Extract of Malt with Pancreatin.

Pancreatin,	saccharated	gr. 64
Extract of	malt fl	.oz. 16

Triturate the pancreatin and a small portion of the extract to a smooth paste and add the remainder of the extract.

### Extract of Malt with Pepsin.

Pepsin, pure.,gr.	64
Hydrochloric acidfl.dr.	
Glycerinfl.oz.	1
Water fl.oz.	1
Extract of maltfl.oz.	14

Add the acid to the water and in this dissolve the pepsin, then add the glycerin and finally the malt extract.

# Extract of Malt with Compound Syrup of Phosphates.

Compound	syrup of	phosphates,	
		fl.oz.	8
Extract of		fl oz	

# Extract of Malt with Pancreatin and Pepsin.

Pancreatin, saccharated gr.	64
Pepsin, saccharatedgr. 1	128
Simple syrupfl.oz.	1
Extract of malt, enough to	
make fl.oz.	16

Triturate the pepsin and pancreatin to a smooth paste with the syrup, then add the malt extract.

# Extract of Malt with Phosphate of Iron and Quinine.

Elixir of ir	on phosphate	and
		fl.oz. 8
Extract of :	malt	fl.oz. 8

# Extract of Malt with Phosphate of Iron, Quinine and Strychnine.

Elixir of iron phosphate,	qui-
nine and strychnine	fl.oz. 8
Extract of malt	fl.oz. 8

### Extract of Malt with Yerba Santa.

Fluid ex	tract	of yerba	santa	fl.oz.	1
Extract					

#### Extract of Manaca, Fluid.

Manaca root,	fine powder	av.oz. 171/2
Glycerin	<del>.</del>	fl.oz. 3
Alcohol.		

Water ..... of each, sufficient

Moisten the drug with a mixture of the glycerin and 4½ fluidounces of alcohol, then pack in a percolator and exhaust by the usual process of percolation, using as a menstruum a mixture of 3 parts by measure of alcohol and 1 of water. Reserve the first 11 fluidounces of percolate, evaporate the weak percolate to 5 fluidounces and mix with the reserve percolate.—D. modified.

#### Extract of Mezereon.

Exhaust mezereum in coarse powder by means of percolation or any other suitable process, using alcohol as a menstruum. Evaporate the percolate on a water bath to thin extract.—Germ. Phar.

The yield of extract is about 10 per cent.

#### Extract of Opium, Aqueous.

This extract of opium of the German Pharm, differs from that of the U. S. Pharm, in not containing an addition of milk sugar to standardize the morphine strength and keep it in the powder form.

This addition of a foreign substance is,

however, not necessary, if the process is modified, in shaking the concentrated opium infusion with an equal part of petroleum benzin; allow to stand together until separation has taken place; decant the benzin and evaporate the infusion to dryness. The extract so produced is easily reduced to powder and will remain in this form, is readily soluble in water, producing therewith a clear solution.

### Extract of Opium, Aqueous Deodorized Fluid.

The article sold generally under this title is the deodorized tincture of opium of the U. S. pharmacopœia.

# Extract of Opium, Camphorated Fluid. (Concentrated Paregoric.)

Opium, fine powder gr. 240	)
Benzoic acidgr. 240	)
Camphor	)
Oil of anisefl.dr. 4	Į
Alcoholfl.oz. 19	9
Waterfl,oz.	}

Dissolve the camphor, oil and acid in the alcohol, triturate opium to a smooth paste with the water, mix the two liquids, macerate for 7 days, agitating frequently, and filter.

This is similar to the "convenience" preparations put up by manufacturers which may be used for making paregoric. To make the latter mix 2 fluidounces of the concentrated preparation with 1 fluidounce of glycerin and 13 fluidounces of diluted alcohol.

### Extract of Pink Root and Senna, Fluid.

Fluid extract of pink rootfl.oz. 10 Fluid extract sennafl.oz. 6
Oil of caraway
Oil of anise

#### Extracts, Powdered.

Most extracts now appear on the market, not only in the old-style plastic form, but also in the form of powder. Inasmuch as almost all plant extracts are injured by the application of heat, the powdered extracts do not very well represent the drug, less so in fact than almost any preparation made from a crude drug. However, there is a demand for these powdered extracts, and this demand must and will be supplied. Extracts

like those of opium and kino can be evaporated to dryness without any appreciable injury to the principles present, but others again, like hyoscyamus, valerian, etc., are very susceptible to change, either because of the presence of easily decomposable principles or because of the presence of volatile matter. These latter extracts should never be prepared in the powdered form.

In making powdered extracts of drugs containing volatile or easily decomposable principles, the temperature employed in drying should be quite low, say not to exceed 55 degs. C. The drying of the extract may be facilitated by spreading out in a thin layer and warming in a drying room or closet at the specified temperature, driving a current of warm, dry air through the chamber if this be convenient. Even then it may not be advisable or possible to reduce the extract to such dryness that it can be powdered, and then it becomes necessary to add a small amount of some substance, such as milk sugar or some of the powder of the drug itself. If the drug contains considerable fixed oil, the extract cannot be dried thoroughly and the intervention of such a powder is necessary. It is also to be remembered in this connection that in making powdered extracts, glycerin must not enter into the menstruum employed in the extraction of the drug, as this will prevent the extract from drying.

After an extract has been reduced to dryness, either with or without the intervention of the added powder, it should be reduced to fine powder and then be preserved in well-stoppered, wide-mouthed bottles.

#### Extract of Rhubarb, Aromatic Fluid.

Rhubarbav.oz.	131
Cinnamonav.oz.	23/4
Cloves av.oz.	23/
Nutmeg av.oz.	$1\frac{1}{4}$
Diluted alcohol sufficie	

Reduce the drugs to moderately coarse powder and extract by the usual method of percolation, so as to make 16 fluidounces of product, using diluted alcohol as a menstruum.—N. F., 1st edition.

If 1 fluidounce of this be mixed with 15 fluidounces of syrup, the product is practical-

ly identical with the aromatic syrup of rhubarb of the pharmacopæia.

### Extract of Rhubarb, Compound.

Extract of rhubarb av.oz.	3
Extract of aloesav.oz.	
Resin of jalapav.oz.	1/2
Soap, powderedav.oz.	2

All of the above, if not in fine powder, should be reduced to this condition and then should be well mixed.—Germ. Phar.

### Extract of Quebracho.

Exhaust finely powdered quebracho by any suitable process, using as a menstruum a mixture of 4 parts of alcohol and 5 of water by volume, then evaporate the tincture on a water bath either to thick extract or to dryness.

The yield of thick extract is about 11 per cent, of dry extract 9 to 10 per cent.—D. modified.

### Extract of Rose, Inspissated.

Rose le	aves,	cut	moder	ately		
coarse					.av.oz.	83/
Alcohol,						, ,
Water,						
Glycerin			of e	ach.	sufficier	nt

Mix 30 fluidounces of alcohol and 10 of water, pour this mixture on the rose leaves, allow it to stand for 24 hours, express and evaporate the resulting fluid to 4 fluidounces. Allow the evaporated extract to stand for 24 hours at the room temperature, filter and evaporate the filtrate to a syrupy consistence. Then add sufficient glycerin to bring the whole up to 2 fluidounces. Of this extract, which is clearly soluble, 4 fluidrams is sufficient to make 16 fluidounces of honey of rose.

However, according to the U. S. pharmacopecia, honey of rose should be made from fluid extract of rose.

#### Extracts, Saccharated.

These are a class of powdered extracts which represent the drug, weight for weight. They are prepared by exhausting the powdered drug with a suitable menstruum, evaporating the tincture to thick extract, adding some sugar of milk, continuing the evaporation to dryness, powdering and adding enough milk sugar to make up the weight of the original drug.

With one exception, these extracts are

seldom or never prescribed or used, the exception being saccharated extract of coto.

### Extract of Sarsaparilla.

Exhaust this drug in fine powder by the process of percolation or any other suitable process, using as menstruum a mixture of 6 parts of alcohol and 5 of water by volume; evaporate the tincture on a water bath to thick extract. The yield is about 20 per cent.—D. modified.

# Extract of Sarsaparilla, Compound Fluid.

I.

Sarsaparillaav.oz.	13‡
Licorice root gr.	
Sassafras barkav.oz.	134
Mezereum av.oz.	1/2
Glycerin	13
Alcohol,	
Water of each, suffic	iert

Mix the glycerin with 5 fluidounces of alcohol and 10 of water. Percolate the drugs, mixed and ground to coarse powder, in the usual manner for making fluid extracts, using this mixture as a menstruum, so as to obtain 16 fluidounces of product. If more menstruum be required, a mixture of 1 volume of alcohol and 2 of water should be employed.

—U. S. P.

II. Another preparation of the same name is offered by manufacturers for the rapid preparation of the compound syrup of sarsaparilla. This preparation may be made as follows:

Sarsaparillaav.oz.	143
Licoricegr.	
Senna	
Oil of sassafrasdrops	3
Oil of anisedrops	3
Oil of wintergreendrops	3
Alcohol,	
Waterof each suffici	ent.

Mix the drugs, reduce them to moderately fine powder and extract by percolation or any suitable process to obtain 16 fluidounces of product, using as a menstruum a mixture of 1 part of alcohol to 2 of water by measure. To the product obtained add the volatile oils and shake well.

To make the compound syrup of sarsaparilla, mix 4 fluidounces of this extract with 12 fluidounces of simple syrup.

### Extract of Senega.

Exhaust senega in fine powder by percolation or any other suitable process, using as a menstruum a mixture of 4 parts of alcohol and 5 parts of water by measure; then evaporate the tincture on a water bath to dryness. The yield is about 25 per cent.—D. modified.

### Extract of Senna, Aqueous Fluid.

The preparations sold commercially under this title are usually made by infusing cut senna leaves with enough hot water to just cover them, allowing to macerate for an hour or two, straining and expressing the liquid, infusing the residue again as before, straining and expressing the liquid again in the course of an hour or two, and mixing the two liquids. This liquid usually measures more than a finished fluid extract should measure, and inasmuch as it is not practicable to evaporate this infusion, the fluid extract of the market is prepared from the latter by adding enough alcohol to preserve it, allowing to stand for 24 hours, and filtering the liquid from the precipitated mucilage. The product is, of course, not properly a fluid extract.

A better process for this preparation is the process of the National Formulary for making deodorized fluid extract of senna, which is as follows:

	av.oz. 17½
Alcohol,	of each, sufficient
water	or each, sumclent

Moisten the senna with 5½ fluidounces of alcohol, pack it firmly in a percolator, and percolate it with alcohol until it is practically exhausted by this menstruum. The alcoholic percolate thus obtained is rejected, and the alcohol may be recovered therefrom by distillation. Then take out the moist powder, dry it, and extract by the usual method for making fluid extracts, using diluted alcohol as a menstruum, so as to obtain 16 fluidounces of product.

### Extract of Senna, Compound Fluid.

Senna
Jalap
Corianderav.oz. 2¼
Alcohol,
Water of each, sufficient

Reduce the drugs to fine powder and ex-

haust by percolation or any other suitable process for fluid extracts, using as a menstruum a mixture of 2 parts of alcohol and 1 of water by measure. The product should measure 16 fluidounces.

#### Extract of Senna and Jalap, Fluid.

Senna	83/4
Jalapav.oz.	834
Alcohol,	
Waterof each, sufficier	nt

Mix the drugs, reduce to fine powder, and exhaust by percolation or any other suitable process for fluid extracts, using as a menstruum a mixture of 2 parts of alcohol and 1 of water by measure. The product should

# measure 16 fluidounces. Extract of Squill.

Squill, coarse powder	av.oz. 8
Alcohol,	
337	

Water ..... of each, sufficient

Macerate the drug for 48 hours with 16½ fluidounces of alcohol and 5½ of alcohol and express. Macerate the residue again for 48 hours with 10 fluidounces of alcohol and 3¼ fluidounces of water and again express. Mix the two liquids, filter, and evaporate the filtrate on a water bath to thick consistency.

The yield is about 36 per cent.—D. modified.

#### Extract of Sumbul, Fluid.

Exhaust sumbul in fine powder with alcohol as a menstruum, using percolation or any other suitable process for extraction.

# Extract of Tobacco, Rademacher's.

Tobacco leaves, freshly gathered (green),
Water ......of each, sufficient

Cut the leaf, contuse in a mortar with sufficient water to make a pasty mass, express, and evaporate the liquid to soft extract. The yield is about 4 per cent.

### Extract of Tolu, Fluid.

Tolu balsam......av.oz. 834 Alcohol, sufficient to make...fl.oz. 16

Digest the balsam in a closed vessel with 10 fluidounces of alcohol on a water bath until dissolved, then strain through flannel, and wash the vessel and strainer with enough alcohol to make up the required amount.

This is not rightly a fluid extract; it is

designed for the convenient preparation of the official tincture, which may be prepared by mixing  $2\frac{1}{2}$  fluidounces of the "fluid extract" with  $13\frac{1}{2}$  fluidounces of alcohol.

#### Extract of Tolu, Soluble Fluid.

The preparation which passes under this name may be prepared according to the following process:

Tolu balsamav.oz.	21/2
Magnesium carbonategr.	
Glycerin fl.oz.	
Water,	
Alcohol	ient

Mix 5 fluidounces of the alcohol with the glycerin, add the balsam, and dissolve the latter by the aid of a moderate heat, shaking frequently, and avoiding loss by evaporation. Now add 6 fluidounces of water, allow the mixture to become cold, decant the milky liquid from the resinous precipitate, mix the decantate intimately with the magnesium carbonate in a mortar, filter, and wash mortar and filter with enough of a mixture of 1 part of alcohol and 2 of water by measure to make the filtrate measure 16 fluidounces.

This is designed for the rapid manufacture of syrup of tolu, which may be made by mixing 1 fluidounce of this "fluid extract" with 15 of simple syrup.

It is to be noted that the National Formulary recognizes a soluble tincture of tolu which was intended for the rapid preparation of syrup of tolu; this tincture is, however, not 16 times the strength of the syrup, as stated by the National Formulary.

### Extract of Valerian.

Exhaust valerian root in fine powder by percolation or any other suitable process, using as a menstruum a mixture of 4 parts of alcohol and 5 of water by volume. Evaporate the tincture obtained on a water bath to thick extract. The yield is about 20 per cent.—D. modified.

Extract of Wahoo. (Extract of Euonymus.)

Wahoo, coarse powder ......av.oz. 8 Water, Alcohol ......of each, sufficient

Mix 2 volumes of alcohol and 1 of water and extract the drug in the usual way by percolation, so as to obtain 22 fluidounces of percolate or until the drug is exhausted. Evaporate this percolate on a water-bath to pilular consistence.—U. S. P.

II. The above preparation is in the form of a plastic extract; the British Formulary recognizes a similar preparation called "dry extract of euonymus" or "euonymin," which is prepared as follows:

Euonymus, No. 20 powder ....av.oz. 8 Diluted alcohol,

Milk sugar ......of each, sufficient

Moisten the bark with 8 fluidounces of diluted alcohol, pack in a percolator, and percolate with the same liquid until exhausted. Evaporate the percolate on a water bath to remove the alcohol. While the extract is still soft, incorporate with it enough powdered milk sugar—the actual amount required having been ascertained by first experimenting with a small amount of the extract—so that the final product shall contain 80 per cent of dry extractive. Then continue evaporation until the mixture is brittle when cold. Then powder and put into a well-stoppered bottle.

#### Extract of Wormwood.

Wormwood, fine powder .....av.oz. 8 Alcohol,

Water .....of each, sufficient

Exhaust the drug by percolation or any other suitable process, using as a menstruum a mixture of 3 parts of alcohol and 10 of water by volume. Evaporate the tincture obtained on a water bath to thick extract.—Germ. Pharm. modified.

The yield is about 32 per cent.

# Extract of Yellow Dock, Compound Fluid.

Yellow dockav.oz.	83/
False bittersweetav.oz.	41/2
Figwortav.oz.	21/4
American ivyav.oz.	21/4
Diluted alcoholsufficier	nt

Mix the drugs, reduce to fine powder, and exhaust by percolation or any other suitable process for fluid extracts, using diluted alcohol as a menstruum, the product to measure 16 fluidounces.

#### Extract of Yarrow.

Yarrow,	fine powderav.oz. 8	
Alcohol,		
Water	of each, sufficient	

Exhaust the drug by percolation or any other suitable process, using as a menstruum a mixture of 4 parts of alcohol and 5 of water by measure. Evaporate the tincture on a water bath to thick extract.

The yield of extract is about 22 per cent.—D.

#### Fats.

Fats, or greases, more properly termed Lards, may be found under the latter designation.

### Fuligokali.

Caustic potassaav.oz. 1	
Wood sootav. oz. 5	
Water sufficient	

Dissolve the potassia in a sufficiency of water, about 32 fluidounces, add the soot, boil for one hour, dilute largely with water, filter, evaporate, filtrate to dryness, and place in a well-corked bottle.

The dose is 2 or 3 grains.

### Glycerite of Alum.

Alum,	powder.	 	 			 av.oz.	3
Glyceri	n	 		٠.	 •	 .fl.oz.	141/2

Stir together in a porcelain dish, apply a gentle heat until solution is effected, set aside and decant the clear fluid from any deposited matter.—Brit. Pharm.

#### Glycerite of Borax.

Borax,	powderav.oz	. 4
	nfl.oz	
CD: 1.	1 .11 11 1 1	4

Triturate together until dissolved, or else warm gently, stirring constantly until dissolved.—U. S. P. 1870.

### Glycerite of Boric and Tannic Acids.

Boric acid	
Tannic acid	av.oz. 13/4
Glycerin	fl.oz. 13

Mix the acids with the glycerin, heat on a water bath until dissolved, and strain.

### Glycerite of Carbolic Acid.

Carbolic acid,	crystal	.av.oz.	31/2
Glycerin	crystal	fl.oz.	123/4

Warm the acid, add the glycerin, and stir until mixed.—U. S. P. 1870.

#### Glycerite of Creosote.

Creosote	fl.oz.	13/
Alcohol	.fl.oz.	2
Glycerin	.fl.oz.	51/2
Water	.fl.oz.	63/
Magnesium carbonate	av.oz.	1

Triturate the magnesium carbonate, alcohol

and creosote together in a mortar, add the water and the glycerin, put the whole in a bottle, let stand for several days and filter. The product represents about 10 per cent by weight of creosote, and may be used for making other preparations of this agent.

## Glycerite of Chloroform.

Chloroform	.fl.oz.	11/4
Alcohol	fl.oz.	41/2
Glycerin	.fl.oz.	$10^{'}$
Dissolve tr. e chloroform in the a		
e glycerin, and shake well.		

The product represents 10 per cent by weight of chloroform.-D.

## Glycerite of Gallic Acid.

Gallic acid									av.oz:	31/2
Gallic acid Glycerin	 								.fl.oz.	123/4

Mix well, heat on a water bath until the acid is dissolved, and strain .- Brit. Pharm.

## Glycerite of Guaiac.

Resin of guaiac, powder gr.	640
Solution of potassafl.oz.	1
Glycerin	91/2
Glycerin	16

Mix the solution of potassa with 5 fluidounces of water, and in this liquid macerate the resin for 24 hours. Then filter, and pass enough water through the filter to make the filtrate measure 61/2 fluidounces, and mix this with the glycerin.-N. F.

### Glycerite of Lead Subacetate.

Lead acetateav.oz.	$3\frac{1}{2}$
Lead oxide, powderav.oz. 2; gr.	20
Glycerin	
Distilled waterfl.oz.	

Mix all, boil together for 15 minutes, then filter, and heat again until all the water has evaporated.-Brit. Pharm.

This is of the same strength the solution of lead subacetate U. S. P., and may be employed in making the diluted solution of lead subacetate.

## Glycerite of Starch.

Starchav.oz	. 1
Waterfl.oz	. 1
Glycerinfl.oz. 6½, or av.oz	. 8

To the starch, contained in a porcelain capsule, add the water and glycerin, and stir until a homogenous mixture results. Then apply heat, gradually raising the temperature to a point between 140 and 144 degrees C., stirring constantly until a transparent jelly is an hour, and strain.—Brit. Pharm.

produced. Transfer the product to suitable vessels, provided with well-fitting covers .-U. S. P.

#### Greases.

Greases or fats, more properly termed "lards," may be found under the latter designation.

## Honey of Borax.

1.										
	Borax	po	wder.		 				.av.oz.	2
	Clarifi	ed	honey				٠.	:	.av.oz.	16

Mix and dissolve by the aid of a gentle heat.—U. S. P., 1870.

II.	
Borax	2
Glycerinfl.oz.	1
Honeyav.oz.	
Prepare like the preceding.—Brit. Pha	rm.

## Honey of Rose with Borax.

Honey	of	rose,	U. S	. Р.,	 .av.oz.	10
Borax.					 .av.oz.	1

Mix and dissolve borax by aid of a gentle

## Honey of Rose with Salicylic Acid.

Hone	y of rosea	v.oz. 16
Salicy	lic acid	gr. 140

Triturate the acid intimately with a small portion of the honey of rose, then add the remainder of the honey.

### Honey of Rose with Tannic Acid.

Honey of roseav.oz	. 16
Tannic acidgr	

Triturate the acid intimately with a small portion of the honey of rose, then add the remainder of the rose honey.

## Hydromel.

Honey															
Water.						•	٠	•	•			•	fl.oz.	9	

### Infusion of Buchu.

Buchu			 gr.	510
Distilled	water,	boiling	 .fl.oz.	16

Mix, let stand in a covered vessel for 2 hours, then strain, and pass enough water through the strainer to make 16 fluidounces. -U. S. P., 1870.

11.										
I	Buch	u.						gr	. 30	30
I	Disti	lled	water	, bo	oili	ng		fl.oz		16
1	Mix.	let	stand	in	a	covere	d ves	ssel	for	half

## Infusion of Columbo.

	 	 -	_	 -
Ψ.				
I.				

Columbo, cut small.....av.oz.  $\frac{1}{2}$  Distilled water, cold ......fl.oz. 16

Macerate for 2 hours and strain.—U. S. P. 1870.

H.

Columbo, cut smallgr.	360
Distilled water, coldfl.oz.	

Mix, let stand for half an hour, and strain.

—Brit. Pharm.

## Infusion of Gentian, Compound.

Gentiangr.	230
Bitter orange peel,	
Coriander, of eachgr.	57
Alcohol,	
Water, of each suffice	ient

Extract the drugs in the form of a coarse powder, by percolation, using as a menstruum a mixture of alcohol and water in the proportion of 1 part by volume of the former to 7 of the latter. The product should measure 16 fluidounces.—U. S. P., 1870.

## Infusion of Golden Seal, Compound.

Golden sealgr. 240
Blue cohoshgr. 240
Witch hazel barkgr. 240
Alum gr. 60
Water, boilingfl.oz. 16
Honeysufficient

Infuse the 3 drugs with the boiling water in the usual way, strain, add the alum, dissolve, and then add sufficient honey to sweeten. Used as wash for various forms of sore mouth, and as a gargle.—Eclectic.

### Infusion of Matico.

Maticogr.	360
Distilled water, boiling fl.oz.	

Mix, let stand in a covered vessel for half an hour, and strain.—Brit. Pharm.

## Infusion of Pink Root, Compound.

(Worm	Tea.)
ink root,	bruisedav.oz.

 Senna, cut.
 gr. 144

 Fennel, bruised
 gr. 144

 Manna
 av.oz.

 Water, boiling
 fl.oz.

 16
 16

Mix the above, let stand until cold, and decant the clear liquid.

Dose, 1 to 5 fluidounces.

## Infusion of Sage.

Sage .	 	 	 	gr.	240
Water,					

Mix, let stand in a covered vessel for half an hour, and strain.—U. S. P., 1870.

## Infusion of Sage, Compound.

Sagegr.	240
Hyssopgr.	240
Boraxgr.	30
Water, boilingfl.oz.	

Infuse the drugs in the usual way with the water, strain, and dissolve the borax in the colature. Used as a mouth wash and gargle.

—Eclectic.

## Infusion of Senna.

Sennagr.	480
Coriander, bruisedgr.	60
Distilled water, boiling fl.oz.	16

Mix, let stand in a covered vessel for 1 hour, and strain.—U. S. P., 1870.

## Infusion of Tar. (Tar Water.)

Pine ta	r	 	 	 	.fl.oz.	4
Water						

Mix, shake frequently during 24 hours, decant the supernatant liquid, and filter.— U. S. P., 1870.

## Inhalation of Bromine, Netolitsky's.

Bromine	 	 	.gr.	36
Potassium bromide .				
Water		 	fl.oz.	153/

Dissolve the potassium bromide in a very small portion of the water, then carefully counterbalance the vessel containing the solution on a small balance, then place the proper weights on the opposite pan of the balance, and then drop the bromine from a dropper into the solution until equilibrium of the balance is again restored; now add the remainder of the water.

To inhale, place a small portion of the liquid on a sponge. This is used in cases of croup.

## Injection of Apomorphine, Hypodermic.

Apomorphine hydrochlorate....gr. 2 Camphor water......m. 100

Dissolve and filter. This solution should be prepared only as wanted for use.—Brit. Pharm.

## Injection of Curare, Hypodermic.

Curare ......gr. 5
Distilled water .....sufficient

Reduce the curare to powder in such a way as to prevent it from coming in contact with the naked hand or any other portion of the body, add distilled water to form a thin paste, transfer to a small funnel plugged with absorbent cotton, and gradually pour upon it distilled water until 1 fluidram is obtained.—Brit. Form.

## Injection of Ergotin, Hypodermic.

Dissolve by stirring together.—Brit. Pharm.

## Injection of Morphine, Hypodermic.

Morphine acetate.....gr. 12
Distilled water.....sufficient

Dissolve the morphine salt in 1½ fluidrams of water, and filter, adding through the filter enough water to make 2 fluidrams. If the morphine salt does not completely dissolve, add a drop or two of diluted acetic acid to the liquid before filtering.—Brit. Pharm.

#### Jelly, Currant. (Currant Paste.)

Take fully ripe currants, either red or black, whichever may be wanted; put them into a preserving pan, bruise them and place them on the fire, stirring constantly with a wooden spatula until they become scalding hot. When reduced to a pulp, remove them from the fire and strain and express all the juice from them through a flannel filtering bag or a crash towel. Measure the juice into a very clean and bright copper basin, or, still better, a porcelain-lined basin; place upon the fire and boil for 10 or 15 minutes in order to evaporate some of the water; remove the scum, and add one pound of sugar for each pint of juice. Stir constantly with a wooden spatula until the sugar is dissolved; then remove the scum and immediately fill the jelly glasses, or other suitable vessels.

This is used in making throat lozenges of certain kinds, such as have been used in the London Throat Hospital under the direction of Dr. Morell Mackenzie.

Juice, Elder. (Succus Sambuci. — Roob Sambuci. — Syrupus Sambuci. — Elder Berry Syrup.)

Take any desired quantity of freshly gathered elder berries, heat, with constant stirring, until they burst open, then express through flannel; evaporate the juice to a rather thick extract, and add to this powdered sugar in the proportion of 1 part by weight to every 9 parts by weight of the extract.—Austr. Pharm.

All application of heat must be over a water bath.

## Juice, Huckleberry.

Huckleberriesav.oz.	16
Waterfl.oz.	8
Sugarav.oz.	11/2

Heat the berries in a porcelain or enameled iron evaporating dish, on a water bath, for 1 hour, and express. Heat the residue for an hour with the water, express again, mix the two liquids, add the sugar and heat until dissolved. Strain through a fine cloth and then evaporate to thick extract. This is used sometimes, in domestic practice, in the treatment of the diarrhœa of children.—D.

Juice, Juniper. (Roob Juniperi.—Succus Juniperi.—Juniper Berry Syrup.—
Syrupus Juniperi.)

Juniper berries, fresh, bruised av.oz. 8 Water, hot.....fl.oz. 32

Mix, stir frequently during 12 hours, express, and evaporate the liquid to a thin extract.—Germ. Pharm.

In the absence of fresh berries this preparation may be made from the ordinary dried berries, by the use of a larger proportion of water and a longer period of maceration.

#### Juice, Lemon, Artificial.

I. Citric acidav.oz.	11/2
Potassium carbonategr.	41
Waterfl.oz.	141/2

Mix the acid and potassium with the water, add the sugar when effervescence, then add the peel of a lemon; let stand for 24 hours, and strain.

II.

Citric acidgr. 5	525
Distilled waterfl.oz.	14
Oil of lemon,drops	10
Alcoholfl.oz.	
Dissolve the acid in the water and tl	ne oil

in the alcohol, mix the two solutions and filter.

—H.

#### Kali, Lemon.

Tartaric acidgr. 4	95
Sodium bicarbonateav.oz.	11/2
Sugarav.oz.	51/4
Spirit of lemon sufficient to flav	or

## Kneipp's Remedies, Pastor.

Cough Tea (Hustenthee.)—Coltsfoot leaves, 20 parts; nettle leaves, 10 parts; equisetum, 10 parts; fennel, 5 parts; juniper berries, 5 parts; snake plantain, 5 parts; mallow flowers, 5 parts; linden blossoms, 5 parts; mullein flowers, 2.5 parts; fenugreek, 2.5 parts.

Felon Oil (Malefizoel). — Croton oil, 1 part; oil of sweet almonds, 6 parts.

Blood Purifying Tea (Blutreinigungthee).—Elder flowers, 10 parts; elder leaves, 10 parts; dwarf alder root, 10 parts; sandalwood, 10 parts; buckthorn bark, 10 parts; mistletoe, 10 parts; sloe blossoms, 5 parts, strawberry leaves, 5 parts; nettle leaves, 5 parts; juniper tops, 2½ parts.

Stomach Consoler (Magentrost). — St. John'swort leaves and flowers, 3 parts; millefoil, 1 part; juniper berries, 1 part; dog rose, 1 part; gentian root, 1 part; wormwood, ½ part; buckbean, ½ part; equisetum, ½ part; eye-bright, ½ part; little centaury, ½ part; peppermint oil, 1-10 part; alcohol, 60 p. c., 100 parts.

Wuehlhuberthee.—Aloes, 8 parts; fenugreek, 8 parts; fennel, 25 parts; juniper berries, 25 parts.

Eye-Bright (Augentrost).—Extract aloes, 1-5 part; fennel, 10 parts; eye-bright, 10 parts; alcohol, 20 parts; water, 80 parts.

Dropsy Tea (Wassersuchtsthee).—Equisetum, 40 parts; dog rose, 20 parts; rosemary, 10 parts; elder root, 10 parts; sassafras, 10 parts; rue, 5 parts; buckbean, 5 parts; uva ursi, 5 parts; mistletoe, 5 parts; sandalwood, 5 parts; juniper berries, 5 parts.

Kneipp's Pills.—Rhubarb, 40 grams; extract aloes, 40 grams; soap, 10 grams; juniper berries, 3 grams; fenugreek, 3 grams; dwarf alder, 3 grams; fennel, 3 grams. Make into 600 pills.

#### Lard.

The adipose tissue adhering to the kidneys, mesentery and omentum of the hog is considered the source of the best lard. This is freed from all flesh, then cut into small pieces, removing, as far as possible, all bloody matter and the membranous tissue. Then heat in a tinned copper, porcelain, or enameled iron dish on a water bath until the fat is about all melted, and then strain.

## Lard, Anhydrous.

Lard may be made perfectly anhydrous by heating the preceding on a water bath for about 30 minutes with about one-twelfth its weight of anhydrous or dry sodium sulphate in very fine powder, then filtering through paper, using some system of hot filtration to maintain the lard in a liquid state.

Lard prepared in this manner remains "sweet" much longer than the preceding.—D.

## Lard, Balsamic.

Lard, freshav.oz.	16
Tolu balsamav.oz.	11/2
Etherfl.oz.	1
Sodium sulphate, anhydrousav.oz.	

Melt the lard and add it to the tolu dissolved in ether and the sodium sulphate, heat the mixture on a water bath for 1 hour, stir constantly, and finally filter by hot filtration.—D.

#### Lard, Benzoated.

	acidgr. 7	
Lard	av.oz. 1	6

Melt the lard on a water bath and dissolve the acid in it.—Germ. Pharm.

This is different from the U.S. P. benzoinated lard, which is made by inclosing coarsely powdered benzoin in a piece of muslin, suspending in melted lard, and heating for 2 hours to a temperature not above 60 degrees C.

#### Lards, Factitious.

Quite a number of fats of animals, commonly called "greases," are demanded of pharmacists. Some of these lards or fats cannot be obtained, or at least with great difficulty, and as a rule they are prepared artificially. The formulas given herewith

PHARMACEUTICAL	L PREPARATIONS. 109
will be found acceptable for preparing these	Liniment of Amber Oil, Compound.
" greases."  Goose Grease or Oil.  Olive oil	Rectified oil of amber       fl.oz. 3½         Oil of stillingia       fl.oz. 3½         Oil of lobelia       fl.dr. 10         Olive oil       fl.oz. 7         — Eclectic
Dog's Fat or Grease.	Liniment of Ammonia, Compound.  (Granville's Counter-Irritant.)  Stronger water of ammonia fl.oz. 10 Spirit of camphor fl.oz. 4 Spirit of rosemary fl.oz. 2  Liniment, Anodyne.  Chloral gr. 120 Camphor gr. 120 Ether fl.dr. 2 Chloroform fl.dr. 2 Tincture of opium fl.dr. 1 Oil of sassafras fl.dr. 1 Soap liniment, enough to make fl.oz. 16 Dissolve and mix.
Beef suet	Liniment, Arnica.
Bear's Grease.  Olive oil	Tincture of arnica fl.oz. 4 Soap liniment fl.oz. 12  Liniment, Black.  Olive oil fl.oz. 6 Sulphuric acid fl.dr. 4 Oil of turpentine fl.dr. 2
form.	Add the acid very gradually, and with con-
Other lards may be found under the head of "Oils," and "Suets."	stant stirring, to the olive oil, allow to cool,
Laudanum, Dutchman's.	and add the oil of turpentine.—Eclectic.
Passion flowers	Liniment of Camphor, Compound. (Compound Tincture of Camphor.) (Rheumatic Liniment, Tincture or Drops.)
Liniment, A B C.	Camphor
Liniment of aconite.       fl.oz.       5½         Liniment of belladonna       fl.oz.       5½         Chloroform       fl.oz.       2¾         Camphor       gr. 288         Glycerin       fl.oz.       2¼	Oil of hemlock       fl.oz.       1         Oil of sassafras       fl.dr.       2         Oil of cajeput       fl.dr.       2         Oil of turpentine       fl.dr.       1         Powdered capsicum       av.oz       ½         Alcohol       fl.oz.       15
Liniment of Aconite.	Mix, macerate for 7 days, and filter in a
Fluid extract of aconitefl.oz. 10 Camphorgr. 240 Alcohol, enough to makefl.oz. 16 Mix the fluid extract and the alcohol, and	well-covered funnel.—Eclectic.  The British pharmacopœia also recognizes a "compound liniment of camphor," which is directed to be prepared as follows:
in this mixture dissolve the camphor.—Brit. Pharm. modified.	Camphor
Liniment of Aconite, Homeopathic.	Stronger water of ammonia fl.oz. 334 Alcohol fl.oz. 11½
Tincture of aconite, U. S. Pfl.dr. 4 Alcoholfl.oz. 8	Dissolve the camphor and oil in the alcohol and add the ammonia gradually to this solution with constant shaking.

Liniment of Camphor. (Camphorated	Liniment of Mustard.
Oil.) Camphor, coarse powderav.oz. 6½ Cottonseed oilfl.oz. 28	Volatile oil of mustard fl.dr. 2 Cottonseed oil
Introduce the camphor and oil in a suitable	Liniment of Oils.
flask, apply a gentle heat by means of a	Oil of cedarfl.oz. 4
water bath, loosely stoppering the flask dur-	Oil of cajeput
ing the operation, and agitate from time to	Oil of sassafras
time till the contents of the flask are dis-	—Eclectic.
solved.—U. S. P.	Linimont of Oils Compound
Liniment, Cantharides.	Liniment of Oils, Compound.
Cantharides, powderav.oz. 21/4 Oil of turpentinesufficient	Oil of origanum
Digest the cantharides with 16 fluidounces of oil in a closed vessel on a water bath for	Camphor
3 hours, then strain and add enough oil	Mix, macerate for 7 days, agitating occa-
through the strainer to make the colature	sionally, and filter in a well-covered funnel.—
measure 16 fluidounces.—U. S. P. 1880 and	Eclectic.
N. F. Liniment of Capsicum, Compound.	Liniment of Opium. (Anodyne Lini-
Tincture of capsicumfl.oz. 10	ment.)
Tincture of opiumfl.oz. 2	Tincture of opium
Ammonia water	—Brit. Pharm.
Oil of cinnamonfl.dr. 5	
Spirit of camphorfl.dr. 5	Liniment of Opium, Ammoniated.
—Eclectic.	Soap liuiment
Liniment of Chloroform, Compound.	Tincture of opiumfl.oz. 43/4
Chloroform         fl.oz. 4           Oil of turpentine         fl.oz. 4	Belladonna liniment
Soap liniment	
Liniment of Croton Oil.	Mix and filter.—Brit. Form.  Liniment of Petroleum, Compound.
Croton oil	Crude petroleumfl.oz. 12
Oil of turpentine	Ammonia water
Liniment of Cajeput, Compound.	Tincture of opium
Oil of cajeputfl.dr. 4½	Mix and dissolve.—Eclectic.
Oil of spearmint       fl.dr. 4½         Tincture of opium       fl.dr. 13         Soap liniment       fl.oz. 13½	Liniment of Stillingia, Compound.
Liniment, Diuretic.	Oil of stillingiafl.oz. 1
Soap linimentfl.oz. 2	Oil of cajeput
Tincture of digitalisfl.oz. 2	Alcoholfl.oz. 2
Tincture of squillfl.oz. 2	—Eclectic.
Liniment, Hungarian.	Liniment, Turpentine, Acid. (Brodie's
Cantharides, powdergr. 60	Liniment.)
Garlic, slicedgr. 60	Sulphuric acid fl.dr. 1
Camphor	Olive oil
Black peppergr. 240	Oil of turpentinefl.oz. 1
Diluted acetic acidfl.oz. 6	Add the acid gradually to the olive oil,
Alcohol	stirring in a mortar; when the mixture is

cold, add the oil of turpentine.

Mix all, macerate for 7 days and filter.

PHARMACEUTICAL
Liniment, Turpentine, Camphorated.
Oil of turpentine
Other liniments are mentioned in Parts II and V.
Liniment, Thymol.
Thymol       av.oz.       3/4         Spirit of soap       fl.oz.       133/4         Glycerin       fl.oz.       13/4
Dissolve the thymol in the spirit, add the glycerin, and filter.—D.
Lotion of Borax.
I. Abercrombie's: Boraxgr. 300 Diluted acetic acid
II. Copeland's:
Boraxgr. 80
Rose waterfl.oz. 4
Orange flower water fl.oz. 4
III. Johnson's: Boraxgr. 160
Chalk, precipitatedav.oz. 1 <sup>1</sup> / <sub>4</sub>
Rose waterfl.oz. 4
Alcoholfl.oz. 4
IV. Meig's: Borax
Borax av.oz. 1/2 Morphine sulphate gr. 6
Rose water
Lotion, Bromine, Glover's.
Bromine
Lotion, Capsicum, Griffith's.
Tincture of capsicumfl.oz. 4
Spirit of camphorfl.oz. 4
Water of ammoniafl.oz. 2
Lotion of Ether, Compound.
(Evaporating Lotion.)
Ether
Lotion of Lead Chloride, Tuson's.
Chloride of leadgr. 48 Water, boilingfl.oz. 16
Lotion of Lobelia, Compound.
(Herpetic Wash.)
Bayberry bark       gr. 120         Lobelia herb       gr. 120         Lobelia seed       gr. 120         Yellow dock       gr. 120
Diluted acetic acidsufficient  Extract the mixed drugs in moderately
which the miner diago in moderately

fine powder with the acid by percolation so as to obtain 16 fluidounces of product.— Eclectic.

## Lotion of Myrrh, Compound.

Myrrh, in coarse powder gr. 1	120
Zinc acetategr.	45
Lead acetategr.	15
Watersuffici	ent

Pour 12 fluidounces of boiling water on the myrrh, let stand for 1 hour, stirring frequently; strain, add the salts dissolved in 4 fluidounces of water, and then enough water to make 16 fluidounces of product, and strain again, if necessary.—Eclectic.

Used as a collyrium.

## Lotion, Red. (Red Wash.)

Zinc sulphategr.	40
Compound tincture of lavender, fl.dr.	
Water fl.oz.	16
Cochineal, coloring, N. Fsufficie	ent

## Lotion of Sodium Chlorate, Darling's.

Sodium	chlorategr.	300
Water.	fl.oz.	8

## Lotion of Sulphur, Compound.

(Taylor's Lotion.)

Sulphur, sublimedgr.	360
Borax, powderav.oz.	
Spirit of camphorfl.dr.	
Glycerin fl.dr.	
Water	
*** CLC	1.0

Dissolve the borax in the water, add the spirit and then incorporate the mixture with the sulphur previously triturated to a smooth paste with the glycerin.

#### Lotion of Tin Chloride, Nouche's.

Tin chloride gr.	8
Waterfl.oz.	16

### Lotion, White. (White Wash.)

Sulphurated potassa (sulphuret	
of potash)gr.	60
Zinc sulphategr.	
Waterfl.oz.	

Dissolve each in 2 fluidounces of water and mix the two solutions.

#### Marrow.

Take fresh marrow and heat on a water bath until quite thoroughly melted, then strain with expression, heat the liquid with a small amount of anhydrous sodium sulphate for about 30 minutes, stirring frequently, and filter by hot filtration.—D.

Extract the mixed drugs in moderately A factitious article may be prepared by

melting 1 part of oil of theobroma and adding 2 parts of fresh lard.-H.

## Milk of Magnesia.

Magnesia, calcined, lightgr.	<b>51</b> 0
Glycerinfl.oz.	31/2
Waterfl.oz.	11 1/2

Triturate the magnesia with the water, adding it gradually, and then add the glycerin.-D.

It may also be prepared from magnesium sulphate and alkali solution as follows:

Magnesium sulphate, crystalparts 6	
Waterparts 20	
Solution of potassasufficient	

Dissolve, filter, heat to the boiling point, then gradually add, under constant stirring, solution of potassa enough to produce an alkaline reaction. Transfer the precipitate to a filter and wash thoroughly with hot water. Then, to the washed magma add sufficient cold water, previously deprived of air by boiling, to make the whole weigh 20 parts. One hundred parts contain 5 parts of anhydrous magnesia. It may also be prepared by the use of solution of soda. The alkali solution should be freshly prepared from pure materials.

## Mixture of Bloodroot, Compound.

Syrup of ipecac	.fl.oz. 1
Syrup of squill	.fl.oz. 1
Syrup of tolu	.fl.oz. 1
Tincture of bloodroot	
Paregoric	
9	Eclectic.

#### Mixture, A E. (Chloroformum Mitigatum.)

Alcohol	 	1 by volume
		2 by volume
Ether	 	3 by volume

## Mixture, Castor Oil.

Castor oil	fl.oz.	3
Oil of lemon	m.	40
Oil of cloves	drops	8
Simple syrup		
Solution of potash	fl.dr.	41/2
Orange flower water, enough		, -
	fl 07	8

Mix the oils in a mortar, then incorporate one-third of the solution of potash and afterward the syrup, then an additional third of the solution of potash, then gradually onehalf of the orange flower water, the remain- when there is threatened collapse.

der of the solution of potash, and lastly sufficient solution of potash to produce the required volume.

## Mixture, Cajeput, Compound.

(Hunnis Drops.)

Oil of cajeput	fl.oz.	1
Oil of cloves	fl.oz.	1
Oil of peppermint	fl.oz.	1
Oil of anise	fl.oz.	1
Alcohol	fl.oz.	4

This has been employed in Eclectic practice in the treatment of diarrhœa, cholera, etc.

## Mixture of Camphor, Compound.

Camphor water	fl.oz.	5
Peppermint water	. fl.oz.	5
Spearmint water		
Paregoric		
	_Eclect	

## Mixtures, Cholera or Diarrhœa.

I. Christensen's:

4
1
2
1
10

## II. Ebert's:

Solution of iron nitrate	.fl.dr.	2
Deodorized tincture of opium		
Caraway water	.fl.dr.	4

Dose, from 1/2 to 1 teaspoonful after each evacuation.

#### III. Greenhow's:

Guaiacgr.	240
Clovesgr.	240
Cinnamongr.	240
Brandyfl.oz.	16

Macerate the drugs in moderately fine powder with the brandy for 7 days, and filter.-Eclectic.

The dose is from a teaspoonful to a tablespoonful, in sweetened water, every 15 or 20 minutes until relief is obtained.

#### IV. Hamlin's (1):

Tincture of	opiumfl.oz.	1
Tincture of	rĥubarbfl.oz.	1
Spirit of ca	mphorfl.oz.	1

## V. Hamlin's (2):

Tincture of opium	fl.oz. 1	l
Tincture of ginger	fl.oz. 1	L
Tincture of capsicum	fl.oz. 1	ŀ
Tincture of cardamom	ı fl. oz. 1	L

This is used in the second stage of cholera

VI. Harney's (Gen'l.):	with the Mistura Thielemanni of the Swedish
Chloroform	Pharmacopœia.
Tincture of opiumfl.oz. 1	
Spirit of cinnamon fl. oz. 1	XV. Velpeau's:
Spirit of peppermintfl.oz. 2	Tincture of opiumfl.oz. 2 Compound tincture of catechufl.oz. 2
Camphor, powder gr. 16 Syrup of ginger 2	Spirit of camphorfl.oz. 2
Mix and dissolve.	Mixture, Copper, Rademacher's.
VII. Loomis':	
Tincture of opiumfl.dr. 4	Tincture of copper acetate, (Rademacher's)gr. 60
Tincture of rhubarbfl.dr. 4	Mucilage of acaciafl.dr. 4
Compound tincture of catechu. fl.oz. 1 Oil of sassafras	Cinnamon waterfl.oz. 4½
Compound tincture of lavender	Water fl.oz. 23/4
enough to makefl.oz. 4	Mixture, Iron, Rademacher's.
VIII. Rubini's:	Tincture of iron acetate, Rade-
Camphorav.oz. 1	macher's
Hoffmann's anodynefl.oz. 2	Mucilage of acacia
Dose: 2 to 5 drops on sugar every 20	Mixture of Soda.
minutes until relieved.	Fluid extract of rhubarbfl.dr, 2
IX. Rademacher's:	Tincture of cinnamonfl.dr. 4
Zinc acetategr. 90	Brandyfl.dr. 2
Distilled water	Sodium bicarbonategr. 60 Oil of clovesdrops 2
X. Ruschenberger's:	Oil of cloves
Tiucture of opiumfl.oz. 1	Water
Spirit of camphorfl.oz. 1	Mixture, Starton's.
Tincture of capsicumfl.oz. 1	200
Spirit of peppermint	Magnesium sulphategr. 360 Ferrous sulphategr. 60
Aromatic tincture of rhubarbfl.oz. 1 Dose: 30 to 40 drops in water.	Diluted sulphuric acidfl.dr. 2
•	Syrup of wild cherryfl.oz. 1
XI. Scammon's:	Water, enough to makefl.oz. 4
Tincture of opium	This is given in teaspoonful doses to be
Tincture of capsicumfl.dr. 2	taken after meals.
Diluted alcoholfl.dr. 1	Mixture, Tonic, Compound.
XII. Squibb's:	(Mistura Alterantiæ Composita.)
Tincture of opiumfl.oz. 1	Ferrous sulphategr. 40
Tincture of capsicum fl.oz. 1 Spirit of camphor	Sodium phosphate
Chloroformfl.dr. 3	Quinine (alkaloid)gr. 64
Alcoholfl.dr. 13	Strychnine (alkaloid)gr. 4 Sugarav.oz. 10
XIII. Sun:	Diluted phosphoric acid fl.oz. 9½
Tincture of opiumfl.oz. 1	Distilled water sufficient
Tincture of capsicum	Dissolve the sulphate of iron in 6 fluid-
Spirit of camphor fl.oz. 1	drams of boiling distilled water, also dis-
Spirit of peppermintfl.oz. 1	solve the sodium phosphate in 11/2 fluidounces
Mix them, and filter.	of boiling distilled water, and mix the 2
XIV. Thielemann's:	solutions; collect the precipitate and wash it
Wine of opiumfl.oz. 1	with distilled water until the washings are
Tincture of valerian fl. oz. 1½ Ether fl. dr. 4	tasteless; add this precipitate together with
Oil of peppermint	the quinine and strychnine to the diluted
Fluid extract of ipecac	phosphoric acid, shake until dissolved, add
Alcohol	the sugar, dissolve without heat and strain.
This preparation is practically identical	—Eclectic.

## Mucilage of Irish Moss.

Irish moss......gr. 192
Water, enough to make .....fl.oz. 16

Wash the moss with cold water, then place in a suitable vessel, add 16 fluidounces of water, and heat on a water bath for 15 minutes, frequently strain through muslin, and add enough water through the strainer to make the colature measure 16 fluidounces.—
N. F.

## Mucilage of Linseed.

Linseed,	wholeav.oz.	1
Distilled	water, hotfl.oz.	5

Mix, macerate for 6 hours, stirring frequently, and strain.—D.

## Mucilage of Quince Seed.

(Mucilage of Cydonium.)

Quince seed, whole ......gr. 144
Distilled water ......fl.oz. 16

Mix, macerate for one-half hour, stirring frequently, and strain without expression.—
N. F.

## Mucilage of Starch.

Starch				 			gr.	192
Distilled water			 			.fl.	oz.	16

Triturate the starch with water gradually added until a smooth paste is formed, then boil for a few minutes, constantly stirring.—Brit. Pharm.

Other mucilages will be noticed in Part VII.

## Oil, Angleworm.

Angleworms, freed from adher-
ent dirtpart 1
Olive oilparts 2

Macerate for 3 days in a warm place, then strain or filter.

#### Oil, Anodyne.

Ammonia water	.fl.dr.	31/2
Tincture of opium		
Oleo-balsamic mixture		
Alcohol	.fl.dr.	12
Infused oil of henbane, enough	1	
to make	.fl.oz.	16

## Oil, Baunscheidt. (Compound Oil of Euphorbium.

Baunscheidt was a German charlatan who claimed to cure rheumatic and other diseases by means of what he called a "lebens-

wecker," i. e., "awakener" or "revulseur." This consists of a number of sharp-pointed needles set in a bed of hard rubber. By means of a spiral-spring arrangement these needles are driven into the skin over the seat of pain, not deep enough to draw blood, while into the wounds produced is rubbed the "Baunscheidt oil." This is an irritating substance and produces papular eruptions similar to those produced by croton oil. The effect is that of a powerful counter-irritant. In medicine this treatment is known as acupuncture. Formulas for the oil are as follows:

Euphorbium, powdergr.	160
Cantharides, powdergr.	96
Olive oilfl.oz.	8
Macerate for 7 days and filter.—H.	

II.	
Euphorbium, powdergr	r. 140
Mezereum, cut finegr	r. 280
Cantharides, powdergr	r. 30
Alcoholfl.oz	z. 1
Etherfl.oz	
Olive oilfl.oz	z. 8´~

Mix the two powders, alcohol and ether in a closed vessel, macerate for 7 days, agitating occasionally; then add the oil, macerate again for 7 days, strain, heat the colature gently so as to expel the ether, and filter.—H.

### Oil of Belladonna, Infused.

Prepare from belladonna leaves by the same process as infused oil of chamomile is prepared from chamomile.

### Oil, Carminative. (Colic Oil.)

Oil of spearmintfl.dr. 5
Oil of carawayfl.dr. 2½
Oil of cumin
Oil of fennelfl.dr. 2½
Infused oil of chamomile,
enough to makefl.oz. 16
-Norw. Pharm.

This is what is desired by Scandinavians when green oil is asked for, but Germans understand by this title the infused oil of henbane.

#### Oil of Chamomile, Infused.

Chamomile, coarse powderav.oz.	3
Alcoholfl.oz.	21/2
Ammonia waterdrops	
Olive oilfl.oz.	16

Mix the alcohol and water, and incorporate

quickly with the chamomile, set aside in a closed vessel for 24 hours, then add the oil, heat for 12 hours at a temperature of 50 to 60 degs. C. and strain with expression.

#### Oil, Chloroform.

Chloroformfl.oz.	33/4
Olive oil, enough to makefl.oz.	

### Oil, Cod Liver, Aromatized.

Oil of lemon drops	40
Oil of nerolidrops	8
Oil of peppermintdrops	8
Vanillingr.	1
Coumarin gr.	10
Cod-liver oilfl.oz.	16

Dissolve the coumarin and vanillin in the volatile oils, with the aid of a very gentle heat, and mix the solution with the cod liver oil.—D.

## Oil, Cod Liver, with Iodine. (Iodized Cod Liver Oil.)

Iodinegr.	
Chloroformm.	15
Cod liver oil	16

Triturate the iodine with a few drops of oil, then add the remainder of the oil and the chloroform, transfer the whole to a bottle, and agitate frequently until dissolved.

## Oil, Cod Liver, with Iron. (Ferrated Cod Liver Oil.)

T

٠								
	Iron	benzoate.	 	 	 	 .gr.	68	
		liver oil						

Triturate the iron salt with the oil and warm gently until the former is dissolved. The product contains 1 per cent of the iron salt.

II.

Castile soapgr.	75
Dialyzed ironfl.oz.	
Distilled watersufficie	
Sodium chlorideav.oz.	1
Cod liver oil fl.oz.	

Dissolve the soap in 10 fluidounces of water by the aid of heat, also mix the iron solution with 9 fluidounces of water, add the latter liquid gradually to the soap solution (when cold), stirring constantly. Collect the precipitate without washing; place between folds of filter paper until tolerably dry, place in a porcelain or enameled iron evaporating dish with the salt and iron, heat on a water bath until solution takes place, and filter. The object of the salt is to abstract water

from the iron compound and thus facilitate its solution in the oil.—D.

This contains about ½ per cent of ferric oleate.

#### III.

Ferric chloride,	sublimedgr.	195
Cod liver oil	fl.oz.	16

Triturate in a mortar until the iron salt is dissolved, and, if necessary, filter.

## Oil, Cod Liver, with Iron and Iodine.

(Ferro-Iodized Cod Liver Oil.)

Iodinegr.	
Clean iron wiregr.	
Cod liver oil, enough to makefl.oz.	16

Triturate the iodine, iron and 6 fluidounces of oil in a mortar with some ether until a black mixture results and the iodine and iron have combined; then add the remainder of the oil and filter.

The product contains ½ per cent of ferrous iodide.

## Oil, Cod Liver, with Malt Extract.

For cod liver oil with malt extract, refer to Extract of Malt with Cod Liver Oil.

## Oil, Cod Liver, with Phosphorus.

(Phosphorized Cod Liver Oil.)

This may be made by dissolving 4 grains of phosphorus in 16 fluidounces of cod liver oil by the aid of a gentle heat.

#### Oil of Conium, Infused.

This is to be prepared from conium herb by the same process as infused oil of chamomile is prepared from chamomile.

## Oil, Eel, Factitious.

Oil of sweet almonds fl.oz. 8
Castor oil
Cod liver oil fl.dr. 6
—H.

#### Oil of Eggs.

This is obtained by subjecting hard-boiled yolk of eggs to pressure. About 1 fluid-ounce of oil is thus obtained from 16 yolks. The oil deteriorates very readily and must be preserved in dram-vials in a cool, dark place.

A factitious article may be prepared as follows:

Olive oil fl.oz.	121/2
Cacao butterav.oz.	
Yellow waxav.oz.	1/2

Melt together on a water bath.—H.

It may often be replaced by olive, sweet Ointment, Aconitine. almond or other similar bland fixed oil.

### Oil of Henbane, Infused.

This is to be prepared from henbane leaves by the same process as infused oil of chamomile is prepared from chamomile.

#### Oil of Lilies.

Oil of bergamot	.drops 1	6
Cottonseed oil, bleached	fl.oz. 1	6

## Oil, Rainworm, Artificial.

Oil of birch tar	fl.dr. 5
Benne oil, enough to make f	1.0z 16

## Oil of Rhodium, Factitious.

1.	
Oil of rosefl.dr.	1
Oil of copaibafl.oz.	21/2
II.	,-
Oil of rosedrops	10
Oil of sweet almondfl.dr.	
Balsam of copaibafl.dr.	4
III.	
Oil of rosefl.dr.	1
Oil of sandalwood fl.oz.	21/2

## Oil, Skunk.

Lard oil	fl.oz. 8
Lard	av.oz. 8
Animal oil	drops 5

## Oil of Spike. (British Oil.-Black Oil.) I.

Oil of amber, crudefl.dr.	10
Oil of turpentinefl.oz.	5
Linseed oilfl.oz.	
American petroleumfl.oz.	5
II.	
Olive oil fl.oz.	14

## Oil of Stramonium, Infused.

This is to be prepared from stramonium leaves by the same process as infused oil of chamomile is prepared from chamomile.

Oil of vitriol.....fl.oz. 2

### Oil of Wormwood, Infused.

This is to be prepared from wormwood by the same process as infused oil of chamomile is prepared from chamomile.

### Ointment of Aconite.

Extract of aconite	.av.oz. 1
Simple ointment	.av.oz. 2

Soften the extract with water or diluted alcohol and mix well with the ointment.-Eclectic.

Aconitinegr.	8
Alcohol drops	32
Lard, benzoinatedav.oz.	1

Dissolve the aconitine in the alcohol and add the lard.-Brit, Pharm.

#### Ointment Alkaline.

Potassium carbonategr.	120
Tincture of opiumfl.dr.	1
Simple ointmentav.oz.	1

Triturate the potassium salt to an impalpable powder, mix with the ointment, and then add the tincture.-Eclectic.

## Ointment, Alkaline, Camphorated.

Potassium carbonategr.	
Camphor, powder gr.	6
Lardgr.	420

Warm the lard, add the camphor, stir well till dissolved, and mix the potassium carbonate. - Eclectic.

## Ointment, Ammoniacal.

Lardgr.	180
Suetgr.	120
Oil of sweet almondfl.dr.	1
Stronger water of ammonia (28	
per cent or concentrated)fl.dr.	6

Melt the lard and suet together, add the oil, and when the mixture is tolerably cool, pour it into a wide-mouth bottle, add the ammonia water, cork well, and shake occasionally until cold.—Eclectic.

## Ointment, Antimonial. (Tartar Emetic Ointment.)

Tartar emeticgr.	100
Lardgr.	
—U. S. P. 18	370.

#### Ointment, Astringent, Thompson's.

Mutton suet	.av.oz.	8
Witch hazel or sumach leaves	.av.oz.	1

Melt the suet, boil with the leaves for onehalf hour and stráin.

## Ointment, Bayberry. (Green Salve.)

Bayberry	plaster			 		.av.oz.	4
Olive oil.							1

Melt the plaster, add the oil, and stir until cool.-Eclectic.

This may also be prepared from:

Bayberry wax	av.oz. 2
Gum turpentine	av.oz. 2
Olive oil	fl 07 1

## Ointment, Atropine.

Atropinegr.	8
Alcoholdrops:	3;2
Lard, benzoinated av.oz.	

Dissolve the atropine in the alcohol and add the lard.—Brit. Pharm.

#### Ointment of Boric Acid.

#### I. Lister's formula is as follows:

Boric acid, fine powdergr.	240
White waxgr.	240
Paraffin gr.	480
Oil of sweet almond fl. oz.	

Triturate the acid to a smooth paste with a portion of the oil, melt the paraffin, wax, and remainder of the oil together, and add the previous mixture.

#### II.

Petrolatum, white
Mix thoroughly.—Germ. Pharm.

#### IJ.

Boric acid.					 				.gr.	75
Petrolatum	٠.	•			 			٠	.gr.	450

## Mix intimately.—Brit. Pharm.

## Ointment, Calamine, Rademacher's.

Camphor, powder	gr.110
Lithargeav.o	z. 2
Armenian boleav.o	
Lead carbonateav.o	z. 2
Calamine, preparedav.o	z. 2
Yellow waxav.o	
Lardav.c	

Mix the litharge, lead carbonate, bole and calamine to a smooth paste with a portion of the lard, also melt the wax, to it add the remainder of the lard, add the previous mixture, mix the whole thoroughly, add the camphor and stir occasionally until solid.

## Ointment, Casein, Unna's.

Casein av.oz.	21/
Potassium hydrategr.	$23^{\circ}$
Sodium hydrategr.	5
Glycerinfl.oz.	1
Petrolatum gr. 1	150
Zinc oxidegr.	36
Carbolic acidgr.	
Water, enough to makeav.oz.	16

Prepare the casein as follows: Take milk from which the cream or fat has been entirely removed, curdle it by the addition of rennet extract, at a temperature of 30 to 40 degs. C.; collect the coagulum and wash with

running water or otherwise until the washings are no longer acid; dry carefully and powder.

Dissolve the alkalies in a portion of the water, and in this solution dissolve the casein; add the glycerin and carbolic acid, incorporate the petrolatum and zinc oxide, and finally the remainder of the water.

This is recommended by Unna as a vehicle to be used instead of fatty bodies for inunction.

#### Ointment of Conium.

Extract of coniumgr	. 55
Simple ointmentav.oz	. 1

Soften the extract with water or diluted aicohol and add the ointment.—Eclectic.

## Ointment of Copper Subacetate.

Yellow waxav.oz.	21/4
Lead plasterav.oz.	31/4
Rosinav.oz.	1/2
Olive oilfl.oz.	81/2
Copper subacetate (verdigris)av.oz.	1/2
Olibanum, finest powderav.oz.	11/2

Melt the wax, lead plaster and rosin together and add 7½ fluidounces of the oil. Intimately mix the verdigris with the remaining oil to a smooth paste, add this mixture to the previous one, stir well, add the olibanum, and stir frequently until cool.

Another ointment containing verdigris and known as green ointment is the following:

Verdigris,	powdered	 			.av.oz.	1
Resin cera	te				.av.oz.	15

Add the powder to the ointment, previously melted at a gentle heat, and stir the mixture frequently until it concretes.

### Ointment, Creosote.

Creosote	fl.dr. 1
Lard	av.oz. 2
	-Eclectic.

#### Ointment of Gallic Acid.

Gallic	acid									.av	oz.	1
Benzo	inated	lard								.av	oz.	9
					_	_T	T	-	3	P	188	n

#### Ointment of White Hellebore.

White hellebore (veratrum al-	
bum), powderav.oz.	1
Lardav.oz.	4
Oil of lemondrops	10

Mix well.-Eclectic.

## Ointment of Iodine, Rademacher's.

Iodine	r. 90
Alcohol suffi	cient
Lard	oz. 4

Dissolve the iodine, in a little alcohol and add lard. The product contains about 4 per cent of iodine.

## Ointment of Ipecac.

Ipecac, fine powder	av.oz. 1
Olive oil	, fl.oz. 1
Lard	

Mix well.-Eclectic.

## Ointment, Lead, Hebra's. (Hebra's Diachylon Ointment.)

Lead plaster									av.oz.	1
Olive oil									.fl.oz.	1

Mix at a gentle heat. Prepare only as needed.

According to the United States pharmacopeia lead ointment is scented with oil of lavender flowers. Sometimes the above ointment is made with linseed oil instead of olive oil.

## Ointment of Lead Carbonate, Camphorated.

Camphor, fine powder	gr.	23
Lead ointment	av.oz.	1

Mix the camphor thoroughly with a portion of the ointment and then add the remainder of the ointment.—Germ. Pharm.

## Ointment, Marshmallow. (Althæa Ointment.—Yellow Ointment.)

Turmeric, powdergr.	130
Lardav.oz.	141/2
Yellow waxgr.	420
Resingr.	420

Digest the turmeric in the lard for half an hour over a water-bath, then add the beeswax and the resin previously melted together, melt the whole together, and strain the ointment.—Germ. Pharm.

## Ointment, Lead, Compound. (Mayer's

Olive oilfl.oz.	7
Gum turpentineav.oz.	11/2
Yellow waxav.oz.	3/4
Butter, unsaltedav.oz.	3/4
Red leadav.oz.	3
Honeyav.oz.	2
Camphor, powderav.oz.	11/2

Melt the wax and turpentine together, add | well mixed. - Germ. Pharm.

the butter and oil, heat nearly to boiling, and add gradually, with constant stirring, the red lead; allow to cool, and when nearly cold add the honey and camphor, stirring until the latter is dissolved.—Eclectic.

## Ointment of Red Iodide of Mercury.

Red iodide of mercurygr. 60	0
Simple ointmentav.oz.	
-Rrit Pharm	

## Ointment, Mezereon.

Lardav.oz.	3
Yellow waxgr.	200
Fluid extract of mezereumfl.oz.	

Melt the lard and wax together over a water bath, add the fluid extract and continue heating until all the alcohol has dissipated, meanwhile stirring constantly; then remove the ointment from the source of heat and stir frequently until cool.—U. S. P. 1880.

## Ointment of Black Pepper.

Lard	.av.oz. 4
Suet	.av.oz. 1
Pine tar	
Black pepper, fine powder	.av.oz. 1

Melt the lard and tar together, then add the suet, and finally the pepper; stir frequently until cool.—Eclectic.

### Ointment of Poke.

Extract of	pokegr.	60
	mentav.oz	

Soften the extract with a little water or diluted alcohol and add the ointment.— Eclectic.

## Ointment of Potassium Sulphide.

Sulphurated potash	gr.	55
Sodium carbonate	gr.	55
Lard	av.oz.	1

Triturate the two salts to impalpable powder and then add the lard.—Eclectic.

### Ointment, Rosemary, Compound.

(Unguentum Nervinum. — Aromatic Ointment.)

Yellow waxav.oz. 1
Mutton suet
Lardav.oz. 8
Expressed oil of nutmegav.oz. 1
Oil of rosemaryfl.dr. 4
Oil of juniper berries fl.dr. 4

Melt the wax and suet, add the lard and nutmeg oil, stir till melted, allow to cool, then add the two volatile oils, and stir unti. well mixed.—Germ. Pharm.

Contuse the shepherd's purse to a pulp, add it to the melted lard, and carefully heat the mixture over a direct flame until the moisture has all evaporated.

## Ointment, Sulphur, Alkaline.

Sulphur, washedgr.	560
Potassium carbonategr.	280
Water	
Benzoinated lard av.oz.	4

water, add the other ingredients, and mix well.-U. S. P. 1880.

## Ointment of Sulphur, Compound.

Sulphurav.oz. 1	
White hellebore (Veratrum al-	
bum), fine powdergr. 72	
Potassium nitrate, powderedgr. 10	
Soft soapav.oz. 1	
Ointment of pokeav.oz. 3	
Oil of bergamotdrops 20	
Mix well.—Eclectic.	

## Ointment of Sulphur Iodide.

I.	
Iodide of sulphur gr. 3	0
Benzoinated lardav.oz.	1
Mix well.	
TT	

Petrolatum		av.oz. 1
	-	Brit, Pharm.
Ointment,	Thompson's.	

Sulphur iodide.....gr. 30

~~~~	,	
Yellow	waxav.oz.	4
	av.oz.	
Gum tur	pentineav.oz.	6
Balsam	of firav.oz.	3

#### Ointment of Tobacco.

Extract of tobaccogr.	48
Simple ointmentav.oz.	4

Soften the extract with water or diluted alcohol and add the ointment, - Eclectic.

## Ointment of Wild Indigo.

Fluid extract of wild indigofl.oz.	5
Butterav.oz.	
Beeswax av.oz.	1
Tallow av.oz.	1/2

Melt the fats together, add the fluid ex- Triturate the morphine to fine powder,

tract, and continue heating carefully ur.til all the alcohol and water have evaporated, meanwhile stirring frequently; allow to cool, stirring from time to time till nearly solid .-Eclectic.

Instead of the fluid extract of wild indigo, a corresponding amount of "solid" extract may be used; this should be softened with water or diluted alcohol after which the fats in melted state should be incorporated with it.

### Ointment of Witch Hazel.

Fluid	extract of	witch	hazel		
Simple	ointment			gr.	410
			Deit	Dho	****

## Dissolve the potassium carbonate in the Ointment, Zinc, Compound. (Wilson's Ointment.)

Zinc oxideav.oz.	4
Benzoin, fine powderav.oz.	
Lardav.oz.	15

Rub the zinc oxide and benzoin together, add to the lard, heat the whole on a water bath for one hour, stirring constantly; then strain, and allow to cool with frequent stirring.

## Ointment of Zinc Oxide, Compound.

_		
	Olive oilav.oz.	4
	Spermacetiav.oz.	2
	White waxgr.	320
	Zinc oxideav.oz.	
	Benzoic acidgr.	20
	Morphine sulphategr.	8
	Oil of rosedrops	3

Triturate the zinc oxide, benzoic acid, morphine sulphate and olive oil to a smooth paste, add this to a melted mixture of the wax and spermaceti, stir almost constantly till cool, and then add the oil of rose .-Eclectic.

## Ointment of Zinc Sulphate.

-		-	
	Zinc su	lphategr.	70
	Simple	ointmentgr.	420

Rub the zinc sulphate to an impalpable powder and add the ointment.—Eclectic.

The original contained fresh butter instead of the ointment and the former may be used if desired.

Other ointments will be noted in Parts II and V.

#### Oleate of Morphine.

Morphine	(alkaloid	)	gr.	87
Oleic acid			fl.oz.	4

add to the oleic acid, and dissolve by the aid of a gentle heat.

The above makes a preparation containing 5 per cent of alkaloid. If stronger preparations are desired, relatively greater proportions of alkaloid must be used. A 10 per cent oleate, for instance, would require the use of 175 grains of morphine.

## Oleate of Strychnine.

Strychnine (alkaloid)gr.	35
Oleic acidfl.oz.	4

Triturate together, dissolving by the aid of a gentle heat, if necessary.

The product contains 2 per cent of strychnine.

## Oxymel.

Acetic acidfl.oz. 1
Water
Clarified honeyav.oz. 8
-Brit. Pharm.

## Paste, Carbolic, Lister's.

Carbolic acid, crystal av.oz.	43/
Olive oilfl.oz.	10
Prepared chalksufficie	nt

Dissolve the acid in the oil and add sufficient chalk to make a soft paste.—D.

#### Paste, Lassar's.

Salicylic acidgr.	72
Zinc oxideav.oz.	1
Starchav.oz.	
Petrolatumav.oz.	

## Paste, Wax.

Yellow waxav.		
Cocoanut oilav.	oz.	11/4
Lanolin		
Borax, powder	gr.	70
Distilled waterfl.	oz.	91/2

Melt the wax, add the oil, and then the lanolin, and after allowing to cool somewhat incorporate the borax dissolved in the water. This is used as an ointment vehicle.—D.

## Physic, White Liquid. (Dow's Physic.)

Sodium sulphateav	oz. 4
Waterfl	
Nitric acidfl	.dr. 4
Hydrochloric acid fl	.dr. 4
Alum	

Dissolve the sodium sulphate in the water, filter, and add the other ingredients.— Eclectic.

## Pills, Anti-Canker, Thompsonian.

Each pin should contain:	
Iron subcarbonategr.	21/2

## Pills, Anti-Catarrhal, Andrews'.

(Anti-Grippe or Yellow Pills.)

Each pill should contain:

Quinine salicylategr.	1
Arsenious acidgr.	T25
Extract of belladonna leavesgr.	4

## Pills, Anti-Constipation, Carson's.

Each pill should conta
------------------------

Extract of cascara sagrada gr.	1
Extract of rhubarbgr.	1
Extract of nux vomica gr.	
Extract of aloingr.	1/4

## Pills, Anti-Constipation, Fothergill's. Each pill should contain:

Strychnine		 .gr.	24
Aloes		 .gr.	11/
Black pepper			
Extract of cascara sagrada.	•	 or	12/
Dittiact of cascara sagrada,		 . 5	-/3

## Pills, Arsenic, Hebra's.

Each pin should contain:	
Arsenious acidgr.	$\frac{2}{25}$
Extract of licorice, dried gr.	3/4
Licorice wood, powdergr.	3/4
Mucilage of acacia sufficient to form ma	ass.
Roll in lycopodium.—D.	

## Pills of Camphor, Compound.

(Cholera Pill.)
Each pill should contain:

Camphorgr. 1	
Opium gr. 1	
Kinogr. 1	
Capsicumgr.	1
Confection of rose, sufficient to form	5
a mass	

### Pills, Cook's.

Each pill should contain:		
Rhubarb	gr.	1
Aloes	gr.	1
Calomel	.gr	3
Soap, powdered	.gr	1
Date 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		

## Pills, Gout, Lartigue's.

Each pill should contain:

-				
Extract of c	olocynth, o	compound.	gr.	4
Extract of	digitalis		gr.	è
Extract of o	olchicum r	pot	gr.	2

## Pills, Tonic, Aiken's.

Each pin should contain.	
Quinine sulphategr.	1
Arsenious acidgr.	30
Reduced irongr	. 2/3
Strychninegr	

Pills of Zinc	Acetate,	Rademacher's.
Each pill sho	uld contain	:

Zinc acetate.										.gr.	3
Licorice root										.gr.	1

## Plaster, Bayberry. (Green Plaster.)

Gum turpentine							.av.oz.	4
Bayberry wax							.av.oz.	4

Melt together, strain, and stir until cool.— Eclectic.

## Plaster, Logan's.

Lead oxideav.oz.	
Lead carbonateav.oz.	
Soapav.oz.	11/2
Butter, freshgr.	240
Olive oilfl.oz.	
Masticgr.	20

Mix the soap, oil and butter, add the lead oxide, and boil the whole gently for an hour and a half or until it has acquired a pale brown color, stirring constantly meanwhile; the heat may then be increased and the boiling continued till a portion of the melted plaster dropped on a smooth board is found not to adhere; then remove the vessel from the fire and add the mastic to the mixture.

## Plaster, Menthol.

I.	
Lead plasterav.oz.	71/2
Yellow waxav.oz.	1
Resin av.oz.	1/2
Mentholav.oz.	1

Melt the plaster, wax and resin, strain and as the mixture cools, add the menthol. Spread the composition on cloth or leather.—D.

II.		
Menthol Yellow wax	av.oz.	31/4
Yellow wax		11/2
Resin	av.oz.	111/

Melt the resin and wax together, and as the mixture cools, stir in the menthol until dissolved.—Brit. Pharm.

The first plaster contains 10 per cent of menthol, the second 20 per cent.

## Plaster, Miraculous, Rademacher's.

Red leadav.oz.	4
Olive oilav.oz.	9
Amber, powder gr.	96
Camphor, powdergr.	60
Burnt alumgr.	30

Boil the lead and oil until a black plaster

is formed, allow this to cool somewhat, add the other ingredients and mix well.

## Plaster, Resin, Compound. (Adhesive and Strengthening Plaster.)

Resin av.oz.	9
Beeswaxav.oz.	1
Camphor gr.	60
Oil of hemlockfl.dr.	2
Oil of sassafrasfl.dr.	2
Oil of turpentinefl.dr.	1
Olive oilfl.dr.	2

Melt the resin and wax together, allow to cool, and while still liquid add the camphor dissolved in the oils.—Eclectic.

## Plaster, Thapsia.

Yellow wax av.oz.	61/2
Resinav.oz.	21/2
Gum turpentineav.oz.	21/2
Venice turpentineav.oz.	
Glycerinfl.dr.	5
Honeyfl.dr.	5
Resin of thapsiagr. 5	510

Melt the first three ingredients and add the others, and stir until a plaster mass is formed.—Codex.

## Pomade, Juniper.

Lardav.oz.	61/2
Paraffin waxgr.	240
White waxgr.	60
Oil of juniper berries fl.dr.	3
Fowler's solutionfl.dr.	

Melt the two waxes, then add the lard, remove vessel from the fire, add the other ingredients, and stir well until cold.—Eclectic.

## Powder, Antimonial, Tyson's.

	Antimony oxideav.oz. Calcium phosphateav.oz.	
	II.	Ü
1	Antimony oxideav.oz.	1
	Calcium phosphateav.oz.	41/2

## Powder, Arsenical, Come's.

Red sulphide of mercury av.	oz. 6
Arsenious acidav.	oz. 2
Dragon's blood	gr. 260
Charcoal, animal	gr. 175

Triturate together to a very fine powder.

Powder,	Black.	(Emmenagogue Powder.)	)
---------	--------	-----------------------	---

Sulphurav.oz. 1
Myrrhav.oz. 1
Steel filingsav.oz. 1
Loaf sugarav.oz. 1
White winefl.oz. 4

Mix, and, by aid of a gentle heat, evaporate till nearly dry; pulverize the mass when cold.—Eclectic.

This is the formula as originally given. The substance is best dispensed in pill form.

## Powder of Camphor, Compound.

Tannic acid	 	gr. 120
Kino	 	gr. 120
Camphor		
Opium		

Each should be in fine powder and the whole should be well mixed.—Eclectic.

## Powder of Charcoal, Compound.

Wood charcoalav.oz. 4
Rhubarbav.oz. 2
Sodium bicarbonateav.oz. 1

All should be in fine powder and the whole well mixed.—Eclectic.

### Powder, Cough, Thompsonian.

Lobelia herbav.oz. 2
Licorice rootav.oz. 2
Skunk cabbageav.oz. 2
Sugarav.oz. 2

All should be in fine powder and be well mixed.

#### Powder of Cypripedium, Compound.

Refer to Compound Powder of Ladies' Slipper.

### Powder, Dover's, Camphorated.

(Beach's Diaphoretic Powder.)

Opiumgr.	55
Ipecac	
Camphorgr.	220
Potassium bitartrate av.oz.	2

All should be in powder and should be well mixed.—Eclectic.

### Powders, Effervescent.

See Effervescent Salts.

## **Powder, Effervescing.** (Pulvis Aerophorus.)

Sodium bicarbonateav.oz.	21/2
Tartaric acid	21/4
Sugarav.oz.	43/4

All should be in fine powder and perfectly

dry, and should then be well mixed.—Germ. Pharm.

Preserve in well-stoppered bottles in a dry place.

## Powder, Effervescing, with Magnesia.

L.		
	Tartaric acidav.oz.	1
	Oil-sugar of lemonav.oz.	2
	Sugar, moderately fine powderav.oz.	3
	Magnesium carbonateav.oz.	4
	—Germ. Forn	n.
ſ١	T	

II.	
Tartaric acid	.av.oz. 1
Citric acid	.av.oz. 2
Sugar	av.oz. 3
Magnesium carbonate	.av.oz. 4
9	

Prepare and preserve like the preceding.

—D.

## Powder of Glycyrrhizin, Aromatic.

Ammoniated glycyrrhizingr.	120
Oil of carawaydrop	
Oil of coriander drop	
Oil of anisedrops	
Oil of cassiadrops	

Adapted for disguising taste of quinine in powder.

## Powder of Golden Seal, Compound.

Golden seal		.av.oz.	2
Blue cohosh		.av.oz.	2
Helonias		.av.oz.	2
Reduce to fine powder	r.—Eclect	ic.	

## Powder, Gun. (Pulvis Pyrius.—Pulvis

•	
Potassium nitratev.oz.	15
Potassium nitrateav.oz. Charcoalav.oz.	3
Sulphur	2

When gunpowder is required for veterinary recipes, it may be prepared extemporaneously according to this formula.

## Powder, Gun, White.

Potassium ferrocyanide	av.oz.	1
Sugar	av.oz.	1
Potassium chlorate	av.oz.	S

Reduce each to powder separately, then mix well, but very carefully, using a bone or wooden spatula.

## Powder of Ladies' Slipper, Compound. (Nerve Powder.)

Reduce to fine powder. - Eclectic.

Powder of Ipecac, Compound.	Powder of Pleurisy Root, Compound.
Ipecac	Pleurisy root
Powder, Laxative, Gregory's.	Podophyllingr. 10
Ginger	Cream of tartargr. 450  Mix well.—Eclectic.  Powder of Podophyllum, Compound.
Magnesium carbonateav.oz. 8  Each should be in fine powder, and the whole should be mixed intimately.  Dose, one-half to one teaspoonful.  Powder, Nephritic, Rademacher's.	Blue flag av.oz. 2 Podophyllum av.oz. 2 Dogsbane (or bitter root) av.oz. 2 Swamp milkweed (Asclepias incarnata) av.oz. 2 Blood root av.oz. 1
Cochineal, powder	Reduce to fine powder.—Eclectic.
Powder, Nerve, Thompsonian.	Powder of Quinine, Compound.  Ouinine sulphateav.oz. 1
This is powdered cypripedium or ladies' slipper root.	Iron ferrocyanide
Powder of Opium, Compound.	Powder of Rhubarb, Compound.
Opium.       av.oz. 3/4 (gr. 328)         Black pepper.       av.oz. 1         Ginger.       av.oz. 2½         Caraway.       av.oz. 3         Tragacanth.       gr. 110	(Neutralizing Powder.)  Rhubarb
All should be in powder and should be	whole should be well mixed.—Eclectic.
well mixed.—Brit. Pharm.	Powder of Senna, Compound.
	-
Powder of Phosphates, Compound.  Sodium phosphate	Senna powder
Sodium phosphate	Senna powder
Sodium phosphate	Senna powder

All should be in fine powder and the whole about one hundred years ago in Germany.

weli mixed.—Eclectic.

He entertained peculiar ideas regarding the

practice of medicine, as well as the preparation and action of drugs. Many of his peculiar preparations are prescribed in this country by the older German physicians, and more recently some of them have been employed by physicians partial to the eclectic school.

This class of preparations includes the following:

- 1. Drops, Dysmenorrhæa.
- 2. Extract of Tobacco.
- 3. Mixture, Copper.
- 4. Mixture, Diarrhœa.
- 5. Mixture, Iron.
- 6. Ointment, Calamine.
- 7. Ointment, Iodine.
- 8. Ointment, Shepherd's Purse.
- 9. Pills, Zinc Acetate.
- 10. Plaster, Miraculous.
- 11. Powder, Nephritic.
- 12. Solution of Calcium Chloride.
- 13. Solution of Sodium Nitrate.
- 14. Solution, Anodyne Turpentine.
- 15. Tincture of Celandine.
- 16. Tincture of Cochineal.
- 17. Tincture of Colocynth.
- 18, Tincture of Copper Acetate.
- 19. Tincture of Golden Rod.
- 20. Tincture of Hips.
- 21. Tincture of Iron Acetate.
- 22 Tincture of Mary Thistle.
- 23. Tincture of Mugwort Root.
- 24. Tincture of Nux Vomica.
- 25. Tincture of Shepherd's Purse.
- 26. Water, Acorn.
- 27. Water, Castor.
- 28. Water, Nux Vomica.
- 29. Water, Quassia.
- 30. Water, Tobacco.

Formulas for making these preparations may be found elsewhere in Part I.

#### Resinoids.

Refer for these to Concentrations.

### Salts, Effervescent.

The effervescent salts or powders, for which formulas are here given, are most conveniently and efficiently dispensed in the form of fine powders, because in this condition they can be made extemporaneously and with an assurance of their freshness and efficiency. The popular demand, however, seems to be

for granular effervescent salts, the preparation of which requires certain modifications of the formulas, important only in so far as they enable the dispenser to granulate the powder in a convenient and expeditious manner.

Effervescent powders or salts are composed of the medicinal agent in admixture with an alkaline bicarbonate, an organic acid, and sugar. The proportion of the medicinal agent is dependent upon its dose, that of the alkaline bicarbonate and of the organic acid is dependent upon their molecular relation to each other, while the proportion of sugar is dependent upon the quantity necessary as a sweetening agent and diluent.

The ingredients for making the fine pulverulent form of effervescent powders are: The medicinal agent, sodium bicarbonate, tartaric acid and sugar, and it is necessary that these be well dried before mixing them.

To make the granular form of effervescent salts the ingredients need not be dried, unless specially directed, and the ingredients are the same as for the pulverulent form, with the single exception that one-half the molecule of tartaric acid is replaced by one-half a molecule of powdered citric acid.

In order to facilitate the manufacture of effervescent salts, the new National Formulary recognizes three new preparations, viz.: saccharated citric acid, saccharated tartaric acid, and saccharated sodium bicarbonate.

Saccharated citric acid is prepared by mixing 5 av. ounces of citric acid with 3 av. ounces of sugar, each to be in very fine powder.

Saccharated tartaric acid is prepared by mixing  $6\frac{3}{4}$  av. ounces of tartaric acid with  $3\frac{1}{4}$  av. ounces of sugar, each to be in very fine powder.

Saccharated sodium bicarbonate is prepared by mixing 6 av. ounces of sodium bicarbonate with 2 av. ounces of sugar, each to be in very fine powder.

The ingredients of these preparations should be intimately mixed, and should be preserved in well-stoppered bottles.

The proportion of sugar in these saccharates is so adjusted that when either of the acid saccharates is mixed with an equal weight

of the alkaline saccharate, the acid and alkali are in molecular relation to each other, and, when dissolved in water, will form the neutral tartrate and citrate of sodium respectively.

With these three saccharates in stock, it becomes possible to make effervescent powders or salts quickly with any medicinal agent that may be prescribed, their use being exemplified by the following general formulas:

#### I. Fine Powder:

Medicinal agent, fine powderav.oz.	2
Saccharated sodium bicarbon-	
ateav.oz.	19
Saccharated tartaric acidav.oz.	

Triturate the ingredients until a uniformly mixed powder is obtained. In order to insure permanence of the product, the ingredients should be well dried before mixing.

## II. Granular Salt:

Medicinal agent, fine powder av.oz.	2
Saccharated sodium bicarbon-	
ateav.oz.	19
Saccharated tartaric acid av oz	91/

Saccharated citric acid.....av.oz. 91/2

Mix the ingredients (not dried) in a mortar, transfer them to an evaporating dish, and heat on a water bath, kept at 60 to 71 degrees C., under constant stirring with a wooden spatula, until dry and uniformly granular.

The saccharated citric acid, being made from crystallized citric acid containing one molecule of water of crystallization, supplies the moisture necessary to cause the powder, when heated, to cake and adhere together. If the somewhat pasty mass is then stirred with the spatula, small granules are readily formed. and these become firm when completely dried.

Another method of making the granular effervescent salts is to add to either of the above mixtures just enough alcohol to form a pasty mass, then rubbing this through a No. 20 porcelain colander, and drying the product in a drying room at a gentle heat. alcohol used should be just sufficient to form a pasty mass, as an excess would make the drying require a longer period of time, and at the same time would prove a serious waste.

cent salts is recognized by the U.S.P., the heat method by the N. F.

It is, of course, not always that the proportion of the ingredients is the same as in the above general formulas. There may, for example, be more or less of the medicinal agent, and it may be necessary to add an extra amount of sugar. It is not necessary, either, to use any of the saccharates mentioned, the alkaline bicarbonate, acid, sugar and medicinal agent or agents.

Throughout the process of making these "salts," contact with metals should be carefully avoided. The mixtures should always be preserved in well-stoppered, wide-mouthed bottles.

The following formulas for effervescent salts are taken from standard works of reference:

### Caffeine, Citrated.

Caffeinegr.	110
Citric acidgr.	110
Sodium bicarbonateav.oz.	81
Tartaric acidav.oz.	71/2
Tartaric acidav.oz. Sugarav.oz.	834

To be granulated (according to U. S. P.) by the alcohol process. It may also be made according to the N. F., or heat, process. may also be dispensed in powder form.

### Carlsbad Salt, Artificial,

Artificial Carlsbad saltav.oz.	41/
Saccharated sodium bicarbon-	
ateav.oz.	$10_{\frac{1}{4}}$
Saccharated tartaric acid av. oz.	101

Mix the ingredients, previously well dried, and triturate them until a uniform powder is obtained. If desired in the granular form, substitute saccharated citric acid, not dried, for one-half of the saccharated tartaric acid. and prepare the granulated compound as directed under the general formula.-N. F.

A solution of about 87 gr. of this preparation in 6 fluidounces of water represents an equal volume of Carlsbad water (Sprudel) in its essential constituents.

#### Iron Citrate.

Iron pyrophosphate, solubleav.oz.	2
Citric acidav.oz.	
Sodium bicarbonateav.oz.	5
Sugarav.oz.	10

Triturate the iron salt to powder, add the other ingredients, mix thoroughly in a The alcohol method of making efferves- porcelain mortar, and gently heat the whole

(with mortar) on a water bath, triturating constantly until a moist mass is formed, which agglutinates sufficiently so it may be passed through a sieve. Then dry and bottle the product in the usual manner.-Germ. Form.

Iron cit	rate, solubleav.oz.	1
Sodium	bicarbonateav.oz.	10
Tartario	acidav.oz.	7
Citric a	eid av.oz.	1
Sugar.	av.oz.	8
Alcohol	fl.oz.	7

Reduce the iron salt to very fine powder, add the other solids in fine powder, mix well, warm slightly in an evaporating dish, moisten with the alcohol, and sift dry, and bottle in the usual manner. - D.

Properly speaking, the first "iron citrate" is not a citrate, but a pyrophosphate.

## Iron Citrate with Magnesia; or iron and magnesium citrate.

Iron citrate, solubleav.oz.	1
Magnesium carbonategr.	220
Sodium carbonateav.oz.	10
Tartaric acidav.oz.	8
Citric acid,av.oz.	11/2
Sugarav.oz.	8
Alcoholfl.oz.	7

Prepare like the preceding.—D.

#### Iron and Quinine Citrate.

Soluble citrate of iron and quinine.gr.	97
Saccharated sodium bicarbon-	
ateav.oz.	11
Saccharated tartaric acidav.oz.	

Mix the ingredients, previously well dried, and triturate them until a uniform powder is obtained. If desired in the granular form, saccharated citric acid may be substituted for one-half of the tartaric acid, then follow the rules of the general formula.-N. F.

Ninety gr. (or about a heaped teaspoonful) of this preparation represent about 1 gr. of citrate of iron and quinine.

#### Iron Phosphate.

Iron phosphate, soluble, very fine powdergr.	215
Saccharated sodium bicarbon-	
ateav.oz,	
Saccharated tartaric acidav.oz.	10

Mix the ingredients, previously well dried, and triturate them until a uniform powder is obtained. If desired in the granular form, follow the rules of the general formula, substituting saccharated citric acid for one-half ate, and the remainder of the citric acid,

the saccharated tartaric acid, heating, etc.-

Ninety gr. (about a heaped teaspoonful) represent about 2 gr. of iron phosphate.

## Iron Pyrophosphate.

See Iron Citrate above.

## Kissingen Salt, Artificial.

Artificial Kissingen	saltav.oz. 7
Saccharated sodium	bicarbonate av. oz. 9
Saccharated tartaric	acidav.oz. 9

Mix the ingredients, previously well dried, and triturate them until a uniform powder is obtained. If desired in the granular form, follow the general formula by substituting saccharated citric acid for one-half the saccharated tartaric acid, heating, etc.-N. F.

A solution of about 80 gr. of this preparation in 6 fluidounces of water represents an equal volume of Kissingen water (Rackoczi springs) in its essential constituents.

#### Lithium Carbonate.

Lithium carbonateav.oz. %	9
Sodium bicarbonateav.oz. 6	
Tartaric acidav.oz. 4	Ŀ
Sugarav.oz. 8	3
Alcohol. fl.oz. 9	•

Mix the solids, and moisten and sift in the regulation manner. Dry first at 20 degrees C., then increase the temperature to 40 degrees C., until perfectly dry.-D.

#### Lithium Citrate.

Lithium citrateav.oz.	2
Sodium bicarbonateav.oz.	6
Tartaric acidav.oz.	4
Milk sugarav.oz.	4
Sugarav.oz.	
Alcoholfl.oz.	9

Prepare like the preceding.—D.

## Magnesium Citrate.

Magnesium carbonateav.oz.	21/2
Citric acidav.oz.	11 1/2
Sodium bicarbonateav.oz.	
Sugarav.oz.	2
Alcohol	

Distilled water.....of each, sufficient

Mix the magnesium carbonate with 71/2 av. ounces of citric acid and 1 fluidounce of water, so as to form a thick paste. Dry this at a temperature not exceeding 30 degrees C., and reduce to fine powder. Then mix this intimately with the sugar, sodium bicarbondampen with alcohol, and granulate and dry.
—U. S. P.

This preparation may also be prepared according to either of the N. F. processes, by mixing the powdered magnesium citrate, obtained in this formula, with the saccharates, and heating, if desired.

## Magnesium Sulphate.

Magnesium sulphate, crystalav.oz.	
Sodium bicarbonateav.oz.	18
Tartaric acidav.oz.	91/2
Citric acidav.oz.	$6\frac{1}{4}$
Sugarav.oz.	5‡

Dry the magnesium salt at a temperature of about 55 degrees C., until it has lost nearly one-fourth (23 per cent) of its weight; powder the product, and mix with the sugar and other ingredients, all in fine powder. Then granulate the mixture by the heat method.—Brit. Pharm.

## Pepsin.

Pepsin, pure, powdered	gr. 150
Citric acid	.av.oz. 534
Tartaric acid	.av.oz. 5
Sodium bicarbonate	av.oz. 121
Sugar	

Make by the U. S. P. process, or it may be prepared by either of the N. F. processes.

### Pepsin and Bismuth.

Pepsin, pure, powderedgr.	150
Citrate of bismuth and ammo-	
niumgr.	150
Citric acidav.oz.	53/
Tartaric acidav.oz.	41/2
Sodium bicarbonateav.oz.	
Sugargr.	
8	

Prepare like the preceding.

#### Potassium Bromide.

Potassium bromide, very fir	
Saccharated sodium bicarbon	1-
ate	av. oz. 12
Saccharated tartaric acid	av. oz. 12

Mix the ingredients, previously well dried, and triturate them until a uniform powder is obtained. If desired in the granular form, follow the general formula by substituting saccharated citric acid for one-half the saccharated tartaric acid, heating, etc.—N. F.

Ninety gr. (or about a heaped teaspoonful) of this preparation represent about 10 gr. of potassium bromide.

Potassium Bromide, with Caffeine.—N. F. Prepare this like the preceding, adding 131 gr. of caffeine to the above mixture.

#### Potassium Citrate.

Citric acidav.oz.	7
Potassium bicarbonateav.oz.	10
Sugarav.oz.	$-5\frac{1}{4}$

Powder the ingredients separately, and mix them thoroughly in a warm mortar. A pasty mass will be produced which may be granulated as described in the general formula.— U. S. P.

### Sodium Citro-Tartrate.

Sodium bicarbonateav.oz. 8	31/2
Tartaric acidav.oz.	41/2
Citric acidav.oz.	3~~
Sugarav.oz.	

Prepare according to the general formula.

-Brit. Pharm.

## Sodium Phosphate.

Sodium phosphate, crystal av.oz.	121/2
Sodium bicarbonateav.oz.	
Tartaric acidav.oz.	$6\frac{3}{4}$
Citric acidav.oz.	41/2

Dry the sodium phosphate until it has lost 60 per cent of its weight, then powder, add the other ingredients, and complete the process as according to the preceding formula.—Brit. Pharm.

#### Sodium Sulphate.

Sodium sulphate, clear crystals.av.oz.	121/2
Sodium bicarbonateav.oz.	
Tartaric acidav.oz.	634
Citric acidav.oz.	41/2

Dry the sodium sulphate until it has lost rather more than one-half (56 per cent) of its weight, then add the other ingredients, and prepare the salt according to the preceding formula.—Brit. Pharm.

## Vichy Salt, Artificial.

Artificial Vichy saltav.oz.	6
Saccharated sodium bicarbonate.av.oz.	91/2
Saccharated tartaric acid	91%

Mix the ingredients, previously well dried, and triturate them until a uniform powder is obtained. If desired in the granular form, follow the general formula by substituting saccharated citric acid for one-half the saccharated tartaric acid, heating, etc.—N. F.

A solution of about 57 gr. of this preparation in 6 fluidounces of water represents an equal volume of Vichy water (Grande Grille spring) in its essential constituents.

Vichy Salt, Artificial, with Lithiun	n.
Artificial Vichy saltav.oz.	4
Lithium citrate, very fine powder.gr.	650
Saccharated sodium bicarbon-	
ateav.oz.	
Saccharated tartaric acidav.oz.	10

Mix the ingredients, previously well dried, and triturate them until a uniform powder is obtained. If desired in the granular form, follow the general formula by substituting saccharated citric acid for one-half the saccharated tartaric acid, heating, etc.—N. F.

Ninety gr. (or about a heaped teaspoonful) of this preparation represent 14 gr. of artificial Vichy salt, and 5 gr. of lithium citrate.

## Salt of Lemon, Artificial.

Oxalic acid				.av.oz.	4
Potassium carbonate				.av.oz.	2
Cream of tartar				.av.oz.	6

## Salts, Mineral Water, Artificial.

See Part IV.

## Salt, Sea, Artificial. (Sal Marinum.)

I.		
Sodium chlorideav.o		
Magnesium chlorideav.o	z.	11
Calcium chlorideav.o	z.	2
Potassium bromide	r.	130
Potassium iodideg	r.	88
Magnesium sulphateav.o	z.	61/
	_	-D.
II.		

I.		
Sodium chloride	av.oz.	80
Magnesium sulphate	av.oz.	16
Calcium chloride	av.oz.	$3\frac{1}{4}$
Potassium iodide		
Potassium bromide	gr.	35
	_	H.

## Silk, Carbolated.

#### I. Lister's:

White waxgr. 44	
Carbolic acid, crystalav.oz. 1	
Silk thread, strong (not dyed)sufficient	

Mix the acid and wax by fusion, place into the mixture as much of the thread as may be desired and allow it to remain until the mixture is cold. Then wipe off the excess of liquid from the thread by means of a cloth, and then preserve the medicated fiber in a mixture of:

Carbolic acid, crystal	gr. 7	5
Glycerinfl.	.dr.	91/2
Alcoholfl.	oz.	2
	r	`

#### II. Czerny:

Boil strong or thick silk thread (uncolored) in 5 per cent carbolic acid water, for from 10 minutes to  $1\frac{1}{2}$  hours, according to the thickness of the thread. For at the end of every half hour of boiling the water should be renewed. The fiber should be preserved in 2 per cent carbolic acid water.—D.

## Silk, Iodoform, Partsch.

Wind strong silk thread (uncolored) upon a glass spool, or other similar suitable object, and macerate for 2 days in a 10 per cent, solution of iodoform in ether. Then dry the fiber by exposing for a moment to the atmosphere, and preserve in well-closed glass bottles or jars.—D.

## Silk, Sublimated.

I.

Macerate strong uncolored silk thread for 24 hours in a 1 per cent solution of mercuric chloride in distilled water, then preserve in a solution of:

Mercuric	chloride.	 gr. 1
Glycerin.		 fl,dr. 3
Alcohol		 fl.oz. 5
		_D

#### II. Schede-Kuemmell:

Boil strong uncolored silk for 2 hours in a 1 per cent aqueous solution of mercuric chloride, and preserve in a one-tenth aqueous solution of the same agent.—D.

## Snuff, Catarrh.

Morphine hydrochlorate	.gr.	3
Acacia, fine powder		
Bismuth subnitrate	.gr.	540

Mix them intimately by trituration.—N. F.

### Snuff, Cephalic.

Sageav.o	z. 1
Lavender flowersav.o	
Marjoramav.o	
Wild gingerav.o	
White helleboreg	

All should be in fine powder, and be well mixed.

### Snuff, Menthol.

Mentholgr.	45
Sugar of milkav.oz.	
Sugarav.oz.	1/2
Coffee, roastedav.oz.	11/2
Boric acidgr.	90

All should be in fine powder and be well mixed.—H.

## Snuff, Schneeberger.

н	u	٠	

White helleboreav.oz.	2
Orris root,	
Bayberry bark of each, av.oz.	1
Starchav.oz.	12
Oil of cloves drops	20

The first four ingredients should be in the finest powder and the whole should be well mixed.

#### II.

Rice powder	av.oz.	5
Canada snake root	av.oz.	1
White hellebore	av.oz.	3/1
Orris root		
Oil of bergamot		
8	1	

All should be in fine powder and should be well mixed.—H.

## III.

Starch powderav.oz.	5
Orris rootav.oz.	3
Canada snake rootav.oz.	1
White helleboreav.oz.	1
Oleobalsamic mixturedrops	10

All should be in fine powder and should be well mixed.—H. modified.

## Solution of Acid Phosphates. (Compound Solution of Phosphoric Acid.)

I. \_

Bone ash, fine powder	av.oz.	171/2
Sulphuric acid (sp. gr., 1.830)	av.oz.	131/4
Water	.fl.oz.	$64^{'}$

Mix the bone ash with 16 fluidounces of water, add the sulphuric acid, diluted with 32 fluidounces of water, and mix thoroughly with a porcelain or glass stirrer. Now add the remainder of the water and set the mixture aside for 24 hours, stirring occasionally. Then transfer the mixture to a strong muslin strainer, and subject this to a gradual pressure (avoiding contact with metals), so as to express as much of the liquid as possible. Lastly, filter this through paper.—N. F.

#### H.

Calcium carbonate, precipitatedgr.	
Calcined magnesia gr.	116
Potassium carbonategr.	151
Iron phosphategr.	64
Phosphoric acid, 85 per centfl.oz.	31/2
Water, enough to make fl.oz.	16

Mix the acid with 8 fluidounces of water, add the calcium carbonate gradually, with constant stirring; when effervescence has

ceased, add the calcined magnesia in the same way and then add the potassium carbonate; finally add the remainder of the water, allowing the liquid to stand for several days, if possible, and filtering.

## Solution of Aluminium Acetate. (Burow's Solution).

Acetate of leadav.o	z. 16
Potassa alumav.o	
Sulphate of sodaav.c	oz. 1
Water gall-	on 1

Dissolve the alum and soda in half the water; the lead in the other half; mix, filter.

Solution of Aloes and Soda, Met-

tauer's. (Mettauer's Laxative Aperient.)

Macerate for 2 weeks, agitating occasionally, and filter.

#### Solution of Aluminum Chloride.

Aluminum sulphateav.oz. 5
Barium chlorideav.oz. 5
Distilled watersufficient

Dissolve the aluminum salt in 8 fluidounces of hot water, and the barium salt in 10 fluidounces of hot distilled water; mix the solutions, and heat the mixture on a water bath to about 70 or 75 degs. C.; then allow to cool, filter and pass enough water through the filter to make the product weigh 20 av.ounces.—D.

Instead of using barium chloride, calcium chloride may be used. In this case, however, the mixture of the two salts should be set aside in a cold place, for at least one week, in order that the excess of sulphate of calcium, not retained in solution, may separate. And when filtering the solution, it will hardly pay to wash out the small amount retained by the precipitate. Of course, when a solution of a definite strength is required, the first-mentioned formula should be used. But when the liquid is wanted as a disinfectant, the second method may be employed.

### Solution of Ammonia, Anisated.

(Liquor Ammonii Anisatus.)

Oil of anise						
Alcohol Ammonia water				 	.fl.oz.	131/2
Ammonia water	٠.	٠.	٠.	 	.fl.oz.	21/2

-Germ. and Austr. Pharm.

#### Solution of Ammonium Benzoate.

Ammonium carbonate gr.	500
Benzoic acidgr.	1165
Distilled water, enough to	
makefl.oz.	16

Reduce the ammonium salt to powder, mix it and the acid in a capacious vessel, add the water, stir frequently until effervescence has ceased and solution is complete, and filter.

Each fluidram contains 10 gr. of ammonium benzoate.

#### Solution of Ammonium Valerianate.

Ammonium valerianate gr. !	240
Borax, powdergr.	384
Ammonia watersuffici	ent
Distilled water, enough to make fl.oz.	16

Mix the ammonium valerianate with 2 fluidounces of distilled water and add ammonia water, drop by drop, until a clear and slightly alkaline solution is produced; then add 4 fluidounces of water and the borax, stir the whole well, and when all or almost all has dissolved, add enough water to make 16 fluidounces, and filter.

This makes a tasteless and odorless preparation.

## Solution of Annatto. (Extract of Annatto.)

Annattoav.oz. 4	
Potassium carbonate av.oz. 4	
Water sufficient	

Boil the annatto and potassium carbonate with 16 fluidounces of water until the annatto is dissolved, then strain, and add enough water through the strainer to make the colature measure 16 fluidounces.

This preparation is employed for coloring purposes.

## Solution of Antimony, Chloride.

(Butter of Antimony.)

Black	sulphide	of	antimony,	
pure			av.oz.	63/4
Hydro	chloric ac	id	fl.oz.	32

Place the antimony compound in a porcelain or enameled-iron dish, add the acid, apply to the mixture at first a gentle heat which must be gradually increased, as the evolution of gas slackens, until the mixture boils. Continue boiling for 15 minutes, then remove the vessel from the fire, and filter the liquid through calico, returning that which passes through at first until a perfectly clear liquid is obtained. Concentrate this by evaporation to 16 fluidounces, and preserve in a glass-stoppered bottle.—Brit. Pharm.

Owing to the fact that most of the black antimony of the market is very impure, great care should be exercised in the selection of a suitable article. This black antimony should be in fine powder. The boiling of the liquid should either be done under a good flue or in the open air, to avoid tainting the atmosphere of the room with the disagreeable odorous sulphuretted hydrogen gas.

## Solution, Antiseptic, Seiler's.

Sodium bicarbonategr.	240
Boraxgr.	
Sodium benzoategr.	10
Sodium salicylategr.	10
Eucalyptol m.	5
Thymolgr.	5
Mentholgr.	21/2
Oil of wintergreendrops	3′˜
Glycerinfl.oz.	41/4
Alcoholfl.oz.	1
Distilled water, sufficient to	
makepints	8

Dissolve the salt in 64 fluidounces of water by the aid of heat, also the eucalyptol, thymol, menthol, and oil in the alcohol, mix the two solutions, add the glycerin and the remainder of the water, allow to stand for 24 hours, and filter.

## Solution, Antiseptic, Lister's. (Lister's Antiseptic Fluid.)

•	Benzoic acidgr.	64
	Borax	64
	Boric acidgr. 1	128
	Thymolgr.	20
	Oil of eucalyptusdrops	5
	Oil of wintergreen 'drops	5
	Oil of peppermintdrops	3
	Oil of thyme (white)drop	1
	Fluid extract of wild indigo drops	20
	Alcoholfl.oz.	6
	Distilled watersufficie	ent

Dissolve the two acids and borax by the aid of heat in 8 fluidounces of water, also dissolve the thymol and oils in the alcohol, mix the two solutions, agitating frequently during mixing, add the fluid extract, and then enough water to make 16 fluidounces; set aside for 24 hours, and filter through purified talcum.

1.	
Boric acidgr.	128
Thymol gr.	20
Eucalyptoldrops	5
Oil of wintergreendrops	5
Oil of peppermintdrops	3
Oil of thyme, whitedrop	1
Fluid extract of wild indigom.	30
Alcoholfl.oz.	3
Distilled water, sufficient to	
make	16

Dissolve the acid in some of the water, add the other ingredients to the alcohol, dissolve, mix the two solutions, add the remainder of the water, let stand for 24 hours, and filter through purified talcum.

#### III.

Boric acidgr.	128
Thymolgr.	16
Mentholgr.	16
Oil of eucalyptusdrops	4
Oil of wintergreendrops	4
Oil of horsemint drops	4
Water fl.oz.	12
Alcoholfl.oz.	4
Caramel drops 1	or 2

Dissolve the boric acid in the water and the other ingredients in the alcohol and mix the solutions; let stand for a day or two, shaking frequently, and filter.

#### Solution of Borax Comp. (Dobell's.)

Borate and bicarb, of soda, each, gr.	120
Carbolic acid crystgr.	24
Glycerinfl.oz.	1/2
Add water, fl. oz. 16 and mix.	

## Solution of Bromine.

Brominegr.	70
Potassium bromide gr.	
Waterfl.oz.	16

In order to avoid inhaling the intensely acrid bromine vapor, the potassium bromide should be dissolved in a small portion of the water and this solution contained in a bottle be balanced on the "scales," and then the bromine may be dropped into the solution from a medicine dropper; the balance of the water should then be added.

This solution must not be confounded with the N. F. preparation, which is very much stronger. The latter is prepared as follows:

Brominegr. 48	30
Potassium bromide gr. 24	10
Waterfl.oz.	4

water contained in a bottle, add the bromine, and shake the mixture until this is dissolved. Keep the solution in glass-stoppered vials in a dark place.

## Solution of Calcium Chloride, Rademacher's.

Calcium	chloride av.oz.	51/2
Water.		10

Solution of Coal Tar. (Liquor Carbonis Detergens.—Liquor Picis Carbonis.)

۰		
	Coal tarav.oz.	4
	Tincture of quillaja, N. Ffl.oz.	
	Alcohol fl.oz.	8

Digest for 2 days, decant the clear liquid and filter.

#### II.

Quillaja bark, coarse powdergr.	450
Alcoholsuffici	ent
Prepared coal tarav.oz.	2

Extract the quillaja by percolation with the alcohol so as to obtain 10 fluidounces of product; to this add the tar, digest at a temperature of about 50 degs. C. for 2 days, allow to become cold, and decant the clear liquid or filter.-Brit. Pharm.

Prepared coal tar is made by placing commercial tar in a shallow vessel, and heating to a temperature of 50 degs. C. for 1 hour, stirring frequently.-Brit. Form.

## Solution of Cocaine Hydrochlorate.

-	
Cocaine hydrochlorate	gr. 66
Salicylic acid	
Distilled water, enough to	

Boil the water, add the acid, then the cocaine, cool, and add enough water to produce the requisite volume.—Brit. Pharm.

This solution is intended to contain approximately 10 per cent of cocaine hydrochlorate.

## Solution of Gold and Arsenic Bromides.

Arsenous acid	 	gr. 19
Tribromide of gold		
Bromine water,		Ŭ

Distilled water......of each, sufficient

Introduce the arsenous acid and about 18 fluidrams of bromine water into a flask and heat gently until all free bromine has disappeared. Then add bromine water, 20 to 30 Dissolve the potassium bromide in the drops at a time, until it will be present in

slight excess, or until the solution does not become colorless after some time. Transfer the solution to a porcelain capsule, expel the excess of bromine with the aid of gentle heat, dilute it with water to about 14 fluidounces, and dissolve in this the tribromide of gold, adding enough water to make 16 fluidounces.

Ten minims of this solution contain 1-32 grain of gold tribromide and the equivalent of 1-16 grain of arsenic tribromide.—N. F.

#### Solution of Gutta Percha.

(Traumaticin.)

Gutta percha	1
Chloroformfl.oz.	
Lead carbonate, fine powder av.oz.	1

Add the gutta percha to 5 fluidounces of chloroform contained in a closed bottle, and shake occasionally until solution has taken place. Then add the lead carbonate previously mixed with the remainder of the chloroform, and, having several times shaken the whole together at intervals of one-half hour, set the mixture aside until the insoluble matters have subsided and the solution has become perfectly clear. Lastly, decant the clear liquid and preserve in small, cork-stoppered bottles.—U. S. P. 1880 and N. F.

Purified gutta percha only should be employed in making this preparation.

## Solution of Hydrastis, Colorless.

(Glycerite of Hydrastine.)

•	Hydrastine hydrochlorategr.	25	
	Aluminum chloridegr.	23	
	Calcium chloridegr.		
	Magnesium chloridegr.		
	Potassium chloridegr.	1/2	
	Diluted hydrochloric acidm.		
	Distilled waterfl.oz.	6	
	Glycerin, enough to makefl:oz.	16	

Dissolve the salts in the water, add the acid, then the glycerin, and filter.

H.

Hydrastis, fine powder		
Glycerin		
Ether		
Diluted sulphuric acid	.fl.dr.	1
Distilled water	.fl.oz.	8

Exhaust the hydrastis with ether, recover the ether by distillation, to the residue add the water previously mixed with the acid, let stand 7 days, agitating frequently, decant

the aqueous solution, and mix it with the glycerin.

## Solution of Hypophosphites, Acid.

Calcium hypophosphitegr.	384
Sodium hypophosphitegr.	
Potassium hypophosphite gr.	64
Iron phosphate, solublegr.	64
Hypophosphorous acid, diluted fl.oz.	3
Water, enough to makefl.oz.	16

Triturate the hypophosphites to fine powder and dissolve in 12 fluidounces of water; dissolve the iron salt in 1 fluidounce of hot water and add to previous solution; then add the acid, and after standing 24 hours, filter, adding enough water through the filter to make 16 fluidounces of product.

## Solution of Iodine, Caustic, Churchill's.

Iodinegr.	1825
Potassium iodideav.oz. Waterfl.oz.	81/4
Water fl.oz	16

Mix and dissolve.—N. F.

## Solution of Iodine, Caustic, Lugol's.

Iodinegr.	480
Potassium iodidegr.	480
Waterfl.oz.	2

Dissolve the potassium iodide in the water and add the iodine.

This is for application as a caustic.

## Solution of Iodine, Compound or Lugol's.

Iodinegr. Potassium iodidegr.	$\frac{360}{720}$
Distilled water, enough to	120
makefl.oz.	16

Mix and dissolve. Keep the solution in glass-stoppered bottles.—U. S. P.

#### Solution of Iodine, Magendie's.

Iodinegr.	2
Potassium iodidegr.	240
Peppermint waterfl.oz.	$\epsilon$

Dissolve the potassium iodide in the water and add the iodine.

## Solution of Iodine, Rubefacient, Lugol's.

Iodinegr. 2	40
Potassium iodidegr. 4	80
Waterfl.oz.	

Dissolve the potassium iodide in the water and add the iodine.

This is for external use.

## Solution of Iron and Ammonium Citrate.

## Solution of Iron and Quinine Citrate.

Citrate of iron and quinine,

soluble......av.oz 8
Distilled water, enough to make fl.oz. 16

## Solution of Dialyzed Iron.

Dissolve and filter.

Mix 3 fluidounces of the iron chloride solution with 20 fluidounces of water and stir into the mixture sufficient ammonia water to impart a distinct ammoniacal odor. Collect the precipitate on calico or muslin, wash it with distilled water and squeeze the strainer to remove superfluous water. Add the precipitate to the remainder of the solution of iron chloride, stir thoroughly, warm gently, and when complete, or nearly complete, solution is effected, filter if necessary, place the liquid in a dialyzer, and dialyze in the usual manner until the liquid on the dialyzer is almost tasteless. Then add to this liquid enough water to make it measure 14 fluidounces.-Brit. Pharm.

Most of the so-called "dialyzed iron" of the market is not prepared by dialysis, but is made by a process the same or very similar to the one given under "solution of iron oxychloride."

## Solution of Iron Oxychloride.

Mix the iron chloride solution with 5½ fluidounces of water, and the ammonia water with 10½ fluidounces of water, add the iron solution gradually, with constant stirring, to the ammoniacal liquid, wash the precipitate thoroughly with water, collect it, press somewhat to remove excess of water, add it to the hydrochloric acid contained in a suitable vessel (a bottle), agitate frequently during 3 days, then warm until complete solution

has taken place, and then add enough water to reduce the solution to a specific gravity of 1,050.—Germ. Pharm.

## Solution of Iron Phosphate.

Iron phosphate, soluble .....av.oz. 8 Distilled water, enough to make fl.oz. 16

Solution of Iron "Protoxide." (Solution of Iron Protocitrate.—Solution of Ferrous Citrate.)

Simple syrup . . . . . of each, sufficient

Dissolve the two salts separately in 32 fluidounces of water, mix by adding the iron solution to the sodium solution with constant stirring, collecting the precipitate, washing it quickly with more water, until the washings are tasteless, then dissolve by the aid of a gentle heat in 4 fluidounces of water containing the citric acid and add enough simple syrup to make 16 fluidounces.

## Solution of Iron Pyrophosphate.

Iron pyrophosphate, soluble ..av.oz. 8 Distilled water, enough to make fl.oz. 16

## Solution of Iron Salicylate.

Ferrous sulphate, puregr.	192
Sodium salicylategr.	
Sodium acetateav.oz.	334
Distilled waterfl.oz.	87
Mix and dissolve.	

## Solution of Magnesium Borocitrate.

Magnesium carbonate, powdergr	. 230
Citric acidgr	. 460
Borax, powder gr	. 460
Water, enough to make fl.oz	. 16

Dissolve the citric acid in 15 fluidrams of water at a boiling temperature, then add the magnesium carbonate and afterward the borax; filter, and then add the remainder of the water.

The solution contains about 10 gr. of the dry magnesium borocitrate in each fluid-ounce.

## Solution of Magnesium Citrate.

٠.	
	Magnesium carbonate gr. 225
	Citric acidgr. 450
	Syrup of citric acidfl.oz. 2
	Potassium bicarbonategr. 38
	Distilled watersufficient
	Dissolve the acid in 4 fluidounces of water,

add the magnesium carbonate, and stir until dissolved. Filter the solution, add the syrup and enough water to make 12 fluidounces, introduce the whole at once into a bottle, add the bicarbonate, and at once cork the bottle and tie it over securely with a stout twine.

II.

4	
Citric acidgr.	360
Magnesia, calcinedgr.	
Syrup of citric acid fl.oz.	1
Potassium bicarbonate, crystalgr.	40
Distilled water, enough to	
makefl.oz.	12

Mix the acid, magnesia and 4 fluidounces of water, stir or agitate until dissolved, add the syrup and the remainder of the water, filter, introduce the clear filtrate into a suitable bottle, add the potassium salt, and cork and tie over the bottle immediately.

The above is intended for 1 bottle.

## Solution of Mercury Albuminate.

Egg albumen, freshgr.	900
Mercuric chloride, puregr.	60
Sodium chloridegr.	240
Distilled waterfl.oz.	10

Beat the egg albumen to foam, allow this to become liquid again by standing, and then add to it a solution of the two salts in the water. Set the liquid aside for 2 days in a cool and dark place, and filter.

This preparation must be kept in the dark.-Germ. Form.

## Solution of Mercury and Arsenic Iodides. (Donovan's Solution.)

Arsenious acid, C. P g Mercury g	
Iodineg	r. 102
Alcoholfl.o	

Triturate the arsenic, mercury, iodine, and alcohol together until a dry mass is obtained; then triturate with this mixture 141/2 fluidounces of water gradually added, transfer to a flask, heat to boiling, allow to cool, and filter, adding through the filter enough distilled water to make 16 fluidounces of filtrate.

This formula may be used by pharmacists who do not desire to carry a quantity of arsenic iodide in stock.

It is to be observed that under no consideration should the ordinary commercial im- in the form of citrate.-N. F.

pure powdered arsenious acid, so-called "arsenic," be used.

## Solution of Mercury Chloride.

Mercury chloridegr. 2
Ammonium chloridegr. 2
Distilled waterfl.oz. 4
Dissolve and filter -Rrit Pharm

## Solution of Mercury and Potassium

Tartrate. (Liqueur de Pressavin.) Mercury oxide, freshly precipitated .....gr. 175 Potassium bitartrate.....gr. 175 Distilled water, hot.....fl.oz. 16

Mix, dissolve and filter.

In using, mix 1 fluidounce of this with 32 fluidounces of water, and give a wineglassful 3 or 4 times daily.

## Solution of Morphine.

I. Magendie's Solution:

Morphine sulphategr.	16
Distilled water, warmfl.oz.	1

Dissolve the morphine sulphate in the water, and filter the solution through a small pellet of absorbent cotton. When the solution is cold, pass a little distilled water through the cotton, if necessary, to make the filtrate measure 1 fluidounce. Keep the solution in well-stoppered vials, in a dark place.-N. F.

II. Morphine sulphate ......gr. 1 Distilled water.....fl.oz. 1 -- U. S. P. 1870.

## Solution of Morphine Acetate.

Morphine acetategr.	9
Acetic acid, dilutedm.	
Alcohol	
Distilled water fl.dr.	
-Brit. Phar	
-Brit. Phar	111.

## Solution of Morphine Citrate.

Morphine (alkaloid)gr.	16
Citric acidgr.	
Cochinealgr.	
Alcohol	
Distilled water, enough to make fl.oz.	1

Triturate the solids with the alcohol and 10 fluidrams of water; filter and pass the remainder of the distilled water through the filter.

This solution should not be kept on hand, but prepared only when required.

Each fluidram contains 2 gr. of morphine

Solution of Morphine Bimeconate.	add the alcohol, and macerate in this mixture
Morphine (alkaloid)gr. 13½	the rennet (or the washed mucous membrane
Meconic acid gr. 12	of the fresh stomach of a suckling calf), dur-
Alcohol fl.oz. 1 Distilled water, enough to make fl.oz. 4	ing 3 days, under frequent agitation; then
	filter.—N. F
Add the morphine and acid to the alcohol, then add the water, dissolve by agitation,	II.
and filter through white paper.—Brit. Pharm.	Rennet, fresh
modified.	Salt
Solution of Morphine Hydrochlorate.	Diluted alcohol
The state of the s	Sherry winefl.oz. 16
I. Morphine hydrochlorate gr. 19	Cut the rennet, knead together with the
Distilled water	salt, and set aside for a day, then add the
—Codex.	water and diluted alcohol, let macerate for
II. Morphine hydrochlorategr. 9	several weeks, add the sherry wine and filter.
Hydrochloric acid, dilutedm. 18	
Alcoholfl.dr. 4	Solution of Saccharin.
Distilled waterfl.dr. 12	Saccharingr. 512
—Brit. Pharm.	Bicarbonate of sodiumgr. 240
Solution of Potassa. (Liquor Potassa.)	Alcoholfl.oz. 4 Water, sufficient to makefl.oz. 16
Potassium hydrateav.oz. 1	Mix and dissolve. Each fluidram repre-
Distilled water	sents 4 grains of saccharin.
Mix and dissolve.—U. S. P.	Galatia and Galatia
Solution of Potassium Acetate.	Solution of Strychnine Acetate. (Hall's
Acetic acid	Solution of Strychnine.)
Potassium bicarbonate, Waterof each, sufficient	Acetate of strychninegr. 1
To the acid add $2\frac{1}{4}$ fluidounces of water,	Diluted acetic aciddrops 15 Alcohol
add gradually 6½ av. ounces of potassium	Comp. tincture of cardamomdrops 5
bicarbonate, heat the liquid to boiling, then	Water, sufficient to makefl.oz. 1
neutralize by the further addition of the	Each fl. dr. has 1/8 gr. acetate strychnine.
potassium salt, and then add enough water	
to make 16 fluidounces.—Germ. Pharm.	Solution, Turpentine, Anodyne,
Solution of Potassium Permanganate.	Rademacher's.
I.	Spirit of ether
Potassium permanganategr. 128	Species, Alterative. (Swedish Blood-
Distilled waterfl.oz. 16	Purifying Tea.—Compound Species of
Mix and dissolve. Preserve the solution	Guaiac.—Species ad InfusumLignorum.)
in a glass or rubber-stoppered bottle.	Licorice root, cutav.oz. 1½
II.	Saponaria, cut
Potassium permanganategr. 77	Juniper root, cutav.oz. 6
Distilled water	Guaiacum wood, raspedav.oz. 9
	—Swed. Pharm.
Solution of Rennet. (Liquor Seriparus.  -Liquid Rennet. — Rennet Wine.—	Species, Aromatic. (Aromatic Tea.—
Essence of Rennet.)	Species Resolventes.)
I.	Peppermint, cut fine
Calves' rennet, freshav.oz. 13/4	Wild thyme, cut fine
Sodium chloride	Lavender flower,av.oz. 2
Alcohol	Cloves, cut fine
Dissolve the sodium chloride in the water,	Cubebs, coarse powderav.oz. 1 —Germ. Pharm.
- 1301.0 the boarding officerate in the water,	, Ooimi Lilainii

Species, Bitter. (Bitter Tea.)	Species, Long Life. (Species ad Longam
I.	Vitam.)
Wormwood, cut	Aloes av.oz. 6 Rhubarb av.oz. 1 Gentian av.oz. 1 Zedoary av.oz. 1 Galangal av.oz. 1 Myrrh av.oz. 1 Agaric av.oz. 2 Theriac av.oz. 1 Reduce the first seven ingredients to small pieces, then rub the agaric to coarse powder, triturate the theriac with it and mix the whole.—D.  Species, Marshmallow. (Marshmallow Tea.—Species Althæa.)
-Swed. Pharm Species, Carminative.	Marshmallow root, cutav.oz. 10
Anise	Marshmallow leaves, cutav.oz. 5 Licorice root, cut
Species, Diuretic. (Diuretic Tea.)	Althæa, peeledav.oz. 8
I.  Lovage root, cut	Coltsfoot leaves       av.oz. 4         Licorice root, Russian, peeled. av.oz. 3         Anise       av.oz. 2         Mullein flowers       av.oz. 2         Orris root       av.oz. 2         Cut, bruise and mix them.—N. F.
II. An older formula is this:	Species, Saxon. (Saxon Tea.—Species
Licorice root, cut.       av.oz. 4         Lovage root, cut.       av.oz. 3         Rest harrow root, cut.       av.oz. 3         Pansy herb, cut.       av.oz. 3         Juniper berries, crushed.       av.oz. 3         Parsley seed.       av.oz. 2         Anise seed.       av.oz. 2         D.	Laxantes Schrammii.)  Senna, cut
Species, Gargle. (Species ad Gargarisma.)	Guaiac wood
Elder flowers	Rest harrow, cut
Species, Laxative. (St. Germain Tea.)	Spirit of Angelica, Compound.
Senna, cut	I.   Angelica root, cut   .av.oz.   2½   Valerian, cut   .gr.   270   Juniper berries, bruised   .gr.   270   Alcohol   .fl. oz.   13   Water   .fl. oz.   18   Camphor   .gr.   120   .g
water; then sprinkle over it, as uniformly as possible, the potassium bitartrate. When it	Macerate the roots and berries in the alco- hol and water for 24 hours, then distill off 16
has become dry, mix it lightly and uniformly	fluidounces, and dissolve the camphor in the
with the other incredients. N. E.	distillate Com Phorm

distillate.—Germ. Pharm.

with the other ingredients.-N. F.

II.	Spirit of Formic Acid, Compound.
Oil of angelica rootdrops 12 Oil of valeriandrops 4	Oil of lavender flowers m 80
Oil of juniper berriesdrops 4	Oil of turpentine
Camphor gr. 120	Spirit of formic acid, enough to make
Alcohol, enough to makefl.oz. 16 —H.	'  n
Spirit of Almond, Bitter.	Spirit of Horse-Radish, Compound.
Oil of bitter almond	Horseradish root, fresh, scraped, av. oz. 2
Alcohol	. Ditter orange peer, cut sman and
Dissolve the oil in the alcohol and a	27
enough water to make 16 fluidounces	Alcohol flor 8
S. P.	Mix and distill 16 fluidounces.—Brit,
Spirit of Ammonia, Succinic.	Pharm.
Oleobalsamic mixturefl.oz.	}
Alcohol fl.oz.	
Ammonia water	On or lavender nowers
—D.	
Spirit of Cardamom, Compound.	—U. S. P.
Oil of cardamomdrops 15	Spirit of Lemon.
Oil of caraway drops & Oil of cassia	
Alcoholfl.oz.	Lemon peel, freshly gratedgr. 390
Glycerin	,
Dissolve the oils in the alcohol, add	Dissolve the oil in 14 fluidounces of alco-
glycerin, and then the water.—N. F	hol, add the peel, macerate for 24 hours, filter, and through the filter add enough di-
Spirit of Cinnamon.	luted alcohol to make 16 fluidounces.—
Oil of cinnamonfl.dr. 15	U. S. P.
Alcohol, enough to makefl.oz. 10	Guinit of Molines (C. i.i. of P. 1)
—U. S. P	This may be prepared by distilling 4 av.
Spirit of Cloves.	auman of molines how with 14 duidous and
Oil of cloves	0/2
Spirit of Ether, Camphorated. (Ne	16 fluidounces of product.—D. modified.
Drops.)	It may also be prepared by dissolving 24
Camphorav.oz.	drops of oil of melissa in 12 fluidounces of alcohol and adding 4 fluidounces of water.
Alcoholfl.oz. 10	03/4
Ether	Spirit of Menssa, Compound. (Carmelite
Spirit of Formic Acid. (Spiritus For	Spirit.—Aromatic Spirit.—Karmeliter
carum.—Spirit of Ants.)	
Formic acidfl.dr.	I. Melissa herbgr. 450
Distilled waterfl.oz.	Lemon peelgr. 384
Alcohol, enough to makefl.oz. 1	Cevlon cinnamongr. 96
Mix the formic acid with the distil	Clovesgr. 96
water, and add the alcohol.  Formic acid for this preparation sho	337
have a specific gravity of 1.060 to 1.063	
N. F.	Germ. Pharm.

A quicker and more convenient process is this:

Oil of	melissadrops 1	8
	lemondrops 1	
	nutmegdrops 1	
Oil of	clovesdrops 1	$^2$
Oil of	Ceylon cinnamondrops 1	2
	ol, enough to makefl.oz. 1	
	н	

## Spirit of Mastic Compound. (Spiritus Matriculis.—Mutter Spiritus.)

Mastic av.oz.	1
Olibanumav.oz.	1
Myrrh av.oz.	1
Alcoholfl.oz.	
Waterfl.oz.	8

Digest the gums with the alcohol, add the water and distill one pint.

## Spirit of Rosemary.

Oil of rosemaryfl.dr.	$2\frac{1}{2}$
Oil of rosemaryfl.dr. Alcohol, enough to makefl.oz.	16
—Brit, Phar	m.

## Spirit of Rosemary, Compound. (Aqua Hungarica.)

Spirit of rosemaryfl.oz.	91/2
Spirit of lavender flowers fl.oz.	
Spirit of sagefl.oz.	

The spirits may be made from the respective oils and alcohol so as to contain 10 per cent of the former.—D. and H.

## Spirit of Soap.

Castile soap, shavingav.oz. Alcoholfl.oz.	23/
Alcoholfl.oz.	91/2
Water, enough to makefl.oz.	16

Introduce the soap into a bottle, add the alcohol and  $3\frac{1}{2}$  fluidounces of water, cork the bottle, and immerse in hot water, frequently shaking. When the soap is dissolved, allow the bottle and contents to become cold, add the remainder of the water and filter.—N F.

## Spirit of Soap, Camphorated. (Liquid Opodeldoc.)

Mix and filter.-Germ. Pharm.

## Spirit of Thyme.

Oil of thyme, white	45
Alcoholfl.oz.	
Waterfl.oz.	4
Mix and filter -D	

## Sponge, Burnt, Artificial.

Mix and reduce to fine powder.-H.

Prepared oyster shell is made by thoroughly cleaning the shell, then reducing in a mortar, separating the finer particles by elutriation and drying the latter.

## Sponge, Carbolized.

Carbolic acid, crystal gr.	450
Alcoholfl.oz.	
Waterfl.oz.	18

Bleached sponges are allowed to remain in this solution for 24 hours, when an equal volume of water is added. The sponges are to remain in the fluid.—D.

## Starch, Iodized.

Starch	 	 	 gr.	475
Iodine				
Water	 	 	 suffic	cient

Triturate the iodine with a little distilled water, add the starch gradually and continue trituration until a uniform blue-black product is obtained. Dry this at a temperature not exceeding 40 degrees C., rub to fine powder, and preserve in well-stoppered bottles.—U. S. P. 1880 and N. F.

## Stone, Medicinal. (Lapis Medicamentosus.—Lapis Mirabilis.)

Alumav.oz.	3
Lithargeav.oz.	
Armenian boleav.oz.	
Sulphate of iron, driedav.oz.	11/2
Diluted acetic acidav.oz.	2

Mix and evaporate to dryness.

#### Suet. (Tallow.)

Mutton snet is official in the United States pharmacopœia, but beef suet is also used. Either may be prepared by taking the fat from the vicinity of the kidneys of the sheep or cow, cutting into small pieces, heating on

a water bath until the fat is quite melted and then straining with expression through flannel. Dieterich recommends adding to the fatty matter in the dish 1-20 its weight of dried sodium sulphate in fine powder, continuing the heat for 15 minutes after thorough fusion has occurred, stirring frequently and filtering by hot filtration. The sodium sulphate removes moisture and assists in separating the membranes.

Beef suet has a slightly lower fusing point than mutton suet, otherwise the two are practically alike, so that one or the other may be selected, depending upon the use the product is to serve.

## Suet, Benzoated.

Mutton suetav.oz.	10
Benzoin, coarse powder av.oz.	1
Sodium sulphate, driedav.oz.	1

Heat the three ingredients together on a water bath for one hour, stirring frequently, then strain through flannel or filter. The suet to be used should be such as has not already been treated with the dried sodium sulphate.—D.

## Suet, Deer.

Beef tallow is usually dispensed for this. It is generally sold either in the form of flat cake or of cylinders about 1 inch in diameter.

## Suet, Salicylated.

Salicylic a	cid	0
	etav.lb.	

Melt the suet on a water bath, add the acid, and dissolve by stirring.—Germ. Pharm.

## Sugar, Coumarin. (Elæosaccharum Cumarini.)

Coumarin .		 	gr.	31
Sugar, fine	powder	 	av.oz.	8

Mix well and keep in well-closed bottles.

—D.

## Sugar, Vanilla. (Elæosaccharum Vanillæ.)

mmæ.)	
Vanillaav.oz.	1
Alcohol fl.oz.	1
Sugar of milk, crystalav.oz.	2
Sugar, crystal (i. e. rock candy). av. oz.	7
Sugar, fine powder, enough to	
makeav.oz.	10

Cut the vanilla into very small pieces by means of a shears or sharp knife, add the alcohol, macerate for 30 minutes, add the milk sugar, contuse until tolerably well reduced, add one-half of the rock candy, contuse and triturate until a tolerably fine powder is produced, sift through a No. 50 sieve, return the residue to the mortar, add the remainder of the rock candy, contuse, triturate, and sift as before, return the coarser particles to the mortar, and continue the trituration and sifting until nearly all has passed through the sieve; finally add the powdered sugar, mix well, and preserve in well-stoppered bottles.

For general sale the above should be mixed with nine times its weight of powdered sugar.—D.

## Sugar, Vanillin. (Elæosaccharum Vanillini.)

Va	anilli	n							 	gr.	75
St	ıgar,	pov	vd	er	ed				 :	av.oz.	8

Mix well and preserve in well-stoppered bottles.—Codex.

This has about the same relative strength as vanilla sugar and may be employed in place of the latter.

Flavored sugars made with volatile oils should be prepared by intimately mixing 1 drop of oil with 30 grains of powdered sugar.—N. F.

They are properly known as oil-sugars or oleosaccharates (Latin: elæosacchara).

## Syrup of Ammonium Chloride.

Ammonium chlorideav.oz.	
Sugarav.oz.	11
Water, enough to makefl.oz.	16

Dissolve the ammonium salt in 10 fluidounces of water, add the sugar and the remainder of the water, dissolve by agitation and strain if necessary.

## Syrup of Apomorphine Hydrochlorate.

Apomorphine hydrochlorategr.	4
Hydrochloric acid, dilutefl.dr.	11/2
Alcohol	$5\frac{1}{2}$
Distilled waterfl.dr.	51/2
Simple syrupfl.oz. 1	41/2

Mix the alcohol and water, add the apomorphine to it, dissolve by agitation, add the acid and the syrup.—Brit. Form.

Syrup	of	Aral	ia,	Compound.	(Com-
pot	nd	Syrup	of	Spikenard.—A	lterative
Syr	up.	)			

Spikenard rootgr. 300
Burdockgr. 300
Yellow dockgr. 300
Guaiacum woodgr. 300
Sassafras bark gr. 240
Prickly ash barkgr. 240
Elder flowersgr. 240
Blue flag rootgr. 240
Glycerin,
Diluted alcoholof each, sufficient
Sugarav.oz. 10

Mix the drugs and reduce to a coarse powder, extract in the usual way by percolation, with diluted alcohol; obtain 10 fluid-ounces of percolate in which dissolve the sugar by percolation and to this last percolate add, if necessary, enough glycerin to make 16 fluidounces.—Eclectic.

## Syrup of Asafetida.

Asafetida, select gumgrs.	240
Boiling waterfl.oz.	
Sugarav.oz.	131/2

Rub the gum with a portion of the water to a smooth paste, add the remainder of the water and sugar, dissolve by aid of gentle heat and strain.—Eclectic.

## Syrup of Bayberry, Thompsonian.

Bayberry barkav.oz.	8
Diluted alcoholfl.oz.	
Sugarav.oz.	8

Macerate the bark with the diluted alcohol in a warm place for two days, strain, evaporate the colature to 8 fluidounces and in this dissolve the sugar.

## Syrup of Belladonna, Compound.

Fluid extract of belladonna root, fl.dr.	2
Fluid extract of chestnut leaves.fl.dr.	4
Syrup of wild cherry bark, enough	
to makefl.oz.	16

This is an effective mixture for whooping cough.

## Syrup of Blackberry, Aromatic.

Blackberry root barkav.oz.	21/2
Cinnamongr.	
Nutmeggr.	
Clovesgr.	60
Allspicegr.	60
Sugarav.oz.	11
Diluted alcohol,	
Blackberry juiceof each, suffic	ient

Reduce the drug to moderately coarse powder, and percolate in the usual manner with diluted alcohol until 4 fluidounces of percolate are obtained. To this add 7 fluidounces of the juice and the sugar, dissolve by agitation, and strain.—N. F.

## Syrup of Buckthorn Bark. (Syrup of Frangula.)

Fluid extract of frangula.....fl.dr. 10 Simple syrup, enough to make. fl.oz. 16

## Syrup of Buckthorn Berries.

(Syrupus Spinæ Cervinæ.)

Dissolve the sugar in 7 fluidounces of the juice, with the aid of a gentle heat, allow the syrup to cool, then add enough of the juice to make 16 fluidounces and strain if necessary.—N. F.

## Syrup of Butyl Chloral.

See Syrup of Croton Chloral.

# Syrup of Calcium Hypophosphite. Syrup of Calcium and Sodium Hypophosphite.

See Syrups of the Hypophosphites.

## Syrup of Calcium Phosphate.

## I. Wiegand's:

Calcium	phosphate,	precipi-	
tated		av.oz.	11/4
Hydrochle	oric acid	fl.dr.	41/2
Sugar		av.oz.	141/2

Dissolve the calcium phosphate in the acid previously mixed with 12 fluidounces of water, filter, add the sugar and the remainder of the water, dissolve by agitation and strain.

#### II. Durand's:

Calcium phosphate, precipitated.gr. 2 Phosphoric acid, glacialgr. 2	
Sugarav.oz.	
Distilled waterfl.oz.	8
Spirit of lemondrops	24

Mix the calcium phosphate with the water, heat moderately, gradually add the acid until all the calcium salt is dissolved, replace the water lost by evaporation, filter, dissolve the sugar in the filtrate, strain, if necessary, and add the spirit.

### Syrup of Cascara Sagrada.

Fluid extract of cascara sagrada,	
Brit. Formfl.oz.	31
Fluid extract of licoricefl.oz.	21/2
Carminative tincturefl.dr.	11/2
Simple syrup, enough to makefl.oz.	
Brit For	

## Syrup of Cherries. (Syrupus Cerasorum.)

Crush black, sour cherries with the stones to a pulp, set aside in a covered vessel so that it will be at a temperature of about 20 degrees C., and stir occasionally. When 2 parts by measure of clear or filtered liquid and 1 of alcohol no longer become cloudy, the juice is to be strained with expression and filtered. To every 8 fluid-ounces of juice, add 15 av. ounces of sugar and 16½ fluidounces of simple syrup, dissolve by agitation and strain.—Germ. Pharm.

## Syrup of Chloral.

I.	
Chloral hydrategr.	320
Distilled waterfl.dr.	
Simple syrup, enough to make fl.oz.	

Dissolve the chloral in the water and add the syrup.—Brit. Pharm.

U		
	Chloral hydrategr.	120
	Distilled waterfl.dr.	
	Simple syrupfl.oz.	31/2
	Spirit of peppermintdrops	4

Dissolve the chloral in the water, add the syrup, and then the spirit.—Codex.

#### Syrup of Cinchona.

Tincture of cincho	na 3
Simple syrup	fl.oz. 13
	—Н.

# Syrup of Cochineal. (Syrupus Coccionellæ.)

Cochineal powderav.oz.	11
Alcoholfl.oz.	$3\frac{1}{2}$
Simple syrupfl.oz.	121/2

Mix, let stand for several days, and strain through flannel.—H.

## Syrup of Codeine.

Codeine sulphategr.	
Simple syrupfl.oz.	4

Reduce the codeine sulphate to a fine powder and dissolve it in the syrup previously warmed.

Each fluidram contains about ½ gr. of codeine sulphate.—N. F

H.		
Codeine (alkaloid)	gr. 4	$^{4}$
Diluted alcohol	fl.dr.	2
Distilled water	fl.dr. 2	2
Simple syrup, enough to make	fl.oz. 4	4

Dissolve the codeine in the diluted alcohol and add the other ingredients.—Brit. Form.

#### III.

Codeine (alkaloid.) gr. 4	
Alcoholfl.dr. 2	
Simple syrup, enough to makefl.oz. 4	

Dissolve the codeine in the alcohol and add the syrup.—Codex.

### Syrup of Corydalis, Compound.

Corydalis rootgr. 600
Twin-leaf rootgr. 300
Blue flag rootgr. 150
Sheep laurel leavesgr. 150
Sugarav.oz. 10
Simple syrup,
Alcohol,
Water of each, sufficient

Reduce the mixed drugs to powder and extract by percolation in the usual way so as to obtain 10 fluidounces of product, using as a menstruum a mixture of alcohol and water in the proportion of 1 of the former to 2 of the latter; in the percolate dissolve the sugar by agitation or percolation, and add to the solution enough simple syrup to make 16 fluidounces.—Eclectic.

#### Syrup of Croton Chloral.

Croton chloral hydrate	gr. 256	
Simple syrup, enough to r		

Dissolve the croton chloral in the syrup previously made hot.—Brit. Form.

### Syrup of Creosote.

Glycerite of Simple syrup	creosote.	 	 .fl.oz.	21/2
Simple syrup	· · · · · · · · ·	 	 .fl.oz.	131/2

This contains 12 per cent of creosote.

#### Syrup of Cubeb.

Fluid extract of cubebfl.oz	2
Magnesium carbonategr.	240
Sugarav.oz.	13
Oil of bitter almonddrop	
Orange flower water fl.oz.	2
Water, enough to makefl.oz.	16

Triturate the fluid extract with the magnesium carbonate, then add slowly, with constant trituration, 2 av. ounces of the sugar in small portions; when thoroughly mixed, add

gradually first the orange flower water, and then 7 fluidounces of water, triturating the mixture until the sugar is dissolved; filter and add sufficient water to make the filtrate measure 11 fluidounces in which the sugar is to be dissolved without heat; to the saccharine solution add the oil dissolved in a little alcohol, and then enough water to make 16 fluidounces.

## Syrup of Digitalis.

Tincture of digitalis, U. S. P. . . fl.dr. 6 Simple syrup, enough to make . . fl.oz. 16 —Codex.

#### Syrup of Ergotin.

Extract of ergotgr. Simple syrupfl.oz.	
* * *	-H.

### Syrup of Ether.

Ether	.fl.dr.	4
Alcohol		
Distilled water	.fl.oz.	41/
Simple syrup, enough to make.		
	—Cod	ex

#### Syrup of Eucalyptus.

Fluid extract of eucalyptusfl.dr.	
Magnesium carbonategr.	360
Waterfl.oz.	
Sugarav.oz.	15

Triturate the fluid extract with the magnesium carbonate, add the water gradually, let stand 1 hour, filter, and in the filtrate dissolve the sugar without heat.

## Syrup of Fox Lungs. (Fuchs Lungen Saft.—Syrupus Pulmonum Vulpium.)

Pectoral elixirfl.dr.	01
Syrup of sennafl.dr.	22
Simple syrupfl.dr.	
Glycerin	1

## Syrup of Garlic, Artificial, Thomp-

Tincture of asafetida	.fl.dr.	1
Acetic acid, concentrated	.fl.dr.	6
Simple syrup	.fl.oz.	16

#### Syrup of Glycyrrhizin.

Ammoniated glycyrrhizin gr.	180
Glycerinfl.oz.	1
Waterfl.oz.	
Sugar av.oz.	12

Heat the water, add the glycyrrhizin, stir tion.—Eclectic.

until dissolved, filter, add the sugar and glycerin, shake until dissolved, and strain.

### Syrup of Guaiac.

Red root

Guaiac, powderav.oz.	
Potassagr.	60
Sugarav.oz.	13
Water	

Dissolve the potassa in the water, add the guaiac, macerate for 7 days, filter, add the sugar, dissolve, and strain.

or 320

### Syrup of Horehound, Compound.

Neu 1001gi-	020
Elecampanegr.	320
Spikenardgr.	
Comfrey gr.	
Wild cherry barkgr.	
Horehoundgr.	
Blood root gr.	
	100
Glycerin,	
Alcohol,	
Waterof each, suffic	
Sugarav.oz	. 10

Mix the drugs, reduce to fine powder and extract by percolation in the usual way so as to obtain 10 fluidounces of percolate, using as a menstruum a mixture of 2 parts of water and 1 of alcohol by measure; in the percolate dissolve the sugar by agitation or percolation, and to this solution add enough glycerin to make 16 fluidounces.—Eclectic.

## Syrup of Horseradish, Compound.

(Cough Elixir.—Vegetable Elixir.)

Fresh root of horseradish,	
gratedav.oz.	1
Bonesetgr.	240
Canada snake rootgr.	120
Sugarav.oz.	14
Boiling water,	

Diluted acetic acid ...of each, sufficient

Add the horseradish to 4 fluidounces of diluted acetic acid, macerate for 2 days, express, and add enough of the acid to the expressed marc so that the liquid obtained by again expressing latter, added to the previous liquid, will make 4 fluidounces, express again, mix the two liquids and filter.

Infuse the boneset and snake root in the usual way so as to obtain 4 fluidounces of product.

Mix the two liquids and in this mixture dissolve the sugar by agitation or percolation.—Eclectic.

### Syrup of Hypophosphites.

(Churchill's Syrup of Hypophosphites.)

Calcium hypophosphite gr.	345
Sodium hypophosphitegr.	115
Potassium hypophosphite gr.	115
Diluted hypophosphorous acidm.	15
Sugarav.oz.	83
Spirit of lemonfl.dr.	11/2
Water, enough to makefl.oz.	16

Triturate the hypophosphites with 7 fluidounces of water until they are dissolved, add the spirit, and the acid, and filter. In the filtrate dissolve the sugar by agitation or percolation and add enough water through the filter to make 16 fluidounces. Strain, if necessary.—U. S. P.

#### II. Parrish's Formula:

Calcium hypophosphitegr.	288
Sodium hypophosphitegr.	
Potassium hypophosphitegr.	96
Sugarav.oz.	
Distilled water, hot fl.oz.	9
Orange flower waterfl.dr.	4

Make a solution of the hypophosphites in the hot water, filter, dissolve the sugar in the filtrate, strain, and to the colature add the orange flower water.

## Syrup of Hypophosphites, Compound.

I.
Calcium hypophosphite gr. 256
Potassium hypophosphite gr. 128
Sodium hypophosphitegr. 128
Iron hypophosphite gr. 16
Manganese hypophosphitegr. 16
Potassium citrate gr. 40
Citric acid gr. 15
Quinine hydrochlorate gr. 8
Tincture of nux vomicam. 160
Sugarav.oz. 13
Watersufficient

Rub the hypophosphites of iron and of manganese with the potassium citrate and citric acid to powder, add 1 fluidounce of water, and warm the mixture a few minutes until a clear greenish solution is obtained. Introduce the other hypophosphites and the quinine hydrochlorate, previously triturated together, into a bottle, next add the sugar, the iron and manganese solution first prepared, the tincture of nux vomica, and, lastly, enough water to make up the volume, as soon as the sugar is saturated by the liquid, to 16 fluidounces. Agitate until solution

has been effected, and strain, if necessary.
—N. F.

## Syrup of Hypophosphite of Calcium.

Calcium hypophosphite gr. \$	256
Citric acidgr.	10
Sugarav.oz.	
Water, enough to makefl.oz.	$16^{'}$

Dissolve the calcium hypophosphite and citric acid in 8 fluidounces of water, filter the solution, add the sugar to the filtrate, and pass enough water through the filter to make the product, after the sugar has been dissolved by agitation, measure 16 fluidounces.

Each fluidram contains 2 gr. of calcium hypophosphite.—N. F.

## Syrup of Hypophosphite of Calcium, Manganese and Potassium.

Calcium hypophosphitegr.	256
Manganese hypophosphitegr.	
Potassium hypophosphitegr.	128
Distilled water, boiling fl.oz.	31/4
Simple syrup, enough to make fl.oz.	16

Triturate the hypophosphites with the water, filter, and add the syrup.

## Syrup of Hypophosphite of Calcium and Sodium.

Calcium hypophosphite gr.	256
Sodium hypophosphitegr.	256
Citric acidgr.	
Sugarav.oz.	
Water, enough to makefl.oz.	

Dissolve the two hypophosphites and citric acid in 8 fluidounces of water, filter the solution, add the sugar to the filtrate, and pass enough water through the filter to make the product, after the sugar has been dissolved by agitation, measure 16 fluidounces.

Each fluidram contains 2 gr. each of calcium and sodium hypophosphites.—N. F.

#### Syrup of Hypophosphites with Iron.

Ferrous lactate, in crustsgr.	72
Potassium citrategr.	72
Syrup of hypophosphites, enough	
to makefl.oz.	16

Triturate the two salts with a small quantity of syrup gradually added, until they are dissolved, then add the remainder of the syrup.

This preparation should be freshly made when wanted.—U. S. P.

## Syrup of Hypophosphite of Iron.

Iron hypophosphitegr.	128
Potassium citrategr.	
Orange flower waterfl.oz.	
Simple syrup, enough to make fl.oz.	

Dissolve the iron hypophosphite with the aid of the potassium citrate in the orange flower water, and add the syrup.

Each fluidram contains 1 gr. of hypophosphite of iron (ferric).—N. F.

## Syrup of Hypophosphite of Manganese.

Manganese sulphategr.	120
Calcium hypophosphite, gr.	80
Sugarav.oz.	
Orange flower waterfl.dr.	
Watersuffic	

Dissolve the hypophosphite and sulphate in separate portions of water, mix the two solutions, filter, washing the precipitate in the filter with fresh distilled water; evaporate the filtrate to 8 fluidounces, dissolve the sugar in the latter, strain, and add the orange flower water. Each fluidounce contains  $2\frac{1}{3}$  gr. of manganese hypophosphite.

## Syrup of Hypophosphite of Sodium.

Sodium hypophosphitegr.	256
Citric acidgr.	
Sugarav.oz.	13
Water, enough to makefl.oz.	16

Dissolve the sodium hypophosphite and the citric acid in 8 fluidounces of water, and filter the solution. In this dissolve the sugar by agitation, and pass the remainder of the water through the filter.

Each fluidram contains 2 gr. of sodium hypophosphite.—N. F.

## Syrup of Iron and Sodium Albuminate.

White of egg no. 4
Sugarav.oz. 2
Tinct, chloride of ironfl.oz. 2
Solution of soda, .
Water of each, sufficient

Mix the white of egg with the sugar and add enough water to effect complete solution; add the tincture of iron, and then just enough of the solution of soda to dissolve the coagulated albumen; finally make up to 16 fluidounces with water.

## Syrup of Iron (Ferric) Chloride.

1.	
Solution of iron chloride	fl.dr. 2
Simple syrup, enough to make.	
	-Codex.

••	
Tincture of chloride of ironfl.oz.	1
Sodium citrateav.oz.	2
Waterfl.oz.	6
Sugarav.oz.	10
Syrup, enough to makefl.oz.	

Mix the tincture of ferric chloride with the water and dissolve in this mixture the sodium citrate and the sugar with the aid of heat; when cold add sufficient syrup to make 16 fluidounces.

## Syrup of Iron (Ferrous) Chloride.

See Syrup of Iron Protochloride.-N. F.

## Syrup of Iron Citrate.

Iron citrate, solublegr.	240
Distilled water, hotfl.dr.	4
Simple syrup, enough to make fl.oz.	
—Co	

## Syrup of Iron Hypophosphite.

See Syrup of Hypophosphites.

## Syrup of Iron and Quinine Iodides.

## I. Bouchardat's formula:

Iodinegr. 4	2 .
Iron, in powdergr. 1	7
Simple syrupfl.oz. 1	
Quinine sulphategr.	
Diluted sulphuric acidsufficier	nt
Distilled waterfl.dr.	

Digest the iodine, iron, and 3 fluidrams of the water until the red-brown color of the iodine has disappeared; filter through a small filter into the syrup. Then dissolve the sulphate of quinine in  $1\frac{1}{2}$  fluidrams of water with the aid of diluted sulphuric acid and mix this solution with the previously prepared syrup.

II.

Quinine sulphate	gr. 20
Hypophosphorous acid, diluted	
Potassium iodide	
Simple syrup, enough to make	
Syrup of iron iodide (U. S.)	
Syrup of fron louide (0. 3.)	11.02. 0

To the quinine sulphate add about 10 drops of commercial solution of hypophosphorous acid and then a small amount of syrup; when the quinine salt is dissolved,

add the remainder of the syrup and after- Syrup of Iron Pyrophosphate. wards the potassium iodide dissolved in a few drops of water; mix well. Now add the syrup of iron iodide and mix. Should any cloudiness appear, clear it up by a few drops of the hypophosphorous acid.

A fluidram of this syrup contains about 4 gr. of dry iodide of iron and about 6 gr. of hydriodide of quinine.

## Syrup of Iron and Ammonium Phosphate.

Iron sulphategr. 635
Sodium phosphategr. 820
Glacial phosphoric acid, C. Pgr. 900
Ammonia watersufficient
Sugar
Distilled watersufficient

Dissolve the sodium phosphate and the iron sulphate separately in distilled water, mix the solution, and wash the resulting precipitated iron phosphate. Then to one-half of the phosphoric acid, dissolved in 21/4 fluidounces of water, add ammonia water until exactly neutral. To the remainder of the phosphoric acid, dissolved in a like amount of water, add the moist iron phosphate and dissolve by the aid of a gentle heat; then add the solution of ammonium phosphate and the sugar, dissolve the whole, strain, and evaporate to 16 fluidounces.

Each fluidram contains 4½ gr. iron phosphate, 43/4 gr. ammonium phosphate, and 31/2 gr. of phosphoric acid.

## Syrup of Iron and Ammonium Tartrate.

Tartrate of iron and potassium. gr. 225 Distilled water, hot......fl.dr. Simple syrup, enough to make fl.oz. 16

Dissolve the iron salt in the water and add the syrup.—Codex.

### Syrup of Iron Iodohydrargyrate.

Syrup of iodide of ironfl.oz.	15
Red iodide of mercurygr.	3

## Syrup of Iron and Potassium Tartrate.

Tartrate of iron and potassium..gr. 225 Distilled water, hot.....fl.dr. Simple syrup, enough to make fl.oz.

Dissolve the iron salt in the water and add the syrup.-Codex.

Iron pyrophosphate, solublegr.	90
Distilled waterfl.dr.	
Simple syrup, enough to makefl.oz.	
—Cod	

#### Syrup of Liquidambar.

Sweet-gum bark, coarse	
dered	
Sugar	av.oz. 14
Water	sufficient

Moisten the bark thoroughly with water, macerate in a close vessel for 24 hours, pack in a percolator, and pour on water until 8 fluidounces of percolate are obtained. this dissolve the sugar by agitation or percolation. - Eclectic.

This has been recommended for the bowel complaints of children, also for chronic cough and mucous affections.

### Syrup of Lobelia.

Vinegar of lobeliafl.oz.	8
Sugarav.oz.	

Dissolve by aid of heat not exceeding 82 degs. C.; continue heat for 3 hours, removing any scum that may form, and strain while hot,—Eclectic.

## Syrup of Lobelia, Thompsonian.

Lobelia seed	.av.oz.	1
Water		
Vinegar		
Sugar		
Tincture of Jobelia	fl oz	4

Boil the lobelia with the water and vinegar for one-half hour, occasionally replacing the water lost by evaporation, then strain, add the sugar, dissolve, and add the tincture.

#### Syrup of Maidenhair. (Syrupus Capilli Veneris.)

Maidenhairgr.	320
Distilled water, hotfl.oz.	10
Sugarav.oz.	11

Macerate the fern with the water for 6 hours, strain, add the sugar and dissolve. -Codex.

#### Syrup of Manganese Iodide.

Manganese sulphate.	gr. 960
Potassium iodide	gr. 1140
Sugar	
Distilled water,	
Simple syrup	of each sufficient

Dissolve the two salts each in 3 fluidounces of water to which 2 fluidrams of syrup have been added, mix them, place in a cool location for at least one-half hour, filter, allowing the filtrate to pass into a bottle containing the sugar; add sufficient water through the filter to make the whole measure 16 fluid-ounces, dissolve the sugar by agitation, and filter if necessary.

Each fluidram contains about 7½ gr. of manganese iodide.

### Syrup of Manganese Hypophosphite.

See Syrup of Hypophosphite of Manganese.

### Syrup of Manganese Phosphate.

Manganese sulphate	gr.	920
Sodium phosphate.av.oz. 31/2		
Hydrochloric acid		
Sugar		
Water, enough to make		

Dissolve the salts separately in 10 fluidounces of water, and add solution of sodium phosphate to the solution of manganese sulphate as long as it produces a precipitate, which wash with cold water, and then dissolve by means of the hydrochloric acid; dilute this solution till it measure 8¾ fluidounces, and in this dissolve the sugar.

Each fluidram contains 5 gr. of manganese phosphate.

## Syrup of Mercury Iodide. (Syrup of Gibert.)

Red iodide of mercurygr.	3
Potassium iodidegr.	120
Water fl.dr.	3
Simple syrup, enough to make fl.oz.	10

Dissolve the mercuric and potassium iodides in the water and add the syrup.

## Syrup of Mitchella, Compound.

(Compound Syrup of Partridge Berry.
—Mother's Cordial.—Compound Syrup of Squaw Vine.)

Mitchella
Helonias rootgr. 240
Cramp barkgr. 240
Blue cohosh gr. 240
Oil of sassafrasdrops 4
Sugarav.oz. 10
Water,
Alcoholof each, sufficient

Mix the drugs, reduce to powder, add the oil, and percolate in the usual manner so as to obtain 11 fluidounces of product, using as Germ. Form,

a menstruum a mixture of 1 part of alcohol by measure and 2 of water; in this percolate dissolve the sugar.—Eclectic.

### Syrup of Nickel Bromide.

Nickel bromidegr.	320
Glycerinfl.oz.	
Water	
Sugarav.oz.	16

Dissolve the nickel bromide in the water, filter, add the glycerin, and in this mixture dissolve the sugar by agitation or percolation.

## Syrup Opiated. (Syrupus Opiatus.)

Extract of opiumgr.	191
Waterfl.dr.	2
Simple syrup, enough to makefl.oz.	16

Dissolve the extract in the water and add the syrup.—Codex.

### Syrup of Osmunda, Compound.

Osmunda	gr. 600
Burdock	gr. 300
Yellow dock	gr. 300
Turkey corn	gr. 300
Comfrey	gr. 300
Stillingia	gr. 300
Prickly ash berries	gr. 300
Calamus	gr. 75
Sugar	.av.oz. $8\frac{1}{2}$
Diluted alcohol	sufficient

Grind drugs to coarse powder, extract them by percolation with diluted alcohol to make 11<sup>1</sup>/<sub>4</sub> fluidounces of product, in which the sugar is to be dissolved.—Eclectic.

#### Syrup of Peppermint.

Peppermint, herb, cutgr.	324
Alcohol	31/2
Water fl.oz. Sugar av.oz.	41/2
Sugarav.oz.	61/4
Simple syrupfl.oz.	8

Moisten the drug with the alcohol, add the water, let macerate for 24 hours, express, add the sugar and syrup and dissolve by agitation.—Germ. Pharm.

#### Syrup of Peru Balsam.

Peru balsam	av.oz. 1
Sugar,	
Waterof each	h, sufficient

Upon the balsam pour 10 fluidounces of hot water, set aside for 24 hours, occasionally agitating, filter, and in 8 fluidounces of filtrate dissolve 13 av. ounces of sugar,—Germ. Form,

## Syrup of Poke Root, Compound.

Poke root gr.	640
American ivy bark gr.	640
Black cohoshgr.	320
Sheep laurelgr.	320
Oil of sassafras drops	
Oil of wintergreendrops	
Sugarav.oz.	
Alcohol,	
Water,	

Simple syrup ......of each, sufficient

Mix the drugs, reduce to fine powder, add the oils, and extract by percolation so as to obtain 10 fluidounces of percolate, using as a menstruum a mixture of 1 part of alcohol by measure and 2 of water. In this dissolve the sugar, and then add enough simple syrup to make 16 fluidounces.—Eclectic.

## Syrup of Quinine Sulphate.

Quinine sulphategr. 96	3
Diluted sulphuric acid	
Distilled waterfl.dr.	
Simple syrup, enough to makefl.oz. 16	;

Dissolve the quinine in the acid and water and add the syrup. -- Codex.

## Syrup, of Restorative, Thompsonian. (Restorative Cordial, "Number five.")

American poplargr.	144
European poplargr.	
Bayberry root barkgr.	
Waterfl.oz.	
Sugarav.oz.	

Boil the drugs for a few minutes with the water, strain, add the sugar, then 150 gr. of peachmeat reduced to fine condition and finally 4 fluidounces of brandy (or tincture of myrrh).

#### Syrup of Rhubarb and Potassium, Compound. (Neutralizing Cordial.)

I.	40
Rhubarbgr. 2	
Hydrastis gr. 1	
Cinnamongr. 1	
Potassium carbonategr. 2	
Oil of peppermintdrops	
Sugarav.oz.	
Alcoholfl.oz.	8
Waterfl.oz.	16

Dissolve the potassium carbonate in a portion of the water and mix in a suitable sized container with the rhubarb, hydrastis and cinnamon, the last three being in fine powof the water and allow to stand for 48 hours, agitating the whole briskly at frequent intervals. Decant the clear portion, and filter the remainder through absorbent cotton, adding sufficient water through the filter to make the whole measure 24 fluidounces. In this dissolve the sugar by agitation, and add the oil of peppermint.—Eclectic.

II.

Fluid extract of rhubarbfl.dr. Fluid extract of golden sealfl.dr.	4½ 2¼
Potassium carbonategr.	128
Simple syrupfl.oz.	4
Tincture of cinnamonfl.oz.	1
Spirit of peppermintfl.dr.	1
Diluted alcohol, enough to	
make	· 16

Dissolve the potassium carbonate in the syrup, and add the solution to the fluid extracts, tincture and spirit, previously mixed with 10 fluidounces of diluted alcohol. well, add the remainder of the diluted alcohol, and filter, if necessary.

## Syrup of Saccharin.

Saccharin gr.	150
Sodium carbonate, puregr.	165
(Or sodium bicarbonate, puregr.	180)
Distilled waterfl.oz.	32

Dissolve by the aid of a gentle heat.

This may be employed as a substitute for simple syrup.

## Syrup of Senna with Manna. (Syrupus Mannatus. -- Compound Syrup of Manna.)

Syrup of senna, U. S. P.....fl.oz. 4 Syrup of manna, N. F........fl.oz. 4
—Germ. Pharm.

## Syrup of Starch Iodide.

	gr. 30 gr. 250
Ether,	8
Water,	
Sugar	of each sufficient

Dissolve the iodine in ether, pour the solution on the starch and triturate until all the ether has evaporated. Then transfer the mixture to a porcelain capsule and heat on a water bath for one-half hour, stirring very frequently. At first considerable iodine vapor is evolved, but this soon ceases. From der. Now add the alcohol and the remainder the soluble starch iodide thus formed, the syrup may be prepared by dissolving 84 gr. in 7½ fluidounces hot water, and in the solution dissolving 14 av. ounces of sugar.

This syrup represents 1-10 per cent of iodine.

### Syrup, Strengthening, Thompsonian.

Comfreyav.oz.	2
Elecampaneav.oz.	1
Hoarhoundav.oz.	1/2
Waterfl.oz.	
Beth root, powdergr.	120
Brandyfl.oz.	8
Sugarav.oz.	8

Boil the first three drugs with the water until 24 fluidounces of liquid can be obtained, strain, add the remaining ingredients, and shake occasionally until the sugar is dissolved.

## Syrup of Strychnine Sulphate.

Strychnine sulphategr.	
Distilled waterfl.dr.	1/2
Simple syrup, enough to makefl.oz.	

Dissolve the strychnine in the water and add the syrup.—Codex.

## Syrup of Superphosphate of Iron.

Add freshly precipitated iron phosphate (see manner of preparation under Syrup of Iron and Ammonium Phosphate) to saturation to a boiling solution of glacial phosphoric acid. On concentrating and cooling, the product forms a soft mass, which is freely soluble in water in all proportions and is free from inky taste.

The syrup may be prepared by dissolving 5 gr. of this substance in a fluidram of simple syrup.

## Syrup of Sweet Gum.

See Syrup of Liquidambar.

## Syrup of Tar, Compound.

Fluid extract of licoricefl.oz.	2
Paregoric fl.oz.	
Syrup of ipecacfl.oz.	2
Glycerite of tarfl.dr.	
Syrup of tolu, enough to make . fl.oz.	16

#### Syrup of Tartaric Acid.

Tartaric acidgr.	100
Distilled waterfl.dr.	2
Simple syrupfl.oz.	
	7.7

Dissolve the acid in the water and add to the syrup.

If this solution be flavored with spirit of

lemon or spirit of orange, it may be called syrup of lemon or syrup of orange.—Codex.

#### Syrup of Tolu.

The following is employed very largely and furnishes an excellent product:

Tincture of tolu	.fl.oz.	2
Magnesium carbonate	gr.	120
Sugar	.av.oz.	281/2
Water	fl.oz.	16

Triturate the tincture with the magnesium compound and 2 ounces of sugar to a smooth paste, gradually add the remainder of the sugar, stirring constantly meanwhile, filter, and in the filtrate dissolve the remainder of the sugar by agitation or percolation.—U. S. P. 1870.

### Syrup of Trifolium, Compound.

(Syrup of Red Clover Blossom.—Syrup of Red Clover, Compound.)

Fluid extract of trifolium blos-	
somsfl.oz.	1
Fluid extract of berberis aqui-	
foliumfl.dr.	4
Fluid extract of poke rootfl.dr.	4
Fluid extract of burdock fl.dr.	4
Fluid extract of cascara amarga fl.dr.	4
Fluid extract of stillingiafl.dr.	4
Fluid ext. of prickly ash bark.fl.dr.	1
Potassium iodidegr. 1	128
Simple syrup, enough to make fl.oz.	16

#### Syrup of Turpentine.

-	
Gum turpentinegr.	320
Magnesium carbonate gr.	
Tincture of tolufl.dr.	21/2
Glycerin	12
Gum arabicav.oz.	
Sugarav.oz.	13
Water, enough to makefl.oz.	16

Triturate the first three together, add 6 fluidounces of water, filter, add the other ingredients, shake until dissolved, and strain.

## Syrup of Uva Ursi, Compound.

Fluid extract of uva ursifl.oz. 1
Fluid extract of buchufl.oz. 1
Fluid extract of cubebs fl.oz. 1
Sweet spirit of nitrefl.oz. 1
Simple syrupfl.oz. 8

## Syrup of Valerian, Compound.

, ·=	
Fluid extract of valerian fl.oz.	4
Fluid extract of scullcapfl.oz.	3
Tincture of hyoscyamusfl.oz., &	ટ
Spirit of wintergreenfl.dr.	1
Simple syrup, enough to make. fl.oz. 16	å

### Syrup of White Pine, Compound.

Reduce the vegetable drugs to moderately coarse powder, moisten the powder with a menstruum composed of 1 volume of alcohol and 3 volumes of water, and macerate for 12 hours. Then percolate with the same menstruum until 8 fluidounces of tincture have been obtained, in which dissolve the sugar and the morphine sulphate; lastly, add the chloroform, and sufficient syrup to make 16 fluidounces, and strain.—N. F.

### Syrup of Wild Cherry, Compound.

Wild cherryav.oz.	. 2
Spikenardav.oz.	. 1/3
Ipecacgr.	. 160
Bloodroot gr.	
Tincture of opiumfl.dr.	
Sugarav.oz.	
Alcohol,	
Water of each suffic	cient

Mix the drugs and reduce to tolerably fine powder and extract them by percolation in the usual way, using a menstruum consisting of alcohol and water in the proportions of 1 of the former and 2 of the latter by measure, obtaining 10 fluidounces of percolate; in this dissolve the sugar by agitation or percolation.

## Syrup of Wild Cherry and Hoar-hound.

Wild cherry bark, coarse pow-	
derav.oz.	4
Hoarhoundav.oz.	1
Glycerinfl.oz.	1
Alcohol fl.oz,	
Sugarav.oz.	
Watersufficie	

Mix the glycerin and alcohol with 8 fluidounces of water; moisten the wild cherry and hoarhound with 2 fluidounces of this mixture, pack in a cylindrical percolator tightly covered; after 24 hours' maceration proceed with percolation, using the remainder of the

menstruum, and afterward sufficient water to make 10 fluidounces of percolate; in this dissolve the sugar by agitation, without heat, and strain.

## Syrup of Wintergreen.

Wintergreen leavesav.oz.	2
Boiling watersufficie	nt
Sugarav.oz.	12
Oil of wintergreendrops	5

Infuse the leaves in enough water to complete 8 fluidounces of infusion; in it dissolve the sugar with the aid of heat, and when the syrup is nearly cold add the oil previously dissolved in a small amount of alcohol.

## Syrup of Wormwood. (Syrup of Absinthium.)

Wormwoodav.oz.	1
Boiling waterfl.oz.	8
Sugarav.oz.	14

Infuse the drug with the water for 12 hours, strain with expression, and filter, and in the filtrate dissolve the sugar by agitation or percolation.

## Syrup of Yellow Dock, Compound.

(Compound Syrup of Rumex.—Scrofulous Syrup.)

Yellow dockgr. 480
False bittersweetgr. 240
American ivy barkgr. 120
Figwort
Alcohol,
Waterof each, sufficient
Sugar

Mix the drug, reduce to fine powder and extract by percolation so as to obtain 10 fluidounces of product, using a menstruum composed of 1 part of alcohol by measure to 2 of water; in this percolate dissolve the sugar by agitation or percolation.—Eclectic.

#### Syrup of Yerba Santa.

Fluid extract of eriodictyonfl.oz.	
Calcined magnesiaav.oz.	1/2
Calcined magnesiaav.oz. Waterfl.oz.	71/2
Sugarav.oz.	

Mix the fluid extract with the calcined magnesia and add the water gradually, with constant stirring; let it stand 24 hours and filter; add the sugar and dissolve with the aid of gentle heat.

## Syrup of Yerba Santa, Aromatic.

(Aromatic Syrup of Eriodictyon.)

Fluid extract of yerba santafl.dr.	4
Solution of potassa fl.dr.	3
Compound tincture of carda-	
momfl.oz.	1
Oil of sassafrasdrops	4
Oil of lemondrops	
Oil of clovesdrops	
Alcoholfl.dr.	
Sugarav.oz.	
Water, enough to makefl.oz.	
, 3	

Mix the fluid extract and solution of potassa, then add 12 fluidrams of water previously mixed with the compound tincture of cardamom, and afterwards add the oils dissolved in the alcohol. Shake the mixture thoroughly, then filter it, and pour enough water through the filter to obtain 6 fluid-ounces of filtrate. Pour this upon the sugar contained in a bottle, and dissolve it by placing the bottle in hot water, frequently agitating. Lastly, cool the product and add enough water, passed through the filter previously used, to make 16 fluidounces.—N. F.

Other syrups are mentioned in Parts II, IV and V.

## Tablets, Antiseptic, Dr. C. M. Wilson's.

Each tablet should contain:

Corrosive sublimate.						.gr.	7.7
Ammonium chloride							

If added to 16 fluidounces of water, the product will represent a solution of 1 in 1000.

#### Tallows.

Refer to Suets.

#### Teas.

Refer to Species.

#### Thompsonian Remedies.

- "Number one" is lobelia.
- "Number two" is capsicum.
- "Number three" is bayberry, sumach and capsicum.
- "Number four" is the "bitters." See Bitters.
- "Number five" is restorative syrup or cordial. See Syrups.
- "Number six," hot drops or rheumatic liquid drops. See Tincture of Capsicum and Myrrh. latter.

" Third	Preparation	:''
---------	-------------	-----

Lobelia seed	gr. 480
Capsicum	
Cypripedium, powder, an	even teaspoonful.
Number six	fl.oz. 12

Macerate and retain liquid on dregs.

Brown lobelia is lobelia seed.

Green lobelia is lobelia herb.

Camphor julep. See Camphor.

Conserve of hollyhock. Refer to Confections.

Ointment, astringent. See Ointments. Pills, iron, compound. See Pills. Poultice, charcoal:

Charcoalav.oz. 3
Gingerav.oz. 1
Bayberryav.oz. 1
Slippery elmav.oz. 3
Water, hotsufficient

Powder, composition. See N. F.
Powder, cough. See Powders.
Powder, nerve. See Powders.
Salve. See Ointments.
Syrup of bayberry. See Syrups.
Syrup of garlic, artificial. See Syrups.
Syrup of lobelia. See Syrups.
Syrup or cordial, restoratives. See Syrups.
Syrup, strengthening. See Syrups.

#### Tincture of Aconite Leaf.

Aconite	leaves,	powder .	 	.gr. 960

Extract the drug by percolation so as to obtain 16 fluidounces of product.—U. S. P. 1870.

This preparation must not be confounded with tincture of aconite (root) U. S. P. or Fleming's tincture of aconite N. F.

#### Tincture of Asarum.

Canada snake-root, powder...av.oz. 4
Alcohol, sufficient to make....fl.oz. 16

#### Tincture of Arbor Vitæ. (Tincture Thuja.)

Arbor vitæ, fresh tops.......av.oz. 3½
Alcohol, enough to make.....fl.oz. 16

Macerate the arbor vitæ with 16 fluidounces of alcohol for 7 days, then decant the liquid, express the residue, treat the latter with fresh portions of alcohol, expressing forcibly each time, until 16 fluidounces of liquid are obtained, and finally filter the latter.

## Tincture of Asafetida, Compound.

Asafetidagr.	200
Lupulingr.	
Stramonium seedgr.	200
Valerian root gr	
Alcohol	

Mix the drugs, reduce to coarse powder, add the alcohol, macerate for 14 days, strain, express and filter.—Eclectic.

## Tincture of Avena Sativa, Homeopathic. (Tincture of Oats.)

Oats, unhuskeda	v.oz. 8
Potassium carbonate,	
Water,	
Alcohol of each su	fficient

Grind the oats to moderately fine powder, moisten with a 5 per cent aqueous solution of potassium carbonate—first warmed to 45 degs. C.—macerate for 3 hours, pack in a percolator and add alcohol until 16 fluid-ounces of product are obtained.

## Tincture of Bees, (Honey.) (Tinctura Apis Mellificæ.)

Collect quantity of living honey-bees in a bottle, agitate the latter so as to irritate them and then cover them with alcohol; after a few days, the liquid is ready for use.— Eclectic.

## Tincture of Black Cohosh, Compound. (Compound Tincture of Cimicifuga.)

Tincture of black cohosh . . . . . fl.oz. 8¼
Tincture of blood root, U. S. P. fl.oz. 6
Tincture of poke root . . . . . fl.oz. 1¾
—Eclectic.

## Tincture of Bloodroot, Compound.

(Emetic Tincture.)

Blood rootav.oz.	1
Lobelia herbav.oz.	
Skunk cabbageav.oz.	1
Diluted alcoholsufficien	ıt

Extract the mixed drugs in fine powder by percolation with diluted alcohol so as to obtain 16 fluidounces of product.—Eclectic.

#### Tincture of Blue Cohosh.

Blue cohosh, fine	powder	av.oz. 31/4
Alcohol		sufficient

Extract the drug by percolation with alcohol so as to obtain 16 fluidounces of product.—Eclectic.

## Tincture of Blue Cohosh, Compound.

Blue cohosh, fine powdergr.	640
Ergot, fine powdergr.	
Water pepper, fine powdergr.	
Oil of savinfl.dr.	21/2
Alcoholsuffic	ient

Extract the mixed drugs by percolation with alcohol, so that the percolate with the oil added will make 16 fluidounces.—
Eclectic.

#### Tincture of Blue Flag.

Blue flag,	fine powder	av.oz. 31/4
Alcohol	s	ufficient

Extract the drug by percolation with alcohol, so as to obtain 16 fluidounces of product.—Eclectic.

#### Tincture of Buchu.

Buchu,	coarse	powderav.oz.	2
Diluted	alcoho	ol sufficier	at

Extract the drug by percolation so as to obtain 16 fluidounces of product.—Brit. Pharm.

#### Tincture of Burdock Seed.

Burdock see	d, ground	av.oz. 41/2
Water,		,-
Alcohol	of eac	h sufficient

Mix the liquids in the proportion of 1 by measure of the water to 3 of the alcohol, and percolate the drug in the usual way, until 16 fluidounces of percolate are obtained.

## Tincture of Cacao. (Tincture of Theobroma.)

Cacao beans, freshly roastedav.oz.	
Cinnamon	
Diluted alcohol, enough to make fl.oz.	

Reduce the cacao beans and the cinnamon to moderately fine powder, add 16 fluid-ounces of diluted alcohol, macerate for 4 or 5 days, agitating occasionally, transfer to a glass percolator and percolate, adding sufficient more of the diluted alcohol to make the percolate, including the tincture of vanilla, measure 16 fluidounces.

#### Tincture of Cactus Grandiflorus.

	flowers				
cact	us grand	ifloru	s	av.oz.	41/2
Alcoho	ol			fl.oz.	16

Macerate for 14 days, occasionally agitating, express and filter.—Eclectic.

## Tincture of Cajuput, Compound.

Oil of cajuputfl.oz.	4
Oil of peppermint fl.oz.	
Oil of clovesfl.oz.	
Alcohol fl.oz.	

#### Tincture of Calamus.

Calamus, coarse powder av.oz.	3
Water fl.oz. Alcohol fl.oz.	41/2
Alcoholfl.oz.	13

Mix, macerate for 7 days, agitating occasionally, strain with expression and filter.—Germ. Pharm.

## Tincture of Carduus Mariæ, Rademacher's. (Tincture of Mary Thistle.)

Carduus Mariæ fruit, wholeav.oz.	10
Alcohol fl.oz.	12
Distilled waterfl.oz.	10

Macerate for 8 days, then filter. The fruit is used whole, owing to its highly mucilaginous character.

### Tincture, Carminative.

Cardamom seed, bruisedgr.	480
Tincture of gingerfl.oz.	
Oil of cinnamonm.	80
Oil of clovesm.	80
Oil of carawaym.	80
Alcohol, enough to makefl.oz.	16

Macerate the cardamom with 12 fluidounces of alcohol for 7 days, decant the liquid, express the residue, filter the entire liquid, add the oils to the filtrate, and finally add the remainder of the alcohol.—Brit. Form.

### Tincture of Cascara Sagrada.

Cascara sagrada, coarse powder av.oz.	3
Water	
Alcohol fl.oz.	111/

Mix, macerate for 10 days, agitating occasionally, express, and filter.—Codex.

#### Tincture of Castor.

Castor (	Russian	preferred)	av.oz. 11/4

Reduce the castor to as fine a condition as possible, macerate with the alcohol for 14 days, occasionally agitating, express, and filter, adding enough alcohol through the filter to make the liquid measure 16 fluid-ounces.—Eelectic.

### Tincture of Castor, Ammoniated.

Castorgr.	480
Asafetidagr.	240
Spirit of ammoniafl.oz.	16

Reduce the drugs to coarse powder, add the spirit, macerate for 7 days, agitating occasionally, and express.—Eclectic.

## Tincture of Celandine, Rademacher's.

(Tincture of Chelidonium.)

Fresh	herb	of	chelidonium	
maji	ıs		av.oz.	10
			fl.oz.	

Contuse the herb tea to a pulp, add the alcohol, macerate for 8 days, express, and filter.

#### Tincture of Chinoidin.

Chinoidin		av.oz.	11/2
Alcohol			
Hydrochloric ac			
Water			
	erm, and Swed		

## Tincture of Chloroform, Compound.

Chloroform	fl.oz. 1
Alcohol	
Compound tincture of	carda-
mom	
	-Brit, Pharm.

#### Tincture of Cinnamon, Compound.

Cinnamongr. 24	0
Cardamom	
Prickly-ash berriesgr. 9	0
Gingergr. 9	0
Diluted alcohol sufficier	ıt

Extract the mixed drugs in fine powder by percolation so as to obtain 16 fluidounces of product.—Eclectic.

## Tincture of Cocculus Indicus, Homeopathic.

Cocculus Indicus, powderav.oz.	3
Alcoholfl.oz.	16
Waterfl.dr.	10

Place all in a well-corked bottle, and macerate for 8 days, shaking the bottle well twice a day; then press out and filter. This makes the mother tincture.

#### Tincture of Cochineal.

Cochineal, powder.......av.oz. 2 Diluted alcohol, enough to make fl. oz. 16

Extract the drug by percolation or maceration.—Brit. Pharm.

The product may be used for coloring elixirs and other preparations.

## Tincture of Cochineal, Rademacher's.

(Tinctura Coccionellæ.)

Cochineal, coarse powder.....av.oz. 1 

Macerate for 3 days, agitating occasionally, and filter.

## Tincture of Colchicum, Compound.

Tincture of colchicum seed . . . . fl.oz. 8 Tincture of black cohosh.....fl.oz. 8 -Eclectic.

## Tincture of Colocynth.

Colocynth, with seeds, cut Alcohol.....sufficient

Percolate alcohol through the drug so as to obtain 16 fluidounces of tincture. —Germ. Pharm.

## Tincture of Colocynth Seed, Rademacher's.

Colocynth seed......av.oz. 3 Alcohol ......sufficient

Wash the seed with water, dry it, reduce to coarse powder, add 161/2 fluidounces of alcohol. Macerate for 14 days, agitating, express, filter, and add enough alcohol to the filtrate to make 16 fluidounces.

#### Tincture of Conium.

Diluted alcohol ......sufficient

Extract the drug by percolation so as to obtain 16 fluidounces of product, adding the acid to that portion of the diluted alcohol which is used for moistening the drug.-U. S. P. 1880.

## Tincture of Convallaria. (Tincture of Lily of the Valley.)

Lily of the valley flowers and stalks, dried, coarse powder..av.oz. 2 Diluted alcohol .....sufficient

Extract the drug by percolation, so as to make 16 fluidounces of product.—Brit. Form.

## Tincture of Copper Acetate, Rademacher's.

Copper sulphate, puregr. Lead acetate, puregr.	
Distilled water fl.oz. Alcohol fl.oz.	8 1/2

smooth paste is formed, transfer this to a copper vessel, add the water, heat to boiling, allow to cool, add the alcohol, set aside for 4 weeks, agitating frequently, and filter.

A more expeditious process is the following:

Copper acetate, crystallized....gr. 480 Distilled water.....fl.oz. Alcohol . . . . . . . . . . . . . . . . . fl.oz.

Dissolve the acetate in the water previously warmed and filter.

#### Tincture of Corydalis. (Tincture of Turkey Corn.)

Turkey corn, fine powder.....av.oz. 31/4 Diluted alcohol ......sufficient

Extract the drug by percolation with diluted alcohol so as to obtain 16 fluidounces of product.—Eclectic.

## Tincture of Culver's Root. (Tincture of Leptandra.)

Culver's root.....av.oz. 3¼ Diluted alcohol . . . . . . . . sufficient

Extract the drug in moderately fine powder by percolation so as to obtain 16 fluidounces of product.-Eclectic.

#### Tincture of Digitalis, Ethereal.

Digitalis, cut fine.....av.oz. 11/4 Spirit of ether.....fl.oz. 16

Mix, macerate for 7 days, and filter.

#### Tincture of Ergot.

Ergot, powder ......av.oz. 4 Diluted alcohol .....sufficient

Percolate the drug so as to obtain 16 fluidounces of tincture.-Brit. Pharm.

#### Tincture of Eucalyptus.

Eucalyptus powder .....av.oz. 3 1/4 Alcohol.....sufficient

Extract the drug by percolation so as to obtain 16 fluidounces of product. - Brit. Form.

## Tincture of Golden Rod, Rademacher's. (Tinctura Virgæ Aureæ.)

Fresh herb of golden rods ....av.oz. 10 

Contuse the herb to a pulp, add the alco-Triturate the two salts together until a hol, macerate for 8 days, express, and filter-

## Tincture of Golden Seal, Compound.

Tincture	of	golden	seal,	U.		
S. P					.fl.oz.	91/2
S. P Tincture	of	lobelia			fl.oz.	$6\frac{1}{2}$
				1	Eclecti.	r´¯

## Tincture of Guaiac, Aromatic.

Refer to Greenhow's Cholera Mixture.

## Tincture of Hips, Rademacher's.

(Tinctura Cynosbati.)

Fresh rose	hips,	cut fine.	 av. oz. 2
Alcohol			 .sufficient

Macerate the hips with 12 fluidounces of alcohol, agitating frequently, express, filter and add enough alcohol to the filtrate to make 12 fluidounces.

### Tincture of Iodine, Compound.

Iodinegr.	240
Potassium iodidegr.	480
Alcoholfl.oz.	

Mix and dissolve.-U. S. P. 1870.

This must not be confused with the compound solution of iodine of the present pharmacopæia.

## Tincture of Iron, Compound.

Tincture of ferrated extract of
apples, N. Ffl.oz. 8
Vinous tincture of rhubarbfl.oz. 8
Tincture of nux vomicafl.oz: 1
See also next formula.

# Tincture of Iron, Athenstædt, Compound. (Aromatic Tincture of Iron. —Athenstædt's Tincture.)

Soluble oxide of irongr.	330
Distilled waterfl.oz.	19
Simple syrupfl.oz.	6
Alcoholfl.oz.	61/2
Citric acidgr.	30
Tincture of orange peelm.	50
Aromatic tincturedrops	12
Tincture of cinnamondrops	12
Tincture of vanilladrops	12
Acetic etherdrop	1
•	

Dissolve the iron salt in the water, then add the other ingredients and filter.

The iron oxide used for the above should represent 10 per cent of metallic iron. If it be weaker, a proportionately larger amount should be employed, and slightly decreasing the amount of syrup subsequently added.

## Tincture of Iron Acetate, Rademacher's.

Iron sulphate, puregr.	656
Lead acetate, puregr.	
Diluted acetic acidfl.oz.	6
Distilled waterfl.oz.	
Alcoholfl.oz.	

Triturate the two salts together to a pasty mass, introduce this into an iron vessel, add the water and acid, heat to boiling, allow to cool, transfer to a large flask, add the alcohol, set the flask, loosely stoppered, aside for several months, agitating occasionally until the liquid has acquired a light red tint, and finally filter.

A more expeditious process for making this preparation would be by the use of solution of iron tersulphate, as follows:

Solution of iron tersulphatefl.oz.	21/2
Distilled waterfl.oz.	
Lead acetate, pureav.oż.	
Diluted acetic acidfl.oz.	
Alcoholfl.oz.	

Dissolve the lead acetate in the acid, add the iron solution previously mixed with the water, then gradually add the alcohol, set aside for one or two weeks and decant the clear liquid which is the finished product.

#### Tincture of Jaborandi.

Jaborandi, powder.	 	 ٠.	av.oz.	4
Diluted alcohol	 	 	sufficie	nt

Extract by percolation so as to obtain 16 fluidounces of product.—Brit. Pharm.

## Tincture of Kalmia. (Tincture of Sheep Laurel or Mountain Mint.)

Sheep la	aurel leaves,	ground	av.oz.	31/4
Diluted	alcohol		.sufficier	nt

Extract the drug by percolation so as to obtain 16, fluidounces of product.—Eclectic.

## Tincture of Lobelia, Compound. (King's Expectorant Tincture.)

,	
Lobelia (herb)gr.	120
Bloodrootgr.	120
Skunk cabbagegr.	120
Canada snake rootgr.	120
Pleurisy rootgr.	120
Water,	
Alcoholof each, suffice	ient

Mix the drugs and reduce to fine powder; mix the alcohol and water in the proportion of 3 of the former to 1 of the latter, and

extract the mixed drugs by percolation with this menstruum so as to obtain 16 fluidounces of product.—Eclectic.

#### Tincture of Lemon Peel.

Lemon peel, fresh, sliced thin..av.oz. 2 Diluted alcohol.....sufficient

Mix the peel with 16 fluidounces of diluted alcohol, macerate for 7 days, agitating occasionally; strain, express and filter, adding enough menstruum to make a total filtrate of 16 fluidounces.—Brit. Pharm.

## Tincture of Lobelia and Capsicum, Compound. (Antispasmodic Tincture.)

Lobeliaav.oz. 1
Capsicumav.oz. 1
Skunk cabbageav.oz. 1
Diluted alcoholsufficient

Mix the drugs in moderately fine powder and extract by percolation with diluted alcohol so as to obtain 16 fluidounces of product.

—Eclectic.

## Tincture of Lupulin.

T 11														0.4
Lupulin	٠	•	•	٠	•	•	•		•		•	•	av.oz.	23/4
Alcohol								ı					.fl.oz.	16

Macerate for 7 days, shaking occasionally, and filter, adding enough alcohol through the filter to make 16 fluidounces.—Eclectic.

#### Tinctures, Mother.

The "mother tinctures" of the homeopathists are the basic preparations of crude drugs from which the attenuations or "potencies" are formed. They should be made from freshly gathered drug, which is to be contused and then macerated for 14 days with a mixture of alcohol and water in such proportion that the percentage of spirit in the product shall be the same as in the diluted alcohol, and the drug shall bear to the finished product the relation of 1 to 10, the former being calculated as dry drug. In other words a portion of the drug must first be dried, then bruise remainder to a pulp, add 5 times its weight (as dry drug) of alcohol and then enough water to make up twice the weight of the alcohol used, strain and express at the end of 14 days, add enough of diluted

alcohol through the strainer to make up the weight of drug and liquid just expressed, and finally filter the whole.

Inasmuch as fresh drugs are not generally obtainable, dry drugs are usually employed in their stead.

Mother tinctures may be conveniently prepared from ordinary tinctures by dilution with diluted alcohol.

Only distilled water should be used for homeopathic preparations; also the best alcohol should be used for these. Good "cologne spirit" is generally dispensed "homeopathic alcohol."

## Tincture of Mugwort Root, Rademacher's. (Tincture of Artemisia.)

Mugwort root, cut fine.....av.oz. 3 Diluted alcohol.....fl.oz. 15

Mix, macerate for 3 days, express, and filter.

#### Tinctures, Normal.

The use of the term "normal" as applied to tinctures is confined to Eclectic pharmacy, and it signifies that the product, the "normal tincture," represents the dry drug, weight for weight. They may be prepared from fresh or from dry drugs or by the use of a greater or less proportion of alcohol in the menstruum, but the strength of the product is always based upon the drug in a dry condition. Good fluid extracts may always be dispensed in place of the corresponding "normal tinctures."

Normal tinctures are designated by Eclectics for brevity's sake, as "tinctures ( $\frac{n}{l}$ )." Tinctures one-half the strength of normal tinctures are called semi-normal ( $\frac{n}{l}$ ); one-fifth as quinti-normal ( $\frac{n}{l}$ ); one-tenth as decinormal ( $\frac{n}{l}$ ), etc.

## Tincture of Nux Vomica, Rade-

macher's. (Tinctura Nucum Vomicarum.—Tinctura Strychni.)

The Rademacher tincture is to be prepared from grated or rasped nux vomica by maceration with diluted alcohol. Inasmuch as it is approximately of the same relative strength as the preparation of the U. S. P., the latter should always be dispensed for it.

#### Tincture of Opium, Ammoniated.

Opiumgr.	80
Spanish saffrongr.	
Benzoic acidgr.	
Oil of anise	
Stronger water of ammoniafl.oz.	31/
Alcoholsuffic	

Mix the first five ingredients with 13 fluidounces of alcohol, macerate for 7 days, agitating occasionally, express, filter, and add enough alcohol to the filtrate to make 16 fluidounces.—Brit. Pharm.

## Tincture of Opium, Benzoated.

(Anisated Tincture of Opium.—Compound Tincture of Camphor.)

These are various terms used to designate paregoric or camphorated tincture of opium.

A rapid process for making this preparation is as follows:

Tincture of opiumfl.dr.	5
Spirit of camphorfl.dr.	5
Oil of anisefl.dr.	
Benzoic acidgr.	
Diluted alcohol, enough to	
makefl.oz.	16
M: 121 1 C1	

Mix, dissolve, and filter.

## Tincture of Opium Compound.

Tincture of opiumfl.oz.	3
Tincture of capsicumfl.oz.	3
Spirit of camphorfl.oz.	
Chloroformfl.dr.	
Alcohol, enough to makefl.oz.	

#### Tincture of Opium, Crocated.

(Tinctura Opii Crocata.—Tincture of Opium and Saffron.—Sydenham's Laudanum.—Compound Wine of Opium.)

Opium, powderav.oz.	11/2
Spanish saffronav.oz.	1/2
Cloves, bruisedgr.	90
Cassia bark, coarse powdergr.	90
Alcoholfl.oz.	634
Waterfl.oz.	91/4

Mix all, macerate for 7 days, agitating occasionally, and filter.—Germ. Pharm.

### Tincture of Phosphorus, Compound.

Phosphorusgr. 8	3
Chloroformfl.dr. 14	1
Alcohol, enough to makefl.oz. 10	)

Place the phosphorus in a stoppered bottle, apply the heat of a water bath until

dissolved, and then add the alcohol, then shake well.

This tincture should be protected from the light, in accurately stoppered bottles.

Each fluidram contains is grain of phosphorus.—Brit. Pharm.

## Tincture of Poison Oak. (Tincture of Poison Ivy.)

Fresh leaves of .rhus	toxico-	
dendron	av.oz. 9	,
Alcohol	fl.oz. 6	;

Macerate for 14 days, express and filter in a well-covered funnel—Eclectic.

## Tincture of Podophyllum.

Podophyllum,	fine	powder	av.oz. 31/4
Alcohol			sufficient

Extract the drug by percolation with alcohol so as to obtain 16 fluidounces of product.

—Eclectic.

#### Tincture of Poke Root.

Poke root,	fine powder	 av.oz. 31/2	1
Diluted ald	cohol	 .sufficient	

Extract the drug by percolation with diluted alcohol so as to obtain 16 fluidounces of product.—Eclectic.

### Tincture of Poke Root, Compound.

Fluid extract of poke fl.oz. Fluid extract of cardamom fl.dr.	
Diluted alcohol, enough to make.fl.oz.	16
Mix and filter.	

#### Tincture of Prickly Ash Berries.

Prickly ash berries, fine powder.av.oz. 4½ Diluted alcohol.....sufficient

Extract the drug by percolation with diluted alcohol so as to obtain 16 fluidounces of product.—Eclectic.

#### Tincture of Pulsatilla.

Pulsatilla herb,	fresh	 .av.oz. 81/2
Absolute alcoho	1	 sufficient

Cut the herb into small pieces and add absolute alcohol enough so that the product will measure 16 fluidounces; macerate for 14 days, express and filter.—Eclectic.

#### Tincture of Quinine.

Quinine sulphategr.	128
Tincture of bitter orange peelfl.oz.	
-Brit. Pha	

## Tincture of Quinine, Ammoniated.

Quinine sulphategr.	128
Water of ammoniafl.oz.	2
Diluted alcoholfl.oz.	14

Dissolve the sulphate of quinine in the alcohol with aid of a gentle heat and add the ammonia.—Brit. Pharm.

## Tincture of Quillaja. (Tincture of Soap Bark.)

Quillaja, coarse powderav.oz. Alcoholfl.oz.	31/4
Alcoholfl.oz.	534
Watersufficier	ıt

Boil the quillaja with 13 fluidounces of water for 15 minutes, strain, wash the residue on the strainer, with 1½ fluidounces of water, boil the strain liquid down to 10 fluidounces, allow to cool, add the alcohol, filter, and through the filtrate add enough water to make the filtrate measure 16 fluidounces.—U. S. P.

#### Tincture of Rhubarb, Compound.

Rhubarbgr. 384
Dogsbane (apocynum androsæm).gr. 192
Golden sealgr. 192
Gentiangr. 192
Prickly ash berriesgr. 192
Diluted alcoholsufficient

Mix the drugs, reduce to fine powder, and extract with diluted alcohol, by percolation so as to obtain 16 fluidounces of product.— Eclectic.

## Tincture of Rhubarb, Koehlreuter's.

Rhubarb, cut fineav oz.	21/2
Bitter orange peel, cut finegr.	
Centaury, cut finegr.	180
Fennel, crushedgr.	110
Distilled waterfl.oz.	9
Alcoholfl.oz.	71/2
THEORIE	. /2

Mix and macerate for 8 days, strain and filter.

## Tincture of Saffron. (Tincture of Crocus.)

	•	
Saffron	av.oz. í	13/
D'1-1 1 1 1 1	CC 1	- /4
Diluted alcohol	sufficien	t

Extract the drug by percolation so as to obtain 16 fluidounces of product.—U. S. P.

#### Tincture of Savin.

Savin, coarse	powder,	.av.oz. 2
Diluted alcoh	ol	sufficient

Extract the drug by percolation so as to obtain 16 fluidounces of tincture.—Br.c. Pharm.

## Tincture of Savin, Compound.

Fluid extract of savinfl.oz.	L
Tincture of castorfl.oz.	71/2
Tincture of myrrhfl.oz.	71/2

## Tincture of Senna, Compound. (Elixir Salutis.—Elixir of Health.)

Alexandria senna, cutgr.	480
Jalap, finely powderedgr.	
Coriandergr.	120
Raisins, deprived of seedsav.oz.	
Diluted alcoholfl.oz.	16

Macerate for 7 days, shaking occasionally, and filter.—Eclectic.

## Tincture of Serpentaria, Compound. (Sudorific Tincture.)

Serpentariagr.	160
Ipecacgr.	
Spanish saffrongr.	
Camphorgr.	
Opiumgr.	
Diluted alcoholfl.oz.	

Macerate the finely powdered drugs with the diluted alcohol for 7 days, agitating occasionally, and filter.—Eclectic.

## Tincture of Shepherd's Purse, Rademacher's. (Tinctura Bursæ Pastoris.)

Shepherd's purse herb, freshly	
gatheredav.oz. 1	0
Alcoholfl.oz. 1	2

Contuse the herb to pulp, add the alcohol, macerate for 7 days, express and filter.

#### Tincture of Skunk Cabbage.

Extract the drug in fine powder by percolation with diluted alcohol so as to obtain 16 fluidounces of product.—Eclectic.

#### Tincture of Stavesacre.

Stavesacre seed, fine powder...av.oz. 11
Absolute alcohol.........sufficient

Percolate the drug with the absolute alcohol so as to obtain 16 fluidounces of product.— Eclectic.

#### Tincture of Stillingia.

Jullingia, fine	powder	av.oz. 3
Diluted alcoho		

Extract the drug by percolation with diluted alcohol so as to obtain 16 fluidounces of product.—Eclectic.

## Tincture of Strychnine.

Strychnine (alkaloid)gr.	12
Alcoholfl.oz.	4

Agitate occasionally until dissolved.—Brit. Pharm.

#### Tincture of Strychnine, Compound.

Strychnine (alkaloid)gr.	16
Acetic acidfl.dr.	4
Compound tincture of carda-	
momfl.dr.	4
Waterfl.oz.	
Alcoholfl.oz.	71/2
	,-

Dissolve the strychnine in the alcohol and acetic acid, add the remaining ingredients, and filter.—Eclectic.

#### Tincture of Sulphur.

Washed s	ulphur	290
Absolute	alcoholfl.oz.	16

Mix, macerate for 4 days, agitating occasionally, and filter.—H.

## Tincture of Sulphur, Homeopathic.

Washed	sulphur.	 	 .av.oz.	11/2
Alcohol		 	 fl.oz.	16

Mix, macerate for 8 days, shaking twice a day, decanting the clear liquid, and filtering. This is considered equal to the first centesimal potency.

#### Tincture of Tolu.

Tolu	enough to makefl.o	z. 13/
Alcohol,	enough to makefl.o	z. 16

Mix, agitate occasionally until dissolved, and filter.—U. S. P.

#### Tincture of Valerian, Ethereal.

Valerian	powder	av.oz.	21/2
Spirit of	ether	sufficie	nt

Mix the drug with 15 fluidounces of spirit, macerate for 7 days, agitating occasionally, express, add enough spirit of ether to make 15 fluidounces, and filter in a closely covered funnel.—Garm, Pharm.

## Tincture of Wahoo. (Tincture of Eu-

337-1 11-		0.7
wanoo bark,	powder	av.oz. 314
Alcohol		sufficient

Extract the drug by percolation so as to obtain 16 fluidounces of product.—Brit. Form.

## Tincture of Water Pepper.

Water-pepper	.av.oz. 41
Diluted alcohol	sufficient

Extract the powdered drug by percolation with diluted alcohol so as to obtain 16 fluid-ounces of product.—Eclectic.

#### Tincture of Witch Hazel Bark.

Hamamelis bark, No. 20 pow-	
derav.oz.	13/4
Diluted alcohol, enough to make, fl.oz.	16

Extract the drug by percolation.—Brit. Form.

## Tincture of Wormwood. (Tincture of Absinthium.)

Wormwoodav.oz.	3
Waterfl.oz.	31/2
Waterfl.oz. Alcoholfl.oz.	121/2

Mix, macerate for 7 days, agitating occasionally, express, and filter —Germ. Pharm.

This may also be prepared by extracting the powdered drug by percolation.

## Tincture of Wormwood, Compound.

(Bitter Stomach Drops.)

Wormwoodgr.	520
Blessed thistle gr.	
Galangal rootgr.	
Orange berriesgr.	
Diluted alcohol suffic	

Mix the drugs, reduce to powder, and extract by percolation with diluted alcohol so as to obtain 16 fluidounces of product..

#### Tisanes.

This is the French appellation for the class of preparations known to us as "infusions."

#### Transfusion Fluid.

#### A. (Billroth's.)

Sodium phosphate	gr.	3
Sodium carbonate		20
Ammonium carbonate	.gr.	20
Sodium chloride		
Alcohol	m.	160
Distilled water, enough to		00
make	fl.oz.	20

### B. (Little's.)

Codium phosphate	or.	3
assium chloride	gr.	6
.iodium carbonate		
Sodium chloride	gr.	60

Listilled water, enough to make.fl.oz. 20

C. (Weber's.)	
Sodium bicarbonategr.	6
Calcium chloridegr.	6
Potassium chloride gr.	6
Sodium chloridegr.	480
Distilled water, enough to	
makefl.oz.	- 8

For use, dilute 1 fluidounce of this solution with water at 50 degrees C., so as to make 16 fluidounces.

#### Turpentine, Canada.

This is Balsam of Fir.

#### Turpentine Venice, Factitious.

Gum turpentineav.oz.	10
White resinav.oz.	3
Oil of turpentine fl. oz.	10

Melt the resin, add the turpentine, allow it to melt, then add the oil, and strain.

## Vinegar of Bloodroot. (Vinegar of Sanguinaria.)

Bloodroot,	powderav.oz.	13/
Diluted ac	etic acidsufficier	nt

Extract the drug by percolation so as to obtain 16 fluidounces of product.—U. S. P. 1880 and N. F.

### Vinegar of Cantharides.

Cantharides, powdergr. 70	0
Glacial acetic acid fl.dr. 1	
Acetic acidsufficier	ıt

Mix 10 fluidounces of acetic acid with the glacial acid, add the drug, maintain at a temperature of 94 degrees C., for 2 hours, allow to cool, and transfer to a percolate; after the liquid has drained off, add enough acetic acid through the percolator to make the liquid measure 16 fluidounces.—Brit. Pharm.

## Vinegar of Ipecac.

Ipecac in No. 20 powdergr.	360
Diluted acetic acid, enough to	
make fl.oz.	16

Extract the drug by percolation.—Brit. Pharm.

## Vinegar of Lobelia.

Lobelia,	powderav.oz.	13/
Diluted	acetic acidsufficier	at

Extract the drug by percolation so as to obtain 16 fluidounces of product.—U. S P. 1880 and N. F.

#### Vinegar of Sabadilla.

	a, powderav.oz.	
Diluted	acetic acid fl.oz. 1	4
Alcohol		$^2$

Macerate for 7 days, then filter.

This preparation is useful for the extermination of vermin upon the body.

## Vinegar, White Wine, Imitation.

Acetic acidfl.oz.	16
Tartaric acidav.oz.	1
Acetic etherfl.dr.	4
White winefl.oz.	16
Waterfl.oz.	32

## Water, Acorn, Rademacher's. (Aqua Glandium.—Aqua Quercus.)

Acorns,	depriv	ed of	the h	ard	
outer h	ulls, in	coarse	powder	av.oz.	91
Alcohol.				fl.oz.	3
Water				fl. oz.	32

Mix, macerate for 24 hours, and then distill off 16 fluidounces.

### Water, Bromine.

—H.

## Water, Caraway.

Mix 1½ av. ounces of bruised caraway seed with 32 fluidounces of water and distill off 16 fluidounces.—Brit. Pharm.

A quicker process is to triturate 15 drops of oil of caraway with  $\frac{1}{2}$  av. ounce of purified talcum until well mixed, then add 16 fluid-ounces of distilled water, macerate for 6 hours, and filter.

#### Water, Carbolized.

Glycerite of carbolic acid, U. S. P. 1870fl. oz. 11/4
1870fl.oz. 1¼
Water
—U. S. P. 1870.

If it is not desired to keep the glycerite on hand, the above may be prepared (approximately) from 2 fluidrams of carbolic acid in the fluid form, as it is usually kept on hand by pharmacists for dispensing purposes; 1 fluidounce of glycerin, and enough water to make 16 fluidounces.

II.
Carbolic acid, liquefied ......fl.dr. 4
Distilled water .........fl.oz. 14½
—Germ. and Austr. Pharms,

## Water, Carminative. (Aqua Carminativa. —Wind Wasser.)

Oil of orangedrops	7
Oil of carawaydrops	
Oil of lemondrops	7
Oil of fenneldrops	7
Oil of corianderdrops	
Oil of spearmint drops	7
Alcohol fl.oz.	2
Chamomile waterfl.oz.	14

Mix the oils with the alcohol, add the water and filter.—H.

## Water, Castor, Rademacher's. (Aqua Castoreum.)

Canadian castor, cut fine, av.oz.	
Alcohol fl.oz.	3
Waterfl.oz.	

Mix, macerate for 12 hours, and distill off 16 fluidounces..

#### Water, Chamomile.

Chamomile							.av.oz.	11/2
Water							fl.oz.	48

Mix, macerate for 24 hours, and distill off 16 fluidounces.—Austr. Pharm..

A quicker process is to triturate 15 drops of oil of chamomile with  $\frac{1}{2}$  av. ounce of purified talcum until well mixed, then add 16 fluidounces of distilled water, macerate for 6 hours, and filter.

## Water, Cherry. (Aqua Cerasorum Nigrorum.)

Bitter almond water......fl.dr. 7 Distilled water, enough to make fl.oz. 16

#### Water, Chlorine.

A rapid process and one that is entirely satisfactory is as follows:

Potassium chlorategr.	40
Hydrochloric acid, C. Pfl.dr.	
Distilled water, enough to make fl.oz.	

Mix the salt with the acid in a bottle and when vapors begin to be evolved add I fluidounce of water, stopper the bottle, shake well until the crystals are dissolved, and then add the remainder of the water.

## Water, Crystal. (Aqua Crystallina.)

Potassium	bitartrate.	 gr.	120
Sugar			
Water		 .fl.oz.	15

Mix, dissolve and filter. This is a pleasant beverage for use in febrile disease.

## Water, Dill. (Aqua Anethi.)

Dill fruit (seed)av.oz. 1	1
Water	Ē

Mix and distill off 16 fluidounces.—Brit. Pharm.

This may be prepared by triturating 15 drops of oil of dill with  $\frac{1}{2}$  av. ounce of purified talcum until well mixed, add 16 fluid-ounces of distilled water, macerate for 6 hours, and filter.

#### Water, Linden.

Linden	flowers,	dry	.av.oz. 1

Mix and distill off 16 fluidounces.

Linden flowers are derived from our common basswood tree. Instead of one-half ounce of the dry leaves,  $2\frac{1}{2}$  ounces of the fresh leaves may be employed.—D. modified.

## Water, Nux Vomica, Rademacher's.

(Aqua Nucum Vomicarum.—Aqua Strychni Seminis.)

	vomica,			
pie	ces	 	av.oz.	10
Alcol	nol	 	fl.oz.	2
			fl.oz.	

Mix, macerate for 24 hours and distill off 16 fluidounces.

#### Water, Pimento.

Pimento, bruisedav.oz.	11
Waterfl.oz.	32

Mix and distill off 16 fluidounces.—Brit. Pharm.

It may also be prepared by triturating 15 drops of oil of pimento with ½ av. ounce of purified talcum until well mixed, adding 16 fluidounces of water, macerating for 6 hours and filtering.

## Water, Quassia, Rademacher's.

Quassia	barkav.oz.	11
	woodav.oz.	
	fl.oz.	
Water .	fl.oz. 3	2

Mix, macerate for 24 hours, and distill off 16 fluidounces.

#### Water, Tar.

See Infusion of Tar.

## Water, Tobacco, Rademacher's.

(Aqua Nicotianæ.)

	freshly gath-	
ered	 av.oz.	16
	fl.oz.	
Water	 fl.oz.	64

Concise the leaves, bruise in a mortar, add the alcohol and water, macerate for 12 hours, and then distill off 16 fluidounces.

## Water, Tolu.

Tincture of	tolu	.fl.dr. 1
Magnesium	carbonate	gr. 60

Triturate the tincture with the magnesium carbonate until well mixed, add the water gradually with stirring, and filter.

This is used to some extent in lieu of distilled water for making mucilage of acacia.

## Wax, Yellow, Filtered.

Yellow	waxav.oz.	16
Sodium	sulphate, dried, in fine	
powd	er, gr.	350

Melt the wax at the lowest possible temperature, add the sodium sulphate, maintain the wax at the same temperature for 4 hours, stirring frequently, and filter by hot filtration.

This is an excellent product suitable for white ointments and cerates provided the temperature employed in preparation was not too high.—D.

### Wine of Aloes.

Purified aloesav.oz. 1
Cardamomgr. 75
Gingergr. 75
Alcohol,
White wine of each sufficient

Mix the three drugs, reduce to coarse powder, add 2 fluidounces of alcohol and 14 of wine, macerate for 7 days, agitating occasionally, and filter; add through the filter enough of a mixture of one part of alcohol to 7 of wine by volume to make the filtrate measure 16 fluidounces.—U. S. P. 1880.

## Wine of Beef and Iron. (Beef, Wine and Iron.)

Extract of beefgr. 5	256
Tincture of citro-chloride of	
ironfl.dr.	41/4
Water, hotfl.oz.	
Sherry wine, enough to make fl. oz.	16

contained in a mortar or other suitable vessel, and triturate until a smooth mixture results. Then gradually add, while stirring, 12 fluid-ounces of sherry wine. Next add the tincture and the remainder of the wine. Transfer the mixture to a bottle, set this aside for a few days in a cold place, if convenient, filter, and pass enough sherry wine through the filter to restore the original volume.

#### Wine of Beef, Iron and Cinchona.

(Beef, Wine, Iron and Cinchona.)

Extract of beefgr. Tincture of citro-chloride of	256
ironfl.dr.	41/2
Quinine sulphategr.	16
Cinchonidine sulphategr.	8
Citric acidgr.	6
Water, hotfl.oz.	1
Angelica wine. enough to make.fl.oz.	16

Dissolve the citric acid and the quinine and cinchonidine sulphates in the hot water, and pour the solution upon the extract of beef contained in a mortar or other suitable vessel. Triturate the liquid with the extract, until they form a smooth mixture, then gradually add, while stirring, 12 fluidounces of angelica wine, and afterwards the tincture of citro-chloride of iron. Transfer the mixture to a bottle, set this aside for a few days in a cold place, if convenient; filter, and pass the remainder of the angelica wine.

#### Wine of Beef, Iron and Coca.

Extract of beefgr.	256
Tincture of citro-chloride of ironm.	
Water, hotfl.oz.	1
Simple syrupfl.oz.	1
Fluid extract of cocafl.dr.	101/2
Sherry wine, enough to makefl.oz.	16

Triturate the extract of beef with the water until dissolved, add 10 fluidounces of wine, then the tincture, syrup, fluid extract, and the remainder of the wine, and filter.

Each  $\frac{1}{2}$  fluidounce represents 8 gr. of beef extract, 8 m. of tincture of iron, and 20 m. of coca.

#### Wine of Cinchona.

Yellow cinchona, coarse powderav.oz. 3 Port wine:	4
Port wine:fl.oz. 16	

Sherry wine, enough to make . fl.oz. 16 Macerate for 8 days and then filter.—
Pour the hot water on the extract of beef Germ. Pharm.

### Wine of Cinchona, Compound.

Yellow cinchonaav.oz.	13/
Bitter orange peelgr.	
Chamomilegr.	
Alcoholfl.oz.	
White winefl.oz.	

Bruise the drugs, macerate in the mixed alcohol and wine for 10 days, agitating occasionally, and filter.—Codex.

#### Wine of Cinchona and Cacao.

Fluid extract of yellow cinchonafl.oz.	1
Tincture of cacao. fl.oz. Simple syrup. fl.oz.	1
Angelica winefl.oz.	$\tilde{12}$
Mix and filter, if necessary.	

#### Wine of Cinchona and Coca.

Fluid	extract of cocafl.oz	. 1
Wine	of cinchona and cacaofl.oz	. 15

## Wine of Cod Liver Oil.

Gaduolgr.	64
Alcohol	
Simple syrupfl.oz.	2
Fuller's earthgr.	
Port wineenough to make fl.oz.	16

Mix the gaduol with the alcohol, and add the fuller's earth, rub well together, and then add the syrup and 13 fluidounces of wine, let stand a day or two, shaking occasionally, then filter, passing the remainder of the wine through the filter.

A preparation of the same character but of a more distinctive taste and appearance may be compounded as follows:

Gaduolgr. 64	
Alcoholfl.dr. 4	
Fuller's earthgr. 240	1
Port wine, claret wine, equal parts	
of each, enough to make fl.oz. 16	
Proceed as before.	

Compounds prepared as above contain 25 per cent of the active medicinal principles of cod liver oil.

#### Wine of Columbo.

		av.oz.	
Red wine	 	fl. oz.	16

Mix, macerate for 10 days, agitating occasionally, express, and filter.—Codex.

## Wine of Creosote.

Glyceritefl.c	z. 4
Simple syrupfl.c	
Waterfl.c	
White winefl.c	z. 51/
	2. 0/2

This contains  $2\frac{1}{2}$  per cent of creosote.

### Wine of Creosote, Compound.

Creosotefl.dr.	2
Alcoholfl.dr.	4
Tincture of gentianfl.oz.	5
Sherry wineenough to make fl.oz.	16

## Wine of Damiana. (Wine of Turnera.)

Fluid extract of damiana fl.oz.	3
Simple elixirfl.oz.	3
Sherry winefl.oz.	10

Mix and filter, if necessary.

### Wine, Diuretic, Hufeland's.

Fluid extract of	bryony	 fl.oz.	1
Sherry wine		 fl. oz.	11

## Wine of Elder Bark. (Hydragogue Tincture.)

Elder barkgr. 4	80
Parsley rootgr. 4	
Sherry winesufficie	nt

Extract the mixed drugs by percolation with the wine so as to obtain 16 fluidounces of product.—Eclectic.

## Wine of Golden Seal, Compound. (Wine Bitters.)

Golden seal	gr. 20
Tulip tree bark	
Bitter root (dogsbane)	gr. 20
Prickley ash berries	
Sassafras bark	
Capsicum	
Sherry winesu	

Extract the mixed drugs in coarse powder by percolation with the wine so as to obtain 16 fluidounces of product.—Eclectic.

#### Wine of Iron, Bitter.

Iron and quinine citrate, solublegr.	360
Tincture of sweet orange peelfl.oz.	
Simple syrup fl.oz.	5
White wineenough to make fl.oz.	16

Dissolve the iron and quinine citrate in 8 fluidounces of wine, add to this the tincture, syrup, and remainder of the wine, set aside for several days, and filter.—U. S P.

#### Wine of Kola.

Elixir of kolafl.oz.	
Claret winefl.oz.	3
Mix; let stand for 24 hours, and filter.	

#### Wine of Iron, Sweet.

Cinchona, powdergr.	60
Bitter orange peel, powdergr.	30
Citric acidgr.	30
Citrate of iron, solublegr.	120
Waterfl.oz.	
Sherry wine fl.oz.	7
Tincture of sweet orange peel.fl.oz.	$3\frac{1}{2}$
Simple syrupfl.dr.	$14^{\circ}$

Mix the tincture with the water and with this percolate the mixed cinchona and orange peel, adding enough more of the same menstruum to make 7 fluidounces, add to this the citric acid and iron salt dissolved in the wine, then add the syrup and filter.

#### Wine of Iron and Quinine Citrate.

Citrate of iron and quininegr.	48
Water, hotfl.oz.	2
Syrup of lemonfl.oz.	2
Sherry wineenough to make fl.oz.	16

Dissolve the iron and quinine citrate in the water, add the other ingredients, and filter if necessary.

#### Wine of Iron and Potassium Tartrate.

Tartrate of iron and potassiumgr. 10	
Water, hotfl.dr.	4
Water of ammoniasufficie	
Angelica wine, enough to make fl.oz.	16

Dissolve the salt in the water, carefully neutralize the acid in the wine with ammonia, mix the two liquids, and filter.

### Wine of Licorice with Opium.

(Vinum Liquiritiæ Thebaicum.)

Opium powdergr. 145
Spanish saffron, coarsely pow-
deredgr. 145
Extract of licorice, powdergr. 145
Malaga wine

Macerate the drugs for 5 days with 16 fluidounces of wine, and filter, adding enough wine through the filter to make 16 fluidounces.

The product contains 2 per cent of each of the drugs.

#### Wine of Coca.

Elixir of cocafl.oz. 8	
Claret winefl.oz. 8	
Mix; let stand for 24 hours, and filter.	

#### Wine of Iron Citrate.

Iron citrate, soluble gr.	288
Tincture of sweet orange peel.fl.oz.	
Simple syrupfl.oz.	13
White wineenough to make fl.oz.	16

Dissolve the iron salt in 12 fluidounces of wine, to this add the tincture, syrup, and remainder of the wine, set the mixture aside for several days, and filter.—U. S. P.

### Wine of Orange.

Oil of orange	5
Alcohol fl.dr.	4
Magnesium carbonate gr.	240
Simple syrupfl.oz.	2
Sherry winefl.oz.	

Dissolve the oil in the alcohol, triturate with the magnesium carbonate, add the other ingredients and filter.

#### Wine of Pancreatin.

	atin, pure gr.	
Simple	elixirfl.oz.	5
Sherry	winefl.oz.	11

Macerate the pancreatin in the elixir for 24 hours, then add the wine and filter.

#### Wine of Quinine.

Quinine sulphategr.	16
Citric acidgr.	24
Orange winefl.oz.	

Mix, let stand for 3 days, agitating occasionally, and filter.—Brit. Pharm.

## Wine of Wafer Ash. (Wine of Ptelea.)

Wafer	ash,	powderav.oz. 2-	ļ
		sufficient	

Extract the drug by percolation so as to obtain 16 fluidounces of product.

## Wine of Wormwood. (Wine of Absinthium.)

Wormwood,	cutav.oz.	1/2
Alcohol	fl.oz.	1
Sherry wine	fl.oz.	16

Mix, macerate for 7 days, agitating occasionally, and filter.—Codex.

#### PART II.

## HOUSEHOLD REMEDIES.

While this part of the work is not intended in any sense as a treatise upon the manufacture of proprietary medicines, it is proposed to impart to the pharmacist such information as he requires in preparing remedies for ordinary ailments, for popular use and sale, and thereby to replace "patent" medicines. These remedies may be offered by the pharmacist for sale under his own name or that of a pseudonym, if the latter be preferred.

#### Ague Cures.

Remedies for the relief and cure of fever, ague, and chills, or, more properly, malaria, are not of such frequent use as they were some years ago; however, there is still a demand in some sections, and the pharmacist must be prepared to supply the want. Most of these remedies are made to contain cinchona or one or more of its alkaloids, sometimes also arsenic and frequently some iron or a cathartic.

These preparations may assume the form of liquid or pills, or even some other form, but the two mentioned are the most common. They may be entitled "Peruvian Ague Cure," "Ague Tonic, " "Ague Tonic Syrup" "Ague Tonic Cure," "Chill Cure," "Chill Tonic," "Ague Pills," "Anti-Chill Pills,"

Cholagogue," "Ague Remedy," "Ague Bitters," "Ague Specific," "Fever and "Febrifuge," "Ague Tonic," Specific," etc.

1.	
Tincture of eucalyptus (1 in 10) fl.oz.	2
Tincture of serpentariafl.oz.	4
Tincture of capsicumfl.dr.	5
Tincture of myrrhfl.dr.	5
Tincture of nux vomicafl.dr.	2
Quinine sulphategr.	
Elixir of licorice, enough to make fl.oz.	16
II.	

L.	
Cinchonidine sulphateav.oz.	21/2
Aromatic sulphuric acidfl.oz.	1
Tincture of chloride of ironfl.oz.	3
Tincture of nux vomica fl.oz.	2
Syrup of ginger, enough to make fl. oz.	64

III.	
Quinidine sulphateav.oz.	2
Solution of arsenous acidfl.dr.	4
Fluid extract of sennafl.oz.	1
Syrup of licorice enough to make flor	

Owing to the absence from the above of the bitter taste of quinine, cinchonidine or cinchona bark, it may be termed "Tasteless Chill Cure," "Tasteless Ague Tonic," or "Tasteless Chill Tonic."

IV.	
Tincture of valerianfl.oz.	41/2
Tincture of cinchonafl.oz.	41/
Elixir of licorice, enough to make fl. oz.	64
V.	
Fluid extract of cinchonafl.oz.	6

	Fluid extract of cinchonafl.oz.	6
	Aromatic sulphuric acidfl.oz.	2
	Tincture of gingerfl.oz.	3
	Fluid extract of clovesfl.oz.	1
	Fluid extract of rhubarbfl.dr.	4
	Fluid extract of cinnamon fl.dr.	4
	Fluid extract of podophyllumfl.dr.	1
	Alcoholfl.oz.	
	Simple syrupenough to make fl.oz.	64
<b>(</b> 7:	T .	

		-	1	0				
VI.								
Ci	ncho	na,	yello	w, powdei	r	.a	v.oz.	4
C1	oves,	po	wder			.a	v.oz.	1/2

Dose: A teaspoonful every 3 hours. This preparation has been termed "Dutch. Ague Remedy."

#### VII.

Make pills, each containing	
Chinoidine gr.	1
Iron ferrocyanidegr.	1
Oil of black pepper drop	1
Arsenous acidgr.	20
VIII	

Make pills, each containing	
Cinchonidine sulphate gr.	3
Podophyllingr.	
Gingergr.	

#### Asthma Remedies.

The number of titles employed to designate asthma remedies is comparatively small; the following are appropriate: "Asthma Remedy," "Asthma Elixir," "Antasthmatic Powder," "Antasthmatic Remedy," "Asthma Cure," "Asthma Mixture," etc.

Asthma remedies are of about three kinds, one for internal administration, one for inhalation, and one to be ignited, the vapor to be inhaled. Favorite ingredients of the first kind are the alkali iodides, ammonium salts, grindelia, wild cherry, lobelia, belladonna, Hoffman's anodyne, etc.

The liquid preparations for inhalation contain amyl nitrite and oil of mustard. Asthma remedies to be used by ignition are usually composed of stramonium, potassium nitrate, belladonna, etc.

I.

Ammonium iodide gr.	120
Tincture of lobeliafl.dr.	2
Tincture of belladonnafl.dr.	2
Fluid extract of grindelia ro-	
busta fl.dr.	4
Fluid extract of licorice fl.dr.	4
Syrup of tolu, enough to make fl.oz.	4

Label: A teaspoonful three times a day. Extra doses to be given during a paroxysm.

TT.

L+	
Potassium iodidegr.	160
Fluid extract of grindeliafl.dr.	$2\frac{1}{2}$
Fluid extract of belladonna fl. dr.	1/2
Tincture of gelsemiumm.	50
Waterfl.oz.	1
Simple elixirenough to make fl.oz.	4

Give one and a half teaspoonfuls every 2 or 3 hours. After about three doses are taken the difficulty of breathing will have disappeared.

#### III.

Ammonium bromidegr.	120
Ammonium chloridegr.	120
Tincture of lobeliafl.dr.	3
Fluid extract of grindeliafl.oz.	1
Compound spirit of ether fl. oz.	1
Waterfl.oz.	1
Syrup of licorice, enough to	
makefl.oz.	4

IV.

Stramoniumav.oz.	2
Cannabis indic aav.oz.	1
Lobeliaav.oz.	1
Eucalyptusav.oz.	1
Teaav.oz.	
Aniseedgr.	
Potassium nitrateav.oz.	11

Reduce all to powder and pass through a No. 30 sieve; then dry thoroughly. If the potassium nitrate be mixed with 3 ounces of water, and the vegetable powder be

moistened with the solution, and then thoroughly dried, it burns better; but if all the ingredients are well dried, and the niter is thoroughly mixed, this is not necessary.

V.

Potassium iodidegr. 60	
Sodium nitrite gr. 60	
Fluid extract of quebracho fl.dr. 3	
Fluid extract of coffee fl.dr. 3	
Fluid extract of sanguinariadrops 15	
Syrup of ipecacfl.dr. 8	
Chloroform waterfl.oz4	

This is to be used like Nos. I, II, and III.

VI.

•	1.	
	Tincture of stramoniumfl.dr.	2
	Tincture of lobelia, etherealfl.dr.	1
	Potassium nitratefl.dr.	1
	Spirit of nitrous etherfl.dr.	
	Aromatic tincturefl.dr.	4
	Chloroform waterfl.oz.	2

Directions: Two tablespoonfuls at bed time, and when difficult breathing comes on.

#### VII.

Ammonium iodidegr.	120
Ammonium bromidegr.	180
Syrup of tolufl.oz.	3
Tincture of lobeliafl.oz.	5

Teaspoonful every 1, 2, 3 or 4 hours.

This is known as "Fothergill's Asthma Mixture."

VIII

Hoffman's anodyne	fl.oz. 1
Syrup of tar	
Syrup of wild cherry	
Syrup of lactucarium	fl.oz. 2

Dose: A teaspoonful every 2 or 3 hours as needed.

IX.

Amyl nitritefl.dr.	2
Oil of mustard (essential)drops	10
Etherfl.dr.	
Alcohol fl.oz.	4

Put a few drops of the liquid upon some cotton or sponge and inhale from the latter.

X.

Fluid extract of grindelia ro-	
busta	2
Potassium iodidegr.	
Tincture of opium fl.dr.	
Hoffman's anodynefl.dr.	
Syrup of wild cherry, enough	
to makefl.oz.	4
to make	-

Dose: Two teaspoonfuls.

X1.	
Stramonium leavesav.o.	z. 1
Belladonna leavesav.o.	z. 1
Saltpetreg	r. 72
Opiumg	

Mix the drugs in powder. This is to be burned and the vapors inhaled.

#### XII.

Cubebav.oz.	$\frac{1}{2}$
Digitalisav.oz.	$\frac{1}{2}$
Jaborandiav.oz.	1
Štramoniumav.oz.	
Potassium nitrateav.oz.	
Grindeliaav.oz.	
Eucalyptusav.oz.	
Cascarilla gr.	60

Mix when perfectly dry, and burn from 1 to 2 drams, inhaling the fumes.

#### XIII.

Potassium nitrateav.oz.	$\frac{1}{2}$
Aniseedav.oz.	
Stramoniumav.oz. 1	L

Mix all in powder, and use like the preceding.

#### XIV.

Stramonium, cutav.oz.	10
Alcohol fl.oz.	2
Potassium nitrateav.oz.	11
Sodium nitrategr.	60
Potassium carbonate gr.	8
Water fl.oz.	

Moisten the stramonium leaves with the alcohol, allow it to remain tightly packed in a well-covered vessel for 24 hours; dissolve the salts in the water, and impregnate the leaves with the solution; again pack them tightly as before, and allow to remain for 24 hours, then remove them and dry carefully. XV.

The Paper of Potassium Nitrate of the U. S. P. may be used. It is prepared by immersing strips of white unsized paper in a solution of potassium nitrate in 4 parts of water, and then drying them.

In using they are to be ignited and the vapor inhaled.

## XVI.

Cigarettes may be prepared from a mixture of equal parts of stramonium, lobelia, and belladonna herbs. These may be cut to a very fine condition, then rolled in suitable paper, and sealing the ends to prevent the drug from falling out by turning up the ends of the paper.

## Barber's Itch, Remedies for.

Resorcinav.oz.	1
Glycerinfl.oz.	1
Waterfl.oz.	1
Colognefl.dr.	4
Lac sulphurav.oz.	11/2
Alcoholfl.oz.	4
Apply several times daily.	

#### III.

Shave off the hairs or cut them very short; then apply, once or twice a week, an ointment composed of:

Prepared chalkav.oz.	1
Coal targr. 45 to	180
Glycerinfl.dr.	
Simple cerate av.oz.	

#### III.

Shave the affected part closely every day and rub in this ointment twice daily:

Tannic acidgr.	90
Lac sulphurgr.	180
Zinc oxideav.oz.	1
Starchav.oz.	1
Petrolatumav.oz.	2

#### T 3.7

During the inflammatory stage the following may be applied:

Ichthyol gr.	20
Salicylic acid gr.	10
Oleate of mercury, 10 per centgr.	120
Oil of lavenderdrops	
Lanolingr.	

This to be kept constantly applied to the affected parts.

#### V

Ichthyol gr.	30
Salicylic acidgr.	
Mercury oleategr.	
Zinc oxidegr.	
Starchgr.	210
Petrolatumgr.	420
 *	

#### VI.

Tannic acid gr.	15
Sulphur, precipitatedgr.	
Petrolatum or.	

#### VII.

Other remedies may be found under the head of "Eczema Remedies," "Remedies for Itch," and "Ointments or Salves."

#### Bitters.

Suitable names for these preparations are "Wahoo Bitters," "Jaundice Bitters,"

IV.

"Stomach Bitters," "Hop Bitters," "Planta-		
tion Bitters," "American Plant Bitters,"		
"Prickly Ash Bitters," "Tonic Bitters,"		
"Iron Tonic Bitters," "Blood Bitters,"		
"Liver Bitters," "German Bitters," "Spring		
Bitters," "Burdock Bitters," etc.		

These preparations differ greatly from one another. All are stomachies and tonics; some are cathartics, others diuretics; some are also alterative.

Some "bitters" are simply disguised "drinks."

I.		
	Gentian av.oz.	1 3/
	Bitter orange peelgr.	240
	Calamus gr.	420
	Tincture of citro-chloride of	
	iron fl.dr.	6
	Alcohol,	
	Waterof each, suffic	ient
	Syrupy glucosefl.oz.	

Extract the drug with a mixture of alcohol and water in the proportion of 1 of the former to 2 of the latter, to obtain 12 fluidounces of product, to which add the iron tincture and glucose.

## II.

Aloes av.oz.	6
Sassafras av.oz.	
Hopsav.oz.	
Gentianav.oz.	
Chamomileav.oz.	2
Acetic acidav.oz.	1
Alcoholfl.oz.	32
Waterenough to make gal.	1

Mix the acid and alcohol with 2 quarts of water, also mix the drugs, reduce to coarse powder, pack and macerate in a percolator, pass the previous mixture through it, and then enough hot water to make 1 gallon.

#### III.

Fluid extract of gentianfl.oz.	1
Fluid extract of yellow cin- chonafl.oz.	1
Quinine sulphategr.	_
Iron citrate, solubleav.oz.	2.
Spirit of orangefl.oz.	1
Simple syrupfl.oz.	32
Water, hotfl.oz.	8
Sherry wineenough to make gal.	1

Dissolve the iron salt in the water, add the other ingredients, and filter.

Cascara sagradaav.oz.	
Dandelion av.oz.	
Gentian av.oz.	
German chamomil e 3v.oz.	
Stillingiaaoz.	
7.1	

 Bitter orange peel
 av.oz.
 1

 Cloves
 gr. 180

 Spirit of orange
 fl.oz.
 1

 Sugar
 av.oz.
 16

 Alcohol,

Water.....of each, sufficient

Mix the seven drugs and reduce to moderately coarse powder, extract by percolation with a mixture of 1 part of alcohol to 3 of water so as to obtain 120 fluidounces of product. To this add the spirit of orange and the sugar; dissolve the latter by agitation.

V.

V •	
Wahoo barkav.oz.	31/2
Gentianav.oz.	11/2
Tamarack barkav.oz.	11/2
Fraseraav.oz.	11/4
Dogwood barkav.oz.	1
Golden sealav.oz.	3
Canada snake rootav.oz.	$1\frac{3}{4}$
Angelica rootav.oz.	2
Serpentariaav.oz.	7
Prickly ash berriesav.oz.	1/2
Podophyllumav.oz.	1
Canella barkav.oz.	2
Buckthornav.oz.	1
Sugarav.oz.	30
Alcohol,	
Waterof each, sufficie	ent

Mix the drugs, reduce to moderately coarse powder, and extract by percolation with a menstruum consisting of 1 part of alcohol to 3 of water by measure so as to obtain 112 fluidounces of percolate; in this dissolve the sugar by agitation or percolation.

VI.

٧	1.	
	Hopsav.oz.	4
	Dandelion av.oz.	4
	Gentianav.oz.	4
	German chamomileav.oz.	
	Stillingiaav.oz.	4
	Sugarav.oz.	32
	Water,	
	Alcoholof each, sufficient	ent

Mix the drugs, reduce to coarse powder and extract by percolation so as to obtain 110 fluid-ounces of percolate, using a menstruum composed of 1 part of alcohol to 3 of water, by measure; in this percolate dissolve the sugar by agitation or percolation.

#### VII.

Sugarav.oz.	6
Calamus rootav.oz.	
Bitter orange peelav.oz.	6
Peruvian barkav.oz.	6
Gentianav.oz.	6
Calumbaav.oz.	1/2
Rhubarbav.oz.	2
Cinnamon av.oz.	1
Clovesav.oz.	1
Diluted alcoholgal.	1
. 0	

Reduce all the drugs to a coarse powder, and macerate two weeks with the menstruum; then strain, express and filter.

#### VIII.

Orange peel, groundav.oz. 6
Gentian root, groundav.oz. 4
Roman chamomile flowersav.oz. 2
Rye whiskeygal 1

Macerate for 7 days, occasionally shaking the mixture; then express and filter through paper.

#### IX.

Gentian root, groundav.oz.	$2\frac{1}{2}$
Bitter orange peelav.oz.	$1\frac{1}{2}$
Canella, groundav.oz.	
Cochineal, bruisedgr.	30
Alcohol fl.oz.	16
Waterfl.oz.	16

Macerate for 7 days in a suitable vessel, occasionally agitating; express and filter through paper.

#### X.

Gentianav.oz.	4
Peruvian barkav.oz.	2
Roman chamomile flowersav.oz.	1
Quassiaav.oz.	1/2
Bitter orange peelav.oz.	1
Diluted alcoholgal.	
9	

Mix the drugs, reduce to coarse powder, mix with the diluted alcohol, macerate for 7 days, agitating occasionally, then express and filter.

#### XI.

Orange berries	.av.oz. 5
Orange peel, bitter	gr. 200
Calamus	gr, 100
Pimpinella	
Hops	gr. 50
Simple syrup	
Diluted alcohol	sufficient

Mix the drugs, reduce to fine powder, extract by percolation with the diluted alcohol so as to obtain 124 fluidounces of product, and to this add the syrup.

#### Blood Purifiers.

These preparations may be put up under the names of "Sarsaparilla" "Sarsaparilla Syrup," "Compound Extract of Sarsaparilla," "Compound Sarsaparilla with Burdock and Iodide of Potassium," "Compound Syrup of Red Clover Blossoms," "Blood Purifying Tea," "Alterative Mixture," "Blood Cleanser," or whatever other title may be appropriate, or desirable.

The ingredients of this mixture are sarsaparilla, stillingia, burdock, yellow dock, red clover, any one or more of these combined, sometimes with potassium iodide, often with some laxative. Blood-purifying mixtures are in fact mainly cathartics.

Formerly all blood purifiers were prepared in the liquid form; latterly some are prepared in the form of "teas," or species. Examples of both kinds are given among the recipes that follow. The "teas" may be termed "Blood-Purifying Tea," "Sarsaparilla Tea," etc.

### Τ.

Potassium iodidegr. 24	0
1 Otassium Todidegi. & x	
Waterfl.oz.	2
Fluid extract of burdockfl.oz.	$^2$
Compound fluid extract of sar-	
saparillafl.oz.	8
Syrupy glucose, enough to makefl. oz. 1	6

Mix. Dose, 1 to 4 teaspoonfuls, according to age.

#### II.

Buckthorn barkav.oz. 20
Rochelle saltav.oz. 8
Potassium bicarbonategr. 240
Sugarav.oz. 24
Alcohol fl.oz. 24
Spirit of lemonfl.dr. 6
Tincture of gingerfl.dr. 6
Oil anisedrops 15
Watersufficient

With sufficient water make 6 pints of decoction from the buckthorn bark, which may be deprived of bitterness by the addition of calcined magnesia (see Bitterless Extract of Cascara Sagrada, Part I) and dissolve in it the Rochelle salt, potassium bicarbonate and sugar. After allowing to stand for some time, clarify by straining through flannel. Then mix the remaining ingredients and incorporate with the

decoction. The dose is from 1 to 2 table-	IV
	IX.
spoonfuls morning and evening, some time	Stillingia
after meals.	Burdockav.oz. 3
III. ·	Blue flag
Cream of tartarav.oz. ½ Potassium bicarbonategr. 150	Podophyllum
Fluid extract of podophyllumfl.dr. 1	Senna
Compound fluid extract of sar-	Potassium iodidegr. 480
saparilla	Diluted alcoholsufficient
Compound tincture of carda- momfl.dr. 2	Mix the drugs, except the iodide, reduce
Glycerin	to coarse powder, extract with diluted alcohol
Waterenough to make fl.oz. 16	so as to obtain 1 gallon of percolate, and in
Dissolve the two potassium compounds in	this dissolve the potassium iodide
8 fluidounces of water by the aid of a gentle	-
heat, add the remaining ingredients, set	X.
aside for about 12 hours, and filter.	Sarsaparillaav.oz. 6 Burdock rootav.oz. 3
IV.	Dandelion rootav.oz. 3
Sodium sulphovinateav.oz. 1	Prickly ash barkav.oz. 3
Compound syrup of sarsaparilla, fl.oz. 4	Chamomile, Romanav.oz. 3
Fluid extract of dandelionfl.oz. 2	Sassafras barkav.oz. 4 Potassium iodideav.oz. 1
Syrup of orange	Sodium salicylate
	Glycerin
Mix and filter.	Alcohol
V.	Water, enough to makegal. 1
Sodium sulphate	Grind all the drugs to No. 20 powder.
Water	Mix the glycerin and alcohol with 2 quarts
Fluid extract of sennafl.oz. 1	of water. Macerate 24 hours and percolate.
Essence of sarsaparilla (Part	When the liquid has ceased to drop, pour in
V1.)fl.dr. 1 Syrupy glucose, enough to make.fl.oz. 16	hot water until a gallon altogether has been
VI.	obtained. Add the potassium iodide and
Rochelle saltav.oz. 1½	sodium salicylate and dissolve. If not suffi-
Compound fluid extract of sar-	ciently dark to suit the eye, add 1 fluidounce
saparilla	of caramel.
Fluid extract of dandelion $fl.oz$ . $1\frac{1}{2}$ Glycerin $fl.oz$ . $2$	XI.
Syrupfl.oz. 2	Sodium sulphateav.oz. 1
Spirit of wintergreen fl.dr. 1	Water
Water, enough to makefl.oz. 16	Fluid extract of burdock fl. oz. 2  Fluid extract of senna fl. oz. 1
VII.	Compound syrup of sarsaparilla fl.oz. 8
Rochelle salt	Syrupy glucose, enough to
Fluid extract of sarsaparillafl.oz. 1½	makefl.oz. 16
Fluid extract of dandelionfl oz. 5	XII.
Compound syrup of sarsaparilla.fl.oz. 14 Syrup of wild cherryfl.oz. 14	Stillingia
Water, enough to makefl.oz. 16	Blue flag
Mix and filter.	Prickly-ash barkav.oz. 2
VIII.	Coriander seed
Fluid extract of cascara sagrada fl.oz. 4	Licorice
Glycerin fl.oz. 4	Yellow dockav.oz. 4
Syrup of gingerfl.oz, 4	Potassium iodideav.oz. 2
Peppermint waterfl.oz. 4	Alcohol, Waterof each, sufficient
Sodium sulphitefl.dr. 1	
Mix. Dose, from $\frac{1}{2}$ to 1 tablespoonful as required.	Mix the ground drugs, and moisten with
· · · · · · · · · · · · · · · · · · ·	menstruum (3 parts water, 1 part alcohol by

measure); pack in a percolator and allow to macerate for 48 hours, then percolate to  $2\frac{1}{2}$  pints; in this dissolve first the potassium iodide, and then, by cold percolation, 3 avpounds of sugar.

#### XIII.

Sarsaparillaav.oz.	61/2
Stillingiaav.oz.	61/2
Dandelionav.oz.	31/4
Pipsissewaav.oz.	31/4
Poke rootav.oz.	$3\frac{7}{4}$
Sennaav.oz.	314
Licoriceav.oz.	$3\frac{7}{4}$
Sodium sulphateav.oz.	2
Sugar	48
Alcohol,	
Water, enough to makegal.	1

Mix the drugs, except sodium sulphate, reduce to coarse powder, percolate with a mixture of 1 of alcohol to 3 of water, until 100 fluidounces are obtained. In this dissolve the sugar and sodium sulphate.

#### XIV

Sennaav.oz.	20
Sugarav.oz.	20
Fennel seedav.oz.	
Caraway seedav.oz.	2
Juniper berriesav.oz.	6
Celery seedav.oz.	
Couch grassav.oz.	
Sassafras barkav.oz.	
Rochelle saltav.oz.	

All should be in coarse powder and should be well mixed. The mixture is a cathartic, alterative and diuretic.

#### XV.

Sennaav.oz.	10
Couch grassav.oz.	
Chicoryav.oz.	1
Juniper woodav.oz.	1
Rest harrow rootav.oz.	1
Dandelionav.oz.	
Guaiac woodav.oz.	1

Mix and make into a species.

This preparation is known as "Koeller's Blood-Purifying Tea."—H.

#### XVI.

Sennaav.oz.	
Corianderav.oz.	2
Manna av.oz.	4
Cream of tartargr.	140

Make into a species.

#### XVII

Aloesav.oz. 5	
Agaricgr. 240	
Gentiangr. 240	
Galangalgr. 240	
Rhubarbgr. 240	
Zedoarygr. 240	
Myrrhgr. 240	
Opiumgr. 2	
Saffron gr. 5	
Cardamomgr. 5	
Cassia gr. 5	
Sugarav.lbs. 3	
Diluted alcoholsufficient	

Macerate the ground drugs with 100 fluidounces of diluted alcohol for 7 days, agitating frequently, then filter, and in the filtrate dissolve the sugar by agitation.

This may be sold as "Blood-Purifying Drops." It resembles "elixir of long life" or "Swedish Bitters."

#### Bunion Cures.

The remedies recommended for the relief and cure of corns are usually also recommended for the removal of bunions. While these remedies often do afford relief, the two maladies are almost as distinct as they could possibly be. Corns are inflammations of the skin, whereas bunions are inflammations of the synovial membrane, which connects the great toe with the foot proper. Nothing less than a surgical operation will absolutely and permanently cure bunions.

Relief to bunions is often afforded as stated by the application of corn cures, assisted by frequent bathing in hot water. Frequently anointing with petrolatum, the application of tincture of iodine or of iodine ointment, or the wearing of a rubber protector will prove beneficial. A warm flaxseedmeal poultice at night often eases a great deal.

Dr. Shoemaker, in his admirable work entitled "Heredity, Health and Personal Beauty," recommends the following paint:

Carbolic acid	.fl.dr. 2
Tincture of iodine	. fl.dr. 2
Glycerin	

To be applied with a camel's hair pencil every day. Copper oleate plaster is also advised by the same authority.

Burns and Scalds, Applications for
I. Calution of ablantic of the A
Solution of chloride of ironfl.dr. 4 Petrolatumav.oz. 4
II.
Europhengr. 50
Olive oil gr. 100
Lanolinav.oz. 1
Petrolatumav.oz. 2
III.
Salolav.oz. 1
Olive oilfl.oz. 6
Lime waterfl.oz. 6

Dissolve the salol in the oil, add the lime water, and agitate well.

IV.

It is also recommended first to wash the burns with saturated solution of boric acid and then to apply absorbent gauze saturated with the following:

Lime waterfl.oz.	8
Linseed oil, rawfl.oz.	8
Thymolgr.	75

Dissolve the thymol in the oil, add the water, and agitate thoroughly.

In about 10 days substitute this ointment:

Bismuth subnitrategr.	150
Boric acidgr.	75
Olive oilfl.dr.	5
Lanolinav.oz.	2

Lanolinav.oz. 1
Benzoated lardav.oz. 2
Lime waterfl.oz. 3
VI.
Mentholgr. 20
Iodoform gr. 20
Glycerin

Mix well.

VII. Some of the preparations under the heading "Ointments or Salves," may also be utilized.

## Catarrh and Cold in the Head, Remedies for.

Catarrh (or coryza) manifests itself in two general forms, one being acute, which is usually known as "cold-in-the-head;" the other, the chronic form, which is known by the popular designation "catarrh." Both forms may be treated in the same, or, at least, in a similar, manner.

The remedies for catarrh are multiform. Mix and Some are snuffs, others are to be used by as a snuff.

inhalation; some by spraying, others by insufflation or by injection with syringe or a douche. Others again are in the form of an ointment, which is to be applied to the interior of the nostrils, then to be drawn up; and finally others again are to be taken internally, the latter containing tonics combined with some agent which diffuses itself through the system and thus attacks the local manifestations of the disease.

The snuffs contain various agents, the most conspicuous being cocaine, the indiscriminate or careless use of which cannot be too severely condemned. Purchasers of such snuffs should invariably be warned that the cocaine habit is, of all forms, probably the most deadly, and that great danger attends its use in catarrh snuffs. It should never be recommended in cases of chronic catarrh where its use would be liable to prove continuous.

The catarrh remedies used by inhalation contain either menthol or iodine combined with carbolic acid. Those used by spraying (with nasal atomizers) contain liquid petrolatum, having in solution menthol, thymol and various antiseptic volatile oils. The catarrh ointments contain ingredients similar to those of the preceding. They are usually made with thymol, menthol, or some oil, together with boric acid or bismuth salt and petrolatum.

The combinations for insufflation contain potassium iodide, or other iodide, salt, potassium chloride, golden seal, carbolic acid, camphor, etc.

Good titles for catarrh remedies are "catarrh remedy," "catarrh snuff," "cream balm," "catarrh balm," "catarrhine," "catarrh cure," "menthol snuff," "mentholin," etc.

I.		
	Boric acidgr.	60
	Iodoformgr.	60
	Bismuth subnitrategr.	
	Elm barkgr.	700

Mix and use as a snuff.

II.	
Mentholgr.	5
Sugarav.oz.	1

Mix and reduce to very fine powder. Use as a snuff.

THE STATE	D I CKIM C BAIK I.
III.	X.
Sodium salicylategr, 75	Cocaine hydrochlorategr. 10
Boric acid, fine powderav.oz. 1	Oil of eucalyptusgr. 3
Cocaine hydrochlorategr. 20	Iodoform
Use like the preceding.	XI.
IV.	Boric acid, powderav.oz. 1
Cocaine hydrochlorate gr. 2 Bismuth subnitrate gr. 90	Sugar, powdered
Quinine bisulphate gr. 6	Mentholgr. 30
Orris rootgr. 30	XII.
All should be in fine powder and should	Betolgr. 150
be well mixed.	Menthol         .gr. 15           Cocaine         .gr. 6
V. Paria acid	Coffeegr. 30
Boric acidgr. 10 Orris rootgr. 25	Mix and reduce to very fine powder.
Roasted coffee gr. 150	XIII.
Sugar	Bismuth salicylategr. 360
Menthol gr. 15	Camphorgr. 30
Mix the first five ingredients and reduce to	Cocaine hydrochlorategr. 1
an impalpable powder, then incorporate	(VIIX.
thoroughly with the menthol.	XIV. Other catarrh snuffs are mentioned
VI.	in Part I.
Sodium bicarbonategr. 8 Magnesium carbonategr. 12	XV.
Menthol gr. 4	Eucalyptol
Cocaine hydrochlorate gr. 16 Milk sugar	Menthol gr. 16
	Liquid petrolatumfl.oz. 15
Mix and reduce to impalpable powder.	XVI.
VI a. White hellebore, powdergr. 120 Orris root, powdergr. 60	Oil of eucalyptusdrops 8 Thymolgr. 8
Rice powderav.oz. 134	Thymolgr. 8 Mentholgr. 4
Oil of lavenderdrop 1 Oil of cassiadrop 1	Oil of wintergreendrops 4
Oil of lemondrop 1	Liquid petrolatum
Reduce the drug to fine powder and mix	The last two above are to be used in a
with the oils.	nasal atomizer.
VII.	XVII.
Sugar	Boric acidgr. 5 Zinc sulphategr. 1
Borax	Distilled waterfl.oz. 1
Oil of peppermint drops 4	To be used like the preceding.
Reduce the sugar, borax and salt to fine	XVIII.
powder and add the oil.	Oil of eucalyptusdrops 20
VIII.	Carbolic acid drops 2
Bismuth subnitrategr. 90	Liquid petrolatumfl.oz. 1
Benzoingr. 90	Useful in all stages of nasal catarrh, in-
Boric acidgr. 60 Mentholgr. 3	cluding the grip. To be used by spraying.
Take a pinch 5 or 6 times daily.	XIX.
IX.	Carbolic acid, liquefiedfl.dr. 7 Oil of turpentinefl.dr. 3
Tannic acidgr. 2	Alcoholfl.dr. 15
Orris rootgr. 90	Ammonia waterfl.dr. 8
Sugargr. 90	This solution is dropped upon absorbent
Mix and reduce to an impalpable powder.	cotton which should be kept in a tightly

stoppered bottle, when not in use. The vapor is to be inhaled frequently. The vapors are also recommended for diphtheria.

The above is the well-known and largely used "olfactorium anticatarrhoicum" of Hager.

#### XX.

Compound tincture of iodinem.	30
Carbolic acid, crystal gr.	
Glycerin	
Water, enough to makefl.oz.	4

Mix the glycerin with the acid liquefied at a gentle heat, add the tincture and the water and expose the mixture to sunlight until it has become colorless.

This is the "Carbolized Solution of Iodine" of the N. F. and is used by inhalation.

XXI. Menthol is an excellent agent for treatment of catarrh by inhalation. It is commonly put up in vessels more or less resembling tubes, which may be opened at both ends to permit free circulation of air through the tube when in use, and which may be closed tightly when not in use.

#### XXII.

Boric acid, powder gr.	120
Mentholgr.	60
Thymolgr.	24
Eucalyptol drops	
Bismuth subcarbonategr.	120
Cold creamgr.	
Petrolatum gr.	300

Dissolve the menthol and thymol in the petrolatum, which has been melted by the application of a gentle heat, allow the solution to solidify, add the other ingredients and make to an intimate mixture.

This mixture is to be applied into the nostrils several times daily.

#### XXIII.

Thymolgr.	- 3
Oil of wintergreendrops	2
Bismuth subcarbonategr.	
Petrolatumav.oz.	1

Prepare and use like the preceding.

#### XXIV.

Mentholgr. 1 to	5
Boric acidgr.	100
Petrolatum av.oz.	11/2

Prepare and use like the preceding.

#### XXV.

Boric acidgr.	60
Boraxgr.	60
Sodium chloridegr.	30
Lister's antiseptic solutionfl.oz.	
Waterfl.oz.	

This may be used by insufflation, but preferably by means of a douche.

#### XXVI.

Borax	120
Sodium bicarbonate gr.	
Carbolic acidfl.dr.	1/2
Carbolic acidfl.dr. Glycerinfl.oz.	1
Water, enough to makefl.oz.	16

This is to be used like the preceding.

#### XXVII

22. 7 11.	
Sodium chloridegr.	240
Potassium chlorate gr.	
Ammonium iodidegr.	4
Carbolic aciddrops	16
Camphor, powder gr.	16
Golden seal, powdergr.	40

Mix well.

This is to be treated with water to make an infusion, which latter is then to be used by insufflation or injection.

#### XXVIII.

Potassium	iodidegr.	60
Compound	tincture of cardamom fl.oz.	4
Compound	tincture of gentianfl.oz.	12

#### Cathartics.

The class of remedial substances most commonly in demand partake of the form of cathartics. In offering a preparation of this kind for sale, it is customary, as well as advisable, to recommend it for chronic constipation or as a liver invigorator, a blood purifier, a remedy against biliousness or headache, an anti-dyspeptic, etc.

This class of pharmacal remedies may take the form of an elixir, syrup, infusion, tincture, pastille or troche, effervescent salt, pill, confection, powder or species.

Ι.

L٠		
	Buckthorn barkav.oz.	8
	Licorice rootav.oz.	4
	Butternut barkav. oz.	4
	Fennelav.oz.	2
	Carawayav.oz.	2
	Gingerav.oz.	
	Fluid extract of senna fl.oz.	
	Watersufficier	nt

Mix the drug, reduce to coarse powder, introduce into a vessel containing 64

duidounces of boiling water, continue boiling for 15 minutes, strain and express, adding enough water to make up the measure of 64 fluidounces. Allow this to stand a few hours, decant 56 fluidounces of clear liquid, add the fluid extract and flavor the whole, if thought advisable, by the addition of oil of anise, orange or peppermint.

#### II.

Sennaav.oz. 5
Licorice rootav.oz. 5
Buckthornav.oz. 5
Anise
Fennelav.oz. 1
Carawayav.oz.
Gingerav.oz.
Rochelle saltav.oz. 3
Alcohol,
Water of each, sufficient
Oil of wintergreendrops 8
Oil of sassafrasdrops 8

Reduce the drugs to moderately fine powder and percolate with diluted alcohol until 48 fluidounces are obtained. Then dissolve the Rochelle salts in 16 fluidounces of water, add this to the percolate and flavor the whole with the oils.

III. Compound cathartic elixir or compound elixir of cascara sagrada may be sold or dispensed if desired.

#### IV.

Fluid extract of cascara sagrada (tasteless)	1,
Fluid extract of berberis aquifo-	
lium fl.dr.	3
Senna, coarse powderfl.dr.	6
Prunesav.oz.	2
Figsav.oz.	2
Oil of fenneldrops	10
Oil of cinnamondrops	10
Oil of clovesdrops	
Sugarav.oz.	
Water, enough to makefl.oz.	16

Chop the figs and prunes, without stones, to a fine hash, mix with senna and steep in 12 fluidounces of water for 3 hours, and strain through a No. 4 wire sieve. To this liquid add the sugar, dissolve by agitation, add the fluid extracts and oils, and make up to one pint with hot water, which has been poured over the fruit on sieve.

Four fluidrams of fluid extract of licorice or 180 grains of powdered nutmegs may be substituted for the oils as a flavor.

V.

Bitterless fluid extract of cascara fl.oz.	2
Fluid extract of rhubarbfl.oz.	1
Fluid extract of sennafl.oz.	1
Oil of fenneldrops	5
Oil of carawaydrops	5
Potassium carbonategr.	10
Saccharingr.	
Simple syrup, enough to make .fl.oz.	16

Add the potassium carbonate to the fluid extract of rhubarb and dissolve, then add the other fluid extracts, the oils and saccharin, and finally the syrup.

#### VI.

Extract of senna, deresinified.av.oz.	4
Pulp of purging cassiaav.oz.	2
Pulp of tamarinds av.oz.	1/2
Extract of licoricegr.	180
Resin of scammonygr.	180 *
Tartaric acidgr.	60
Sugarav.oz.	5

Make into oval lozenges of 80 grains each. These may be dusted with powdered sugar or they may be dipped in melted chocolate and afterwards covered with a coating of granulated sugar.

VII. Wash some dried prunes (a pound, for instance), place them in a saucepan over a dull fire or on a sand-bath, with just sufficient water to nearly cover them; when they have boiled long enough to become quite soft, and the greater part of the water has been evaporated, allow them to cool, and rub them in a large mortar so as to crush the fruit, but not the kernels. Transfer them to a coarse straining cloth and squeeze the pulp through it. This should be about the consistence of honey in the water. If not, it can be made so by evaporating it over a water bath. Now make a mixture of:

Senna, powder	4
Sugar, powderav.oz.	4
Jalap, powderav.oz.	1/2
Gum arabic, powderav.oz.	3/4
Aromatic powderav.oz.	3/4

Add sufficient quantity of the prune paste to make a mass and divide into troches of convenient size.

VIII. Confection of senna of figs, and of prunes in Part I, may be sold or dispensed either as they are in paste form or they may be molded into pastilles like the preceding.

HOUSEHOLD REMEDIES. ' 175	
IX.	XV.
Jalap, powder	Buckthorn bark, cut       av.oz.       10         Dande ion, cut.       av.oz.       10         Senna, cut.       av.oz.       10         Licorice root, cut.       av.oz.       2         Coriander, bruised       gr. 300         Anise, bruised       gr. 300
X.  Senna, powder	XVI. The following may be dispensed as "Grape Salt" or "Fruit Salt" or "Fruit Saline:"  Sodium bicarbonate
XI. Formulas for cathartic pills might be given by the score. The compound cathar-	XVII. Artificial Carlsbad salt (Part IV)
tic or vegetable cathartic pills may be supplied as cathartic pills. Other formulas which may be used are the following:	or the same salt in effervescent form may be dispensed for cathartic purposes; also the effervescent magnesium citrate (Part I) may be used for the same purpose.
Aloin       gr. 10         Podophyllin       gr. 10         Capsicum       gr. 10         Extract of nux vomica       gr. 20	XVIII. Among other preparations that may be recommended for cathartic purposes are some of the liver remedies, blood purifiers and bitters.
Make 100 pills.	Chilliblain Group
Aloin       gr. 10         Jalap       gr. 10         Extract of henbane       gr. 5         Extract of nux vomica       gr. 5         Oleoresin of capsicum       gr. 5         Podophyllin       gr. 20         Make 100 pills	Chilblain Cures.  Pharmacists are often called upon for some simple remedy for the relief of the annoying affection known as chilblains. The formulas which follow may be recommended for their relief and cure.
These latter two form very small pills and may be called "Little Liver Granules," "Little Cathartic Pills," or some similar name.	I.   Creosote.
XII. Laxative species, Part I., may be dispensed when a cathartic tea is demanded. The next two formulas may also be utilized:	II.  Zinc oxidegr. 60 Camphor, powdergr. 30
XIII.       senna, cut.       av.oz. 8         Manna       av.oz. 3         Coriander       av.oz. 1	Myrrh, powdergr. 30 Opium powdergr. 30 Lardgr. 480 Dissolve the camphor in the lard, which
XIV.	has been melted at a gentle heat, allow this
Senna         av.oz.         8           Couch grass, cut         av.oz.         8           Fennel, bruised         gr. 160           Elder flower         gr. 160	to cool, add the other ingredients, and mix well.  III.  Opfum, powderedgr. 30
Mix well. This may be sold as "German Herb Tea," or "Cathartic Tea."	Camphorgr. 40 Nutgall ointmentav.oz. 1

Camphorgr. 110 Carbolic acidfl.dr. 1
Yellow waxav.oz. 1/2
Melt the wax and add the petrolatum; when nearly cold, add the other ingredients pre-
viously well mixed.
v.
Zinc soziodol
VI.
Beef marrowgr. 300 Marshmallow ointmentav.oz. 2½
Venice turpentinegr. 150
Hydrochloric acidfl.dr. 2½ Camphorgr. 90
Extract of opium:gr. 50
Melt the fats together and dissolve the
camphor in the fluid, soften the extract of
opium with a few drops of water in a mortar, and rub up with about one-half ounce of the
fats, then add more of the fats, so as to have
the extract equally distributed; mix with the
remainder, and incorporate the hydrochloric
acid by constant stirring, as in making cold cream.
VII. Oil of rosemarydrops 15
Camphor
Oil of turpentine
,-
VIII. Citrin ointmentgr. 480
Camphor gr. 60
Oil of turpentine       fl.dr.       2         Olive oil       fl dr.       4
To be applied with gentle friction before
the chilblains break.
IX.
Ammonium chloridegr. 30
Water
Alcohol
Apply morning and evening.
X.
Zinc oxide gr. 240
Tannic acid

Peru balsam ......fl.oz.

Oil of eucalyptus.....fl.dr.

XI.
Carbolic acid       drops 15         Camphor       gr. 60         Eucalyptus oil       fl.dr. 1         Alcohol, enough to make       fl.oz. 4
XII.
Resorcin       gr. 35         Ichthyol       gr. 35         Tannin       gr. 35         Water       fl.oz. 3
To be painted on at night.
XIII.
Carbolic acidgr. 6 Liniment of aconitefl.dr. 2 Liniment of belladonnafl.dr. 2 Collodion, flexiblefl.oz. 1
To be painted on once or twice daily.
XIV.  Solution of lead subacetatefl.dr. 2 Camphorgr. 120 Oil of turpentinefl.oz. 1
XV. Other suitable preparations may be
found under the head of "Ointments or

#### Cholera Remedies.

See Diarrhœa and Dysentery Remedies.

Salves," and "Liniments," Part II.

### Corn Eradicators.

Remedies for the removal of corns are very numerous and assume divers forms. One of the most common and most popular is a preparation of collodion containing extract of cannabis indica and salicylic acid, sometimes also lactic or acetic acid. Another popular preparation is an ointment or cerate containing about 10 per cent of salicylic acid. These are usually to be applied for 4 or 5 nights consecutively, followed by a hot foot bath, when the corn can be picked out. If this first treatment does not produce satisfactory results, it should be repeated. Better results are obtained if a hot foot bath be taken, also, before application of the remedy, then scraping or cutting off the calloused tissue as far as possible.

Dissolve the extract in the alcohol, and the salicylic acid in about 5 ar. ounces of flexible

collodion contained in a tared bottle. Then add the former solution to the latter, and finally add enough flexible collodion to make 10 av. ounces.—N. F.

II.

Salicylic acidgr.	30
Lactic acid, concentratedgr.	20
Collodion, enough to makefl.oz.	1

Mix and dissolve. Apply like the preceding.

III.

Extract of cannabis indica,	
(Squibb's)gr.	30
Salicylic acidgr.	
Oil of turpentinefl.dr.	21/2
Collodionfl.oz.	
Acetic acid, glacialfl.dr.	1

Mix the first three ingredients intimately, add the collodion, dissolve, and then add the acetic acid.

IV.

Salicylic acid.										.av.oz.	1/2
Simple cerate	•		•			•				.av.oz.	41/2
J.											

Yellow wax. av.oz. 6
Venice turpentine av.oz. Resin av.oz. Salicylic acid av.oz. Peru balsam av.oz. Petrolatum av.oz. 1

Melt the resin and wax and add the other ingredients; stir until cold.

VI.

Potassium carbonateav.oz. 1
Simple cerateav.oz. 2
Verdigrisenough to color sufficiently

To be applied on a cloth.

VII.

Lead plasterav	.oz.	3
Resinav	.oz.	$^2$
Verdigrisav	.oz.	1

Melt together and spread upon leather.

VIII.

4	***	
	Resinav.oz	. 3
	Yellow waxav.oz.	3
	Gum turpentineav.oz.	1
	Elemiav.oz.	1
	Beef tallowav.oz.	1
	Wood charcoal, very fine pow-	
	derav.oz.	1
	Monochloracetic acid gr.	108
	Glycerinfl.dr.	
	•	/ ·

Melt the first five ingredients, also mix the advisable to have two preparations, one withouther ingredients, and incorporate thoroughly out morphine or opium; this to be entitled

with the first mixture, when the latter begins to cool, and stir frequently until cool.

This plaster may be formed into pills which, when used, may be warmed in the hand and then spread out flat on a piece of silk; the latter then to be applied to the corn.—D.

If this plaster be warmed and spread on cloths, the latter may be sold as "Spread Corn Plaster."

IX.

Soap plaster ......av.oz. 6
Salicylic acid ......av.oz. ½

Melt the plaster, add the salicylic acid, and stir frequently until cool. This may be spread upon cloth like the preceding.

X. The remedies previously mentioned will prove of more benefit to "hard" corns then to "soft" corns. It has been recommended to treat the latter by painting with a solution of silver nitrate in 8 parts of distilled water every fourth or fifth day, in the meantime keeping the toes apart by means of a pledget of cotton smeared with petrolatum, zinc ointment or other bland fatty substance.

The cure or removal of corns is facilitated by frequent washing of the feet, followed by removal of all dead tissue.

Lately dry tannin placed between the toes where the corn is located is recommended as a cure for "soft" corns.

### Cough and Cold Remedies.

Suggested titles for these remedies are "Cough Remedy," "Cough Mixture," "Cough Syrup," "Syrup of Tar and Wild Cherry," "Children's Cough Cure" (if intended for children exclusively), "White Pine Syrup," "Honey of Hoarhound and Tar," "Tar, Tolu and Wild Cherry," "Tar Hoarhound Cough Syrup," "Lung Balsam," "Expectorant," "Cough Cordial," "Cough Balsam," etc.

Opium in some form or a salt of morphine are constituents of almost every cough mixture. This should not be administered to small children or infants and hence it may be advisable to have two preparations, one without morphine or opium; this to be entitled

"Children's Cough Cure," or "Infant	VIII.
Cough Mixture," and the other with mor-	Hoarhoundgr. 120
phine or opium.	Irish mossav.oz. 1½
Other remedies are noticed under the head	Ammonium chlorideav.oz. 2½
of Remedies for Throat Affections.	Water, boiling       fl.oz.       64         Oil of tar       fl.dr.       2
	Chloroform fl.dr. 4
I.	Fluid extract of sennafl.oz. 2
Syrup of tolu	Compound syrup of squillfl.oz. 16
Syrup of wild cherryfl.oz. 4 Tincture of henbanefl.oz. 4	Mix the hoarhound, Irish moss and am-
Hoffman's anodynefl.oz. 4	monium chloride, add the water, allow to
Water	stand for 12 hours, and strain. Dissolve
II. White Pine "Expectorant" is an ex-	the oil of tar in the chloroform, add the fluid
cellent remedy. For formula, see Part I.	extract and compound syrup, and add to the
The morphine may be omitted if desired.	previous mixture
III.	IX.
Extract of licorice, purifiedav.oz. 1	Syrup of tar
Syrupy glucoseav.oz. 10	Syrup of squillfl.oz. 16
Ammonium chlorideav.oz. 1 Paregoricfl.oz. 1	Fluid extract of lobeliafl.dr. 6
Wine of antimonyfl.dr. 4	Tincture of opiumfl.oz. 4
Spirit of nitrous etherfl.dr. 2	Fluid extract of ipecacfl.dr. 4
Water, enough to makefl.oz. 16	X.
Mix. Dose, 1 to 4 teaspoonfuls.	Syrup of tar
This mixture is known as "Improved	Syrup of tolu
Brown Mixture.''	Syrup of wild cherryfl.oz. 50 Syrup of squillfl.oz. 16
	Syrup of senegafl.oz. 6
IV.	Ammonium chlorideav.oz. 3½
Tincture of tolu fl.dr. 4 Paregoric fl.oz. 2	Morphine sulphategr. 15
Syrup of squill	Tartar emeticgr. 30
Honey, enough to makefl.oz. 16	Water
Dose. One teaspoonful.	
-	Dissolve the ammonium chloride, mor-
V.	phine salt, and tartar emetic in the water,
Terebene	filter the solution, and add the other ingre-
Sugar	dients.
Yolk of egg 4	Either of these last two formulas may be
Anise waterfl.oz. 4	used for the preparation of Syrup of Tar and
Camphor water	Wild Cherry.
Distilled water, enough to make fl.oz. 16	XI.
Triturate the acacia, sugar, and terebene in	Ammonium chlorideav.oz. 4
a mortar, beat the egg yolk with the flavored	Chloroform
waters, make an emulsion by rubbing this	Syrup of tolu
with the contents of the mortar, and add the	Mix all and dissolve by agitation.
distilled water. Lemon juice may be sub-	
stituted for the distilled water.	XII. See Yerba Santa Cough Mixture, or
VI.	Compound Syrup of Yerba Santa.
Oil of tarfl.dr. 2	XIII.
Fluid extract of hoarhoundfl.dr. 4	Wild cherry barkgr. 240
Oil of anisedrops 5	Senega
Jamaica rumfl.oz. 8	Extract of coniumgr. 15
Honey, enough to makefl.oz. 16	Ginfl.dr. 1
Dose: A half to 1 teaspoonful.	Ginfl.dr. 1 Compound tincture of carda-
VII.	momfl.dr. 1
Fluid extract of hoarhoundfl.oz. 6	Watersufficient
Syrup of wild cherryfl.oz. 18	Mix the drugs, reduce to coarse powder,
Syrup of tarfl.oz. 36	extract by percolation with water so as to

obtain 8 fluidounces of percolate and to this add the other ingredients.

Two teaspoonfuls in water constitute the usual dose to relieve cough.

This is known as "Dr. Pancoast's Cough Mixture."

#### XIV.

Tincture of capsicumf	l.oz. 1
Syrup of wild cherryf	1.oz. 2
Mucilagef	
Syrup of tarf	
Syrup of hydriodic acidf	

Label: A teaspoonful four times daily for persistent, dry, hacking cough, which resists usual treatment.

#### XV.

Raw linseed oil	oz. 2
Oil of cassia	dr. 1/2
Oil of wintergreenfl.	dr. 1/2
Oil of sassafrasfl.	dr. ½
Acacia, powder	gr. 240
Mucilage of Irish moss, N. Ffl.	oz. 2
Glycerinfl.	oz. 1
Simple syrupfl.	oz. $2\frac{1}{2}$
Morphine sulphate	
Chloral hydrate	gr. 240
Diluted hydrocyanic acid fl.	dr. ½
Water, enough to makefl.	oz. 16
36.1	

Make an emulsion in the usual way.

This may be sold as a "flaxseed or linseed cough syrup or balsam."

#### XVI.

Spirit of chloroform	.drops	20
Hydrobromic acid	.drops	30
Syrup of squill	.fl.dr.	1
Water, enough to make	.fl.oz.	1

Mix. To be given in 1 dose for an adult; for children, the quantity to be reduced according to age.

This is known commonly as "Fothergill's Hydrobromic Acid Cough Mixture."

XVII. See also Emulsion of Linseed Oil, Part I., which may be employed.

#### XVIII.

Tincture of red spruce gumfl.oz.	2
Sugarav.oz.	26
Waterfl.oz.	16
Caramel fl.dr. 1 or gr.	60
Fuller's earthgr.	120

Mix 2 ounces of the sugar with the tincture of spruce and fuller's earth, rub well and add the water in divided portions; then filter, returning the filtrate until it comes will form a very satisfactory article. Or the

through clear; add the caramel and sugar, which dissolve with a gentle heat, and strain while warm.

A syrup of a different and richer appearance may be made by mixing equal parts of the syrup prepared as above and Syrup of Wild Cherry of the U. S. Pharmacopæia.

The tincture of red spruce gum directed for use in the above is best prepared as follows:

Red spruce gum, fine powder. .av.oz. 2 Alcohol, enough to make . . . . . fl.oz. 16

Macerate until dissolved and filter.

The above syrup is the formula for what is known as "Spruce Gum Syrup," or "Syrup of Red Spruce Gum."

#### XIX.

Ammonium chloridegr.	180
Tartar emeticgr.	2
Morphine sulphategr.	3
Syrup of licorice fl.oz.	4

In teaspoonful doses.

This preparation has been known as "Davis' Cough Mixture."

#### XX.

Syrup of squill	.fl.dr. 2
Wine of ipecac	.fl.dr. 1
Paregoric	
Simple syrup	.fl.dr. 4
Water	.fl.oz. 3

This preparation has been known as "Dr. Child's Cough Mixture."

#### XXI.

1		
	Tincture of tolufl.oz.	4
	Fluid extract of lobeliafl.oz.	2
	Fluid extract of cannabis indica.fl.oz.	2
	Chloroformfl.oz.	1
	Morphine sulphategr.	32
	Tartar emeticgr.	32
	Spirit of peppermint fl.dr.	11/2
	Simple syrupgal.	1

Dissolve the morphine and tartar emetic in a little water; mix the two fluid extracts, tincture, chloroform and spirit, shake well, add a portion of the syrup, shake again, add the remainder of the syrup and then the solution previously prepared.

This makes a turbid preparation such as is commonly sold as "Chlorodyne Cough Cure."

XXII. If a lozenge is desired, the Troches of Glycyrrhiza and Opium of the U.S.P.

lozenges mentioned under the heading "Remedies for Throat Affections" may be employed.

### Cough (Whooping) Remedies.

Ammonium picrategr.	1
Ammonium chloridegr.	24
Purified extract of licoricegr.	
Vaterfl.oz.	

Dose: A teaspoonful for children up to 2 years of age and 2 teaspoonfuls for children 3 to 5 years of age.

II.
Ether
Chloroformfl.oz. 3
Oil of turpentinefl.oz. 1

Hold to the mouth on cloth or sponge, and allow the child to inhale the vapor.

III.	
Creosote, puredrops	10
Paregoric	
Syrup of glucosefl.oz.	
Caramelsufficient to co	

IV.
Butyl-chloral hyrate gr. 15
Potassium bromide dr. 60
Etherdrops 20
Tincture of belladonnadrops 15
Tincture of hyoscyamusdrops 25
Syrup of tolu, enough to make. fl.oz. 4
Label: A teaspoonful 4 times a day.

v.	
Terpin hydrategr.	15
Antipyringr.	15
Acacia	
Syrup of orangefl.dr.	
Linden flower waterfl.oz.	2

## Diarrhœa and Dysentery Remedies.

These preparations may be put under the titles "Blackberry Balsam," "Blackberry Cordial," "Blackberry Elixir," "Diarrhœa Cordial," "Diarrhœa Remedy," "Diarrhœa and Cholera Cure," etc.

The ingredients of these preparations number among the following: Blackberry root bark, blackberry juice, rhubarb, nutgall, witch hazel, catechu, kino, peppermint, opium, capsicum, ginger, aromatics and syrups. The blackberry juice and syrup are introduced for the purpose of disguising the styptic taste of the blackberry root bark, object of employing the aromatics, these of its age.

latter also serving as carminatives. Capsicum is introduced with the view of utilizing its stimulant properties. Opium should be introduced with some misgivings, for, as a rule, these "cordials" or "balsams" are given to children. In fact, it may be advisable to have two preparations, one containing opium and intended for older children and adults, the other, without opium, for infants and younger children.

- I. Compound Elixir of Blackberry, Part I.
- II. Aromatic Syrup of Blackberry, Part I.
- III. Compound Elixir of Dewberry. See Part I.

IV.

Fluid extract of blackberry root fl.oz.	2
Tincture of vanillafl.dr.	4
Compound elixir of taraxaçumfl.oz.	4
Simple elixir, enough to makefl.oz.	16

V.	
Fluid extract of blackberry root fl.oz.	2
Simple elixirfl.oz.	7
Compound elixir of taraxacumfl.oz.	

VI.				
Fluid	extract of b	lackberry	root fl.oz	. 10
Fluid	extract of g	alls	fl.oz	. 2
Arom	atic tincture		fl.oz	. 4
Simpl	e elixir, eno	ough to m	akegal	l. 1
Set as	side a few .da	ays and fi	lter.	

V	II.	
	Camphorav.oz.	$2\frac{1}{4}$
	Fluid extract of rhubarbfl.oz.	31/2
	Oil of peppermintfl.oz.	1
	Tincture of capsicumfl.oz.	
	Tincture of opiumfl.oz.	
	Chloroformfl.dr.	4
	Sodium bicarbonateav.oz.	8
	Alcohol	64
	Simple syrup, enough to makegal.	1

Dissolve the camphor and oil in the alcohol, and add the chloroform and two tinctures. Mix the fluid extract, sodium bicarbonate, and a portion of the syrup, let stand in an open vessel for several hours, then add to the previous mixture, add the remainder of the syrup, and filter in a well-covered funnel.

#### VIII.

Fluid extract of blackberry rootfl.oz.	3
Aromatic syrup of rhubarbfl.oz.	8
Fluid extract of hamamelisfl.oz.	
Tincture of opiumfl.oz.	

A teaspoonful every 2, 3, or 4 hours; a nutgall or other astringent. This is also one child should be given 5 drops for every year

Tripe bidemoetites, trittini, pint	
	1
Blackberry root gr. 48	30
Mace	30
Cloves	30
Allspicegr.	30
Cassia	30
Gingergr.	30
Port wine fl.oz.	4
Alcoholfl.oz.	2
Watersufficier	nt

Express the juice from the berries and add sufficient water through the residue to make the liquid measure 12 fluidounces; add the wine and alcohol. Mix the drugs and grind to tolerably fine powder, moisten with the liquid before mentioned, pack lightly in a percolator, soak with menstruum, macerate for 24 hours, and then percolate, passing the remainder of the liquid through the drug. If the percolate is less than 16 fluidounces, add enough menstruum consisting of alcohol and water in the proportion of 1 of the former to 4 of the latter to make up this amount. Known as "Blackberry Cordial."

7		
	Fluid extract of ipecac fl.dr.	6
	Tincture of opiumfl.oz.	3
	Aromatic tincture of rhubarbfl.oz.	3
	Simple syrupfl.oz.	3
	Alcoholfl.oz.	3
	Fluid extract of logwood fl.dr.	12
	Fluid extract of blackberry root fl.dr.	12

Dose: A teaspoonful every 3 hours.

XI.			
F	'luid extract of blackberry root	fl.oz.	32
F	luid extract of ginger	fl.oz.	10
C	compound tincture of catechu	fl.oz.	48
P	aregoric	fl.oz.	15
S	imple syrup, enough to make .	.gal.	1
		_	

١.	11.	
	Fluid extract of blackberry root fl.oz.	4
	Tincture of opiumfl.oz.	4
	Tincture of gingerfl.oz.	4
	Tincture of catechufl.oz.	4
	Tincture of kinofl.oz.	4
	Tincture of capsicumfl.oz.	
	Sugarav.oz.	
	Alcohol fl.oz.	36
	Water, enough to make gal.	1
	-	

Mix all and dissolve the sugar by agitation or percolation.

XIII. Some of the preparations in Part I, known as "Cholera Mixtures," might be utilized.

XIV. See also the Liniments in Part II.

#### XV.

Tincture of opium, deodorized...fl.dr. 2 Diluted sulphuric acid......fl.dr. 2 Tincture of cardamom comp....fl.oz. 1½ Camphor water, enough to make.fl.oz. 6

Directions: A tablespoonful, undiluted, every 3 hours until relieved.

#### XVI.

Tincture of opium, deodorized...fl.dr. 4
Tincture chloride of iron ......fl.dr. 4

Mix; 10 to 15 drops in some water, after each movement of the bowels.

These are useful for acute and chronic dysentery.

### Dyspepsia Remedies.

Owing to the prevalence of dyspepsia, remedies for this complaint are in considerable demand. Most of the "bitters" and the liver remedies, and many of the blood purifiers are usually recommended for dyspepsia.

The remedies recommended especially for dyspepsia contain some stomachic tonic like golden seal, columbo, gentian, bitter orange, etc., combined with a laxative like aloes, sodium phosphate, rhubarb, etc.; sometimes an alkali like sodium bicarbonate, as well as a carminative or stimulant like peppermint, capsicum, elixir, etc.

I.		
	Fluid extract of rhubarbfl.dr.	5
	Fluid extract of columbo fl.dr.	5
	Fluid extract of chamomilefl.dr.	5
	Fluid extract of bitter orangefl.dr.	10
	Fluid extract of life everlasting.fl.oz.	5
	Sodium phosphate av.oz.	2
	Water, hotfl.oz.	8
	Simple elixir, enough to makefl.oz.	64

Mix the fluid extracts with a portion of the elixir, dissolve the sodium salt in the water, add to the previous mixture, then incorporate the remainder of the elixir, and filter.

ΙI		
	Rhubarbav.oz.	3
	Golden sealav.oz	3/4
	Cape aloesgr.	
	Peppermint herbav.oz.	3
	Potassium carbonate av.oz.	1
	Capsicumgr.	15
	Sugarav.oz.	24
	Alcohol,	
	Waterof each, sufficient	ent

Mix the rhubarb, golden seal, aloes, peppermint and capsicum, reduce to coarse powder, extract by percolation with a mixture of 3 volumes of alcohol and 10 of water, so as to obtain 50 fluidounces of percolate, having first dissolved the potassium carbonate in the water. In the percolate dissolve the sugar, either by agitation or percolation, and then add enough more of the menstruum to make 64 fluidounces.

#### III.

Sodium bicarbonateav.oz.	1
Sodium sulphateav.oz.	2
Tincture of gentian compoundfl.oz.	4
Fluid extract of sennafl.dr.	2
Fluid extract of rhubarbfl.dr.	
Oil of carawaydrops	20
Water, sufficient to makefl.oz.	

Dissolve the sodium sulphate and bicarbonate in the water, add the oil of caraway to the tincture and fluid extracts and mix together. Dose: A tablespoonful after meals and at bedtime, in some water.

#### IV.

Compound tincture of gentianfl.oz.	1
Tincture of columbofl.dr.	4
Tincture of nux vomica fl.dr.	1/2
Nitromuriatic acidfl.dr.	
Simple syrup, enough to make fl. oz. 4	4

A teaspoonful 3 times daily.

#### V.

Carbolic aciddrops	18
Tincture nux vomicafl.dr.	11/2
Nitrohydrochloric acid, diluted.fl.dr.	$1\frac{7}{2}$
Pepsin elixir	
Brandyfl.oz.	
•	

Direction: A teaspoonful 3 times a day before meals. This is used in fermentative dyspepsia.

#### VI.

Cocaine hydrochlorate gr.	12
Hydrochloric acid, diluted fl.dr.	1
Elixir of Garus or aromatic	
elixir fl.oz.	121/2
Waterfl.oz.	21/2

Dose: One tablespoonful after eating, for dyspepsia complicated with gastralgia.

This is known as "Huchard's Elixir."

#### VII.

Infusion of rhubarb	fl.oz.	141/2
Resorcin	gr.	120
Sodium bicarbonate	gr.	320
Peppermint water	fl.oz.	1

A tablespoonful every hour. Useful for catarrh of the stomach.

VIII. Sometimes elixir of pepsin or compound powder of pepsin or tablets, soda and hot.

peppermint, or soda and pepsin are recommended for dyspepsia.

IX. Recently dyspepsia tablets have appeared on the market. Something very similar may be prepared according to this formula:

Sodium bicarbonate	.gr.	5
Resin of jalap	.gr.	1
Extract of hydrastis, powder	.gr.	3
Menthol	.gr.	1/8

Make 1 tablet. Pepsin may be added if desired, but it will not serve any purpose whatever in the mixture.

One, 2 or 3 of the tablets may be taken after each meal.

This mixture may conveniently be compressed by one of the hand compressors now so readily obtained.

#### Ear Medicines.

Medicines for the ear are of two kinds, one for earache, which may be called "Earache Drops," "Earache Remedies," or "Earache Oils," the other being intended for the improvement of the hearing, the latter kind being known as "Ear Oils" or "Acoustic Oils." Of course no medicine will actually improve the hearing when impaired, unless it be due to a waxy concretion present, which may be softened and dissolved.

### I. Earache Remedies:

71.											
(	Olive	oil				 				.fl.dr.	4
(	Chlor	oform		٠.		 				.fl.dr.	4
ъ											

 Camphor-chloial
 ...m. 40

 Glycerin
 fl.dr. 4½

 Oil of sweet almonds
 fl.dr. 2¾

Three drops of this mixture on absorbent cotton to be placed in the ear twice a day, some also being rubbed behind the ear.

#### II. Acoustic Oils: '

A.	
Oil of turpentinefl.oz.	1
Oil of sweet almondsfl.oz.	
One to 2 drops on cotton in the ear.	

В.

Garlic, fresh							.av.oz.	1/2
Bay leaves							.av.oz.	1/2
Olive oil					 		fl. oz.	8

Boil together 15 minutes and filter while ot.

C.		
	Olive oil fl.dr.	2
	Oil of cajeputfl.dr.	2
	Oil of sassafrasfl.dr.	2
	Oil of rosemaryfl.dr.	2
	Camphorgr.	120
	Mix and dissolve.	

While all of the above will soften ear wax, the following is also employed for this purpose:

Boric :	aci	d.			 						.gr.	15
Glycer	in										fl.dr.	4
Water												

Mix and dissolve by the aid of heat. Warm 5 or 10 drops and put into the ear twice daily.

# Eczema or Salt Rheum, Remedies for.

Eczema is one of the numerous class of parasitic skin diseases like itch, ringworm, barber's itch, etc., and the treatment is consequently very similar.

Ι.		
	Rice powdergr.	240
	Talcum gr.	120
	Zinc oleategr.	60
	Bismuth subnitrate gr.	30
	8	

This should be dusted freely on the surface, and repeated every 2 or 3 hours. Sometimes, on account of the intense pruritus or itching, it is necessary to apply a cooling application; a solution of thymol, 1 to 1,000, is very good. Salicylic acid and menthol are also useful.

II.	
Diluted alcoholfl.oz.	6
Glycerinfl.dr.	
Cologne fl.dr.	
Tincture benzoinfl.dr.	3
Salicylic acidgr.	60
Mentholgr.	60

After applying this, the above powder should be dusted on freely. Continue the cooling treatment (thymol) as long as the active stage continues, and then use a mild ointment containing substances that influence the inflammation, such as the next formula.

11.	
Cold creamav.oz.	1
Petrolatumav.oz.	
Zinc oxideav.oz.	
Bismuth subnitrategr.	75
Ichthyolgr.	
Carbolic aciddrops	15
Apply twice a day. Used in mild sta	

IV.	
Zinc oxideav.oz. 4	
Chalk, powderav.oz. 2	
Lead waterfl.oz. 2	
Linseed oil	

Mix the chalk and zinc oxide; emulsify the lead water and linseed oil by shaking together. Finally mix the powders with the emulsion, rubbing constantly until a smooth paste is obtained.

V.		
	Zinc oxideav.oz.	2
	Sulphur, powderav.oz.	2
	Chalk, precipitatedav.oz.	2
	Linseed oilfl.oz.	2
	Lime waterfl.oz.	2

Mix the powders together, emulsify the oil and lime water by shaking together, and finally incorporate the mixtures by rubbing until homogeneous.

V1.	
Zinc oxide	.gr. 15
Talcum powder	
Tar	
Petrolatum	

Make into an ointment, and apply to the part morning and night.

VII.	
Tar ointmentgr.	60
Cold creamav.oz.	3
Zinc oxidegr.	120
Spread on lint and apply.	

VIII. Lassar's paste. See Part I.

IX. For other remedies, see under headings "Remedies for Barber's Itch," "Remedies for Itch," and "Ointments or Salves."

#### Eye, Remedies for Diseases of the.

The public demand for preparations for the eye is not large, but it is quite steady, and the formulas here given will suffice for the ordinary affections. If the condition of the eye seems at all serious or is long continued, the patient should invariably be advised to consult a physician who makes a specialty of ophthalmology.

The remedies for the eye may be either lotions or ointments. Suitable names are the following: "Reliable Eye Salve (or Water)," "White Eye Salve (or Water)," "Standard Eye Salve (or Water)," "Imperial Eye Salve (or Water)," etc.

I. Zinc sulphategr. 20	IX. See also Calamine Ointment and Calamine Cerate, Part I.
Morphine sulphate      gr. 20         Rose water       fl.oz. 4         Distilled water       fl.oz. 4	Feet, Remedies for Perspiration and Fetor of the.
Mix, dissolve and filter.	Perspiration and fetor of the feet is a tol-
II.       Boric acid, C. P.       gr. 80         Zinc sulphate       gr. 8         Morphine sulphate       gr. 8         Glycerin       fl.oz. 1         Rose water       fl.oz. 7         Mix, dissolve and filter.	erably common complaint, but does not appear to the public to be of sufficient importance to make it necessary to consult a physician. Most remedies recommended by pharmacists are in powder, by reason of the majority of them containing talcum with
III.	salicylic acid. These powders should always
Mercuric chloride         gr. 2           Ammonium chloride         gr. 12           Cochineal         gr. 3           Alcohol         fl. dr. 2           Water, enough to make         fl. oz. 16	be exceedingly fine, as otherwise they will prove irritating to the already sensitive integument.  I.
Mix, and filter after 12 hours.  This is known as "Mackenzie's Eye	Salicylic acid
Lotion:"	This is the "Salicylated Powder of Tal-
IV.	cum' of the N. F. The corresponding
Glycerite of hydrastis fl.dr. 1 Zinc sulphate gr. 8 Morphine sulphate gr. 8	preparation of the German pharmacopœia contains powdered starch instead of the boric
Borax	acid.
Glycerin	II.  Mercuric chloridegr. 1 Sodium salicylateav.oz. 1 Prepared chalkav.oz. 1
V. Zinc oxide gr. 5 Morphine sulphate	Dust a little of the powder in the socks
Morphine sulphategr. 2 Camphorgr. 1	every morning.
White wax	III.  Zinc oleate, powderav.oz. 1 Boric acid, fine powderav.oz. 2
Melt the wax, add the lard, allow to cool,	Talcum, powderav.oz. 3
when nearly cold add the camphor, allow it to dissolve, and then incorporate with the	IV. Salicylic acidav.oz.
other ingredients so as to make a thoroughly smooth ointment.	Alum, powder.       av.oz.       ½         Starch.       av.oz.       2         Oil of bergamot.       drops 54
VI. Burnt alumgr. 90	Alcohol
Opium, powdergr. 60	Dissolve the acid and oil in the alcohol,
Olive oil	and rub in mortar with the other ingredients until the alcohol is dissipated.
VII. Barium chloridegr. 6	V.
Calomel      gr. 10         Simple ointment       .av.oz. 1         Oil of rose      drop 1	Orris, powder
Use in scrofulous ophthalmia.	VI.
VIII. Yellow mercuric oxide gr. 5 Petrolatum	Salol
Reduce to a very smooth ointment.	_D.

VII.
Zinc oxideav.oz. 2
Starchav.oz. 234
Salicylic acidgr. 120
Talcum, powderav.oz. 15
Oil of wintergreendrops 12
—D.
VIII.
Potassium permanganategr. 8
Thymol gr. 16
Distilled water

### Female Disorders, Remedies for.

Remedies for female disorders are of several kinds. Many are uterine tonics (such as Nos, I and II below), these being intended to strengthen or "tone up" organs of gestation to fit woman to endure child bearing with comfort, to prevent, relieve, and cure distressing pains which occur from excessive or tardy menstruation, and to relieve the nervous disorders accompanying these complaints. These preparations are known by such names as "Catholicon," "Woman's Friend," "Female Remedy," "Mother's Friend," "Female Tonic," etc.

No III below is used as a galactagogue to increase the flow of milk in nursing women; No. IV as an anti-galactagogue or anti-lacteant to suppress or retard the flow of milk; and Nos. V and VI, as emmenagogues to assist the functions of the womb during or just preceding menstruation.

T.

Fluid extract of squaw vine fl. oz	. 4
Fluid extract of cramp barkfl.oz	
Fluid extract of blue cohoshfl.oz	
Fluid extract of damiana fl. oz	
Fluid extract of helonias fl.oz	
Fluid extract of cinchonafl.oz	
Sherry winefl.oz	z. 50
· ·	
II.	
Fluid extract of life root fl.oz	. 1
Fluid extract of helonias fl.oz	
Fluid extract of black hawfl.oz	
Fluid extract of cascarilla fl. oz	
Fluid extract of cascara sagrada fl.di	. 2
Fluid extract of Jamaica dog-	
woodfl.dı	. 6
Fluid extract of rhubarbfl.di	
Alcoholfl.oz	
Simple syrupfl.oz	
Simple elixir, enough to make, fl.oz	

L L .	I.	Į.					
-------	----	----	--	--	--	--	--

Fluid extract of castor oil plant	
leavesfl.oz.	12
Fennelav.oz.	4
Anise av.oz.	4
Wintergreen herbav.oz.	4
Simple elixir, enough to make fl.oz.	48

Mix the 3 drugs, reduce to coarse powder, and extract by slow percolation, using the elixir as a menstruum. When 36 fluidounces of percolate are obtained, add to it the fluid extract.

### IV. Make pills, each containing:

Sodium acetate							.gr.	3
Camphor							.gr.	1
Potassium nitrate							.gr.	1

#### 17

Saccharated carbonate of iron gr.	180
Myrrhgr.	
Ginger	
Nutmeggr.	00

Mix and reduce to fine powder. The dose is 30 gr. taken 3 times daily.

#### VI.

Myrrh		
Saffron	gr. 3	
Oil of clove	drop 1	

Mix and reduce to fine powder. This is sufficient for 1 dose, to be taken 3 times daily.

#### Frost-Bite Remedies.

The continued and repeated application of cold upon those portions of the body not well supplied with blood, such as the finger, toes and ears, is liable to produce chilblains or similar disorder; if this application of cold be severe and long continued, the result is frost bite, more or less severe, followed possibly by gangrene (complete destruction of tissue). If the frost bite is only moderately severe, any of the stimulant applications mentioned below will prove beneficial. If it be severe enough so that gangrene may supervene, the patient should be directed to place himself immediately under the care of a trustworthy physician.

#### т

Camphor powdergr.	90
Lanolinav.oz.	
Petrolatum av.oz.	1
Hydrochloric acidfl.dr.	1

To be applied evenings.

II.       Iodine      gr. 60         Ether      fl.dr. 10         Collodion      fl.oz. 4         Mix and dissolve by agitation.         Apply with a camel's hair brush.         III.         Tannin      gr. 30         Glycerin or spirit of camphor      fl.oz. 2	s o c t
IV.	1:
Peru balsam         fl.dr.         1½           Oleobalsamic mixture         fl.oz.         1           Cologne         fl.oz.         1	s c s
Paint on the affected parts.	Ι
V. Solution of chloride of iron	s
VI.  Diachylon ointment av.oz. 1 Lanolin av.oz. 1 Cottonseed oil fl.dr. 4 Carbolic acid fl.dr. ½ Oil of lavender drops 15	t I
VII. Other suitable preparations may be found under the head of "Ointments or Salves" and "Liniments," Part II.	I
Gonorrhœa, Gleet and Allied Diseases.	I
Remedies for gonorrhea may be used externally (so-called) or may be exhibited internally. The external remedies are usually in the form of "injections," or "washes," as they are also termed. These usually contain a zinc salt combined with hydrastis or one of its alkaloids, lead acetate, opium, car-	7
bolic acid, or other substance. The remedies for internal use contain copaiba or santal oil or both combined with cubeb, matico, spirit of nitrous ether, gum turpentine, eucalyptus, etc. These internal medicines may take the form of paste, capsules, pills or emulsions.	
The "external" and "internal" remedies may be used simultaneously, although there	7

This is to be taken only in the very first imes daily. II. Berberine hydrochlorate.....gr. 15 Zinc acetate .....gr. 15 Glycerin ......fl.dr. 4 Water, enough to make ......fl.oz. 8 Tincture of hydrastis .....fl.dr. Lime water.....fl.oz. Tannic acid.....gr. 20 Alum .....gr. 20 Zinc sulphate..... Water . . . . fl.oz. Zinc sulphate.....gr. 10 Lead acetate.....gr. 10 Compound tincture of catechu. .fl.dr. Tincture of opium.....fl.dr. Water .....fl.oz. VI. Zinc sulphate.....gr. 15 Extract of opium .....gr. 8 Glycerite of hydrastis ......fl.dr. Glycerin ......fl.oz. Water, enough to make .....fl.oz. 8 VII. Zinc sulpho-carbolate.....gr. 15 to 60 is no benefit to be gained by their conjoint Hydrogen peroxide.....fl.oz. 8 use. VIII. The "external" remedies usually are Balsam of copaiba ......fl.oz. 1 known by some fanciful or odd title, or by a Cubeb, powder ......av.oz. 1 number such as "55," "400," etc. The Matico, powder.....av.oz. 1 "internal" remedies are usually known by Light magnesia sufficient to form a mass.

uch titles as "Sandalwood Pills," "Paste of Copaiba, Cubeb and Santal," "Paste of Copaiba and Santal," "Paste of Copaiba, Cubeb and Matico," "Gonorrhœa Paste." 'Confection of Cubeb and Copaiba,'' "Mass of Copaiba," etc.

Remedies for external use may also be in he form of bougies; one example of these s given among the following formulas.

Every patient suffering with gonorrhœa should be advised to keep the bowels well open and also to refrain from the use of any stimulants during the course of the disease.

I.		
	Potassium citrateav.oz.	1
	Syrup of citric acidfl.oz.	2
	Water enough to make flor	

stages of the disease to render the urine alcaline. A tablespoonful is to be taken 3

IX.	
Resin	
Oil of sandalwoodfl.oz.	
Calcined magnesiaav.oz.	1/2

Melt the resin, add the oil, and stir in the magnesia when nearly cold.

8
4
4
4
2
1/

Melt the wax by the aid of heat, add the copaiba and with the powders form a paste.

#### XI.

Balsam of copaibafl.oz.	21/2
Glycerinfl.dr.	
Sugar, powderav.oz.	
Magnesia (calcined)av.oz.	21/2
Licorice root, powderav.oz.	2

Rub up the copaiba and glycerin intimately together, and then add the remaining ingredients gradually in the order named.-D.

#### XII.

Balsam of copaibaav.oz. 4
Oil of sandalwoodfl.oz. 1
Gum turpentineav.oz. 4
Cubeb, powdersufficient

Melt the turpentine with the copaiba by the aid of gentle heat, add the oil and incorporate in the melted mass as much of the cubebs as will form a suitable paste.

#### XIII.

Alum, powderav.oz.	1
Subcarbonate of ironav.oz.	
Cubeb, powderav.oz.	
Balsam of copaiba, enough to	U
form a suitable mass or paste	
form a suitable mass of paste	

#### XIV.

cubeb.

Balsam of copaibaav.oz. 8
Gum turpentineav.oz. 8
Oleoresin of cubebfl.oz. 1
Alum, powderav.oz. 1
Extract of licorice, powderav.oz. 2
Oil of wintergreen, sufficient to
flavor suitably.
Cubeb powder, sufficient to form

a suitable mass or paste. Melt the turpentine at a gentle heat, add the copaiba, add the oleoresin, alum and extract, allow to cool, and then add the oil and

XV.	
Oil of sandalwood gr.	300
Yellow waxgr.	

Melt the wax at the lowest possible temperature, add the oil, allow to cool, and divide into 100 pills or capsules.

VVI

v
Yellow wax
Balsam of copaibagr. 300
Oleoresin of cubebgr. 100
Iron sulphate, driedgr. 67
Carbolic aciddrops 33
Extract of belladonna, powder gr. 12
Oil of peppermintsufficient to flavor

Melt the wax at the lowest possible temperature, add the balsam, oleoresin, and acid, allow to cool, incorporate the extract, iron salt and oil, and divide into 100 pills or capsules.

Any of the pastes enumerated above may also be converted into pills or capsules.

#### XVII.

F 4 TT+	
Carbolic aciddrop	1
Zinc acetategr.	10
Iron persulphate, powdergr.	4
Extract of hydrastisgr.	
Extract of belladonnagr.	
Cacao butter, grated suffici	
0	

Convert this mixture into 30 bougies, each 2 inches long and about 1/4 inch thick. These may be rolled out on a pill tile or pill ma-See Bougies, Part I, for details. chine. These bougies may be advised for gonorrhea, gleet and spermatorrhæa (nocturnal emissions).

#### XVIII.

Fluid extract of eucalyptusfl.oz.	8
Mucilage of acaciafl.oz.	8
Balsam of copaiba fl.oz.	8
-	

Mix well by agitation.

#### XIX.

Balsam of copaibafl.oz.	2
Oil of sandalwoodfl.dr.	
Acacia, powder gr.	320
Oil of wintergreenfl.dr.	
Simple syrupfl.oz.	4
Water, enough to makefl.oz.	

Mix the balsam and two oils and triturate this mixture intimately with the gum; then add all at once 14 fluidrams of water, triturate rapidly until an emulsion is formed, and add the remainder of the water and the syrup.

#### Gout, Remedies for.

See under Rheumatism and Gout.

### Hay Fever Remedies.

I.
Borax
Capsicum
Ammonium carbonate gr. 30
All in fine powder and to be well mixed.
This is to be insufflated into the nostril
several times daily.
II.
Boric acidgr. 60

Boric acidgr.	
Sodium salicylategr.	75
Cocaine hydrochlorategr.	4

Make a powder.

This powder is to be insufflated into the nostrils frequently during the day. For the eyes, where affected, a wash of zinc or copper sulphate should be ordered.

#### III.

Cocaine hydrochloride gr.	6
Carbolic acidgr.	
Mentholgr.	
Oil of sweet almondfl.dr.	2
Zinc ointmentgr. 5	240

This is to be applied on a cotton pledget. The cocaine makes the above two formulas dangerous without specific warning against the continued use of the preparations.

#### Headache Remedies.

Headache remedies are now quite numerous and almost every pharmacist prepares a remedy to replace the "patented" articles. These remedies are usually put up in the form of powders, capsules, wafers, pills and tablets, but the ingredients are practically alike. The common ingredients of these preparations are acetanilid, phenacetin, caffeine, bromides, sodium bicarbonate (to correct acidity of the stomach), etc.

Some of the headache remedies appear in the form of effervescent salts; the ingredients are, however, similar to those of the other remedies.

Acetanilid gr. 7 Sodium bicarbonate gr. 2 Caffeine gr. 1
Make 1 powder, pill, capsule or tablet.
[.

Phenacetin .....gr. 10 Caffeine . . . . . . . . . . . . . . . . . gr. 1

Make 1 powder, pill, capsule or tablet.

÷	_		_		_	
ı		ı		П		

Acetanilid								.gr.	3
Caffeine								.gr.	1
Sodium bromide								.gr.	7

Make 1 powder, pill, capsule or tablet.

#### IV.

Acetanilidav.oz.	1
Sodium salicylategr.	125
Cerium oxalategr.	65

Mix, make 10 gr. doses, and form into powders, pills, capsules, etc.

#### V.

Caffeinegr.	20
Ammonium carbonate gr.	20
Elixir of guaranafl.oz.	1

One fluidram every hour until relieved. This is suitable for neuralgic headaches.

Oil of lavender flowers fl.dr.	1
Camphor av.oz.	
Water of ammoniafl.oz.	
Alcoholfl.oz.	16

Mix and dissolve. For inhalation and application to the forehead.

#### VII.

Cerium oxalate gr. 192 Sodium bicarbonate gr. 192 Caffeine citrate gr. 48 Magnesium carbonate gr. 48 Licorice root, powder gr. 96 Acetanilid av.oz 1 Oil of coriander drop 1 Oil of orange drops 2	•
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---

Mix intimately, reducing to very fine powder. Divide into 10 gr. powders, wafers or capsules, or make 5 gr. tablets or pills, directing the patient to take 1 of the former or 2 of the latter at a dose.

VIII. The Effervescent Potassium Bromide with Caffeine, Part I, will also be found serviceable.

IX. The latter may also be made with the addition of acetanilid; the product will resemble the different effervescent headache remedies of the market.

#### Itch, Remedies for.

The disease known as itch, or, more properly, scabies, is a very annoying and tolerably common complaint. It is a parasitic skin disease, and for this reason the mode of treatment is practically the same as for the other skin diseases.

HOUSEHOLL
The remedies mentioned below are all to
be applied several times daily.
1.         Red oxide of mercury       av.oz. 1         Burgundy pitch       av.oz. 1         Oil of turpentine       fl.oz. 1         Lard       av.oz. 16         Suet       av.oz. 16
Melt the pitch, add the suet and lard, mix well, allow to cool, add the oil, and then in- corporate thoroughly with the mercury oxide.
II.
Sulphur       av.oz. 2½         Birch tar, crude       av.oz. 2½         Prepared chalk       av.oz. 1¾         Green soap       av.oz. 5         Lard       av.oz. 5
This is "Hebra's Itch Ointment."
III.       Potassium nitrate, powder.       gr. 40         White hellebore, powder.       av.oz. 1         Sulphur       av.oz. 2½         Soft soap       av.oz. 2½         Lard       av.oz. 11
IV.
Sulphurated potassa or potassium sulphuret         gr. 300           Sodium carbonate         gr. 120           Lard         av.oz. 2           Soft soap         av.oz. 2           Olive oil         fl.oz. 1
Rub the sulphurated potassa to a very fine
powder and mix intimately with the other
ingredients.
V.
Menthol       gr. 110         Peru balsam       av.oz.       ½         Lanolin       av.oz.       10
Dissolve the menthol in the lanolin melted
at a very gentle heat and incorporate the balsam with this solution.
VI. Sulphurated potassa (sulphuret
of potash)
Make an intimate mixture.  This has been known as "Barton's Lo-
tion."
VII.
Ammoniated mercury.       gr. 10         Lead acetate.       gr. 30         Sulphur.       gr. 60         Carbolic acid       drops 10         Petrolatum       gr. 360

•	
VIII.	
	0
Green soapgr. 12	0.
Waterfl.oz.	8
IX.	
Storaxav.oz.	8
Olive oilfl.oz.	
Alcoholfl.oz.	1
Х.	
Sulphurav.oz.	8
Starch, powderav.oz.	
Oil of bergamotfl.dr.	
XI.	
Crude petroleumav.oz.	2
White way av oz	

Mix the petroleum, wax and alcohol in a flask until solution has taken place. Then add the soap, continue the heat, until the soap is liquefied, allow to cool and cut into bars or else pour into molds before quite hard. This is essentially a "petroleum soap."

Alcohol fl.oz. 2 Castile soap av.oz. 4

XII. Compound Sulphur Ointment, Part I, is an excellent itch remedy.

XIII. Other remedies enumerated under "Remedies for Barber's Itch," "Eczema Remedies" and "Ointments or Salves," may be employed for the troublesome affection.

### Kidney Remedies.

Some years ago preparations for the kidneys were marketed under the name of "Buchus;" there have also been "Diuretic Elixirs," and now these preparations are usually termed "Kidney and Liver Remedies." The latter are therefore to be recommended for affections of the liver, kidney and urinary organs. Many of the remedies for liver complaints may consequently be recommended for kidney affections and conversely many remedies for the kidney may be recommended for the liver.

Kidney remedies need not necessarily be of the liquid form; they may be in the form of coarse powder or species which should be infused with water before using.

I.	
Liverwortav.oz.	1
Hydrangeaav.oz.	1
Scoparius av.oz.	1
Canadian hemp (apocynum)av.oz.	1
Couch grassav.oz.	1
Potassium nitrategr.	
Alcoholfl.oz.	3
Syrupy glucosefl.oz.	3
Watersuffic	ient

Infuse the drugs with hot water so as to make 10 fluidounces of product, and to this add the alcohol and glucose.

Fluid extracts may be substituted for the drugs, in which case the alcohol should be omitted and the amount of water used be reduced to 8 fluidounces.

#### II.

Liverwortav.oz.	4
Jamaica dogwoodav.oz.	1
Couch grassav.oz.	
Gaultheriaav.oz.	2
Potassium nitrateav.oz.	1
Alcoholfl.oz.	32
Glycerinfl.oz.	
Water, enough to makegal.	1

Grind the drugs to coarse powder, percolate with all the glycerin and alcohol mixed with 32 fluidounces of water. When that has all passed add enough hot water to make 1 gallon, add the nitrate of potassium and dissolve.

#### III.

Fluid extract of buchu fl.oz.	3
Fluid extract of dandelionfl.oz.	3
Potassium acetateav.oz.	3
Fluid extract of juniper berries.fl.oz.	$^{2}$
Fluid extract of pareira fl.oz.	2
Fluid extract of stone rootfl.oz.	2
Simple elixir, enough to makefl.oz.	32

### IV.

Poppy heads, bruisedav.oz.	6
Waterfl.oz.	
Potassium nitrateav.oz.	1

Mix the poppy heads and water, boil until the liquid is reduced to about 8 fluidounces, express, adding, if necessary, enough water to make 8 fluidounces and in this dissolve the potassium salt.

Dose: One to 2 teaspoonfuls night and morning.

Buchuav.oz.	6
Juniper berries av.oz.	
Liverwortav.oz.	
Hydrangeaav.oz.	
Potassium acetate av.oz.	2

Mix the drugs, reduce to coarse powder, moisten with a menstruum consisting of a mixture of the above-mentioned amount of alcohol mixed with 32 fluidounces of water. Then extract by percolation in the usual way, using water as a menstruum when all of the above mixture has been consumed. Allow percolation to cease when 58 fluidounces of percolate have been obtained; in this dissolve the sugar and potassium acetate, and then add the spirit of nitrous ether.

VI. Any of the elixirs containing buchu in Part I may be dispensed as kidney remedies.

#### VII.

Buchu	 	 .av.oz. 8
Uva ursi	 	 .av.oz. 8
Juniper berries	 	 .av.oz. 4

Make into coarse powder.

#### VIII.

Chicoryav.oz.	9
Couch grassav.oz.	
Sennaav.oz.	
Red cloverav. oz.	
Rittersweet av.oz.	ŝ

All the drugs should be cut tolerably fine and be well mixed.

#### Liniments.

Liniments are in considerable demand and every pharmacist should have such a preparation ready to offer for sale. Some of these liniments may also be taken internally for cramps, cholera, diarrhæa, etc., and these may be known by such titles as "Pain Cure," "Pain Dispeller," "Pain Expeller," "Rapid Relief," "Pain Killer," etc. Other names which may be employed are "Stimulant Liniment," "Electric Liniment," "Embrocation," "Nerve and Bone Liniment," "Arnica Liniment," "Rheumatic Oil," "Rheumatic Liniment," "Universal Liniment," "Rocky Mountain Liniment,"

HOUSEHOLL	D REMEDIES. 191
"Penetrating Liniment," "Red Oil," "Indian Liniment," "Wizard Liniment," "Wizard Liniment," "Wizard Balm," "Golden Oil," "Knickerbocker Liniment," "Bicycle Liniment," etc. If it be white, it might be known as "White Liniment" or "Cream Liniment."  These liniments are recommended for rheumatism, neuralgia, bruises, chilblains, frost bites, sprains, stings and bites of insects, lameness, etc. Many of the preparations may also be employed for veterinary purposes; see also Part IV.  I.  Capsicum, powdergr. 60 Oil of origanum	VII.  Oil of tar
Kerosene oil	Tincture of arnicafl.oz. 16 Distilled extract of witch hazel fl.oz. 16
strain through muslin.	X. Oil of origanumfl.dr. 3
Spirit of camphor fl.oz. 1 Chloroform fl.dr. 4 Spirit of ammonia fl.dr. 4 Spirit of peppermint fl.dr. 4 Tincture of capsicum fl.dr. 4 Oil of sassafras fl.dr. 4 Oil of turpentine fl.oz. 1 Alcohol, enough to make fl.oz. 32	Oil of sassafras         fl.dr. 3           Kerosene oil         fl.dr. 10           Oil of turpentine         fl.oz. 20           Linseed oil         fl.oz. 25           XI.         Camphor         av.oz. 1½           Oil of turpentine         fl.oz. 23           Liquid petrolatum         fl.oz. 20
III.	Oil of origanumfl.dr. 4 Carbolic acidfl.dr. 4
Kerosene oil. fl.oz. 16 Spirit of ammonia fl.oz. 6 Spirit of camphor fl.oz. 5 Tincture arnica fl.oz. 5 Tincture of opium fl.oz. 4 Tincture of stramonium fl.oz. 4 Oil of origanum fl.oz. 4 Chloroform fl.oz. 3  IV.  Tobacco, rubbed to powder av.oz. 4 Tincture of arnica fl.oz. 24 Soap liniment fl.oz. 24	Ammonia water fl. oz. 5 Capsicum, powder av. oz. 3 Mix, macerate for 3 days, agitating occasionally and strain.  XII. Camphor av. oz. 1 Oil of amber fl. oz. 1 Oil of origanum fl. oz. 2 Crude petrolatum fl. oz. 4 Kerosene oil fl. oz. 10 Oil of turpentine fl. oz. 16 XIII.
Mix, macerate for 2 or 3 days, agitating	Castile soap, powderav.oz. 2
occasionally and strain.  V.  Tobacco	Oil of origanum. fl.oz. 2 Oil of turpentine. fl.oz. 2 Yolks of eggs. 4 Camphorated oil. av.oz. 2 Ammonia water. fl.oz. 8 Mix. Known as "White Oil." XIV. Capsicum, powder. av.oz. 1 Camphorated oil fl.oz. 1½
Tincture of capsicum	Oil of turpentine

X

and yolk—until they are thoroughly mixed, and add:

	Acetic acidfl.oz.	
	Water fl.oz.	16
ζ	V.	
	Tincture of capsicumfl.oz.	20
	Tincture of gingerfl.oz.	
	Tincture of myrrhfl.dr.	10
	Tincture of guaiac fl.dr.	18
	Tincture of opiumfl.oz.	21/2
	Camphor av.oz.	
	Alcohol, enough to makefl.oz.	64

Mix and dissolve the camphor by agitation.

#### XVI.

Oil of camphor (Japanese)fl.oz.	8
Oil of turpentinefl.oz.	
Benzine, deodorizedfl.oz.	16
Cottonseed oil fl.oz.	16
Capsicum, powderav.oz.	1

Macerate the capsicum with the benzine for 7 days, agitating frequently, and strain. Mix the oils of camphor, turpentine and cottonseed and add the previous liquid.

#### XVII.

Tincture of cantharidesfl.oz.	3
Tincture of myrrhfl.oz.	4
Tincture of guaiacfl.oz.	4
Oil of hemlock fl.oz.	2
Oil of turpentine fl.oz.	8
Cottonseed oil	32
Oil of camphor (Japanese)fl.oz.	16
Water of ammonia, strongfl.oz.	4
Solution of potassa fl.oz.	
•	

#### XVIII.

Oil of clove	.fl.dr.	3
Oil of origanum	.fl.dr.	4
Spirit of ammonia	.fl.oz.	4
Ether		
Alcohol	.fl.oz.	32

#### XIX.

The following is similar to certain preparations known by the term "Fluid Lightning:"

Aconitinegr.	2
Oil of mustard, etherealfl.dr.	2
Chloroformfl.dr.	2
Ether, strongerfl.oz.	
Alcoholenough to make fl.oz.	12

The above is a valuable external application for headache, rheumatism, neuralgia, and all nervous pains.

#### XX

Any of the liniments of Part I may also be employed.

#### Liver Remedies.

Remedies for the liver are usually termed "Liver Invigorators" or "Liver Regulators," very frequently "Kidney and Liver Remedies;" in fact most remedies for liver complaints are also recommended for derangements of the kidneys. Some of the liver remedies assume the liquid form, some are in the form of species, some in pill form (see Cathartics), etc. In addition to the formulas here mentioned, some of the blood purifiers, "bitters," cathartics, and kidney remedies may be recommended for assisting the liver in its functions.

#### Τ.

Fluid extract of rhubarbfl.oz. 2 Fluid extract of leptandrafl.oz. 2 Fluid extract of podophyllu m. fl.oz. 2 Compound tincture of gentian. fl.oz. 8 Compound tincture of cardamom fl.oz. 4 Tincture of gingerfl.oz. 2 Simple elixirfl.oz. 12
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### II

Fluid extract of leptandrafl.oz	z. 1
Fluid extract of podophyllumfl.oz	z. 1
Fluid extract of sennafl.oz	z. 5
Fluid extract of serpentariafl.oz	
Diluted alcohol, enough to make fl.o.	z. 64

#### III.

Leptandra, serpentaria, liverwort, senna, butternut, of each...av.oz. 2 Licorice root, anise, of each...av.oz. 1 Mix and reduce to coarse powder.

### Moles, For Removing.

Tartar emetic, fine powdergr.	30
Soap plasterdr.	11/2
Venice turpentinedr.	1/2

Mix intimately, and spread upon adhesive plaster. Apply firmly to the surface of the mole, and when suppuration sets in, remove.

#### Nervous Debility, Remedies for.

Of late it has become quite the fashion for the public in general to believe they are suffering from nervous disorders, and many socalled "nervines" have appeared upon the market. Some of these contain celery, others phosphorus and damiana, the latter also frequently containing kola, nux vomica, gentian, cinchona, or columbo.

These preparations may, according to their form or composition, be known as "Celery Compound," "Celery Nervine," "Celery Cordial," "Nerve Tonic," "Vitalizer,"

"Vitalizing Tonic," "Damiana Compound,"
"Nervous Debility Pills," "Aphrodisiac
Elixir," "Aphrodisiac Pills," "Compound
Damiana Pills," etc.

 I.
 Celery seed
 .gr. 384

 Catnip
 .gr. 640

 German chamomile
 .gr. 384

 Simple elixir
 .fl. oz. 12

 Diluted alcohol
 .sufficient

Percolate the mixed and ground drugs with the elixir and then pass enough diluted alcohol through the drug to make 16 fluidounces of product.

TT.

Fluid extract of celery seed... fl.dr. 10
Fluid extract of catnip.... fl.dr. 13
Fluid extract of chamomile ... fl.dr. 5
Diluted alcohol ... fl.oz 6
Simple syrup ... fl.oz 2
Glycerin... enough to make fl.oz 16

III.

Celery seed av.oz.	2
Red cinchonaav.oz.	
Orange peelav.oz.	1/4
Coriander seedav.oz.	1/4
	1/4
Muriatic acid	
Alcoholfl.oz.	
Glycerinfl.oz.	
Waterfl.oz.	4
Simple syrupfl.oz.	
Diluted alcoholsufficie	ent

Mix all the drugs and grind to a moderately coarse powder. Mix the acid, alcohol, glycerin and water; percolate the drug with this mixture, adding enough diluted alcohol to make 12 fluidounces. Add the syrup and if necessary filter. The flavoring may be altered to suit. Some like rose.

 $\mathbf{IV}$ 

Compound Elixir of Celery. See Part I.

7.		
	Cocaav.oz.	8
	Damianaav.oz.	8
	Gentianav.oz.	8
	Potassium bromideav.oz.	13/4
	Sodium salicylate av.oz.	1
	Dandelion av.oz.	8
	Alcohol fl.oz.	16
	Glycerin,	
	Waterof each, sufficie	ent

Mix 32 fluidounces of alcohol and the glycerin with 80 fluidounces of water. Also mix the coca, damiana, gentian, and dandelion, reduce to coarse powder, extract by lin afterwards.

percolating the previous mixture through it, in the percolate dissolve the salts, and then if necessary pass enough of the mixture of one volume of alcohol and 3 of water through the mixture to make the entire percolate measure one gallon.

 VI.
 Nux vomica
 av.oz. 3

 Damiana
 av.oz. 8

 Gentian
 av.oz. 8

 Columbo
 av.oz. 8

 Phosphoric acid
 fl.oz. 2

 Water,
 Alcohol

 Alcohol
 of each, sufficient

Mix the nux vomica, damiana, gentian, and columbo in ground form, percolate with a mixture of 1 volume of alcohol and 3 of water so as to obtain 62 fluidounces of percolate and to this add the acid

VII.

Make a mixture of drugs as in the preceding instance, percolate in the same manner, obtaining 56 fluidounces of percolate and to this add 7½ fluidounces of spirit of phosphorus.

VIII. Instead of the preceding, use one of the elixirs of Part I, containing phosphorus, damiana, and nux vomica.

IX

2 <b>1</b> .	
Celery seed av.oz.	4
Kolaav.oz.	11/2
Red cloverav.oz.	3
Cascara sagradaav.oz.	3
Simple syrupfl.oz. 1	6
Alcohol, water, of eachsufficier	ıt

Mix the drugs, reduce to coarse powder, percolate with a mixture of 1 volume of alcohol and 3 of water, to obtain 112 fluid-ounces, and to the percolate add the syrup.

This combines the "blood-purifying" laxative, and nerve-tonic properties.

X

۷,	.*	
	Phosphorusgr.	1
	Extract of damianagr.	200
	Extract of nux vomicagr.	
	Make into 100 pills	

Make into 100 pills.

Neuralgia Remedies.

I.					•			
	Menthol	 		 	 	 	.gr.	45
	Cocaine	 		 	 	 ٠.	.gr.	15
	Chloral	 		 	 	 ٠.	.gr.	10
	Petrolatum						or	300

Apply to painful part, covering with muslin afterwards.

II.
The ointment mentioned under Rheuma-
tism and Gout Remedies will be found serviceable as an anti-neuralgic ointment.
III.
Ipecacgr. 60
Quinine sulphategr. 100 Strychninegr. 1
Reduced irongr. 25
Make into 30 pills.
Label: One pill three times a day.
IV.
Atropine sulphategr. 1 Morphine sulphategr. 8
Camphor, powdergr. 120
Chloroform
Alcohol, enough to makefl.oz. 3
Dose: 30 to 40 drops.
V.
Oil of peppermint
Tincture of aconite
Apply every half hour or every hour.
VI.
Arsenic iodide gr. 1
Extract of belladonnagr. 8  Morphine valerianategr. 8
Extract of gentiangr. 5
Fluid extract of aconite rootdrops 5
Make into 60 pills.  Label: Take from 1 to 3 pills in twenty-
four hours.
VII.
Chlor. hydrate, camphor, each, av. oz. 11/2
Morphine sulphategr. 20 Atropine sulphategr. 1
Chloroform II.oz. 1
Mix. Dose: 10 to 20 drops.
Nipples, Cures for Fissured.
(Mammillary Lotions, Ointments, etc.)
I.
Ichthyolgr. 120
Lanolingr. 180
Glycerin
II.
Salicylic acidgr. 30
Tannic acidgr. 8 Boraxgr. 60
White waxgr. 120
Lard, benzoinatedgr. 360

	D FORMULARY.
	III.       Salol       .gr. 60         Cocaine hydrochlorate       .gr. 2         Ether       .fl.dr. 1         Collodion       .fl.dr. 4
	IV.
	Peru balsam         gr. 60           Oil of thyme         fl.dr.           Yolk of egg         1           Water         fl.oz.           Alcohol         fl.oz.
	Triturate the balsam, oil, and egg yoll
	together until an emulsion is formed, then add the water and finally the alcohol.
I	V.
	Lead nitrate
	VI. To prevent fissuring of the nipples, appl lanolin with the onset of labor four times dail till lactation is established. The nipples are then, after each nursing, anointed with the following:  Compound tincture of benzoin.drops 15 Olive oil
	Ointments or Salves.

The ointments mentioned below are useful applications for cuts, burns, ulcers, bruises, bites and stings of insects, frost bites, chilblains, bed sores, etc. Appropriate titles for these preparations are "Household Salve," "Arnica Salve," "Domestic Salve," "Healing Salve," "Carbolic Salve," etc.

Other ointments useful for all parasitic skin diseases may be found under the head of "Remedies for Barber's Itch," "Itch Remedies," and "Eczema Remedies."

Petrolatumav.oz.	16
Yellow waxav.oz.	$1\frac{1}{2}$
Camphorav.oz.	1
Carbolic acid, crystaloz.	
Oil of sassafrasdrops	30

Melt the carbolic acid and while warm add the camphor and oil of sassafras. Melt the wax and add to it the petrolatum, melting them together; while cooling but still liquid add the solution of camphor in carbolic acid, etc., and stir occasionally while cooling. The caustic properties of the carbolic acid are neutralized in this preparation by the camphor.

II.

White waxav.oz.	4
Lardav.oz.	12
Carbolic acid, crystalav.oz.	1/2
Calomelgr.	240
Camphorgr.	

Prepare this like the preceding, thoroughly incorporating the calomel by frequent stirring until the ointment is almost solid.

III.

Solid extract of arnicagr. 1	20
Lardav.oz.	141/2
Yellow waxav.oz.	1 1/2
Hot watersufficie	ent

Dissolve the extract of arnica in the hot water, and thoroughly incorporate it with the lard and beeswax previously melted together.

IV.

Yellow waxav.oz.	
Petrolatumav.oz.	141/2
Arnica flowersav.oz.	4

Melt the wax, add the petrolatum, stir in the flowers, heat moderately for one hour, stirring frequently; strain and allow to cool.

Simple cerate or simple oint-	
mentav.oz.	15
Boric acidav.oz.	1
Make an intimate mixture	

VI.

Yellow wax								.av.oz.	2
Petrolatum								.av.oz.	16
Thymol								.av.oz.	1

Melt the wax, add the petrolatum, and then stir in the thymol.

### Pile Remedies.

Piles, or hemorrhoids, as they are more correctly termed, are a very common and very annoying affection. They are termed "internal" piles when they exist within the IV. sphincter controlling the muscles of the anus, and "external" piles when existing outside of this sphincter. Other terms also are used in describing them: Blind piles which are simply a varicose state of the veins without bleeding; itching piles, bleeding piles, which 2 per cent carbolic acid solution.

are accompanied by loss of blood at every evacuation, and mucous piles, when pus or mucus only is discharged.

Treatment of piles should be both constitutional and local. The constitutional treatment should consist of the taking of compound licorice powder or one of the "bitter waters" at night. For local treatment, a mixture of an astringent like nutgall, tannin, extract of witch hazel, extract of krameria, lead acetate or iron subsulphate, with an anodyne like opium, belladonna, conium, tobacco, stramonium, ergot, or morphine is considered advisable. Other agents sometimes added to this mixture are antiseptics like iodoform, tar. peru balsam, carbolic acid, betanaphthol or salol. This mixture may assume the form of an ointment or of suppositories. The former should be preferred for external, the latter for internal, piles. When the piles are only tolerably severe, these remedies afford prompt relief, but when quite severe, only surgical intervention will effect a cure.

Fluid extract of witch hazelfl.oz.	1
Peru balsamgr.	120
Fenugreekav.oz.	1
Petrolatumav.oz.	
Paraffin av.oz.	4

Melt the petrolatum with gentle heat and macerate therein the fenugreek, for half an hour; then add the paraffin and strain through cloth. When about to solidify, add the extract of witch hazel to which the balsam has been added, stir until cool.

II.

Nutgall, fine powderav.oz.	3/4
Opium, fine powdergr.	90
Lard, freshav.oz.	6

III.

Morphine sulphategr.	2
Olive oilfl.dr.	
Zinc ointmentav.oz.	
Nutgall, fine powdergr.	120

Chrysarobingr.	24
Iodoformgr.	
Extract of belladonnagr.	
Petrolatumor.	480

Before applying wash the parts with a

7	The state of the s
V.	XIII.
Bethanaphthol gr. 10	Tanningr. 36
Extract of ergotgr. 20	Morphine sulphategr. 4
Extract of belladonnagr. 20	Potassium iodidegr. 24
Lead acetategr. 50	Cacao buttersufficient
Opium, powdergr. 50	Make into 12 suppositories.
Simple ointmentgr. 250	
	XIV.
VI.	Iodoformgr. 30
Lead acetategr. 15	Extract of belladonnagr. 3
Extract of coniumgr. 45	Morphine sulphategr. 1½
Crocated tincture of opiumdrops 10	Cacao buttergr. 180
Peru balsamgr. 45	Make into 12 suppositories.
Simple cerategr. 375	
—Н.	XV.
This is known as "Hellmund's Narcotico-	Iodoformgr. 60
	reiu baisamgr. 120
balsamic Ointment.''	Cacao buttergr. 90
VIJ.	White waxgr. 90
Iron persulphate, powdergr. 40	Calcined magnesiagr. 60
Lardav.oz. 1	Make into 12 suppositories.
Oil of bitter almonddrops 2	One of these should be introduced after
VIII.	
	each evacuation.
Extract of belladonnagr. 1½	XVI.
Antipyrin gr. 24	Extract of witch hazelgr. 60
Salolgr. 24	Tannin
Cacao buttersufficient	Opium, powdergr. 4
Make into 12 suppositories.	Cacao buttergr. 180
IX.	Make into 12 suppositories.
	Table 1110 21 Supposition
Cocaine hydrochlorategr. 8	Rheumatism and Gout Remedies.
Oil of eucalyptus	Rheumatic remedies may consist of reme-
Extract of krameriadr. 2	
Cacao buttersufficient	dies for internal or for external use. Those
Make the following into 12 supposi-	for internal use are to be preferred as afford-
tories:	ing better results, but the effects are still
V.	more marked if accompanied by the use of a
X.	1 The second sec
Extract of belladonnagr. 1½	suitable application (liniment).
Iodoform(gr. 12 to) gr. 24	Rheumatism remedies are usually also rec-
Chrysarobin gr. 12	ommended for gout. The remedies for the
Cacao buttersufficient	external treatment of gout may consist of
X a.	liniments or ointments; an example of a
Extract of belladonnagr. 6	suitable gout ointment is mentioned below.
Tannic (or gallic, acidgr. 24°	
Cacao buttersuficient	(See also Liniments, in Part II.)
	I.
XI.	Potassium iodidegr. 240
Hydrastingr. 6	Potassium bromidegr. 240
Ergotingr. 6	Wine of colchicum seedfl.oz. 1
Hamamelingr. 6	Syrup of orange or sarsaparilla fl.oz. 2
Opiumgr. 6	Waterfl.oz. 5
Tannin gr. 12	Directions: A teaspoonful 3, 4 or 5 times
Cacao buttersufficient	
	a day.
XII.	II.
Iron subsulphategr. 36	Salicylic acidgr. 150
Morphine sulphategr. 5	Potassium citrategr. 300
Iodoformgr. 3½	Glycerinfl.oz. 2
Cacao buttersufficient	Simple elixirfl.oz. 4
	Mix and dissolve by agitation.
	man and dissolve by agricultur.

HOUSEHOLD	P REMEDIES. 197
Potassium acetate	IX.  Veratrinegr. 15  Alcoholfl.dr. ½  Lanolinav.oz. 1  Petrolatumav.oz. 1  Oil of bergamotfl.dr. 1  White oil of thymefl.dr. 1
Sodium salicylate	Dissolve the veratrine in the alcohol, add the other ingredients, and mix well.—H.  To be applied night and morning.  X. Many of the liniments may be employed externally for the relief of rheumatic pains; some of the "blood purifiers" and possibly some of the "bitters" may be rec-
Dissolve the oil in the alcohol, add the fluid extracts, then the other ingredients, and dis-	ommended for the cure of rheumatism.
solve by agitation.	Ringworm, Applications for.
V.	I.
Sodium salicylate	Naphthalin
Dose: A tablespoonful three times a day.	Lanolin
VI.	Apply night and morning.
Sodium salicylate	III.  A more effective application than the latter is a saturated solution of salicylic acid in collodion. For a time this application may be quite painful.  IV.  Mercuric chloridegr. 2
Label: A teaspoonful every 3 hours.	Compound tincture of benzoinfl.oz. 1
The following are intended especially for gout:	Mix and dissolve. Paint over the affected parts.
VII.  Mentholgr. 400 Chloroform, enough to makefl.oz. 8	Aromatic sulphuric acidfl.oz. 1 Spirit of nitrous etherfl.oz. 1 Creosotefl.oz. 1 Apply once a day.
Mix and dissolve. To be applied exter-	VI.
nally.	Goa powdergr. 12
VIII.	Lardav.oz. 1
Rhubarb       av.oz. 1         Senna       av.oz. 1         Coriander       av.oz. 1         Fennel       av.oz. 1         Licorice root       av.oz. ½	Apply freely to part affected.  Instead of goa powder a corresponding amount of chrysophanic acid may be employed.
Saffron av.oz. ½ Raisins av.oz. 20	Soothing and Teething Remedies.
Diluted alcoholpts. 8	These preparations usually have a title
Macerate for 14 days, express, and filter.  Dose: 1 to 3 tablespoonfuls daily.	like "Baby Soothing Syrup," "Anise Soothing Drops," "Infant Teething Syrup,"

"Baby Soother," "Teething Powders," etc. Most of the proprietary preparations of this kind depend for their "soothing" effect upon opium; others are simply carminative, depending for their value upon anise, sometimes combined with fennel, ginger, lactucarium, lupulin, etc. Inasmuch as the effects of opium are so pernicious, especially upon infants, it is best to refrain from giving formulas containing this agent or its chief There is also the disadvantage, with the use of opium, that it obscures the real difficulty; the infant may be suffering from a vital or dangerous disease which will result fatally if not properly treated by a competent physician.

In addition to the formulas for soothing remedies to be administered internally, there are given two others for application to the gums during dentition.

I.		
	Anisegr. 5	40
	Fennelgr. 2	30
	Carawaygr. 2	
	Gingergr.	25
	Lactucariumgr.	
	Lupulingr. 1	
	Diluted alcoholsufficie	
	Simple syrupfl.oz.	21

Mix the drugs, reduce to powder, percolate with diluted alcohol so as to obtain 9 fluidounces of product and to this add the syrup.

#### II.

Anise, bruisedav.oz.	11/2
Alcohol fl.oz.	
Simple syrupfl.oz.	6
Anise water, enough to makefl.oz.	

Macerate the anise in the alcohol for 5 days, filter, and to the filtrate add the remaining drugs.

#### III.

Aniseav.oz.	11/4
Fennel av. oz. Lactucarium gr. Hops gr.	1/2
Lactucariumgr.	$25^{\circ}$
Hops gr.	120
Diluted alcoholfl.oz.	
Simple syrup, enough to makefl.oz.	30

Mix the drugs, reduce to powder, percolate syrup.

Anethol	drops	50
Oil of fennel		
Alcohol	fl.oz.	71/2
Water	fl.oz.	31/2
Simple syrup	fl.oz.	14
Purified talcums	ufficie	nt

Dissolve the oil and the anethol in the alcohol, add to a mixture of the syrup and water, let stand a few hours, and filter through

#### V.

Ammonium bromideg	r. 960
Chlorofor mfl.d	r. 2
Fluid extract of coniumfl.d	r. 2
Tincture of henbanefl.d	r. 4
Syrup of glucosefl.o	z. 8
Water, enough to makefl.o	z. 16

1.	
Make powders, each containing	
Pepsin, saccharatedgr.	5
Charcoalgr.	
Magnesium carbonategr.	
Lactucariumgr.	1/6

### Syphilis Remedies.

The recommending of remedies for the use of syphilitics does not come within the province of the pharmacist. The formula below, known as the "Hot Springs Prescription," is given merely because of its more or less extended use. Other remedies which may prove useful are the "blood purifiers" containing potassium iodide.

Potassium iodidegr.	480
Iodine	2
Mercuric chloridegr.	3
Compound tincture of gentian, fl.oz.	2
Fluid extract of sennafl.oz.	1
Compound syrup of sarsaparilla.fl.oz.	8
Water enough to makefl.oz.	16

#### Throat Affections, Remedies for.

Under this heading will be mentioned such preparations as could not conveniently be classed under Cough and Cold Remedies.

1.		
	Morphine sulphategr.	10
	Ipecacgr.	40
	Gingergr.	40
	Tartaric acidgr.	30
	Oil of anisegr.	20
	Sugarav.oz.	16

Mix all the dry ingredients, which should with the diluted alcohol, and add the be in very fine powder, add the oil, mix again, pass through a fine sieve, and convert into a lozenge mass by adding a small amount | IV. of mucilage of acacia or of gum tragacanth, and then sufficient water. Roll the mass out and divide into lozenges of suitable size, spread these out on boards or trays in a warm place and when nearly dry turn them over and allow them to dry on the other

The above forms a good Expectorant Cough Lozenge.

#### II.

Cubebsgr. 140	į
Potassium chlorategr. 280	
Extract of licoricegr. 150	,
Sugarav.oz. 16	,
Pine targr. 20	

Mix the cubeb, potassium chlorate, and sugar, all in fine powder; add the powder, which may be either in powder or plastic form, then the tar, make into a mass like the preceding, divide into lozenges and dry as before.

These lozenges have enjoyed some reputation under the name of "Pine Tree Tar Lozenges." The licorice, cubeb and tar act as an expectorant and the potassium chlorate is intended to relieve soreness of the throat.

#### III.

Fluid extract of pyrethrumm. Pilocarpine hydrochlorategr.	$\frac{\frac{2}{3}}{100}$
Extract of licoricegr.	2100
Glycerinm.	1
Sugar, enough to makegr.	20

This is sufficient for one lozenge, which should be prepared like the preceding.

These lozenges give great relief from the uncomfortable sensations of heat and dryness which characterize many acute and chronic affections of the mucous membrane of the mouth and throat. The lozenge should be allowed to dissolve in the mouth, and one used every two, three or four hours, as necessary. The addition of 2 grains of ammonium chloride will often be beneficial in sub-acute inflammatory conditions of the mucous lining of the respiratory a good purpose.

Cubebsav.oz.	1
Benzoic acidgr.	140
Extract of licoriceav.oz.	21/4
Tragacanthgr.	10
Morphine muriategr.	6
Sugarav.oz.	1
Oil of anisedrops	30
Currant jellyav.oz.	10

Mix all of the solids above, previously reduced to fine powder, add the oil and the jelly, form into a mass (by the addition of water or mucilage if necessary), roll this out like a thin pill pipe, and cut into troches weighing about 10 gr. each.

These lozenges are excellent for hoarseness and for coughs and colds.

•	
Sodium salicylatefl.	dr. 3
Fluid extract of cascara sagradafl.	
Glycerinfl.	dr. 2
Orange flower water, enough	
to makefl.	oz. 2

Label: A teaspoonful every 3 or 4 hours. This is excellent for tonsilitis.

Use 10 drops in a cup of warm water, morning and evening, as a gargle. This is an excellent remedy for quinsy.

#### Tonics.

A variety of preparations known by this title is grouped under this heading. Other tonic preparations may be found under the heading Bitters.

Solution of iron "protoxide"fl.oz.	2
Fluid extract of cinchonafl.oz.	4
Sodium phosphate av.oz.	1
Tincture of nux vomicafl.dr.	$^2$
Fluid extract of sennafl.oz.	1
Water, hotfl.oz.	2
Simple elixir, enough to makefl.oz.	16

Dissolve the sodium phosphate in the tract, while in more chronic affections 2 or 3 water, add the other ingredients, let stand for minims of the oleoresin of cubebs will serve 24 hours, and filter. This may be known as I" Iron Tonic Syrup."

Annual Control of the	
II.	VII.
	Compound ti
Fluid extract of gentianfl.oz. 1 Fluid extract of dandelionfl.oz. 1½	Syrup of coff
Phosphoric acid, diluted fl.dr. 10 Glycerin 8	Simple elixir
Sherry wine	Toe-Nail, In
Compound tincture of cardamom fl.dr. 6	Liquefied chl
Mix and filter.	Apply one dr
III.	Toothache R
Tincture of cinchonafl.oz. 2	Formerly all
Compound tincture of gentian, .fl.oz. 1	pared in the liqu
Tincture of capsicumfl.dr. 1 Fluid extract of hydrastisfl.dr. 2	they were terme
Fluid extract of hydrastisfl.dr. 2 Simple elixir, enough to makefl.oz. 16	pellets, each for
Mix, let stand a few hours, and filter.	of the tooth, ca
	been succeeded
IV.	called "toothacl
Soluble tincture of ginger (Part IV.)fl.oz. 4	which have be
(Part IV.)	consist of a fat
Glycerite of hydrastis	wax or spermac
Sugar         av.oz. 2           Alcohol         fl.oz. 5	carbolic acid, c
Water	phor, etc. Th
Mix, dissolve by agitation, and filter.	sticks or pencils
This may be known as "Ginger Tonic."	saturated with
V.	cut into suital
Hops	similar sticks.
Dandelion $1\frac{1}{2}$	cement is also
Podophyllum	Inasmuch as
Water, boilinggal. 1	proprietary toot
Alcohol	can always "pu
Mix the drugs, reduce to coarse powder,	even find it ackinds of tooth
pour on the water, let stand for 12 hours,	for sale.
decant 7 pints of clear liquid and to this add	Tor saic.
the alcohol.	I.
This preparation has been known as	Morphine
"Hop Tonic."	Acetic acid .
VI.	Alcohol Chloroform .
Cinchona	1
Bitter orange peelav.oz. 1½ Wild cherry barkgr. 100	Dissolve the
Cinnamongr. 60	alcohol, and the
Calamusgr. 30	11.
Simple syrup fl.oz. 11 Alcohol,	Camphor
Water, of each enough to make fl.oz. 32	Oil of cajupu
Reduce the solids to a coarse powder, and	Chloroform . Oil of clove.
percolate with a menstruum consisting of 2	
volumes of alcohol and 1 volume of water	Mix and diss
until 21 fluidounces of percolate is obtained.	III.
Add the syrup, let stand a few days, then	Oil of clove.
filter.	Spirit of ethe

This preparation may be known as "Cali-

saya Tonic."

Compound tincture of gentian	fl.oz.	2
Syrup of coffee	fl.oz.	8
Simple elixir	fl.oz.	6

### growing.

loride of lime. rop at night.

### emedies.

toothache remedies were preuid form-''toothache drops'' ed. Several years ago, pills or r one insertion into the cavity ame into use. These have d lately by pencils or sticks, the wax'' or "toothache gum," ecome very popular. These tty body like yellow or white ceti with which is incorporated reosote, chloral hydrate, camis is then formed into small s, or else absorbent cotton is this mixture, and this is then ble pieces and formed into A formula for an odontalgic given.

s the public rarely asks for a thache remedy, the pharmacist oush " his own article; he may dvantageous to have several ache remedies prepared ready

Morphinegr.	60
Acetic acid fl.dr.	
Alcoholfl.oz.	3
Chloroform fl.oz.	7

alkaloid in the acid, add the en the chloroform.-H.

Camphorav.oz.	2
Oil of cajuputfl.oz.	
Chloroformfl.oz: 5	
Oil of clovefl.dr. 5	2

solve by agitation. -H.

Oil of clovefl.c	z.	2
Spirit of etherfl.c		
Tincture of opiumfl.c		
imetate of optomittions		T

IV.	wax paper or tir
Chloral hydrateav.oz. ½	closed vials.
Camphor	
Morphine sulphategr. 8 Peppermint oilfl.oz. 11	XII.
Rub the solids in a mortar until liquefied	Iodol
and add the oil.	Paraffin oil
	Venice turpenti Yellow wax
V. Creosote	Alkanet root
Chloroformfl.oz, 2	Tuituunta tamati
Alcohol	Triturate togetl
Spirit of soap	tine, and paraffin
VI.	wax colored with substituted for th
Morphinegr. 3 to 6 Oil of peppermintfl.dr. ½	substituted for th
Carbolic acidfl.dr. 2	XIII.
VII.	Mastic Oil of clove
Camphor	Carbon bisulph
Chloroformfl.oz. 1	Amber, powder
Etherfl.oz. 1	Opium, powder Tannin, powde
Tincture of opiumfl.dr. 4	rannin, powde
Oil of thyme       fl.dr. 4         Oil of sassafras       fl.dr. 4         Alcohol, enough to make       fl.oz. 16	Dissolve the r
Alcohol, enough to makefl.oz. 16	carbon, and add t
VIII.	ders previously m
Creosotedrops 60	
Oil of clovesdrops 16	XIV.
Oil of cinnamon drops 16 Alcohol	-Paraffin
	Burgundy pitc Oil of cloves
Directions: Put one drop on a pledget of	Carbolic acid
absorbent cotton and apply.	
IX. Camphorgr. 60	Melt the parat
Camphorgr. 60 Peru balsamgr. 60	add the other in and make mass i
Extract of opiumgr. 60	cotton and cut in
Masticgr. 120	cotton and cut in
Chloroformfl.oz. $2\frac{1}{2}$	XV.
X. White or yellow waxav.oz. 3	Oil of clove
Venice turpentineav.oz. 1¼	Carbolic acid,
Mastic, powderav.oz. 1/2	Yellow beeswa
Opium, powder. gr. 130 Chloral hydrate gr. 110	While still liqu
	absorbent cotton
Melt the first three together, then add the	roll them into the
other ingredients, and stir frequently while	snip off a little
cooling.	introduce into the
XI. Salicylic acidgr. 10	
Salicylic acid	XVI.
Oil of clovegr. 10	White wax or s
Oil of cajuputgr. 20	Carbolic acid, o
Masticgr. 20 Dragon's bloodgr. 20	Chloral hydrate
Venice turpentinegr. 20	Aelt the fat at
Yellow waxgr. 120	and chloral, stir
Melt together and mix at a moderate tem-	merse cotton in
perature, roll out into little rods, cover with	ceding instance.

wax paper or tinfoil, and preserve in well closed vials.

Iodol gr.	15
Paraffin oilgr.	
Venice turpentinegr.	10
Yellow waxgr.	
Alkanet rootsufficient to co	lor

Triturate together the iodol, venice turpentine, and paraffin oil, then add to the melted wax colored with the alkanet. Salol may be substituted for the iodol.

Mastic av.oz.	2
Oil of clovefl.dr.	
Carbon bisulphide fl.oz.	5
Amber, powderav.oz.	
Opium, powder	1
Tannin, powder av.oz.	1/2

Dissolve the mastic in the bisulphide of carbon, and add the oil of clove and the powders previously mixed.

-Paraffingr.	180
Burgundy pitchgr.	
Oil of clovesfl.dr.	
Carbolic acidfl.dr.	1

Melt the paraffin and pitch together and add the other ingredients when nearly cold, and make mass into pills, cones or mix with cotton and cut into strips.

Oil of clovefl.dr.	2
Carbolic acid, crystalav.oz.	6
Yellow beeswaxav.oz.	1

While still liquid immerse thin layers of absorbent cotton and when sufficiently cool roll them into the shape of rods. For use, snip off a little piece, warm it gently, and introduce into the hollow tooth.

White wax or spermacetiav.oz	. 1
Carbolic acid, crystalav.oz.	1/2
Chloral hydrateav.oz	. 1

Aelt the fat at a gentle heat, add the acid and chloral, stir until dissolved, and immerse cotton in the mixture as in the preceding instance.

#### XVII.

Cocaine hydrochlorate	gr.	16
Opium, powder		
Menthol		
Althæa, powdered		
Mucilage of acaciasui		

Make into one-half grain pills and keep in well-stoppered vials. For use, one of these is to be inserted into the hollow tooth.—D.

#### XVIII.

Opium, powdergr.	15
Belladonna root, powdergr.	15
Pyrethrum root, powdergr.	
Oil of clovedrops	
Oil of cajuputdrops	
Oil of sweet almonds drops	
Yellow waxgr.	
8	

After melting together and allowing to cool form the mass into 100 pills which are sprinkled over with clove powder and properly preserved.

#### XIX.

Opium, powdergr. 60	
Pyrethrum root, powdergr. 30	
Creosote sufficient to form a mass	

Make into pellets weighing about one-half grain each.

#### XX.

Tanningr.	40
Opium, powdergr.	80
Ambergr.	80
Masticgr.	160
Oil of clove	40
Carbon disulphidefl.dr.	51/2

After dissolving the mastic in the carbon disulphide add the previously mixed powders.—D.

Chloroform may be substituted for the carbon disulphide, but the latter acts as an instantaneous analgesic, while its odor is marked by the clove oil.

This mixture should be inclosed in a well-stoppered wide-mouth bottle. For use, take out a small portion and insert into the carious teeth.

## Vermifuges.

Pharmacists are very frequently asked to recommend some remedy for the expulsion of worms in children. The various vermifuges are of different forms, such as syrup, solution, lozenge, powder, species and electuary. These may be entitled "Pleasant Worm Syrup," "Santonin Worm Lozenges," "Chocolate Worm Syrup," "Tonic Vermifuge,"
"Ideal Worm Powders," etc. An almost universal ingredient of these preparations is santonin; other common ingredients are oil or fluid extract of wormseed and pink root. These are usually combined with some purgative, such as castor oil, buckthorn, rhubarb, calomel, podophyllin, senna and jalap resin. Another addition sometimes made is some carminative like anise or fennel or the volatile oils of either of these.

The above described remedies are intended only for pin worms and lumbricoid worms. Tapeworms usually infest adult persons and require much different treatment. Formulas serviceable against the latter are also appended.

#### T.

Fluid extract of spigeliafl.oz.	5
Fluid extract of sennafl.oz.	3
Oil of anisedrops	10
Oil of carawaydrops	10
Simple syrupfl.oz.	

Dose, 1 or more teaspoonfuls at intervals until purging commences.

This formula is that of the old and familiar "compound fluid extract senna and spigelia," reduced one-half by the addition of syrup.

#### II.

•	
Santonica, fine powderav.oz.	1
Fluid extract of spigeliafl.oz.	3
Fluid extract of sennafl.oz.	3
Oil of chenopodiumdrops	10
Syrupy glucose, enough to make fl.oz.	
	Fluid extract of spigelia fl.oz. Fluid extract of senna fl.oz. Oil of chenopodium drops Oil of anise drops Oil of caraway drops Oil of fennel drops

#### TIT

4		
	Fluid extract of spigeliafl.dr.	4
	Fluid extract of sennafl.dr.	
	Fluid extract of buckthorn fl.oz.	
	Santonin	
	Alcoholfl.oz.	
	Cacao, powderav.oz	
	Simple syrup, enough to make.fl.oz.	

Make the caoao into a smooth paste with a portion of the syrup, heat to boiling, allow to cool, dissolve the santonin in the alcohol, add the fluid extracts, then the chocolate syrup and the remainder of the syrup, and mix the whole by agitation.

TXT		

Santoningr.	80
Alcoholfl.oz.	
Oil of chenopodiumfl.oz.	1
Fluid extract of chenopodiumfl.dr.	4
Castor oilfl.oz.	

Dissolve the santonin in the alcohol, add this solution to a mixture of the two oils, and to the whole add the fluid extract.

V. Powders may be prepared, each containing santonin, gr. 1, calomel, gr. 1, podophyllin, gr. 1-12 and sugar, gr. 30. The calomel may be omitted and the podophyllin increased to ½ gr., or either may be replaced by resin of jalap.

#### VI.

Santoningr.	50
Sugar, fine powderav.oz.	4
Tragacanth, fine powdergr.	$5\overline{0}$
Orange flower watersufficie	nt

Triturate the santonin to fine powder, add the tragacanth and sugar, mix well, make into a mass with the water, and divide into 100 lozenges. These are the "troches of santonin" of the U. S. P. Plain water may be substituted for the orange flower water.

The dose for a child 1 year old is 1 lozenge night and morning; of 2 years, 2 lozenges; of 4 years, 3; of 8 years, 4; of 10 years or more, 5 to 7 lozenges; in all cases to be taken twice daily, and continuing until worms are evacuated.

VII. To the preceding mass may be added podophyllin, calomet or resin of jalap; they may be colored pink with solution of carmine, cochineal color, or tincture of cudbear; or in place of the latter, powdered cacao or chocolate may be added.

The lozenges may be flavored with anise, fennel, wintergreen, or other flavor, and the gum tragacanth may be replaced by acacia.

#### VIII.

Spigelia	av.oz. 4
Manna	av.oz. 4
Senna	
Fennel	av.oz. 1

Cut the spigelia and senna, bruise the fennel, mix, add the manna and reduce the whole to a uniformly coarse powder.

In using, the above amount is sufficient for 7 pints of infusion or "tea," of which

half a teacupful is to be given to a child 2 years old morning, noon and night before eating.

### IX.

Spigeliaav.oz	. 6
Savinav.oz	. 1
Sennaav.oz	
Cream of tartarav.oz	. 2
Rhubarbav.oz	. 1/2
Fennelav.oz	. 4
Wormseedav.oz	. 2

Mix and reduce to coarse powder.

#### X.

	Extract of licorice, pure or puri-	
	fiedav.oz.	1/2
	Honeyav.oz.	2
- 1	Tamarind pulp, purified av.oz.	21/2
	Jalap, powderav.oz.	1/2
- 1	Santonica, powderav.oz.	2
	Male fern, powderav.oz.	2
	_	

Mix the first 3 substances and add the powders.—D.

The subjoined mixtures are intended for the destruction and expulsion of tapeworms:

#### XI.

Oleoresin of male fern	.drops	30
Pelletierine tannate	gr.	1
Glycerin		

The whole is to be taken in a liberal quantity of sweet milk immediately upon arising in the morning.

#### XII.

Oleoresin							
Calomel.	• •	 	 	٠.	• •	.gr.	5

Divide into 16 capsules.

Early in the morning, 1 capsule every 5 minutes, in a tablespoonful of sweetened water:

#### XIII.

The addition of the chloroform is said greatly to increase the efficacy of oleoresin of male fern. This is of practical value, as grave symptoms of poisoning have been observed after the ingestion of large doses of male fern. The following will prove of benefit:

Oleoresin of male ferngr.	20
Chloroform	40
Castor oilfl.dr.	11/2
Croton oildrops	3

This is sufficient for one treatment.

XIV.

Granatum, coarse powderav.or	z. 2
Watersuffici	ient 🕟
Castor oilfl.or	
Acacia, powderav.os	z. ½
Syrup of licoricefl.oz.	1

Mix the bark with 8 fluidounces of water, macerate for 10 hours, then heat on a water bath for 2 hours, express, heat the residue as before for 2 hours with 7 fluidounces of water, express again, mix the two liquids, evaporate them on a water bath to  $4\frac{1}{2}$  fluidounces, make an emulsion with this and the oil and gum, and finally add the syrup.—D.

An adult is to take one-half of this mixture upon arising, subsequently taking a cup of coffee or tea, and following in one-half hour with the remainder of the mixture.

XV.

Pumpkin seeds, deprived of the
outer membrane av.oz. 1
Water
Honeyav.oz. 1

Beat the seed and water together in a mortar to a uniform paste, then add the honey gradually, mixing the whole well.—D.

The above is recommended for tapeworm in children. No previous fasting is required. In the morning upon arising, the child is to drink a glass of milk; follow this in 1 hour by one-half of the electuary, in 15 minutes the other half, and follow in another 15 minutes with almost a tablespoonful of castor oil.

### Wart Eradicators.

Pharmacists are frequently called upon to recommend or offer some remedy for the eradication of the peculiar excrescences known as warts. These are peculiar in that sometimes a very simple remedy applied but once or twice will cause them to disappear and at other times they persist indefinitely in spite of all treatment.

Every one is familiar, of course, with the application of silver nitrate in sticks, concentrated nitric acid, creosote, carbolic acid, or salicylic acid in any of the forms of corn cures. Glacial, acetic and dichloracetic acid are frequently employed, while some add salicylic acid in the proportion of 1 in 16 to concentrated acetic acid. Another escharotic solution consists of chromic acid 1 part, water

5 parts. Unna recommends mercurial plaster containing 5 per cent of arsenic. It is also recommended to shave off the wart to the quick and then to apply a compress wet with a saturated solution of ammonium chloride. A corrosive collodion for warts consists of 1 part of mercuric chloride dissolved in 20 parts of collodion. Other applications are: Concentrated hydrochloric acid, solution of antimony chloride, solution of mercuric nitrate, potassium bichromate, arsenic oleate and copper oleate. Castor oil and oil of cinnamon are also useful. All of these remedies must be applied once or twice daily until the wart disappears. The outer hard, thick layer of cuticle should always be removed before applying anything whatever.

An ointment sometimes recommended is the following:

Verdigrisgr.	50
Savin, powdergr.	50
Soap cerateav.oz.	1

The following powder is said to be very effective:

Calomelgr.	30
Borie acidgr.	
Salicylic acidgr.	5
Cinnabar gr.	3

Rub into the wart 2 or 3 times a day.

In the case of multiple warts, where a large number appear within a short time, there is some constitutional derangement, and the patients are usually advised to take Fowler's solution in very small doses, or magnesium sulphate in 5 gr. doses 3 times daily. The following application has been recommended:

Sublimed sulphurgr.	120
Glycerinfl.dr.	
Acetic acid	

Apply repeatedly to each wart, continuing the treatment for several days. The warts dry up and then drop off.

Children's warts, appearing principally on the hands, may be removed by applying during several days solution of soda or potassa, and then covering them with collodion containing tannin. The same treatment applies for common warts.

concentrated acetic acid. Another escharotic Erasmus Wilson considers the application solution consists of chromic acid 1 part, water of caustic potassa in stick form to be the

quickest and most reliable eradicator, one treatment being all that is usually required.

Warts appearing on the skin of elderly persons must be looked upon as suspicious (cancer) and should be shown to a reliable surgeon.

#### Worm Medicines.

See Vermifuges.

#### Miscellaneous Remedies.

Under this heading are grouped remedies for such diseases as did not naturally fall into any special chapter:

### Pain Dispeller:

Aromatic spirit of ammoniafl.oz.	1
Compound spirit of etherfl.oz.	1
Spirit of peppermint fl.oz.	1
Compound tincture of lavender.fl.oz.	1
Spirit of camphor fl.oz.	1
Tincture of capsicumfl.oz.	1
Tincture of opiumfl.oz.	1
Tincture of rhubarbfl.oz.	1
Alcohol fl.oz.	3
Water, enough to makefl.oz.	16
,	

This is useful in diarrhœa, dysentery, cramps, etc., and may be known as "Pain Cure," "Pain Dispeller," "Carminative Balsam," "Pain Expeller," etc.

### Bibron's Rattlesnake Poison Remedy:

Brominegr.	100
Diluted alcoholfl.oz.	
Potassium iodidegr.	4
Corrosive sublimategr.	

Mix the bromine and alcohol, place the other ingredients in a mortar and add sufficient of the bromine-alcohol solution to dissolve them.

This has been recommended as an antidote to rattlesnake poisoning and is known as Bibron's antidote.

#### Poison Ivy:

The following has been recommended for poisoning by poison ivy:

Sodium sulphitegr.	60
Glycerinfl.oz.	1/2
Camphor water, enough to make fl.oz.	4

Apply several times daily.

Boil Remedy:

The following are recommended for boils or furuncles:

A.	
Salicylic acidgr.	120
Soap plaster av.oz.	
Lead plasterav.oz.	1

ь.		
Ichthyol	gr.	60
Resin plaster	gr.	60
Lead plaster.	gr. 1	20

These are to be applied daily on a cloth.

#### C. Furuncle Remedy:

Tincture of chloride of ironfl.oz.	1
Liquor potassii arsenitisfl.dr.	11/2
Aquafl.oz.	3

Take a fluidram after each meal.

Use, locally, camphorated carbolic acid and a poultice of linseed meal, when there is much inflammation. Also advise small doses of epsom salts to prevent constipation.

### Remedy for Bruises:

Ammonium chlorideav.oz.	
Alcoholfl.oz.	1/2
Water	5

Dilute acetic acid may be substituted for one-half the water and the alcohol may be replaced by 1 fluidounce of tincture of arnica.

This preparation is a valuable application for bruises or contusions; it is therefore of value in the treatment of ecchymotic conditions, such as "black eye." If applied at once and continuously for a time after the blow has been received, no discoloration will appear.

The following is also useful for bruises and "black eyes:"

Potassium nitrategr.	15
Ammonium chloridegr.	30
Aromatic vinegarfl.dr.	
Water, enough to makefl.oz.	8

#### Insect Bite Remedy:

Olive oil fl.oz.	1
Water of ammoniafl.oz.	
Oil of turpentinefl.dr.	1/2
Tincture of opiumfl.dr.	1/2

The above is recommended as an application to insect bites or stings.—H.

See also Liniments and Ointments in Part II.

#### Delirium Tremens:

Chloral hydrate gr.	90
Potassium bromide gr.	
Hoffmann's dropsfl.dr.	
Tincture of valerianfl.dr.	
Water, enough to makefl.oz.	6

Mix, dissolve and filter if necessary. This mixture is advised against delirium tremens or "jim jams" or the nervous condition bordering upon it,

Croup Remedy:	Erysipelas:
Fluid extract of senegafl.dr. 2	The following are employed for erysipelas:
Fluid extract of ipecacfl.dr. 1/2	I.
Oxymel of squill	Tannic acidgr. 100
Tartar emetic	Camphorgr. 100 Etherfl.dr. 14
Teaspoonful doses of this are recommend-	Mix and dissolve. Apply once or twice
ed for croup.	daily.
Anaphrodisiac Pills:	II.
The following pills are used as anaphrodi-	Sodium salicylategr. 160
siacs or to repress sexual excitement. Each	Sodium bicarbonategr. 80 Water, boilingfl.oz. 16
pill should contain:	
	Apply upon compresses.
Camphor gr. 3	Castor Oil Substitute:
Lactucariumgr. ¼	The following preparation has considera-
Guaranagr. ½ Extract of belladonnagr. ¼	ble sale under such names as "Castor Oil
Denarcotized opiumgr. 1/4	Substitute," "Castroilina," "Castorol,"
2011110011201 opinin 1111111111111111111111111111111111	"Castorbena," etc:
Brown-Sequard's Anti-Epileptic Mixture:	Sennaav.oz. 8
This mixture is in rather extended use, but	Pumpkin seedav.oz. 3
various formulas for it have appeared. The	Anise
following may be employed:	Rochelle saltav.oz. 2
7.	Sodium bicarbonateav.oz. 1
Sodium bromido or 180	Sugarav.oz. 56
Sodium bromidegr. 180 Potassium bromidegr. 180	Oil of wintergreenfl.dr. 2
Ammonium bromidegr. 180	Oil of peppermintfl.dr.
Potassium iodidegr. 90	Alcohol
Ammonium iodidegr. 90	
Ammonium carbonategr. 60 Tincture of columbofl.oz. 1½	Mix the three seeds, bruise to fine powder,
Tincture of columbofl.oz. 1½ Water, enough to makefl.oz. 8	add the senna, pour on 48 fluidounces of hot
,	water, macerate in a warm place for 3 hours,
Mix, dissolve and filter.	pour off the liquid, upon the dregs pour 16
II.	fluidounces of hot water, macerate for 1 hour,
Potassium bromide gr. 30	pour off the liquid, express the residue in a
Potassium iodidegr. 30	cloth, mix the two liquids, evaporate at a
Ammonium bromidegr. 30 Potassium bicarbonategr. 4	gentle heat to 32 fluidounces, add the soda
Infusion of columbo	and salts, dissolve by agitation, add the oils
Prepare like the preceding.	previously dissolved in the alcohol, and per- colate the whole through the sugar-
·	
Brown-Sequard's Neuralgic Pills.	Gross' Neuralgic Pills.
Each pill contains:	Each pill contains: Quinine sulphategr. 2
Extract of hyoscyamusgr. 2	Morphine sulphategr.
Extract of coniumgr. $\frac{2}{3}$	Strychninegr. 30
Extract of Ignatia beangr. ½	Arsenious acidgr. 30
Extract of opiumgr. $\frac{1}{2}$ Extract of aconitegr. $\frac{1}{3}$	Extract of aconite leavesgr. 1/2
Extract of cannabis indgr. \(\frac{3}{4}\)	"Hot Drops," or "No. 6."
Extract of stramoniumgr. 1/5	Tincture of capsicumfl.oz. 8
Extract of belladonnagr. ½	Tincture of myrrh fl.oz. 6 Mix and filter.

#### PART III.

## PROPRIETARY PREPARATIONS.

This department gave rise, in the first edition of the work, to some criticism based upon the argument that, inasmuch as the formulas given in connection with the designated preparations must, by reason of the extreme difficulty, if not the impossibility, of determining exactly the composition of many organic compounds, in numerous instances be more or less conjectural, that therefore all reference to them by their trade titles should be omitted.

This argument wholly mistakes the purpose of this Part and the nature of the formulas given.

The editors hold it to be the paramount duty of every pharmacist to know everything possible concerning the constituents of everything he may be called upon to dispense or sell. In this way only can he discharge his obligations in his professional relations to the physical welfare of his patrons. Secrecy and monopoly have, by the humane command of the professions, no legitimate place in medical science.

The formulas in this department are, therefore, primarily designed not to furnish information for duplicating the various preparations, but to give pharmacists and physicians an approximate idea of their composition and properties. It should be clearly understood that we quote the formulas simply for what they may be worth, considering the credibility of the source and considering also that manufacturers are at liberty to change their formulas at any time.

The reader should bear in mind that the titles of preparations in this department are. in nearly every instance, presumptively the property of their respective manufacturers, and can be legally used (until a court decision may otherwise decree) only to designate the original compounds. Competing preparations made by the retail druggist or others have no right in law or morals to valid trade-marked names; no self-respecting druggist will fraudulently substitute his own products when firmly in a percolator, add more menstruum,

another's is clearly indicated by its specific title. It is, however, the right and the duty of every pharmacist to encourage the use of his own products on their own merits, and in accordance with the recognized principles of honest and open competition.

The source of our information is given in most cases and is believed to be reliable, but should any formula, by reason of any incorrect statement or inference, be regarded as unjust to the original preparation or as impairing its reputation, the correction will be cheerfully made on receipt of the requisite information.

#### REFERENCE ABBREVIATIONS.

A. Dr	American Druggist
A. J. P	American Journal of Pharmacy
Apoth. Ztg	Apotheke Zeitung
B. & C. Dr	British and Colonial Druggist
Can. Dr	Canadian Druggist
D. C	Druggists' Circular
Era Form	Era Formulary
Fenner's Form	Fenner's Formulary
Ind. Ph	Indiana Pharmacist
Kilner's Form	Kilner's Formulary
Nat. Dr	National Druggist
N. I	New Idea
Pharm	Pharmacist
Ph. Central	.Pharmaceutische Centralblatt
Ph. Era	Pharmaceutical Era
Ph. Post	Pharmaceutische Post
Ph. Rec	Pharmaceutical Record
Ph. Rundsch	Pharmaceutische Rundschau
Ph. Ztg	Pharmaceutische Zeitung
	Registered Pharmacist

#### Abernethy's Pills.

## Each pill contains:

Socotrine aloesgr.	2
Extract of henbanegr.	S
Blue massgr.	
Ipecacgr.	-

-Fenner's and Beasley's Formularies.

#### Acetic Cantharidal Vesicant.

Cantharides,	freshly powdered av.oz.	8
	sufficie	

Moisten the drug with the liquid, pack

macerate for a few hours, and then percolate slowly until 16 fluidounces of liquid have been obtained.

Liquor Epispasticus or Blistering Liquid of the British pharmacopeia is one-half the strength of the above.—Pharm.

#### Actina.

Various formulas have been offered for similar preparations as follows:

I.	
Mentholgr.	60
Oil of mustardfl.dr.	2
Alcoholm.	30
Etherfl.dr.	
Sponge, enough to makeoz.	1
-Prof. F. B. Tiffany, K.	C.
II.	
Mentholgr.	60
Oil of mustardfl.dr.	2
Hydrobromic etherfl.dr.	1
Sponge, enough to makeoz.	1

#### Adhæsol.

Copal parts	70
Benzoinparts	6
Toluparts	6
Oil of thymeparts	4
Alphanaphtholparts	6
Etherparts	
-Merck's Bullet	

### Albolene.

According to manufacturers' statements relative to albolene and liquid albolene, these are presumably purified petrolatum and purified liquid petrolatum respectively.

#### Aletris Cordial.

I.

What purports to be a similar preparation is made as follows:

Stargrassav.oz.	1
Blue cohoshav.oz.	1
Cramp barkav.oz.	1
Helonias or false unicornav.oz.	1
Simple syrupfl.oz.	2
Alcoholfl.oz.	
Sherry wine, enough to makefl.oz.	16

Extract the drugs with the 16 fluidounces of wine to which the alcohol has been added, and obtain 14 fluidounces of liquid. Mix this with the syrup and filter if necessary.

II.

The following will make a satisfactory similar preparation (Fenner's Form.):

Unicorn root (aletris)av.oz.	2
Catnipav.oz.	1
Cramp barkav.oz.	1
Mitchellaav.oz.	2
Blue cohoshav.cz.	1/4
Cinnamonav.oz.	1/2
Orange peelav.oz.	
Carawaygr.	60
Sugarav.oz.	
Diluted alcoholfl.oz.	20
Watersufficie	nt

Mix the drugs, grind to coarse powder, moisten the diluted alcohol, pack in a percolator pass the remainder of the diluted alcohol through the drug, and then follow with water until 20 fluidounces of percolate are obtained. In this dissolve the sugar and to the solution add enough water to make 32 fluidounces.

#### Allcock's Porous Plaster.

Hager states it contains India rubber, Burgundy pitch, olibanum, myrrh and a small amount of oil of turpentine.

### Allen's Hair Restorer.

Sulphurgr.	160
Lead acetategr.	
Glycerinfl.oz.	
Water (perfumed)fl.oz.	
—Witts	

Ammonamide. (Ammoniated Phenylace-tamide.)

Probably similar to Ammonol, which see.

## Ammonol (Ammoniated Phenylacetamid).

This is claimed to be a derivative of the amido benzene series— $C_6H_5NH_2$ —containing ammonia in an active form. Dr. Eccles says the following mixture will give like medicinal results:

Acetanilid	parts 4
Sodium bicarbonate	
Ammonium carbonate	. part 1

### Amylocarbol.

Carbolic acidparts	9
Green soapparts	150
Amyl alcoholparts	160
Water, enough to makeparts	1000
Cobl	

#### Anaspaline.

Petrolatum .																	. part 1
Wool fat	•	•	•	•	•	٠	٠	•	٠	۰	•	•	•	•	•	•	parts 3

#### Anæstheto Obtundent.

A dental anæsthetic containing about 1 part of cocaine hydrochlorate in 30 of mixture, also carbolic acid, camphor, glycerin, oils of cinnamon and citronella, and probably alcohol.—Sadtler.

### Analgesin.

This is said to contain ammonium chloride, caffeine, sodium bicarbonate and acetanilid.

It should not be confounded with analgesine, which is the French name for antipyrine.

#### Anderson's Pills.

Т

Barbadoes aloes with some jalap and oil of anise.

#### II.

Aloes												parts 8
Myrrh.												parts 2
Saffron												

Mix and make into a mass with water and oil, and divide into 4-grain pills. This is said to be the original formula.

## Anesthyl.

Ethyl chloride.	 	 		 	parts	5
Methyl chloride	 ٠.	 		 	.part	1

### Angier's Petroleum Emulsion.

This is advertised to contain in each ounce 33½ per cent of purified petroleum and 12 grains of the salts of lime and soda. The following will yield a preparation with these essential ingredients in like proportions:

Liquid petrolatumfl.oz.	16
Acacia, powderav.oz.	8
Glycerinfl.oz.	
Calcium hypophosphitegr.	288
Sodium hypophosphitegr.	
Water, enough to make fl. oz.	

Add the acacia to the petrolatum and mix thoroughly in a large mortar, then add 16 fluidounces of water (all at once) and rub briskly until the emulsion is formed. Dissolve the hypophosphites in 8 fluidounces of water, to which add the glycerin; then add all to the emulsion and rub well together, adding any water necessary to make up the measure of 3 pints of finished product.

### Angostura Bitters.

Angostura barkav.oz.	4
Bitter orange peelav.oz.	2
Canada snake rootav.oz.	2
Cinchonaav.oz.	2
Serpentariaav.oz.	2
Gentianav.oz.	1
Galangalav.oz.	1
Calamusav.oz.	1
Cardamomav.oz.	1/2
Cinnamonav.oz.	1/4
Clovesav.oz.	1/4
Coriander av.oz.	14
Mace av.oz.	1/1
Alkanet rootav.oz.	1/2
Alcohol	/-
Water	
Water	10

Mix the drugs, reduce to fine powder, and extract with a mixture of the alcohol and water.

This makes an extract from which the bitters may be prepared by mixing 1 fluidounce of the extract with 24 fluidounces of alcohol, 40 of water, and 1 fluidounce of glycerin.—Fenner's Formulary.

#### Antidiabetin.

This is said to be a mixture of saccharin and mannite. It is marketed in three strengths—Nos. 70, 10 and 1—these numbers indicating the ratio in sweetening power to cane sugar.—Ph. Ztg.

### Antidiphtherikon.

Oil of birchparts	
Oil of beechparts	3
Alcoholparts	90
Potassium carbonatepart	1
Potassium sulphideparts	
Coblent	tz.

### Antikamnia.

This is claimed to be a "combination of coal-tar derivatives of the series  $C_N$   $H_{2N-6}$ ." One chemist has reported the following composition arrived at by analysis:

Acetanilidparts 47	
Sodium bicarbonateparts 50	
Tartaric acidparts 3	
-	
—A. J. P.	

Another chemist, Goldman, however, gives the following as the composition:

Acetanilid	parts	s 7
Sodium bicarbonate	parts	s 2
Caffeine		

#### Antikol (Anti calor).

The manufacturers state that it contains acetanilid, caffeine citrate, quinine bisulphate, and sodium bicarbonate, each 10 grains, containing 1 grain of caffeine citrate and 1-10 grain quinine bisulphate.

A writer in the Pharmaceutische Rundschau gives the following as the result of an analysis:

Acetanilidparts 30	0
Sodium bicarbonateparts	
Tartaric acidparts	

#### Antikrinin.

See Perl's Antikrinin.

### Arophene.

This contains about 1½ per cent of cocaine hydrochlorate, also carbolic acid, chloral, glycerin, oil of rose and probably alcohol. It is used as a dental anæsthetic.—Sadtler.

# Arsenauro. (Solution of Bromide of Gold and Arsenic: Barclay.)

This, the manufacturers claim, contains, in every 10 drops, 1-32 grain of gold bromide and 1-32 grain of arsenic bromide.

### Aubergier's Syrup of Lactucarium.

According to Procter, the following yields similar results:

Lactucarium, German.	 	gr.	60
Sugar			
Citric acid, powder			
Orange flower water			
Simple syrup			
Diluted alcohol,			
***		cc	

Water..........of each, sufficient Triturate the lactucarium with the sugar until reduced to powder. Put the mixture in a small funnel, and pour on diluted alcohol until the drug is exhausted or until 2½ fluidounces of percolate are obtained. Evaporate this to one-half fluidounce and add it to the syrup previously heated to boiling. Continue the ebullition slowly until the liquid measures 17½ fluidounces, add the citric acid, dissolve, strain, allow to cool, and add the orange flower water.

#### Ayer's (J. C.) Cathartic Pills.

Each pill is said to weigh nearly 4 gr. and consists of aloes, compound extract of colocynth, gamboge, capsicum and oil of peppermint.—Hager.

### Ayer's (J. C.) Cherry Pectoral.

Syrup of wild cherry	fl.dr. 6
Syrup of squill	fl.dr. 3
Tincture of bloodroot	fl.dr. 2
Sweet spirit of nitre	fl.dr. 2
Wine of antimony	fl.dr. 3
Wine of ipecac	
Simple syrup	fl.oz. 1½
Morphine acetate	gr. 2
Spirit of bitter almond	fl.dr. 1
	—N. I

### Ayer's (J. C.) Hair Vigor.

•	
Lead acetateparts	3
Sulphur parts	2
Glycerinparts 1	
Waterparts 8	30
—Popular Hygienic Journa	1.

### Ayer's (J. C.) Sarsaparilla.

Fluid extract of sarsaparillafl.oz.	3
Fluid extract of stillingiafl.dr.	
Fluid extract yellow dock fl.oz.	
Fluid extract podophyllumfl.dr.	21/2
Sugarav.oz.	
Potassium iodidegr.	
Iron iodidegr.	10

### Ayer's Recamier Cream.

According to N. I., its formula is similar to this:

-Pharm.

Zinc oxideav.oz.	2
Glycerinfl.dr.	
Waterfl.dr.	21/2
Spirit of rose (4 fl.dr. to 16 fl.oz.) fl.dr.	1

Triturate together until a perfectly smooth homogeneous mixture results.

In addition to the above, it is said to contain a small amount of corrosive sublimate.

### Ayer's Recamier Moth and Freckle Lotion.

This is said to contain corrosive sublimate dissolved in almond emulsion.—N. I.

#### Ayer's Recamier Powder.

Contains arrow root and zinc oxide. - N. I.

#### Ayer's Recamier Soap.

This is similar to ordinary toilet soap. - N. I.

#### Ayer's Vita Nuova.

Said to be port wine containing a small amount of cocaine.—N. I.

#### Barr's Dental Anæsthetic.

This is an alcoholic solution of oils of clove and peppermint.—Sadtler.

### Barry's Tricopherous.

According to N. I.. a substantially similar preparation is made as follows:

Castor oil	1
Alcoholfl.oz.	
Oil of lavenderdrops	
Oil of bergamotdrops	
Color pink with alkanet root	

## Color pink with alkanet root.

### Beckwith's Hog Cholera Remedy.

This, according to a patent specification, consists of equal parts of mandrake, sulphur, sodium bicarbonate, charred coffee and potassium chlorate.

### Beecham's Pills.

Aloes	)
Rhubarbgr. 90	)
Sodium sulphategr. 24	Ŀ
Saffron, truegr. 24	
—Ind. Ph.	
Mala inte 9 am mille	

## Make into 3 gr. pills.

## Begg's Ague Pills.

Each pill contains
Quinine sulphategr. 3/8
Cinchonine sulphategr. 1/4
Rhubarb gr. 3/8
With a little flavoring.—N. I.

#### Bejean's Gout Cure.

The formula is said to be as follows:	
Sodium salicylategr.	60
Potassium iodidegr.	
Extract of gentiangr.	
Alcoholfl.oz.	2
Oil of wintergreenfl.dr.	11/4
Waterfl.oz.	8

### Benzothymol.

According to the statement of composition by the manufacturers, this contains about the same essential ingredients as Lister's Antiseptic Solution, Part I.

#### Big G.

The following was contributed to the D. C. as making a preparation similar in appearance and effect:

Berberine hydrochlorategr.	15
Zinc acetategr.	
Glycerinfl.dr.	4
Waterfl.oz.	71/2

### Birnev's Catarrh Powder.

Cocaine hydrochlorategr. 19	
Magnesium carbonategr. 18	
Powdered peppermint leavesgr. 5	
Sugar of milk, enough to makeoz. 1	
Divide into 4 equal parts and put into 2-d	r

Divide into 4 equal parts and put into 2-divials.—Dr. Eccles.

# Bishop's Granular Effervescent Citrate of Magnesia.

This contains a mixture of sodium bicarbonate and tartaric acid.—Wittstein.

#### Blancard's Pills.

These are pills of iodide of iron.—Landerer.

## Blatta Insect Powder.

Persian insect powderav.oz. 8
Boraxav.oz. 8
Sulphurav.oz. 4
Oil of eucalyptusfl.dr. 2
II.
Persian insect flowersav.oz. 8
Boraxav.oz. 8
Sulphurav.oz. 4
Eucalyptus leavesav.oz. 4
Mix and reduce to fine powder.

### Boerhave's Bitters. .

Aloes	180
Cinnamonav.oz.	1
Galangalav.oz.	1
Zedoaryav.oz.	1
Angelicaav.oz.	1
Clovesav.oz.	1
Gentianav.oz.	1
Quassia, cutav.oz.	1
Waterfl.oz.	9
Alcoholfl.oz.	6
Sugarav.oz.	3
—Ha	ger.

### Boonekamp Bitters.

Socotrine aloesav.oz.	8
Myrrhav.oz.	4
Galangalav.oz.	4
Saffronav.oz.	4
Clovesav.oz.	1
Wormwoodav.oz.	1
Gentianav.oz.	4
Rhubarbav.oz.	4
Turmericav.oz.	4
Agaricav.oz.	8
Cinnamonav.oz.	1/2
Fennelav.oz.	16
Alcoholpints	12
Waterpints	
In the alcohol before it is mixed	with

In the alcohol, before it is mixed with the water, dissolve:

Oil of	wormwoodm	. 90
	fennelm	
Oil of	curled mintm	. 50
		T

#### Borol.

This is a fused mixture of boric acid and sodium bisulphate.—Ph. Central.

### Borolyptol.

Said to be a combination of 5 per cent of aceto-boro-glyceride, 0.1 per cent of formaldehyde, and pinus pumilio, eucalyptus, myrrh, storax, benzoin. The following may yield a preparation of similar composition:

Glyceride of boroglycerin, U.S.Pdr. 10	
Benzoic acid80	
Oil of eucalyptus	
Oil of pinus pumilio	
Formaldehyde solutionfl.dr. 2	
Water, distilled fl.oz. 16	

#### Borosol.

This is a colorless liquid containing, according to various analyses, aluminium tartrate, boric acid, salicylic acid, and free tartaric acid in aqueous solution.—Coblentz. Borosalyl.

This is said to be similar:

Boric acidparts	25
Salicylic acidparts	32
Water sufficie	

Triturate the acids with a small quantity of water to a smooth paste, dry and reduce to powder.

### Bor-Salicylate.

Appears to be made by triturating together 5 parts of sodium salicylate with 4 of boric acid and a small amount of water, drying and powdering.—Coblentz.

### Boschee's German Syrup.

According to N. I., the following is similar:

41 •	
Oil of tarfl.dr.	. 1
Fluid extract of wild cherryfl.dr.	. 6
Fluid extract of ipecac fl.dr.	. 4
Tincture of opiumfl.dr.	
Waterfl.oz.	
Sugarav.oz.	14
Magnesium carbonategr.	180
Rub the oil thoroughly with the me	

Rub the oil thoroughly with the magnesia in a mortar; mix the fluid extracts with the water, and triturate with the mixture in the mortar; filter and dissolve in the liquid the sugar without heat, and add the tincture.

### Bradycrotine.

According to N. I., the following furnishes a preparation approximating the original:

Caffeine, alkaloidgr. 6
Potassium bromidegr. 20
Sodium bromidegr. 20
Simple syrupfl.dr. 3
Alcohol
Port wine, enough to makefl.dr. 13
Caramelenough to color

### Brandreth's Pills.

Compound extract of colocynthgr. 3 Aloesgr. 13	
Gambogegr. 4	5
Castile soapgr. 3 Oil of peppermintdrops	
Oil of cinnamondrop	1

Glycerin and alcohol to make 100 pills.— Ph. Rundsch.

#### Brassicon.

Oil of peppermint	.fl.dr.	1
Camphor		
Ether	.fl.dr.	21/2
Alcohol	.fl.dr.	7
Essential oil of mustard	.drops	12
0.11 . 1 .		7.

-Suddeutsche Apoth. Ztg.

#### Bromidia.

According to the manufacturers, each fluidram contains 15 grains each of pure chloral hydrate and purified potassium bromide, and ½ grain each of genuine imported extract of cannabis indica and hyoscyamus. The formula below furnishes a preparation containing essentially the same ingredients in about the same proportions:

Extract of cannabis indicagr.	16
Extract of henbanegr.	16
Chloral hydrateav.oz.	
Potassium bromideav.oz.	
Saccharingr.	2
Water, enough to makefl.oz.	
Caramel,	
D '	nen f

Pumice, powder.....of each, sufficient

Triturate the extracts with the pumice to powder, dissolve the chioral, saccharin and potassium bromide in a portion of the water, filter this solution through a filter containing the powder, pass the remainder of the water through the filter and color the liquid with caramel.

### Bromophtharin.

Said to be a mixture of zinc oxide, calcium oxide, carbonate calcium, sodium sulphate, and sand.—Rundsch.

#### Bromo-Seltzer.

The following may yield a similar prepara-

Acetanilidav.oz.	
Tartaric acidav.oz.	21/4
Scdium bicarbonateav.oz.	21/2
Potassium bromideav.oz.	1
Sugar,av.oz.	2

#### Bronchiline.

The following has like ingredients in practically the same proportions as are claimed by the manufacturers for the original (N. I.):

Mulleingr.	64
Hoarhoundgr.	64
Senegagr.	64
Ipecacgr.	64
Sanguinariagr.	64
White pinegr.	64
Wild cherrygr.	256
Chloroformfl.dr.	1
Sugarav.oz.	14
Tar water,	
Alcohol of each, suffic	ient

Mix the drugs, grind to coarse powder, and percolate with a mixture of 3 volumes of alcohol and 8 of tar water, until 8 fluid-ounces of liquid are obtained. Percolate this through the sugar, adding enough tar water through the percolator to make the percolate measure 16 fluidounces, and to the latter add the chloroform.

## Brown's Male Fern Vermifuge.

The following makes a very similar preparation (N. I.):

Fluid extract of male fernfl.oz. 3
Simple syrupfl.oz. 5
Oil of wintergreendrop 1

#### Brown's Bronchial Troches.

This makes a good preparation of the	kind
Extract of coniumav.oz.	1/2
Acacia, powderav.oz.	2
Cubeb, powder av.oz.	2
Extract of licorice, powderav.oz.	8
Sugar, powderav.oz.	12
Oil of peppermint	

Make into a mass with water. Mix thoroughly and divide into lozenges of about 30 grains each.

Some formulas do not mention the oil of peppermint.—W. D.

## Bucklen's Arnica Salve.

Extract of arnicaav.oz.	2
Resin cerateav.oz. 1	6
Petrolatumav.oz.	4
Raisins, seedlessav.oz. 1	6
Fine-cut tobaccoav.oz.	1
Watersufficier	ıt

Boil the raisins and tobacco in 32 fluidounces of water until they are exhausted; express the liquid and evaporate down to 8 av.ounces. Soften extract of arnica with some hot water, mix the preceding liquid

with it, add this to the resin cerate, and petrolatum previously melted together.— Nat. Dr.

#### Bumsted's Gleet Cure.

The following is a favorite prescription:
Zinc sulphategr. 10
Extract of opium, aqueousgr. 60
Glycerite of hydrastisfl.dr. 1
Glycerin
Waterfl.oz. 4
-W. D.

#### Burnett's Cocoaine.

_	
	A similar preparation is as follows:
	Cocoanut oilav.oz. 4
	Cologne waterfl.oz. 4
	· —W D

#### Cactus or Cactine Pills.

An extolled proprietary preparation is claimed to represent in each pill the "active proximate principle of the cactus grandiflora." As neither a process for its isolation or preparation nor a test for identity appears in any standard work, the existence of this active principle in an isolated state has been questioned. Cactus or Cactine Pills may be prepared by saturating homœopathic sugar globules with the concentrated tincture of cactus grandiflora (the so called green drug fluid extract) and the alcohol removed by drying in the air.

#### Calisaya La Rilla.

The following is said (A. D.) to afford a satisfactory compound of similar composition:

atistactory compound or similar	,ompo	5101011
Calisaya, powder	gr.	640
Lime, calcined, powder		
Glycerin		4
Hydrochloric acid		10
Simple syrup		7
Oil of orange	.fl.dr.	3/4
Oil of lemon	drops	10
Oil of coriander	drops	5
Water,	-	
Alcoholof each	, suffic	ient
Fuller's earth		

Mix the calisaya and lime intimately, add 3½ fluidounces of water, stir well and let dry slowly. Percolate with a mixture of the acid and alcohol, adding sufficient alcohol to bring the bulk up to 4 fluidounces. To this add the oils, and, after shaking thoroughly, the glycerin. syrup and sufficient water to make 16 fluidounces; finally add the fuller's earth, shake well and filter.

#### Calolactose.

The following is the composition according to the manufacturers:

Calomelpart 1
Bismuth subnitratepart 1
Milk sugarparts 8

It is important that the ingredients be subjected to prolonged trituration.

## Camphoid.

Collodion	 	part 1
Camphor	 	part 1
Absolute alcohol	 	.part 1

## Campho-Phenique.

Claimed to be a chemical combination of phenol and camphor.

#### Carbolic Smoke Ball.

This is composed of licorice root, flour, white hellebore and some tarry body having the odor of carbolic acid.-N. I.

#### Carter's Little Liver Pills.

The following is said by the N. I. to make a similar preparation:

Podophyllingr. 1	1/2
Aloes, socotrinegr. 3	1/2
Mucilage of acaciasufficient	_
Mix divide into 12 pills and coat with suga	ır.

The entire 12 pills weigh about 71/2 gr.

#### Cascara Cordial.

Cascara sagrada	.av.oz.	3
Berberis aquifolium	gr.	525
Diluted alcohol	.av.oz.	7
Coriander	.av.oz.	1/2
Angelica,	gr.	25
Oil of anise		2
Oil of orange	.drops	2
Granulated sugar	.av.oz.	83/4
Fluid extract of licorice	gr.	180
Tincture of cudbear	suffic	elent
Water, sufficient to make	fl.oz.	32

Mix a decoction of the cascara and water at 212 deg. F. and filter, then dissolve the sugar in resulting liquid. Pack the coriander, angelica and berberis (in coarse powder) in a percolator, and displace with the alcohol, in which the oils have previously been dissolved. Lastly mix the cascara solution, the aromatic tincture and the tincture of cudbear together, and add the fluid extract of licorice.

#### Castoria.

According to the statements of composition made by the manufacturers, the following contains like essential ingredients:

Fluid extract of wormseed f	l.oz.	11/2
Fluid extract of pumpkin seed, i	l oz.	1
Fluid extract of sennaf	l.oz.	1
Fluid extract of rhubarbf	l.dr.	2
Potassium carbonate	.gr.	60
Rochelle salt	.gr.	720
Oil of peppermint		1
Oil of anised	rops	2
Oil of wintergreend	rops	5
Alcohot		1
Sugara	v.oz.	11
Water, enough to make	l.oz.	16
_		

Mix the fluid extracts of wormseed and pumpkin seed with 41/2 fluidounces of water, clarify by filtering through purified talcum, and add enough water through the filter to make filtrate measure 7 fluidounces. To this add the sugar, Rochelle salt and potassium carbonate, dissolve by the aid of gentle heat, add the fluid extracts of senna and rhubarb, and the oils dissolved in the alcohol, and finally enough water, if necessary, to make 16 fluidounces.

Celerina.

See Compound Elixir of Celery, Part I, which contains like essential ingredients as are claimed for the original, which are 5 grains each of celery, coca, kola, viburnum in each teaspoonful, together with aromatics.

#### Centaur Liniment.

The following is said (N. I.) to make a similar preparation:

I. For man:
Oil of pennyroyalfl dr. 4
Oil of thyme (white) fl.dr. 2
Oil of turpentinefl.dr. 2
Soapgr. 130
Caustic sodagr. 10
Water, enough to makefl.oz. 16
II. For beast:
Oil of spearmintfl.dr. 1
Oil of mustard
Oil of turpentinefl.dr. 2
Oil of amber, crudefl.dr. 4
Black oil
Soapgr. 130
Caustic sodagr. 10
Water, enough to makefl.oz. 16
Chamberlain's Relief.

Tircture of capsicumfl.oz. 1
Spirit of camphorfl.dr. 6
Tincture of guaiacfl.dr. 2
Coloringsufficient
—J. J. Pierson.

## Chapman's Alkaline Powder.

Sodium bicarbonate......av.oz. 16 Sodium phosphate, dried.....av.oz. 3

## Chapman's Internal Disinfectant.

This appears to contain sugar.-Eccles.

#### Chevalier's Life for the Hair.

Lead sulphidegr. 1	0
Iron sulphidegr.	1
Lac sulphurgr. 1	9
Glycerin fl.oz.	
Water fl.oz.	
	,

Flavor with rosemary and rose geranium.—Piper.

## Chlor-Lactated Elixir of Pepsin

According to the statements of the manufacturers regarding its composition, this is similar to Compound Elixir of Pepsin, Part I.

## Chlor-Lactated Pepsin Powder.

According to the statements of the manufacturers regarding composition, this is similar to Compound Powder of Pepsin, Part I.

#### Chlorobrom.

A mixture of equal parts of chloralamid and potassium bromide in solution.

#### Chlorol.

This is said (Arch. Med. Belge) to have the following composition:

Mercuric chloridepart	1
Sodium chloridepart	1
Hydrochloric acid part	• 1
Copper sulphateparts	
Waterparts 1	

The sodium chloride is added to render the solution more stable; the hydrochloric acid to prevent the decomposition of the corrosive sublimate in the presence of albuminoid matter; and the copper sulphate for its vomitive effects—in case the chlorol should be taken internally by mistake.

#### Chlorolin.

A solution consisting chiefly of mono and tri-chlorphenols.

#### Chloryl.

See Anesthyl.

#### Churchill's Syrup.

Compound syrup of hypophosphites.

## Christadoro's Hair Dye.

No. 1 contains 60 gr. of pyrogallic acid dissolved in 1 fluidram of alcohol and 4 fluid-ounces of distilled water.

No. 2 consists of 1 av.oz. of silver nitrate dissolved in 1 fluidounce of distilled water and 1 fluidounce of concentrated ammonia to which is added ½ av.oz. of gum arabic cissolved in 3 fluidounces of distilled water.— Era Form.

## Clark's Blood Mixture.

Potassium iodidegr.	128
Spirit of chloroformfl.oz.	
Solution of potassafl.dr.	
Waterfl.oz.	
Caramelsufficient to c	olor
_D	C.

#### Coaline Headache Powders.

Similar	powders a	re mad	e as	follows
	ne bromide			

Mix and divide into 12 powders.

#### Cobb's Pills.

Extract of henbanegr.	30
Extract of coniumgr.	30
Extract of colocynthgr.	40
Extract of nux vomicagr.	4

Mix, make a mass and divide into thirty pills.—Contributed to D. C.

#### Coddington's Asthma Powder.

A similar preparation may be made as follows (N. I.)

Potassium nitrateav.oz. 1
Anise
Stramonium:av.oz. 1
Lobeliaav.oz. 1

#### Coe's Dyspepsia Cure.

Rhubarb, powdergr.	120
Fluid extract of gentian fl.dr.	
Peppermint waterfl.dr.	
Sodium bicarbonategr.	

## Collins' Disinfecting Powder.

Chlorinated	lime,	dry	. parts 2
Burnt alum.			part 1
			-N. I.

#### Condy's Fluid.

•		
Potassiun	permanganategr.	75
Distilled	water fl.oz.	16

-N. I.

## Compound Lithia Tablets.

Calculating from the statement of composition furnished by the manufacturers, the following mixture may furnish a similar result:

Citric acidgr.	1,385
Lithium carbonate gr.	320
Sodium bicarbonategr.	315
Potassium carbonategr.	550
Talcum, purifiedgr	00
Make into 100 tablets.	

#### Conklin's Salve.

Resin.	av.oz. 1	2
Yellow	waxav.oz.	1
Mutton	suetav.oz.	1

#### Corassa Compound.—Kilner's Form.

•		
Sugar, powder	gr.	185
Gentian, powder		
Licorice, powder		
Sodium bicarbonate		
Cochineal, powder	or.	9
-	_	
—A. I	5. Ly	ons

#### Coudray's Eau de Quinine.

•	
Tincture of cinchonafl.oz.	
Tincture of cantharides fl. dr.	4
Spirit of soapfl.oz.	2
Cologne waterfl.oz.	5
Alcoholfl.oz.	5
Peru balsamgr.	192
Oil of bergamotfl.dr.	11/2
Oil of sweet orangefl.dr.	11/2
Oil of rose geraniumdrops	30
Brandy, enough to makefl.oz.	40
Cochinealenough to co	olor

Mix and filter.—Ph. Ztg.

#### Cram's Fluid Lightning.

Contributed by I. L. Fulton to W. D	. :
Oil of mustard, volatilefl.dr.	1
Oil of cajeputfl.dr.	1
Oil of clovefl.dr.	1
Oil of sassafrasfl.dr.	1
Etherfl.dr.	4
Tincture of opiumfl.dr.	6
Alcoholfl.oz.	10

#### Creme Simon.

This is stated (A. D.) to be composed of zinc white and powdered tale suspended in a mixture of glycerin and water and perfumed with oil of patchouly.

#### Cuticura Ointment.

Petrolatum containing boric and carbolic acids, about 2 per cent of the latter, perfumed with oil of bergamot and colored green either with chlorophyll or green aniline.—N. I

#### Cuticura Resolvent.

## Cuticura Soap.

This is toilet soap containing boric and carbolic acids.—N. I.

## Darby's Prophylactic Fluid.

Potassium	permanganate	gr.	5
	sulphate		
	carbonate		13/
	chloride		
	ugh to make		

## Darwin's Liniment.

Oil of wormwood	fl.oz.	1
Oil of thyme (red)		
Stronger water of ammonia		
Wood alcohol	fl.oz.	13

## Davis' (Perry) Pain-Killer.

•
The following is said to be similar (W. D.):
Tincture of myrrhfl.oz. ½
Tincture of guaiacfl.oz. 1½
Tincture of capsicumfl.oz. 1
Spirit of camphorfl. oz. 2
Alcoholfl.oz. 3
Alkanetsufficient to color
Macerate for a day, then filter.

# Davy's Urinal Cakes. (Disinfecting Solid.)

A mixture of resin with sulphates of copper, iron, zinc and sodium and some alum. These salts are probably powdered and mixed with the resin by fusion.—O. J. Bierbach.

## Dean's King Cactus Oil.

The following is suggested as yielding similar results:

Petroleum (neutral) oilfl.oz. 10	8
Kerosenefl.oz.	
Oil of mirbane enough to perfum-	e
Alkanet to colo	r

Filter the kerosene through alkanet root until it is sufficiently colored, add the petroleum oil, and perfume with oil of mirbane.

## Declat's Syrup of Phenic Acid.

The following makes a syrup of phenic acid:

Carbolic acid, pure	gr. 67
Glycerinfl.	
Simple syrupfl.	
Essence of cognacfl.	

#### Dental Anæsthetics.

For the composition of various proprietary dental ansesthetics, see the following in their alphabetical place:

Anæstheto Obtundent, Arophene, Barr's Dental Anæsthetic, Dental Surprise, Dickson's Improved Anæsthetic, Dorsenia, Eureka Dental Anæsthetic, Jessop's Dental Anæsthetic, Odontodol, Odontunder, and Weinman's Dental Anæsthetic.

## Dental Surprise.

This contains about one and one-half per cent of cocaine hydrochlorate and some carbolic acid.—Sadtler.

## Deshler's Salve.

According to N. I., this is similar to Compound Resin Cerate, U. S. P., 1870, which is prepared as follows:

Resinav.oz.	4
Suet av.oz.	4
Yellow waxav.oz,	4
Gum turpentineav.oz.	2
Linseed oilfl.oz.	
20.0	, .

Melt the wax, suet and resin, add the turpentine and then the oil, strain, and stir until cool.

#### Dextro-Quinine.

This is said to consist of quinidine with other alkaloids left after the extraction of quinine from red cinchona.—Fenner's Form.

#### Dextro-Saccharin.

Saccharin	1	p	art 1
Glucose,	crystalparts	1,000	to 2,000
	-		Fischer

## Dickson's Improved Anæsthetic.

This contains 4 per cent of cocaine hydrochlorate with some carbolic acid and chloral hydrate.—Sadtler.

#### Dioviburnia.

According to the manufacturers' statement as to composition, the following may yield a similar preparation:

Fluid extract of blackhawfl.oz. 3
Fluid extract of cramp barkfl.oz. 3
Fluid extract of wild yamfl.oz. 3
Fluid extract of star grassfl.oz. 3
Fluid extract of heloniasfl.oz. 3
Fluid extract of mitchellafl.oz. 3
Fluid extract of blue cohoshfl.oz. 3
Fluid extract of scullcapfl.oz. 3
Simple elixirfl.oz. 8

#### Dorsenia.

This is a dental anæsthetic containing onefifth per cent of cocaine hydrochlorate with some carbolic acid, camphor, and probably alcohol.—Sadtler.

## Dusart's Syrup.

A French proprietary preparation consisting essentially of syrup of lactophosphate of lime flavored with orange flower water.

## Dyche's Compound Capsules of Guaiacol No. 1.

Each capsule contains 5 minims of guaiacol, 10 minims of cod liver oil, and ¼ gr. of codeine.

## Dyche's Compound Capsules of Guaiacol No. 2.

Each capsule contains 5 minims of guaiacol, 10 minims of cod liver oil, and ½ gr. of iodine.

## Dyche's Compound Pills of Potassium Iodide.

Each pill contains 5 grains of potassium iodide and one-twentieth grain of red iodide of mercury.

#### Edison's Polyform.

This is said to make a similar preparation (Ph. Era):

Chloral hydrateav.oz. 1
Camphorav.oz. 1
Morphine sulphategr. 3
Chloroformfl.oz. 1
Etherfl oz. 4
Alcohol fl.oz. 6
Oil of peppermintfl.dr. 1

## Edwards' Alterative and Tonic Bitters.

Fluid extract of hopfl.oz. 1	
Fluid extract of red cinchona, fl dr. 4	
Fluid extract of sarsaparillafl.dr. 3	
Fluid extract of hydrastisfl.dr. 3	
Fluid extract of podophyllumfl.dr. 2	
Oil of wintergreendrops 24	
Oil of sassafrasdrops 12	
Oil of peppermintdrops 8	
Oil of lemondrops 8	
Sugarav.oz. 6	
Alcoholfl.oz. 16	
Water, enough to make fl.oz. 96	

Mix, dissolve the sugar and strain.—Era Form.

## Egyptian Eye Salve.

This is said to be composed of the following (N. I):

White resinav.oz.	12
Burgundy pitchav.oz.	1
Yellow waxav.oz.	1
Mutton suetav.oz.	1
Venice turpentineav.oz.	1
Balsam of firav.oz.	1

## Electric Headache Cures.

These all consist essentially of an alcoholic solution of essential oil of mustard, which is usually sprinkled on a small tuft of cotton inclosed in a wide-mouthed bottle. The twisted wires in these bottles are placed there simply to delude a gullible public.

## Elepizone.

Magnesium bromidegr.	180
Sodium bromidegr.	180
Waterfl.oz.	
Oil of cassiadrops	2
Simple syrup, enough to make fl.oz.	4
N	T

## Elixir of Lactopeptine.

The formulas in Part I, under the heading Compound Elixir of Pepsin, may furnish similar preparations.

#### Elixir Nutrans.

According to the makers' statements, the following and the next have like essential constituents in like proportions:

Fluid extract of cocafl.dr.	51/2
Fluid extract of damianafl.oz.	1
Fluid extract of kolafl.dr.	
Fluid extract of saw palmettofl.oz.	
Extract of beefgr.	256
Simple elixirfl.oz.	131/2
	-

Mix, dissolve the extract by agitation, and filter through purified talcum.

The published formula gives 2 ounces of fresh beef to the fluidounce.

#### Elixir Purgans.

Fluid extract of cascara sagrada fl.oz.	2
Fluid extract of senna, deodor-	
izedfl.oz.	1
Fluid extract of wahoofl.oz.	1
Fluid extract of blue flagfl.dr.	4
Fluid extract of henbane leaves, fl.dr.	4
Simple elixirfl.oz.	11

## Elixir of Salicylic Acid, Compound.

See above in Part I.

## Elixir Six Aperiens.

According to the statement of composition made by the manufacturers, the following yields a preparation having similar essential constituents in about the same proportions:

Fluid extract of caseara sagrada fl.oz	1
Fluid extract of podophyllumfl.oz.	1
Fluid extract of dandelionfl.oz.	
Fluid extract of butternutfl.oz.	
Fluid extract of sennafl.oz.	
Rochelle saltav.oz.	
Simple elixir, enough to makefl.oz.	16

#### Elixir Six Bromides.

According to the statement of composition by the manufacturers, the following contains like essential constituents in about the same proportions, the two last items being given in the advertisements simply as "cannabis indica and aromatics:"

Potassium bromidegr.	640
Sodium bromidegr.	
Ammonium bromidegr.	
Calcium bromidegr.	192
Lithium bromidegr.	64
Iron bromidegr.	64
Tincture of cannabis indicafl.oz.	1
Simple elixir, enough to makefl.oz.	16

## Elixir Six Hypophosphites.

According to the statement of composition by the manufacturers, the following has like constituents in about the same proportions:

Iron hypophosphitegr	. 32
Calcium hypophosphitegr	. 48
Manganese hypophosphitegr	. 16
Potassium hypophosphitegr	. 32
Quinine sulphategr	. 16
Strychnine sulphategr	. 1
Potassium citrategr	. 60
Sugarav.oz	. 4
Alcoholfl.oz	
Compound spirit of orangefl.dr	
Water, enough to makefl.oz	. 16

Dissolve the iron and manganese hypophosphites and potassium citrate in 2 fluidounces of water by the aid of heat, also the
calcium and potassium hypophosphites in the
same amount of water, and also the alkaloidal
salts in a similar amount of water. Mix all
three solutions, let stand for 24 hours, filter,
dissolve the sugar in the filtrate, add the alcohol containing the spirit, and finally add the
remainder of the water.

10

#### Elixir of Six Iodides.

The preparation of the same name in Part I has like essential constituents in about the same proportions.

#### Ely's Cream Balm.

These two widely different formulas have been published (W. D. and N. I.):

Bismuth carbonate.....gr. 15

I.

II

Thymol	$\frac{3}{2}$
I.	
White waxgr.	60
Paraffin waxgr.	30
Sweet almond oilfl.dr.	2
Petrolatumgr.	240
Nitrate of sodiumgr.	

## Eno's Fruit Salt.

Both of the following are said to resemble the original:

Oil of lemon......drops

Oil of orange.....drops

I.

Rochelle saltav.oz.	21/2
Tartaric acidav.oz.	31/4
Sodium bicarbonateav.oz.	$3\frac{3}{4}$
N	T

II.

Magnesium sulphateav.oz. 2
Magnesium citrateav.oz. 2
Potassium bitartrate av.oz. 2
Sodium bicarbonateav.oz. 2
Tartaric acidav.oz. 2
Powdered sugarav.oz. 4

Thoroughly dry, mix and preserve in well-stoppered bottles.—Monthly Mag. Phar.

## Eulyptol.

Carbolic acid	part 1
Oil of eucalyptus	part 1
Salicylic acid	parts 6
	hmelz.

#### Eureka Dental Anæsthetic.

This contains 3¼ per cent of cocaine hydrochlorate with some carbolic acid and oil of rose.—Sadtler.

#### Euthymol.

Lister's Antiseptic Solution, Part I, is of a similar type.

## Esencia de Calisaya.

Calisaya bark,	moderately	
coarse powder.	av.oz.	3
	.,gr.	
Water	fl. oz.	. 2
Cinnamon water.	fl.oz	101/2
Simple syrup	fl.oz.	101/2
Aromatic spirit	fl.oz	1 1/2
	av.oz.	
Alcohol,		
Diluted culphurie	said of such suffic	ciant

Diluted sulphuric acid, of each, sufficient

Slake the quicklime with the water, incorporate the bark, and dry the mixture. Then pack in a percolator and exhaust with hot alcohol so as to obtain 9½ fluidounces of percolate. If the alcohol used be insufficient to secure exhaustion of the drug, more may be added, the excess to be removed subsequently by evaporation or distillation. To the percolate add enough of the acid to precipitate all of the calcium, set aside for ten days, filter, add the remaining ingredients, let stand a few days, agitating frequently, and filter, returning the first portions of the filtrate until the liquid runs through clear.

## Espey's Fragrant Cream.

The following is similar:

Tragacanth, powdergr.	120
Glycerinfl.oz.	
Alcoholfl.oz.	
Waterfl.oz.	4
Oils of lavender, bergamot and	
rosemarysufficient to perf	ume

#### Essence of Pepsin.

See Part I.

## Exodyne.

Acetanilidparts	18
Sodium salicylatepart	1
Sodium bicarbonatepart	
Goldma	

# Febriline. (Tasteless Syrup of Amorphous Quinine—Lyons.)

Advertised as the active principle of amorphous quinine, each teaspoonful equaling 2 grains of the sulphate. This is said to yield a similar result:

Quinidine suspended in syrup, about 2 grains in 1 fluidounce, and flavored with spirit of lemon.—Eccles.

## Fellows' Syrup of Hypophosphites.

The manufacturers state that it contains potash, lime, iron, manganese quinine, strychnine and phosphorus; the whole combined in the form of a syrup with a slightly alkaline reaction. The formula below is based upon analysis and is claimed to furnish a preparation essentially similar to the original:

Calcium hypophosphitegr.	64
Potassium hypophosphitegr.	
Iron sulphategr.	
Manganese sulphategr.	
Quinine sulphategr.	
Strychnine sulphategr.	
Syrupy glucoseav.oz.	
Simple syrupfl.oz.	
Water, enough to make fl.oz.	

Dissolve the calcium and potassium hypophosphites in 2 fluidounces of water. Add to 1 fluidounce of the water 3 fluidrams of the syrup and dissolve in the mixture, by the aid of heat, the remainder of the salts. Mix the solutions, set aside for a few hours, filter into a bottle containing the remainder of the syrup, wash the filter with 1 fluidounce of boiling water, to the liquid add the glucose and then enough water to make 16 fluidounces.

## Flagg's Relief.

Oil of clovefl.oz. 2
Oil of sassafras fl.oz. 4
Spirit of camphorfl.oz. 3
•
-J. J. Pierson.

## Freligh's Tonic. (Phosphorized Cerebro-Spinal Tonic.)

According to the statement of composition by the manufacturers, the following contains the same essential constituents in about the same proportions:

Fluid extract of nux vomicafl.dr.	21/2
Fluid extract of ignatiafl.dr.	11/4
Fluid extract of yellow cinchona fl.oz.	614
Fluid extract of German chamo-	,
milefl.oz.	21/4
Fluid extract of gentianfl dr.	1
Fluid extract of bitter orange	
peel drops	15
Fluid extract of columbodrops	40
Fluid extract of cardamomdrops	
Spirit of phosphorusfl.oz.	
Oil of orangedrops	$2^{2}$
Oil of clovedrop	1
Oil of cassiadrop	1
Simple syrup	

## Frey's Vermifuge.

Castor oil	fl. oz.	1
Aromatic syrup of rhubarb	. fl.oz.	1
Oil of wormseed		
Croton oil		
	r's Forr	

#### Fruit Salt or Saline.

See Tarrant's Aperient, or Eno's Fruit Salt.

## Fluid Hydrastis.

This is a non-alcoholic water-miscible preparation of hydrastis, probably similar to Glycerite of Hydrastis, Part I.

## Ford's Balsam of Hoarhound.

Hoarhoundav.oz.	31/2
Licorice rootav.oz.	31/2
Waterfl.oz.	16
Camphorgr.	75
Opiumgr.	60
Benzoingr.	60
Squillgr.	120
Oil of anisefl.dr.	1
Alcoholfl.oz.	24
Honeyav.oz.	31/2

Macerate the hoarhound and licorice with the water for 12 hours, decant 12 fluidounces, to this add the remaining ingredients except the honey, macerate 7 days, strain, and add the honey.—N. I.

## Fosgate's Anodyne Cordial.

Fluid extract of rhubarb	fl.dr. 5
Fluid extract of rhatany	fl.dr. 2
Fluid extract of ginger	drops 6
Paregoric	
Simple syrup	fl.dr. 1
Diluted alcohol	fl.dr. 5
	_N I

#### Fossilin.

A petroleum product similar to petrolatum.

—Coblentz.

#### Franck's Grains de Sante.

Aloes		parts 4
Jalap		parts 4
Rhubarb		part 1
Syrup of worm		
, .		
Make into 2-gr	ain pilis.—Ca	idet.

#### Gelatol.

An ointment base consisting of a mixture of oil, glycerin, gelatin and water.—Helbing's Mat. Med.

#### Gargling Oil.

 Anmonia water
 fl oz.
 1

 Soft soap
 av oz.
 4

 Benzin
 fl.oz.
 4

 Oil of amber, crude
 fl dr.
 4

 Tincture of iodine
 fl dr.
 2

 Water
 fl.oz.
 20

Mix the petroleum and soap, add the ammonia water, oil of amber and tincture of iodine and aix thoroughly. Then add the benzin and finally the water.

#### Giles' Iodide of Ammonia Liniment.

Iodinegr.	15
Alcoholfl.oz.	8
Camphorgr.	120
Oil of lavenderfl.dr.	
Oil of rosemaryfl dr.	
Water of ammoniafl.oz.	
—D	C.

## Gilt Edge Butter Compound.

This contains 30 per cent of pepsin and 70 per cent of hydrous sodium sulphate, besides a trace of pink coloring matter.—Wiley.

## Gluten Suppositories.

These consist of cacao butter containing 10 per cent of wheat flour.—Vulpius.

## Glycerole Yerbine Compound.

This contains the same essential ingredients in the same proportions as are claimed for a preparation of the same name on the market:

Yerba santaav.oz.	31/2
Licoriceav.oz.	
Grindeliagr	480
Wild cherrygr.	
Potassium bromidegr.	
Pine targr.	45
Potassium carbonategr.	40
Salicylic acidgr.	32
Alcohol.	
Water,	
Glycerin of each, suffic	ient

Mix the yerba santa, licorice, grindelia, and wild cherry to moderately fine powder, add the potassium carbonate, and extract by percolation so as to obtain 16 fluidounces of percolate, using as a menstruum a mixture of equal volumes of alcohol, water and glycerin. To the percolate add the potassium bromide, tar and salicylic acid, set aside for several hours, agitating occasionally, and filter.

## Glycoline.

This is a liquid petrolatum, according to the manufacturers' statement.

#### Glyco-Thymoline. (Kress.)

Each fluidounce is said to contain sodium 24, boric acid 4, benzoin 4, salicylic acid 0.33, eucalyptol 0.33, Thymoline 0.17, betula lenta 0.08, menthol 0.08, pinus pumilio 0.17, glycerin and solvents sufficient. This does not differ materially from Lister's Antiseptic Fluid, Part I, containing, in addition, oil of pinus pumilio. Wild indigo is replaced by red birch.

## Glymol.

This is claimed to be a liquid hydrocarbon obtained from petroleum.

## Godfrey's Cordial.

Oil of sassafrasdrops	6
Tincture of opiumfl.dr.	41/4
Alcoholfl.dr.	
Potassium carbonategr.	60
Molasses	5
Waterenough to make fl.oz.	16

Mix the tincture of opium with the alcohol in which the oil has previously been dissolved. Dissolve the potassium carbonate in 8 fluid-ounces of water, mix this with the molasses, add the mixture first prepared, and then enough water to make 16 fluidounces; allow the mixture to stand until it has become clear, and decant the clear liquid which is to be used.—N. F.

## Gombault's Caustic Balsam.

This is said to consist of about the following (N. I.):

Oil of red thymeparts	3
Oil of amber, rectified parts	8
Oil of rosemaryparts	
Camphorparts	20
Alcoholparts	
Sulphuric acidparts	
Oil of turpentineparts	
Sulphurated oilparts	

Mix the ingredients properly, stirring the acid into the other liquids mixed, and allow to cool.

## Gouraud's Oriental Cream.

This consists of calomel and water, about 39 grains of the former to 1 fluidounce of the latter.—N. I.

## Graham's Cucumber and Elder Flower Cream.

Sweet almond oilfl.oz.	5
Saturated aqueous solution of	
boraxfl oz.	1
Oil of lemondrops	16
Oil of bergamotdrops	
Oil of bitter almonddrop	
Con T	

## Gray's Glycerin Tonic Compound.

Each fluid half-ounce is stated to contain dilute phosphoric acid 12 minims, gentian root 10 grains extract of taraxacum 15 grains, glycerin 80 minims, sherry wine 80 minims, carminatives q. s. An essentially similar preparation is yielded by the following formula:

Compound fl. ext. of gentianfl.oz.	1
Fluid extract of dandelionfl oz.	1
Dilute phosphoric acidm.	384
Glycerinfl.oz.	5
Sherry wine, enough to makefl. oz.	16

## Gray's Specific Pills.

Nearly all of these "specific" pills are composed (W. D.) of asafetida with a little camphor and sometimes hops, or lupulin, e. g.:

nor and sometimes nops, or rupum, c. g.	
Asafetidagr. 2	
Camphorgr. 1	
Lupulingr. 3	8

The "specific action" is in the direction of an anaphrodisiac.

#### Green's August Flower.

The following produces a similar preparation (Coblentz):

Dhuhanh		900
Rhubarb		
Golden seal	.gr.	90
Cape aloes	.gr.	16
Potassium carbonate	.gr.	120
Peppermint		
Capsicum	.gr.	5
Alcoholfl	.oz.	3
Sugarav	oz.	8
Waterff	oz.	10

Macerate the drugs and the potassium carbonate in the alcohol and water for several days, agitating occasionally, then filter, add sufficient alcohol through the filter to make the filtrate measure 16 fluidounces and in the latter dissolve the sugar.

#### Gurania.

A correspondent of the Western Druggist says a mixture of 1 part of caffeine and 2 parts of sodium bicarbonate closely resembles the above.

#### Green's Nervura.

Fisher states (A. D.) that this is similar to the following:

Cocaav.oz.	2
Damianaav.oz.	2
Gentianav.oz.	2
Potassium bromidegr.	210
Sodium salicylategr.	
Dandelionav.oz.	
Alcoholfl.oz.	
Glycerinfl.oz.	
Waterenough to make fl.oz.	
8	

Grind the vegetable drugs to powder; add the alcohol and glycerin with an equal measure of water; macerate 24 hours, then percolate, adding enough alcohol and water in the proportion given to make 32 fluidounces, in which the sodium and potassium salts are to be dissolved.

## Grimault's Matico Injection.

Made by dissolving 8 grains of copper sulphate in 16 fluidounces of infusion of matico, the latter being prepared from 1 ounce of leaves.—Wittstein.

#### Haarlem Oil.

I.	
Balsam of sulphur fl.oz.	1
Oil of turpentinefl.oz.	5
II.	
Balsam of sulphurfl.oz.	3
Barbadoes petroleumfl.oz.	
Crude oil of amberfl oz.	
Oil of turpentinefl.oz.	
Linseed oilfl.oz.	

## Haas' Hog Cholera Remedy.

According to a report of the U. S. Dept. of Agriculture, the following is probably the composition of this remedy:

Powdered soap	parts	10
Potassium carbonate		
Red ochre		
Chalk		
Quicklime		
Calcined magnesia	parts	13

#### Hagan's Magnolia Balm.

The following is said to make a similar preparation:

reparation.		
Zinc oxide	.av.oz.	1/2
Glycerin	fl.oz.	11/2
Water	fl.oz.	2
Carmine	gr.	1/4
Oil of bergamot	drop	1
Oil of lemon	drop	1
	-N. I	

av 07 4

## Haines' Golden Specific.

A formula for producing a preparation said to resemble the original was contributed to the D. C.:

Bayberry root barkparts	16
Gingerparts	8
Capsicumpart	1

Reduce to fine powder and mix thoroughly.

#### Hair's Asthma Remedy.

The following is said to resemble the original (N. I.):

Potassium iodideav.oz.	1
Tar waterfl.oz. 1	6
Caramelsufficient to color light brow	n

#### Hall's Hair Renewer.

The following gives similar results (I	N. I.)
Lead acetategr.	60
Precipitated sulphurgr.	60
Glycerinfl.oz.	8
Sodium chloridegr.	120
Bay rumfl.oz.	2
Jamaica rumfl.oz.	4
Waterfl.oz.	16

#### Hamlin's Wizard Oil.

١.		
	Spirit of camphorfl.oz.	1
	Spirit of ammoniafl.dr.	4
	Oil of sassafrasfl.dr.	
	Oil f clovefl.dr.	2
	Chloroformfl.dr.	4
	Oil of turpentinefl.dr.	4
	Alcoholenough to make fl.oz.	
	—J. J. Pierso	n.

#### Harter's Wild Cherry Bitters.

Wild cherry barkav.oz.	8
Yellow cinchona barkav.oz.	1
Orange peelav.oz.	2
Cardamom seed av. oz.	
Canada snake rootav.oz.	1/2
Diluted alcoholsufficion	ent
Honey	16
Syrup	

Percolate the drugs in moderately fine powder with diluted alcohol, and when 96 fluidounces are obtained, add the honey and syrup.

#### Hartman's Crimson Salt.

Boraxpart 1
Potassium permanganatepart 1
Salt (common)parts 6
Alumparts 8
-Frercksen.

## Hayden's Viburnum Compound.

According to the statement of composition by the manufacturers, the following contains like essential constituents:

Clamp bark	- 4.
Cassia barkav.oz.	2
Scullcapav oz.	1
Wild yamav.oz.	1
Clovesav.oz.	1/2
Alcohol,	
Water,	

Glycerin......of each, sufficient

Mix the drugs, reduce to powder, and extract with a menstruum composed of 1 volume of glycerin, 1 of water, and 2 of alcohol so as to obtain 32 fluidounces of product.

#### Headine.

Cramp bark

	parts 7
Dodium Diodi Bondi	
,	Dr Schneider

## Heiskell's Tetter Ointment.

This is said (W. D.) to be cerate of sub-acetate of lead.

## Helonia Tablets.

According to the statement of composition by the manufacturers, each tablet may be computed to contain essentially the following ingredients:

2
4

The published formula does not specify the amounts of alum, thymol and eucalyptol.

#### Hemicranine.

Phenacetine.										parts 5
Caffeine										part 1
Citric acid										.part 1

#### Henry's Carbolic Salve.

Carbolic acidgr.	5
Resin cerategr.	270
Oil of bergamotdrops	2
Oil of lavenderdrop	

#### Henry's Magnesia.

This is heavy calcined magnesia.

Henry's Tri-Iodides. (Solution of Tri-Iodides—Solution of Triple Hydroiodates.)

The published formula says that each tablespoonful contains as iodides of their alkaloids the active constituents of 30 grains each of colchicum seed, poke-root and bittersweet and 10 grains of sodium salicylate. According to this the formula below approximately represents the preparation:

Fluid extract of colchicum seed.fl.oz.	1
Fluid extract of poke-rootfl.oz.	1
Fluid extract of bittersweet fl. oz.	1
Potassium iodidegr.	64
Sodium salicylategr.	320
Simple elixir, enough to make fl.oz.	16

# Hensel's Tonicum. (Essentia Tonica Henseli.)

Formic acid (sp. gr. 1.200)fl.dr.	3 3/4
Marble dustgr.	140
Ferrous sulphate, crystalgr.	96
Solution of tersulphate of iron fl.dr.	
Glacial acetic acid fl.oz.	31/4
Alcoholfl.oz.	5
Acetic etherfl.dr.	1
Waterfl.oz.	61/2

Dissolve the marble dust in  $3\frac{1}{4}$  fluidounces of water containing the formic acid; also dissolve the ferrous sulphate in the remainder of the water and add the solution of tersulphate of iron and the acetic acid. Mix the two liquids, add the alcohol, allow the calcium sulphate to precipitate, filter, and to the filtrate add the acetic acid.—W. D.

#### Himrod's Asthma Cure.

Lobelia herbav.oz. 2
Stramonium leavesav oz. 2
Potassium nitrateav.oz. 2
Black teaav.oz. 2

Powder, mix and sift.—Contributed to W. D.

## Hinds' Honey and Almond Cream.

This formula G. H. Rose has contributed to D. C. as furnishing a similar preparation:

Cold cream, U. S. Pav.oz.	1
Sweet almond oilfl.oz.	1
Glycerinfl.oz.	1
Boric acidav.oz.	
Solution of sodafl.oz.	
Mucilage of quince seedfl.oz.	5
Water, enough to make fl. oz.	40
Oil of bitter almond,	
Oil of rose, of each, sufficient to perfu	me.

Heat the cold cream, oil and solution of soda together, stirring constantly until an emulsion is formed; then warm together the glycerin, acid, mucilage, and about 30 fluid-ounces of water, mix with the emulsion, stir until cold, and make to 40 fluidounces by adding more water. Lastly add the volatile oils.

#### Hoff's Malt Extract.

The following produces a good preparation of its class:

Althæaav.oz.	8
Corianderav.oz.	8
Star aniseav.oz.	4
Grains of paradiseav.oz.	4
Simple syrupfl.oz.	16
Glycerinfl.oz.	
Oil of lemon,drops	
Oil of orangedrop	1
Caramelav.oz.	8
Water, boilinggal.	11/2

The drugs are mixed, reduced to coarse powder, and infused with the water; to the infusion add the remainder of the ingredients, and mix well; 16 fluidounces of this liquid is added to one barrel of ordinary brewers' beer.

## Holloway's Ointment. An ointment of this

Melt the butter, wax and resin, add the vinegar, allow the whole to simmer for 10 or 12 minutes, allow to cool somewhat, add the remaining ingredients, and stir until cool.

#### Holloway's Pills.

Aloes	.gr. 36
Jalap	
Ginger	
Myrrh	
Mix and make 80 pills.—N. I.	

#### Horner's Rheumatic Lightning.

	-	_	
Fluid extract of colchicum	seed f	l.dr.	1
Fluid extract of black col			
Potassium acetate		.gr.	128
Sodium salicylate		.gr.	256
Alcohol			
Water, enough to make.	. <b></b> . f	l.oz.	16
		-N	. I.

## Hooper's Female Pills.

The following is apparently similar:	
Aloes, socotrinegr. 48	3
Dried sulphate of irongr. 18	
Extract of black helleboregr. 15	3
Myrrhgr. 19	9
Soapgr. 15	3
Powdered canellagr. 6	3
Powdered gingergr. 6	3

Beat them well together into a mass, with syrup or water, and divide into pills, each containing  $2\frac{1}{2}$  grains.

## Hop Bitters.

The following makes a good hop bitters:

Hopsav.oz.	4
Orange peelav.oz.	2
Dandelion av.oz.	2
Buchuav.oz.	1
Mandrakeav.oz.	1/2
Sugarav.oz.	
Alcoholfl.oz.	16
Water, to-makepints	8
•	

Macerate the drugs in coarse powder in the alcohol and 7 pints of water, at a warm temperature, for 8 days. Express and dissolve the sugar in the liquid, to which add water, if necessary, to make 8 pints, and strain.—W. D.

## Horsford's Acid Phosphate.

Solution of Acid Phosphates, Part I, is an excellent preparation containing acid phosphates.

#### Hostetter's Bitters.

The following is said to produce a similar article (D. C.):

Gentiangr.	15
Blessed thistlegr.	15
Calamusgr.	15
Orange peelgr.	60
Oil of orangedrop	1
Sugargr.	320
Alcohol,	

Water, of each enough to make fl.oz. 16

Mix the drugs, extract with a menstruum consisting of 3 volumes of alcohol and 2 of water, and in the liquid obtained dissolve the sugar and oil of orange.

#### Hudson's Honey of Elm.

This is a tooth paste containing precipitated chalk, powdered pumice stone, glycerin, oil of clove, oil of wintergreen, and simple syrup.—Era Form.

#### Hubert's Malvina Cream.

The following will make a preparation resembling the original (N. I.):

White petrolatumav.oz.	6
White waxgr.	
Spermacetigr.	300
Bismuth oxychloridegr.	400
Mercuric chloridegr.	5
Oil of rosedrops	6
Oil of bitter almonddrop	1

Warm together the petrolatum, white wax and spermaceti until melted. While cooling incorporate the bismuth oxychloride and the mercuric chloride, this last previously dissolved in a little alcohol, and when nearly cold stir in the volatile oils.

#### Hubert's Malvina Lotion.

This is said (A. D.) to be an emulsion of almond with rose water and containing oxide of zinc and corrosive sublimate. A formula for such a preparation may be constructed as follows:

Mercuric	chloride		 gr. %	3
Zinc oxid	e		 dr. 8	3
Emulsion	of almond	s	 pin	t
	e emulsion			

Make the emulsion from 120 grains of sweet almonds with rose water.

## Husband's Magnesia.

This is heavy calcined magnesia.

#### Hydroleine.

This contains approximately the same ingredients in the same essential proportions as are claimed for the original, and may be presumed to yield a similar result:

Cod liver oil	fl.oz.	8
Solution of soda	fl.dr.	51/2
Boric acid	gr.	12
Hydrochloric acid	gr.	21/2
Saccharated pancreatin		
Water, enough to make	fl.oz.	12

## Idiaton.

Its approximate composition is given (according to W. D. ) as follows:

Venice turpentinepart 1
Masticpart 1
Chloroformparts 3
Spirit of ammoniaparts 2
Oil of cloveparts 2
Oil of mustard (volatile) a small quantity

This is a toothache remedy sold largely in Germany.

## Injection Brou.

Zinc sulphate	gr. 15
Lead acetate	gr. 30
Tincture of catechu	fl.dr. 1
Crocated tincture of opium	
Water, distilled	
,,	-Hager.

#### Iodia.

This is claimed to be a combination of active principles of stillingia, saxifraga, menispermum, and aromatics, each fluidram containing 5 grains potassium iodide and 3 grains of iron phosphate. Compound Elixir of Potassium Iodide, Part I, appears to be similar in character.

## Iodophenochloral.

	e of iodine	
Carbolic	acid	part 1
Chloral	hydrate	part 1

## Jayne's Tonic Vermifuge.

This contains sodium santoninate, pinkroot, jalap, peppermint, erigeron, wintergreen, sugar and water.—N. I.

## Jayne's Expectorant.

The following will produce a similar preparation (Kilner's Form.):

Syrup of squill	fl. oz. 2
Tincture of tolu	
Tincture of lobelia	fl.dr. 1
Tincture of digitalis	fl.dr. 1
Tincture of opium	fl.dr. 2
Spirit of camphor	
Wine of ipecac	fl.dr. 2
Tartar emetic	gr. 2

Dissolve the tartar emetic in the tincture of lobelia or digitalis, and mix with the remaining ingredients.

#### Jesson's Dental Anæsthetic.

This contains about 2.6 per cent of cocaine hydrochlorate with some carbolic acid and oil of rose.—Sadtler.

## Jewsbury & Brown's Oriental Tooth Paste.

This, according to Nelson's "Handbook," makes a similar preparation:

Carminegr.	60
Water	
Honeyav.oz.	6
Oil of peppermintdrops	
Oil of anisedrops	5
Oil of orangedrops	10
Oil of wintergreendrops	10
Precipitated chalksufficie	nt

Rub together, using enough of the chalk to make a firm paste.

## Kalydor.

Bitter almonds, blanched av.oz	.10
Rose waterfl.oz.	50
Corrosive sublimategr.	5
Ammonium chlorideav.oz.	3/
Cherry laurel waterfl.oz.	11/2
Alcoholfl.oz.	11/2

Mix an emulsion of the almonds and the rose water, strain, add the ammonium chloride and cherry laurel water, and then the corrosive sublimate dissolved in the alcohol.—Nat. Dr.

## Kaputine.

This is chiefly a colored acetanilid.—British Medical Journal.

## Kendall's Spavin Cure.

The following makes a preparation similar to the original (N. I.):

Oil of rosemary	fl.dr.	1
Oil of turpentine	fl.oz.	1
Camphor	gr.	240
Petroleum oil (heavy)	fl.dr.	1/2
Alcohol	fl.oz.	2
Iodine	gr.	125

Dissolve the iodine in a little alcohol and add it to the other ingredients, previously dissolved in the alcohol and water.

#### Kennedy's Medical Discovery.

This makes a similar preparation (N. I.):

Fluid extract of mandrake	fl.dr.	10
Fluid extract of dandelion	fl. oz.	1
F luid extract of leptandra	fl.dr.	3
Alcohol	fl.oz.	5
Water, enough to make	fl. oz.	16

#### Kennedy's Pinus Canadensis (Dark).

The Fluid Extract of Pinus Canadensis, Part I, is presumably similar

## Kennedy's Pinus Canadensis (White).

Zinc sulphatepart 1
Glucoseparts 2
Waterparts 7
Hemlock oila few drops
—Tschenne

#### Kern's Insect Annihilator.

The following is said to resemble this mixture (W. D.):

Deodorized benzin	.fl.oz.	16
Oil of cedar	fl.oz.	1
Oil of wintergreen	fl.oz.	1

This composition is certain destruction to cockroaches, bedbugs, fleas, ants, and other insects.

#### Kidneywort.

The dry preparation is a mixture of drugs, dandelion, hydrangea, etc., with roasted beans. The liquid preparation contains the extract of similar drugs.—Fenner's Form.

#### King's New Discovery.

According to the N. I., a similar preparation is made as follows:

Morphine sulphategr.	8
Sugar, granulatedav.oz.	14
Chloroformdrops	60
Tincture of white pinefl.oz	
Fluid extract of ipecacfl dr.	
Magnesium carbonatedr.	
Water	

Rub one ounce of sugar with ¼ ounce of magnesium carbonate, triturate with the tincture and fluid extract, gradually add the water with continued trituration. Filter, dissolve the morphine and sugar in the filtrate, strain and add the chloroform.

#### Kitchell's Liniment.

Water		parts 3
	water	

Color with caramel.—J. Goldbach.

## Kline's Great Nerve Restorer.

The following is said to make a similar preparation (Dr. R. C. McCann):

Ammonium bromidegr.	180
Potassium bromideav.oz.	3
Potassium bicarbonategr.	70
Tincture of columbofl.dr.	6
Waterfl.oz.	6

## Kenig's Hamburg Breast Tea.

This is a mixture of cut licorice root, althea root, althea flowers, coltsfoot herb, red-poppy petals mallow flowers, calendula flowers, and blind nettle flowers.—Ph. Rundsch.

## Kenig's Hamburg Drops.

This is similar to Elixir of Long Life, Part I.—D. C.

## Kænig's Hamburg Plaster.

Mother plasterav.oz.	8
Suetav.oz.	1/2
Black pitchav.oz.	1/2
Amberav.oz.	1
Peru balsamgr.	45

#### Knox's Disinfecting Yowder.

Chlorina	ated lime.		٠.				.av.oz.	4
Sodium	chloride.						.av.oz.	12

## Lactopeptine.

According to statement of composition by manufacturers, the following are the ingredients of a similar preparation:

Pepsin gr.	48
Pancreatingr.	
Diastasegr.	3
Hydrochloric aciddrops	4
Lactic aciddrops	
Milk sugargr.	240

## Lactopeptine Elixir.

See Elixir of Lactopeptine.

## Lactopeptine Elixir with Bismuth.

The following is an elixir containing lactopeptine and bismuth:

Ammonio-citrate of bismuthgr.	128
Waterfl.oz.	2
Elixir of lactopeptine (prepared	
without acid)fl.oz.	14

Dissolve the bismuth salt in the water by the aid of a small amount of ammonia water, carefully avoiding any excess of the latter, and add this solution to the elixir.

## Lactopeptine Elixir with Calisaya.

The following is an elixir containing lactopeptine and calisaya:

Elixir	cf	lact	op	eptine	 	.fl.oz.	14
				cinchona			

## Lactopeptine Elixir with Calisaya, Iron and Bismuth.

The following is an elixir containing lactopeptine, calisaya, iron and bismuth:

Fluid extract of cinchonafl.dr.	10
Iron pyrophosphate, solublegr.	192
Ammonio-citrate of bismuthgr.	
Waterfl.oz.	
Elixir of lactopeptine (prepared	,-
without acid)fl.oz.	13

Dissolve the bismuth in a small amount of water by the aid of a little ammonia, dissolve the iron salt in the remainder of the water, mix all, and filter.

## Lactopeptine Elixir with Gentian and Chloride of Iron.

The following is an elixir containing lactopeptine, gentian and iron chloride:

Fluid extract of gentianfl.dr.	$^{-2}$
Tincture of citro-chloride of iron fl.oz.	
Elixir of lactopeptine, enough	
to makefl.oz.	16

## Lactopeptine Elixir with Phosphate Laubach's Eclectic Liniment. of Iron, Quinine and Strychnine.

The following is an elixir containing lactopeptine, iron phosphate, quinine and strych-

Quinine hydrochlorategr. 64	
Strychnine sulphategr. 1	
Iron phosphate, solublegr. 64	
Water, distilled	
Elixir of lactopeptine (prepared	
without acid)fl.oz. 14	

## Lactopeptine Elixir with Strychnine and Bismuth.

The following is an elixir containing lactopeptine, strychnine and bismuth:

Ammonio-citrate of bismuthgr.	128
Strychnine sulphategr.	2
Waterfl.oz.	2
Water of ammoniasuffic	ient
Elixir of lactopeptine (prepared	
without acid)fl.oz.	14

Dissolve the bismuth salt in a small amount of water, adding some water of ammonia to insure complete solution, but avoiding any excess of the latter. Dissolve the strychnine salt in the remainder of the water, add this to the elixir, then add the bismuth solution. let the whole stand a few hours, and filter.

## Lactopeptine, Liquid.

Lactopeptinegr. (	600
Distilled waterfl.oz.	10
Glycerinfl. oz.	
D	C

## Lallemand's Gout Specific.

Extract of colchicumgr.	60
Extract of opiumgr.	60
Potassium iodideav.oz.	2
Potassium acetateav.oz.	1
Waterfl.oz.	14
White winefl.oz.	2
—Era Fo	rm.

## Laird's Bloom of Youth.

Zinc oxidegr.	180
Precipitated chalkgr.	180
Oil of bergamotdrops	
Water fl oz.	
	-

#### Lavoline.

This is said to be purified liquid petrolatum .- Reg. Ph.

#### Laxol.

This is said to consist of castor oil sweetened with saccharin and flavored with peppermint.-Ph. Era.

Oil of turpentinefl	.oz,	1
Tincture of arnica flowersfl		
Stronger water of ammoniafl		
Soap linimentfl		
Oil of sassafras		
Oil of thymedr	ops :	16
Alcoholfl	.oz.	4

#### Liquor Sedans.

According to statements of composition by the manufacturers, Elixir of Black Haw Compound, Part I, produces a preparation having the essential constituents of a similar product.

## Liquor Uterans.

According to the manufacturers, each fluidounce represents 55 grains of blackhaw, 28 grains of blue cohosh, 25 grains of Jamaica dogwood, and 55 grains of golden seal, with aromatics. An essentially similar preparation will be obtained by this formula:

Fluid extract of blackhawfl.oz.	13
Glycerite of hydrastisfl.oz.	1 3
Fluid extract of blue cohoshfl.oz.	1
Fluid extract of Jamaica dog-	
wood	1
Simple elixir enough to make fl.oz.	16

#### Listerine.

This is claimed to be the "essential antiseptic constituent of thyme, eucalyptus, baptisia, gaultheria, and mentha arvensis in combination. Each dram also contains 2 grains of refined and purified benzo-boracic acid." Formulas for several good antiseptic solutions are given under "Lister's Antiseptic Solution," Part I.

#### Listol.

This is dithymol diiodide, also known as annidalin and aristol.

#### Lithiated Hydrangea.

Each fluidram is claimed to represent 30 grains of fresh hydrangea and 3 grains of c. p. benzo-salicylate of lithia. The formula below is for a preparation of similar character:

Lithium salicylate	.gr.	240
Lithium benzoate		
Fluid extract of hydrangeaf		
Alcoholf		
Water, enough to make		

## Lloyd's Leontin.

This is a 1-per-cent solution of leontin in an alcoholic menstruum, according to the manufacturers' statement.

#### Lotsil.

Acetanilidgr.	36
Sodium bicarbonategr.	12
Caffeine citrategr.	
Camphor monobromidegr.	6

Dispense either as powder or in 5-grain tablets.

## Low's Magnetic Liniment.

The following makes a similar preparation (N. I.):

Oil of turpentinefl.oz.	1 1/2
Tincture of capsicumfl.oz.	2
Spirit of camphorfl.oz.	16
Stronger water of ammoniafl.oz.	1 1/2
Alcoholfl.oz.	31/2
Oil of sassafras	
Fluid extract of sassafrasfl.dr.	5

#### Luperine.

This is a mixture of powdered columbo, gentian and quassia. - Coblentz.

## Lyon's Kathairon.

This is said (N. I.) to consist of substantially the following:

Alcoholfl.oz.	12
Castor oilfl.oz.	4
Tincture of cantharidesfl.dr.	4
Tannic acidgr.	30
Oil of citronellafl.dr.	1/2
Oil of bergamotfl.dr.	1/2
Oil of clovesfl.dr.	1/2
Oil of rosemarydrops	8
Oil of lavender flowersdrops	8

Mix and filter.

## Lyon's Tooth Powder.

This contains soap, precipitated chalk, pumice stone and oil of wintergreen.-N. I.

## McDade's Succus Alterans. (McDade's Prescription.)

This is claimed to be prepared from the fresh green drugs. The "Extract of Bamboo Brier" Part I, is of similar composition.

## MacDougall's Disinfecting Powder.

This is prepared by adding crude carbolic acid to sodium sulphite.-Frerksen.

#### Maltine.

This is said to be prepared from malted barley, oats and wheat, equal parts, as described in Part I, for making Extract of Malt.

## Maltine with Cascara Sagrada.

Cod liver	oil.							.fl.oz.	3
Maltine .								.fl.oz.	7

## Maltine, Ferrated.

Iron pyrophosphate, solublegr. 128	
Water, hotfl.oz. 1	
Maltine	
Dissolve the iron salt in the water and a	dd

the maltine.

## Maltine with Hypophosphites.

Calcium hypophosphitegr. 4	8
Sodium hypophosphitegr. 4	8
Iron hypophosphitegr. 3	2
Potassium citrategr. 6	0
Waterfl.oz.	
Maltinefl.oz. 1	5

Dissolve the potassium citrate and iron hypophosphite in one-half fluidounce of water by the aid of heat, also the calcium and sodium hypophosphite in the same amount of water, and mix the two solutions with the maltine.

## Maltine with Pensin and Pancreatin.

-	
Pepsin, saccharatedgr.	640
Pancreatin, saccharatedgr.	
Maltine fl.oz.	16
Mix all by trituration	

## Maltine with Iron Phosphate, Quinine and Strychnine.

Iron pyrophosphategr.	64
Quinine hydrochlorategr.	16
Strychnine sulphategr.	12
Distilled water, hotfl.oz.	1
Maltinefl.oz.	15

Dissolve the iron and alkaloidal salts in the water and add to the maltine.

## Mariani Wine of Coca.

The following makes a satisfactory wine of coca:

Coca leaves, coarse powderav.oz.	1 1/2
Alcohol fl.oz.	2
Sugargr.	330
Red winefl.oz.	16

Mix the alcohol and wine, macerate the leaves in 4 fluidounces of this mixture in a moderately warm place for 24 hours, then transfer to a glass percolator, pass remainder of liquid slowly through the drug, and in the percolate dissolve the sugar.

#### Marienbad Reduction Pills.

Potassium bromide10
Sodum bicarbonate
Extract of squill
Guaiac
Senega
Extract of taraxacumsufficient

## Divide into pills weighing 21/2 gr. each.—D.

#### Marrol.

This is said to consist of ox marrow, malt extract, and hop extract.—Coblentz.

## McLean's Strengthening Cordial.

A similar preparation is said to be made as follows (Fenner's Form.):

Gentianav.oz. 8
Columboav.oz. 8
Orange peelav.oz. 2
Corianderav.oz. 1
Serpentariaav.oz. 1
Cardamomav.oz. ½
Whiskeypints 7
Glycerinfl.oz. 1
Water sufficient

Mix the drugs, grind to powder and percolate with the whiskey mixed with the glycerin, and pass through the percolator enough water to make the percolate measure 1 gallon.

## McMunn's Elixir of Opium.

The official deodorized tincture of opium is similar in character.

#### Medicamentum.

See Haarlem Oil.

## Melachol.

Each fluidram, according to the manufacturers, contains 85 grains of the combined sodium phosphate, sodium nitrate and citric acid.

According to W. C. Wescott, the following yields a similar preparation:

Sodium nitrategr.	73
Citric acidgr.	475
Sodium phosphateav.oz. 8. gr.	150
Water, enough to makefl.oz.	8

The mixture of salts and acid will liquefy and solution will be completed upon the addition of water.

## Merrell's Vaginal Discs.

Powdered extract of heloniasgr.	12
Powdered extract of hydrastisgr.	12
Powdered extract of henbanegr.	6
Powdered opiumgr.	6
Thymolgr.	
Gallic acidgr.	
Boric acidgr.	36
Tannic acidgr.	24
Alumgr.	36
Eucalyptolgr.	11/2

Make into 12 compressed tablets.

## Mexican Mustang Liniment.

The following makes a similar preparation (N. I.):

Soapgr.	140
Water, warmfl.oz.	
Oil of turpentinefl.dr.	2
Crude petroleumfl dr.	
Oil of amber. crudefl.dr.	
Oil of thymefl.dr.	2
Kerosenefl.oz.	
Caustic potashgr.	12

Dissolve the soap in the water and incorporate with the other ingredients.

## Micajah Medicated Uterine Wafers.

This formula for a similar preparation has been contributed to a medical journal (W. D):

Mercury bichloridegr. 1
Zinc sulphategr. 5
Bismuth subnitrategr. 15
Acaciagr. 5
Carbolic acidgr. 3
Watersufficient
Water

## Migranin.

Citric acidpart	1
Caffeineparts	9
Antipyrineparts	90

## Mollosin.

Yellow Liquid	wax petrolatum	  	part 1
•			-Ph. Ztg.

## Morrison's Pills.

Aloes	. parts 5
Jalap resin	. parts 5
Jalap root	
Marshmallow root	
Gamboge	. parts 2
Scammony	part 1

Divide into pills, each to contain ½ gr. aloes. Roll in cream of tartar.—Hager.

## Murray's Infallible System Tonic.

Aloes		 	 		<i>.</i>	 .gr.	50
Cinnamon		 	 	٠.	٠	 .gr.	25
Licorice ro	ot.	 	 			 or.	25

Make into a mass with water, divide into 50 parts, and put into gelatin capsules.—
N. I.

## Murdock's Liquid Food.

This is said to consist of defibrinated blood with 15 per cent of whiskey and various tonics and astringents.

## Naphthocresol.

This is a mixture of phenols and cresols dissolved in a solution of resin soap.

# Naphthol Camphor. (Camphorated Naphthol.)

A syrupy liquid said to be prepared by fusing together 1 part of betanaphthol and 2 parts of camphor.

## Nelason's Suppositories.

Calomelgr	12
Extract of belladonnagr.	3
Fluid extract of stramoniumm.	
Cacao buttergr.	
Make into 12 suppositories.	

#### Neurosin.

A French preparation (in syrup or granule form) which contains calcium glycero-phosphate as the active ingredient.—Coblentz.

## Neurosine.

It is claimed that each fluidram contains 5 grains each of c. p. bromides of potassium, sodium and ammonium,  $\frac{1}{8}$  grain bromide zinc,  $\frac{1}{64}$  grain each of extract belladonna and cannabis indica, 4 grains extract lupuli, and 5 minims fluid extract cascara, with aromatic elixirs. The formula below, based on the foregoing is for a similar preparation:

. 0 0	1	
Potassium bromide	.gr.	640
Sodium bromide	.gr.	640
Ammonium bromide	.gr	640
Zinc bromide		
Extract of belladonna		
Extract of cannabis indica	.gr.	2
Extract of hops		512
Fluid extract of cascara sagrada f	dr.	101/2
Simple elixir, enough to make f		

## Nichol's Compound Tasteless Cod Liver Oil.

See Wampole's Tasteless Cod Liver Oil.

#### Nichol's Elixir of Bark and Iron.

This preparation contains, according to the manufacturers, calisaya and protoxide of iron. A formula for a similar preparation is Elixir of Cinchona with Iron Protoxide, Part I.

#### Norton's Chamomile Pills.

Extract of aloes, aqueousgr.	60
Extract of gentiangr.	
Oil of chamomiledrops	20

Make 60 pills.

## Norwood's Tincture.

This is the same as the official tincture of veratrum viride.

#### Oculine.

A solution in ordinary water containing 1 per cent of boric acid and 5 per cent of glycerin.—Ph. Rundsch.

#### Odol.

Saccharingr. 1
Salolgr. 80
Tincture of vanilladrops 24
Spirit of peppermintdrops 36
Spirit of cumin drop 1
Alcohol, enough to makefl.oz. 4

#### Odontunder.

This contains 1.35 per cent of cocaine hydrochlorate with some carbolic acid, glycerin, oil of rose, and probably alcohol.—Sadtler.

#### Osgood's Indian Cholagogue.

Quinine sulphategr.	
Fluid extract of culver's rootfl.dr.	2
Saturated tincture of stillingiafl. oz.	4
Fluid extract of mandrakefl.dr.	3
Oil of sassafrasdrops	10
Oil of wintergreendrops	10
New Orleans molasses, enough	
to makefl.oz.	8
—Kilner's Fo	orm.

## Oxygen Aquæ.

A colorless, odorless and tasteless liquid found to be water.—A. B. Prescott.

#### Oxygen, Compound.

A colorless aqueous solution of ammonium nitrate and lead nitrate, the two salts being in nearly equal proportions, and together forming about three per cent of the solution.—A. B. Prescott.

## Oxygen, Compound, Green's.

An aqueous solution of ammonium nitrate with a very little lead nitrate.—A. B. Prescott.

## Oxygen, Compound, Solid.

This is ammonium nitrate.—A. B. Prescott.

## Oxygen, Compound, O'Leary's.

Contains alcohol, chloroform, bitter almond oil, balsam of tolu and red coloring matter.—
A. B. Prescott.

#### Ozonized Water.

This is said to contain 1 or 2 parts of potassium permanganate dissolved in 500 parts of water.

## Palmer's Invisible Powder.

This contains talcum with coloring .- Snow.

## Palmer's Lily White Tablet.

Precipitated chalkav.oz. 17	
Talcumav.oz. 23	
-Snow.	

#### Palmer's Lotion.

A similar preparation is said to be:	
Corrosive sublimategr.	8
Alumgr.	12
Waterfl.oz.	

## Paine's Celery Compound.

A preparation, said to be similar, may be made by the following formula (D. C.):

Celery seed av.oz.	2
Cerety seed	~
Red cinchonaav.oz.	1
Orange peelav.oz.	1/4
Coriander seedav.oz.	1/4
Lemon peelav.oz.	1/4
Hydrochloric acid	$15^{\circ}$
Alcoholfl.oz.	
Glycerinfl.oz.	3
Waterfl.oz.	
Simple syrupfl.oz.	
0111 111 1	

Grind the solids to moderately coarse powder, mix the acid and the water, add the glycerin and alcohol and in the menstruum so prepared macerate the powder for twenty-four hours; then percolate, adding enough alcohol and water in the proportion given to make 12 fluidounces. Finally add the syrup, and if necessary filter.

#### Pancropepsin.

The Compound Powder of Pepsin, Part I, has the same essential constituents as are claimed for this.

## Pan-Peptic Tablets.

The composition as stated by the manufacturers is 1 grain each of pure pepsin and pure pancreatin, ¼ grain pure caffeine, and acid lactophosphate of calcium and celery. The formula below will furnish a composition essentially similar:

Pepsin	or.	12
Pancreatin		
Caffeine		
Apiol	gı.	0.0
Calcium lactophosphate, soluble.	gr.	30
Make into 12 tablets.		

#### Papier Fayard.

70 4 4 4 4	100
Powdered cantharidesgr.	480
Powdered euphorbiumgr.	
Alcoholfl.oz.	
Venice turpentineav.oz.	3
White resinav.oz.	

Extract the powdered drugs with the alcohol, melt the resin and turpentine, add the extract, and, with a brush, spread the mixture on paper while still warm.—Pharm.

## Papine.

This is said to be the anodyne principle of opium, the narcotic and convulsive elements being eliminated, one fluidram representing the anodyne principle of ½ grain of morphine. The following will yield a preparation conforming to these requirements.

Deodorized tincture of opium...fl.oz. 3½ Simple elixir.....fl.oz. 13

## Parker's Tonic.

The following is similar (N. I.):	
Fluid extract of gingerfl.dr.	2
Fluid extract of sassafrasdrops	15
Fluid extract of capsicumdrop	1
Oil of wintergreendrop	1
Sugarav.oz.	11/4
Waterfl.dr.	
Alcoholfl.dr.	13

#### Parsons' Local Anæsthetic.

6
6
$^2$
1
1
1

Dissolve the camphor in the chloroform, then add oil of clove and then the tinctures.

—Dr. Parsons.

## Pasteurine.

This contains oils of cinnamon, eucalyptus, lemon and wintergreen dissolved in alcohol.

# Peacock's Bromides. (Syrup of Bromides, Compound: Peacock.)

Claimed by the manufacturers to contain in each fluidram 15 grains of the combined c. p. bromides of potassium, sodium, calcium, ammonium and lithium. A compound of a similar character is the following:

Potassium bromidegr.	384
Sodium bromidegr.	384
Ammonium bromidegr.	384
Citric acidgr.	16
Tincture of vanillafl.dr.	4
Compound tincture of cudbear fl.oz.	1 1/2
Simple syrupfl.oz.	3
Water, enough to make,fl.oz.	16

Dissolve, let stand for twenty-four hours and filter.

#### Peckham's Balsam.

White	resinav.oz. 4	Ŀ
Oil of	turpentinefl.oz. 4	Ł

Melt the resin, remove from the fire, add the oil and mix well together.—Fenner's Form.

## Pepsin and Wafer Ash.

Pepsin, puregr.	128
Fluid extract of wafer ash,fl.oz.	
Glycerinfl.oz.	
Water, enough to make fl.oz.	

## Perl's Antikrinin.

Strontium sulphideav.oz.	3
Zinc oxideav.oz.	1
Starchav.oz.	1
Mentholgr.	20
This is employed as a depilatory.—W	v. D

## Peterman's Roach Food.

According to Dr. Eccles, the following is similar:

Boraxav.c	oz. 37
Starchav.c	
Cacao	

#### Petit's Eye Salve.

A similar preparation is the following:

Morphine sulphategr.	1 1
Benzoic acidgr.	4
Ammonia ed mercurygr.	48
Zinc oxidegr.	64
White waxgr.	64
Spermacetigr.	
Olive oilgr.	500
Oil of rosemarydrop	1

## Phenol Sodique.

The following is said to yield a similar preparation (G.M. Beringer):

Coal tarav.oz.	21/4
Soda, causticgr.	120
Water, enough to makefl.oz.	16

Dissolve the soda in 4 fluidounces of warm water, add the coal tar and thoroughly agitate for a few minutes. Then add the remainder of the water and set aside in a covered vessel in a warm place, frequently agitating, for 7 days. Decant and filter.

## Phenosalyl.

Carbolic acidparts 90	
Lactic acidparts 20	
Salicylic acidparts 10	
Mentholpart 1	

Mix by fusing acid and adding other ingredients.

## Phillip's Phospho-Muriate of Quinine, Compound.

According to the manufacturers' statement, each fluidram contains 134 grains of potassa, 1 grain of magnesia, 34 grain of lime, ½ grain of iron, all in the form of phosphates. Further: ¼ grain of quinine hydrochlorate, ½ grain of strychnine, 2 grains of free phosphoric acid. The following formula yields a preparation essentially conforming to these requirements:

Compound solution of phos-	0
phoric acidfl.oz.	3
Sugar av.oz. 1	
Quinine hydrochlorategr.	8
Strychnine sulphategr.	1
Spirit of bitter almonddrops	5
Caramel sufficient to colo	r

Dissolve the sugar in the acid solution, add the alkaloid salts previously dissolved in a small amount of distilled water, then the spirit and caramel, and filter.

#### Phospho-Albumen.

Testicles,	sheep,	sliced	 part 1
Water			 parts 3

Macerate for 12 hours, strain, saturate liquid with sugar and add simple syrup to wake 10 parts. Flavor each 12 fluidounces with 4 fluidrams of spirit of orange. To preserve, a small amount of antiseptic is added.—Stuart.

## Phytoline.

Is the inspissated juice of poke berries after having been touched by frost. Claimed to be prepared by a special process suggested by Dr. W. W. Baxter.

## Pierce's Compound Extract of Smartweed.

The following makes a preparation of smartweed suitable for external application:

Smartweedav.oz.	í
Alcoholfl.oz. 24	Ŀ
Waterfl.oz. 8	3
Camphorgr. 90	)
Oil of hemlock fl.dr.	3
Oil of sassafrasfl.dr.	3

Extract the smartweed with the alcohol and water and to the liquid obtained add the camphor and oils.

# Pierce's Favorite Prescription. Pierce's Golden Medical Discovery.

The formulas quoted from Hager in the previous editions of this work—alcohol and opium appearing in the formulas for both remedies—have been met by an affidavit by Dr. Pierce, received by the editors, declaring that "there is not at the present time, nor has there ever been, any alcohol, opium, chloral, cocaine, or other narcotic, used as ingredients in Dr. Pierce's Golden Medical Discovery or Dr. Pierce's Favorite Prescription" It is evident from this declaration either that Dr. Hager erred in his analyses, or that his results were obtained from imperfect samples of the remedies.

The "Favorite Prescription" was said by Hager to consist of savin, acacia, cinchona, agaric, sugar, cinnamon, tinc digitalis, tinc. opium, oil anise, alcohol, water. The "Medical Discovery" was said to consist of lactucarium, honey, tinc. op um, alcohol and water. Pinapin.

This is essentially a fermented pineapple juice.—Coblentz.

#### Pinaud's Brilliantine.

The following is said to resemble the original (W. D.):

Castor oilfl.oz.	1
Sweet almond oilfl.oz.	
Glycerinfl.dr.	3
Jockey club extractfl.dr.	
Alcohol, enough to makefl.oz.	16

## Pinaud's Eau de Quinine Tonique.

In a suit in the U. S. Circuit court at Boston concerning its tariff classification, the court was satisfied "that this article contains of absolute alcohol substantially 67 per cent by volume, that the solid residuum, amounting to about 18-100 of 1 per cent, consists principally of an odoriferous resin having a fragrance similar to that of benzoin; a minute trace of quinine sulphate, and also a very small percentage of essential oils, the remainder being water." This substantiates essentially an analysis published some years ago by A. Tscheppe. The following is for a somewhat similar preparation:

1 1	
Alcohol	fi.oz. 10
Water	fl.oz. 5½
Yellow cinchona	dr. 4
Cochineal	gr. 30
Fotassium carbonate	gr. 30
Tincture of benzoin	fl.dr. 2
Oil bergamot	drops 30
Oil sweet orange	drops 30
Oil rose geranium	drops 10

Make a decoction of the cinchona and cochineal, strain, then add the potash and alcohol in which have been dissolved the resin and oils. Filter through pumice.

# Pinkham's (Lydia) Vegetable Compound.

This formula was contributed to D. C., and was said to furnish a product resembling the original:

4
4
2
2
2
11/2
$24^{'}$
16
ent

Reduce the first six ingredients to powder, add boiling water enough to cover, let stand till cold, and then percolate with water until 5 pints of liquid are obtained. To this add the sugar, bring to a boil, remove from the fire, strain, and, when cold, add the alcohol.

## Pleis' Fit Powders.

	bromidegr	
Gentian, p	owdergr.	5

Make one powder.—Drug Mill.

## Piso's Consumption Cure.

The following is said to be similar:	
Tincture of tolufl.dr.	4
Fluid extract of lobeliafl.dr.	2
Fluid extract of cannabis indicafl.dr.	2
Sulphate of morphinegr.	4
Tartar emeticgr.	4
Chloroformfl.dr.	1
Essence of spearmintdrops	10
Water, hotfl.oz.	
Sugarav.oz.	14

The fluid extracts, tincture of tolu, chloroform, and essence of spearmint are mixed with sugar, in a bottle. Dissolve the morphine and tartar emetic in the hot water and mix; when thoroughly dissolved, filter.—N. I.

## Plant's Asthma Cigarettes.

Stramonium leaves				 .av.oz	2
Green tea leaves				 .av.oz.	2
Lobelia leaves			 	.av.oz.	13/

Mix, moisten with a saturated solution of potassium nitrate, and dry.—Ph. Post.

#### Platt's Chlorides.

A similar preparation, according to Dr. Tscheppe, is the following:

Aluminum sulphateav.oz.	6
Zinc chlorideav.oz.	
Sodium chlorideav.oz.	2
Calcium chlorideav.oz.	3
Water, enough to makefl.oz.	32

Dissolve the calcium and aluminum salts separately, mix, allow the calcium sulphate to subside, and in the clear liquid dissolve the other ingredients.

## Ponca Compound.

Each tablet is said to contain 3 gr. extract of ponca (?), 1 gr. extract of mitchella, ¼ gr. of caulophyllin, ⅓ gr. of helonin, and ⅓ gr. of viburnin.

#### Potsdam Balsam.

Oleobalsamic mixturefl.oz.	17
Compound spirit of angelicafl oz.	2
Tincture of capsicumfl.dr.	3
Spirit of ammoniafl.dr.	4
-	70

#### Powell's Balm of Anise Seed.

This has a composition similar to paregoric. It has but little camphor, a small amount of rhubarb, and some extract of icorice.—N. I.

## Pozzoni's Complexion Powder.

Γalcum parts 3	5
Calcium carbonateparts 2	0
Bismuth oxychlorideparts	8
—Snow	7.

#### P. P. P.

This is said to contain the fluid extracts of green poke-root, green prickly-ash bark, stillingia, and sarsaparilla, with compound tincture of gentian, potassium iodide, and simple syrup.

## Pyretine.

Acetanilidparts 9
Caffeinepart 1
Calcium carbonateparts 2
Sodium bicarbonateparts 3
Walter.

#### Pyrozone.

This is a solution of hydrogen peroxide.

#### Quickine.

Carbolic acidpart	1
Mercuric chloridepart	50
Alcohol and waterparts	1000
Ph.	Ztg.

#### Quinquinia.

This is a similar preparation, according to a communication by Dr. Lee to the Phila. Med. and Surg. Reporter:

Quinine alkaloid	.15	per cent
Quinidine alkaloid	.15	per cent
Cinchonidine alkaloid	.15	per cent
Cinchonine alkaloid	.25	per cent
Chinoidine	.30	per cent

#### Quina-Laroche.

This is said (Bulletin Societe Royale de Bruxelles) to be prepared as follows:

Red cinchona, coarse powderav.oz.	1
Water, boilingfl oz.	1
Malaga winefl oz. 10	)
Diluted alcohol	
Watersufficien	t
Sugarav.oz.	8

Macerate the drug with the boiling water for 30 minutes, decant the liquid, add the wine, macerate for 8 or 10 hours, again decant the liquid, macerate the dregs with the diluted alcohol, macerate for a few hours, again decant, mix the three liquids, and wash the marc with enough water to make the

entire liquid measure 16 fluidounces. Set this aside for 24 hours, filter, and in the filtrate dissolve the sugar.

The ferruginous preparation is made by adding to the above 8 grains of soluble iron pyrophosphate.

## R. & H. Three Chlorides. (Elixir Ferri, Hydrarg et Arsenicum.)

Each fluidram, according to advertisements, contains  $\frac{1}{18}$  grain of protochloride of iron,  $\frac{1}{128}$  grain of bichloride of mercury,  $\frac{1}{128}$  grain of chloride of arsenic, with calisaya alkaloids and aromatics. The Elixir of Chlorides of Arsenic, Iron and Mercury, Part I, is a similar product.

## Radcliff's Great Seven Seals or Golden Wonder Remedy.

The following is supposed by Nat. Drug. to be similar to the original:

Etherf	l.dr. 6
Chloroformf	
Camphorated oilf	d.dr. 4
Oil of peppermintf	dr. 2
Tincture of capsicumf	
Alcohol	

#### Radway's Pills.

Gambogeg	r.	8
Aloesg		
Jalapg	r.	15
Gingerg	r.	10
Make 30 pills.—Hager.		

## Radway's Ready Relief.

Soap liniment	.fl.oz. 3
Tincture of capsicum	.fl.oz. 1
Ammonia water	
Alcohol	
	Pierson.

#### Radway's Renovating Resolvent.

A similar preparation consists of a vinous tincture of ginger and cardamom, sweetened with sugar.

## Ransom's Hive Syrup and Tolu.

A similar preparation is composed of about the following (N. I.):

Fluid extract of squillfl.dr. 2	
Fluid extract of senegafl.dr. 2	
Soluble tincture of tolufl.dr. 2	
Tartar emeticgr. 4	
Sugarav.oz. 4	
Water, enough to makefl.oz. 4	

## Recamier Preparations.

See Ayer's Recamier Preparations.

#### Resorbin.

An ointment vehicle prepared by emulsifying sweet almond oil with yellow wax, gelatin, and soap.—Coblentz.

## Redlinger's Pills.

Calomel	.gr. 30
Resin of jalap	
Soap	.gr. 30
Gentian, powder	.gr. 30
Fennel, powder	
Mucilage of acaciasufficient for	or mass

Divide into pills weighing 21/2 grains.—D.

## Ricord's Urethral Bougies.

Lead acetate	1
Cacao butter gr. 57	-

Make into 48 suppositories.

#### Richter's Pain Killer.

Dr. Gerhard states that the following will yield a similar preparation:

Tincture of capsicum,	con-	
centrated	fl.oz.	71/2
Soap	gr.	120
Water	fl.dr.	10
Water of ammonia		
Camphor	gr.	180
Oil of rosemary	fl.dr.	1
Oil of lavender		1
Oil of thyme	fl.or.	1
Oil of clove	fl.dr.	1
Oil of cinnamon	drops	10
Caramelsuf	fficient to c	olor

Dissolve the soap in the water, add the solution to the tincture of capsicum, and finally add the other ingredients. Mix thoroughly and filter.

The tincture is made from  $2\frac{1}{2}$  av. ounces of powdered drug extracted with alcohol.

#### Robinson's Elixir of Paraldehyde.

This, according to the manufacturers' statemen's, contains 45 grains of paraldehyde in each fluidounce, dissolved in an aromatic menstruum. The Elixir of Paraldehyde, Part I, furnishes a satisfactory preparation.

## Roseter's Hair Regenerator.

Lead acetategr. 18	
Lac sulphurgr. 24	
Glycerin	
Rose waterfl.oz. 8	
-Ph. Rec.	

#### Rotterin

Zinc chloride	.gr. 10
Zinc sulphocarbonate	.gr. 10
Salicylic acid	.gr. 21/2
Boric acid	.gr. 8
Citric acid	
Thymol	.gr. 1
Sodium chloride	
Distilled waterf	l.oz. 16

The tablets contain one-half the quantities of the solids mentioned above.

## Rourke's Iodine Liniment.

See Giles' Iodide of Ammonia Liniment— D. C.

## Royal Catarrh Cure.

This, according to N. I., contains about the following:

Common saltparts	98
Carbolic acidparts	
Berberine hydrochloratepart	

## Royal Germeteur.

H. R. Slack says the following has the same chemical and physiological properties:

Sulphuric acidfl.oz. 2	
Water saturated with sulphur-	
etted hydrogenfl.oz. 1	
Common water, enough to make. gal. 1	

#### Rubifoam.

A similar preparation is given by the following:

)
1/2
) [
}
}
3
3
í
ó
3
3

Mix the soap, glycerin, syrup and water, stir well, add the alcohol, then the remainder of the ingredients, let stand a few days, and filter at a low temperature (to avoid separation of any soap).

## Ruppert's Face Bleach.

W. Robertson states that the following makes a similar preparation:

Corrosive sublimategr. 8	3
Tincture of benzoinfl.dr. 1	Ĺ
Water, enough to makefl.oz. 8	3

## Sage's Catarrh Remedy.

Powdered golden sealgr.	40
Indigogr.	4
Camphorgr.	16
Carbolic acidm.	
Sodium chloridegr.	

Triturate the camphor to powder by aid of a small quantity of alcohol, and mix with it the salt previously reduced to fine powder; rub the indigo and acid together, mix this with the salt and camphor, finally add the golden seal, and mix intimately in a mortar without much pressure,—Pharm.

## Salicylbromanilid.

See Antinervin.

#### Salubrin.

Acetic acid, glacialp	arts	2
Acetic etherp	arts	25
Alcoholp		
Waterp		

#### Sanitas.

A similar product may be prepared by passing air through warm oil of turpentine, which is in contact with water.—Frerksen.

#### Sanmetto.

This is claimed to be a combination of true santal and saw palmetto in a pleasant aromatic vehicle. The following may yield a preparation similar in therapeutic properties:

Yellow :	sandalwood.	 	av. oz. 1
Saw pal	metto	 	av.oz. 1
	elixir		

Mix the drugs with 16 fluidounces of the elixir, macerate for several days, agitating occasionally, and filter.

#### Saul's Catarrh Remedy.

This is said to be composed of the following (A. D.):

Compound tincture of benzoi	nfl oz. 2
Tincture of tolu	
Chloroform	fl.dr. 1
Sulphuric ether	fl.dr. 1
Aromatic spirit of ammonia.	fl.oz. 1
Oil of tar	fl.dr. 1
Alcohol	fl 07 5

#### Saunder's Bloom of Ninon.

Precipitated chalk	parts 7
Talcum	
Bismuth subcarbonate	parts 2
Zinc oxide	
Starch	
H •W	Snow

## Saw Palmetto Compound.

Fluid extract of saw palmettofl.oz.	2
Fluid extract of corn silkfl.oz.	
Fluid extract of sandalwoodfl.oz.	2
Simple elixirfl.oz.	10

## Schenck's Pulmonic Syrup.

H. M. Wilder has claimed this to be substantially its composition:

Wormwood av.oz. 1
Catnipav.oz. 1
Tansyav.oz. 1
Hyssopav.oz. 1
Hoarhoundav.oz. 1
Hopsav.oz. 1
Chamomileav.oz. 1
Comfreyav.oz. 1
Senegaav.oz. 1
Elecampaneav.oz. 1
Roil with sufficient water to make after

Boil with sufficient water to make, after straining, 2 quarts, then add:

Gum arabicav.oz. 3
Licoriceav.oz. 3
One good-sized turnip and finally:
Sugarlbs. 6
Brandyfl.oz. 16
Juice of 4 lemons.

## Schlotterbeck's Compound Hydrastis Mixture.

According to statement of composition by the manufacturers, the following furnishes a similar compound:

Rhubarbgr. 320
Golden sealgr. 160
Cinnamongr. 160
Potassium carbonate gr. 320
Pancreatingr. 64
Pepsingr. 64
Syrupfl.oz. 2
Water
Simple elixirfl oz. 6
Diluted alcoholsufficient

Mix the rhubarb, golden seal and cinnamon, reduce to powder and percolate with diluted alcohol so as to obtain 6 fluidounces of percolate. To this add the potassium carbonate and pancreatin, agitate occasionally for 24 hours, mix with the pepsin previously dissolved in the water, filter, and lastly add the syrup and elixir.

## Schreyer's Toothache Pellets.

Oil of clovesgr. 1	5
Oil of cassiagr. 1	5
Black peppergr. 6	
Sodium chloridegr. 6	
Acaciagr. 6	

Make into pellets weighing 8 grains each.

—Wittstein.

## Searle's Athlophorus.

Morphine sulphategr.	2
Fluid extract of colchicum seed. fl dr.	1
Fluid extract of guaiacfl.dr.	1
Potassium acetategr.	60
Potassium salicylategr.	60
Diluted alcohel fl.dr.	4
Syrup of squill, enough to make.fl.oz.	6

Make a solution by applying gentle heat.— N. I.

## II.

4	L.		
	Potassium acetategr.		60
	Sodium salicylategr.	4	90
	Sugarav.oz.		4
	Water		14
	Caramelsufficient to o		
	N	T	т

## Scott's Emulsion of Cod Liver Oil.

This is said to contain fifty per cent of cod liver oil, and 6 grains of calcium hypophosphite and 3 grains of sodium hypophosphite to the fluidounce.

## "Schinseng" (or Ginseng) Elixir.

A proprietary ginseng elixir is described as containing the "active proximate principle of the panax schinseng [Chinese ginseng] in an aromatic essence." A satisfactory Ginseng Elixir may be made as follows:

Fluid extract of ginseng.....fl.oz. 2 Simple elixir.....fl.oz. 14

## Seven Sutherland Sisters Hair Grower.

This is said to make a similar preparation (N. I.):

Bay rumfl.oz.	7
Distilled extract of witch hazelfl.oz.	9
Common saltgr.	60
Diluted hydrochloric aciddrop	1
Magnesiasufficie	

Mix the bay rum and witch hazel with some of the magnesia, filter, in the filtrate dissolve the salt, add the acid and filter again if necessary.

#### Sheffield's Dentifrice.

White castile soap, powderav.oz.	1
Precipitated chalkav.oz.	20
Carminegr. 4 or	
Sassafras flavoring (as below)fl.dr.	
Glycerinsuffici	

Rub the solids well together, add the flavoring in small portions during constant trituration and then glycerin enough to form a thin paste.

## (Sassafras Flavoring.)

Oil of sassafrasdr.	51/2
Oil of cinnamondr.	
Oil of wintergreendr.	1/2
Extract of vanilladr.	4
Alcoholdr.	4
~	

Shake before using.—D. C.

## Shiloh's Consumption Cure.

This formula for a similar preparation was contributed to the D. C.:

Chloroformfl.dr.	2
Ålcoholfl.oz.	1
Oil of peppermintdrops	10
Oil of tarfl.dr.	1
Morphine hydrochlorategr.	4
Diluted hydrocyanic acidfl.dr.	1
Extract of licoricegr.	120
Tincture of lobeliafl.dr.	4
Waterfl.oz.	1
Simple syrup, enough to make fl.oz.	16

## Sloan's Condition Powder.

Elecampane root, fenugreek,
flaxseed, juniper berries, pop-
lar bark, resin, mustard,
bran, eachparts 8
Licorice root, ginger, sodium
sulphate, sodium chloride, sul-
phur, iron sulphate, eachparts 6
Gentian, sodium carbonate,
eachparts 4
Black sulphuret of antimony,
potassium nitrate, coriander,
valerian, eachparts 2
Sanguinaria, lobelia, podophyl-
lum, dried alum, eachpart 1

#### Mrs. Smith's Butter Color.

The following is said to resemble the original (N. I.):

Annatto seed, bruisedav.oz.	3
Turmericgr.	120
Ammonium carbonategr.	40
Cottonseed oilfl oz.	7
Lard oilfl.oz.	1

Boil, stirring frequently, until the proper rich color has been attained; then strain and

allow to settle. Only the best material should be used.

## Simmons' Liver Regulator.

The powder consists of:

Liverwort	av.oz. 2	2
Leptandra	av.oz. 🛭	3
Serpentaria	av oz. 🛭	)
Senna	av.oz. 🤅	3

The liquid preparation consists of the above extracted with diluted alcohol.

#### Smith's Electric Oil.

Linseed oil	
Sassafras oil	fl.oz. 1
Chloroform	-Kilner

#### Smith's Tonic Syrup.

Quinine sulphategr. 30	,
Cinchonine sulphategr. 30	)
Fluid extract of podophyllumfl.dr. 2	,
Compound tincture of cardamom fl, dr.	1/2
Soluble citrate of irongr. 64	· -
Sugarav.oz. 10	)
Waterfl.oz.	,
Simple syrup, enough to makefl.oz. 16	,

A few drops of aromatic sulphuric acid may be added, if necessary, to dissolve the alkaloid salts.—N. I.

#### Somnal.

This is said by the Ph. Rundsch. to be an alcoholic solution of chloral hydrate and urethan.

## Solution of Chloro-Phosphide of Arsenic.

Arsenous acid	
Diluted hydrochloric acid	.fl.oz. 3
Distilled water	
Phosphoric acida f	ew drops

Dissolve the arsenous acid in the hydrochloric acid and 7 fluidounces of water by the aid of a gentle heat, add the remainder of the water, and then the phosphoric acid.—Fr. Sieker.

#### Steedman's Soothing Powders.

Opium powdergr.	3
Ipecacgr.	1
Milk sugargr.	8
Rice flourgr.	12

Mix and divide into 8 powders.—N. I.

#### Steresol.

A similar preparation is said to be the following:

Shellacgr.	540
Benzoingr.	75
Telugr.	75
Carbolic acidfl.oz.	
Oil of cinnamonfl.dr.	1′
Saccharingr.	45
Alcohol, enough to makefl.oz.	16

#### St. Jacob's Oil.

Squibb states this contains water, ether, alcohol, turpentine, aconite, and red coloring matter.

The following formula, constructed in conformity with the best information available, may furnish a satisfactory preparation:

Camphoroz.	1
Tincture of aconite rootfl oz.	2
Etherfl oz.	1
Oil of cedar fl.oz.	4
Alcohol enough to makefl.oz.	16
Tincture of alkanet enough to col	

## Stoddart's Peerless Liquid.

Bismuth oxychloridegr.	120
Precipitated chalkgr.	240
Glycerinfl.dr.	
Waterfl.oz.	

Color and perfume.

## Stoughton's Bitters.

Gentianav.oz.	1
Orange peelav.oz.	1
Columboav.oz.	
Chamomileav.oz.	1
Quassiaav.oz.	1
Caramelav.oz.	
Diluted alcoholfl.oz.	80

Extract the coarsely powdered drugs by maceration with the diluted alcohol for one week, agitating occasionally, then filter, and add the caramel.

Compound tincture of gentian is also similar.

## Strong's Arnica Jelly.

See No. VI in Cosmetic Jellies, Part III, for a similar compound.

## Succus Alterans.

See Mc Dade's Succus Alterans.

#### Svapnia.

The following is said to make a similar product (D. S. Dyson):

Deodorized tincture of opium..fl.oz. 16 Powdered gum arabic.....gr. 120

Evaporate the tincture to 4 fluidounces, remove the vessel from the fire, add the gum and triturate thoroughly, replace the vessel on the fire, and continue evaporation until the liquid is of such a density that it will have a syrupy consistence when cold. Now spread the liquid, while still warm, in thin layers on glass or porcelain plates and set aside to dry.

### Swan Down.

Orris root	parts	6
Zinc oxide	parts	13
Talcum	parts	14
	Cma	

## Swayne's Ointment.

Precipitated sulphurav.oz. 2	
Suetav.oz. 3	
Lardav.oz. 3	
—N. I.	

## Syrup of Figs.

I.		
	Senna. washed with alcoholav.oz.	
	Clovesgr.	120
	Cinnamongr.	60
	Nutmeggr.	60
	Sugarav.oz.	
	Water,	
	Alcohol	ient

Percolate the senna and spices with a menstruum composed of 1 volume of alcohol and 3 of water, until 8 fluidounces of percolate are obtained, and in the latter dissolve the sugar.

The senna is "washed" by macerating 4 ounces of the leaves in 16 ounces of alcohol for 2 days; after which they are separated from the liquid, dried and powdered.

 II.
 Senna
 av.oz. 14

 Coriander
 av.oz. 6

 Figs
 av.oz. 24

 Tamarinds
 av.oz. 18

 Cassia pulp
 av.oz. 18

 Prunes
 av.oz. 12

 Extract of licorice
 av.oz. 1½

 Spirit of peppermint
 av.oz. 1½

 Simple syrup
 gal. 1

Make an aqueous extract of the solid ingredients in which the required portion of sugar should be dissolved.

#### Sweet Quinine.

A mixture of cinchonine alkaloid with some ammoniated glycyrrhizin—Procter.

# Syrup of Trifolium Compound. (Compound Syrup of Red Clover Blossoms.)

Fluid extract of red clover	
blossoms fl.oz.	1
Fluid extract of burdockfl.dr.	4
Fluid extract of berberis aqui-	
foliumfl.dr.	4
Fluid extract of stillingia fl.dr.	4
Fluid extract of poke rootfl.dr.	4
Fluid extract of cascara amarga fl.dr.	4
Fluid extract of prickly-ash	
barkfl.dr.	1
Potassium iodidegr.	128
Waterfl.oz.	5
Sugarav.oz.	13

Mix the fluid extracts and the water, let stand for a few hours, filter, and in the filtrate dissolve the sugar and potassium iodide, and strain.

#### Tamar Indien.

According to the Repert de Pharmacie, the ingredients are as follows:

Purified tamarind pulpav.oz. 13	
Sugar, powderav.oz. 1	
Milk sugar, powderav.oz. 13	1/2
Senna, powderav.oz. 1	
Anise, powderav.oz.	4
Tartaric acidgr. 35	•
Essence of lemonfl.dr.	/2
Glycerinsufficient	
Mix and make into troches.	

## Tarrant's Compound Extract of Cubebs and Copaiba.

Nelson's "Handbook" gives the following formula for a compound paste of cubebs and copaiba:

Balsam copaibaav.oz. 1	6
Calcined magnesiaav.oz.	1
Powdered potassium nitrateav.oz.	8
Powdered cubebsav.oz. 4	8
Oil of wintergreenfl.dr.	4
Honeysufficier	ıt

Rub the copaiba and magnesia well together; then add the cubebs and potassium nitrate, using enough honey to form a paste. Lastly add the wintergreen oil.

#### Thymenthol.

Lister's Antiseptic Solution, Part I, has the same essential constituents claimed for this.

## Tarrant's Seltzer Aperient.

The following gives (N. I.) a similar preparation.

Sodium bicarbonateparts 1	7
Tartaric acidparts 1	
Rochelle saltparts 1	

#### Thomas' Electric Oil.

This formula for a similar product has been given by L. L. Briggs:

21 21 21 26	
Camphor	. 240
Oil of wintergreen fl. dr	. 4
Oil of origanumfl.dr	. 4
Chloroformfl.oz	. 1
Tincture of opiumfl.oz	. 1
Oil of sassafrasfl.oz	. 1
Oil of hemlockfl.oz	. 1
Oil of turpentinefl.oz	. 1
Balsam of firav.oz	. 1
Tincture of guaiacumfl.oz	. 1
Tincture of catechufl.oz	. 1
Alcoholfl.oz	
Alkanetsufficient to	color

## Thompson's Eye Water.

Copper sulphate	gr.	5
Zinc sulphate	gr.	20
Rose water		
Spirit of camphor	fl.dr.	2
Tincture of saffron	fl.dr.	2
	_	

-Kilner's Form.

## Thymolyptol.

According to statements of composition by the manufacturers, Lister's Antiseptic Solution, Part I, has about the same essential constituents.

# Tilden's Elixir of Iodo-Bromide of Calcium. Compound.

Compound Elixir of Iodo-Bromide of Calcium, Part I, contains calcium bromide, several iodides, etc.

#### Tobias' Derby Condition Powder.

Tartar emeticav.oz.	1
Black antimonyav.oz.	10
Sulphurav.oz.	
Potassium nitrateav.oz.	5
Fenugreekav.oz.	
uniper berriesav.oz.	
—Hag	er.

## Tolu, Rock and Rye.

Whiskeygal.	1
Rock candy syrupfl.oz.	
Tincture of tolufl.oz.	

Mix, allow to stand for several days, and filter.

## Tonic Aphrodisiac Tablets. (Wayne.)

The following formula contains essentially the same ingredients claimed for the original: makes a satisfactory product:

Extract of damianagr.	100
Extract of saw palmetto gr.	
Extract of cocagr.	100
Extract of nux vomica gr.	10
Phosphorusgr.	1/2
Make into 100 pills.	

## Tongaline.

According to the statement of composition made by the manufacturers, the following contains the essential ingredients in the required proportion for a similar preparation:

Fluid extract of tongsfl.oz.	8
Fluid extract of black cohoshfl.dr.	4
Sodium salicylateav.oz.	
Pilocarpine salicylategr.	
	1/4
Simple elixir, enough to makefl.oz.	16

For all practical purposes the two alkaloids could be replaced by 4 fluidrams of fluid extract of jaborandi and 21/2 fluidounces of fluid extract of colchicum seed.

#### Trousseau's Diuretic Wine.

The following furnishes a satisfactory preparation:

Squillav.oz.	l
Digitalisav.oz.	3
Juniper berriesav.oz. 15	3
Potassium acetateav.oz.	
Alcoholfl.oz. 16	3
White winegal.	Ĺ

## Tyree's Antiseptic Powder.

Formula as published by the manufacturer, in parts: Sodium borate, 50; alum, 50; carbolic acid, 5; glycerin, 5; the crystallized principles of thyme, 5; eucalyptus, 5; gaultheria, 5; mentha, 5. The following will probably represent these conditions:

Borax
Alum
Carbolic acid, crystallizedgr. 20
Thymolgr. 20
Mentholgr. 20
Oil of eucalyptus
Oil of gaultheria

#### Ulyptol.

See Eulyptol.

## Tropic Fruit Laxative.

According to A. Conrath, the following

Jalap, powderav.oz. 1
Senna, powderav.oz. 1
Sugar, powderav.oz. 1
Tamarind pulpav.oz. 6

Make into lozenges weighing 45 gr., coat with chocolate and sugar and wrap in tinfoil.

## Uncle Sam's Nerve and Bone Liniment.

Oil of origanum	.fl.oz.	1
Oil of rosemary	.fl.oz.	1
Oil of amber	.fl.oz.	1
Oil of hemlock	.fl.oz.	1
Oil of turpentine	.fl.oz.	16
Linseed oil		

## Upham's Asthma Remedy.

	_
Stramonium leavesav.oz.	-8
Skunk cabbageav.oz.	8
Lobelia herbav.oz.	
Potassium nitrateav.oz.	4
Waterfl.oz.	16

Mix the three drugs, reduce to powder, add the potassium salt previously dissolved in the water, and dry the whole.-Kilner.

#### Uricedin.

Sodium chloride	parts	4
Lithium citrate	parts	5
Sodium citrate		
Sodium sulphate	parts	68

#### Van Buskirk's Sozodont.

The following resembles the original (N. I.):

#### I. Liquid:

A1 1 1	a	
Alcohol	fl.oz.	1
Water	fl.oz.	11/
Soap		
Oil of wintergreen.	drops	2
Fluid extract of red	sanders sufficie	nt

Dissolve the soap in the mixture of alcohol and water, add the color, perfume with oil of wintergreen, add enough water to make the fluid measure 3 fluidounces.

#### II. Powder:

Infusorial earth (tripoli)gr.	40
Orris rootgr.	125
Precipitated chalk gr.	205

Perfume very lightly with oil of cloves.

#### Van Stan's Stratena.

A good cement of similar nature is the following:

Acetic acidfl.oz.	4
White glueav.oz.	3
French gelatingr.	
Shellac varnishfl.dr.	
Distilled water,fl.oz.	4

Dissolve the glue in the acetic acid with heat, and the gelatin in the water with heat. Mix the two solutions gradually and work until a uniform mixture results; then add the shellac varnish. Mix thoroughly and bottle.

#### Viburnal.

According to the statement of composition by the manufacturers the following contains the essential ingredients of a similar preparation:

Fluid extract of black hawfl.oz.	21/2
Fluid extract of black cohoshfl.oz.	21/2
Compound fluid extract of helo-	, -
nias	5
Simple elixirfl.oz.	6

#### Vin Mariani.

See Mariani Wine of Coca.

#### Vita Nuova.

See Ayer's Vita Nuova.

#### Walker's Vinegar Bitters.

A decoction of aloes, guaiac, sarsaparilla, cinchona, sassafras and golden seal preserved with acetic acid.—Cooley's Cyclopedia.

## Wampole's Asparoline Compound.

Each fluidounce, according to the manufacturers, contains: Diluted alcohol; guaiacum, 30 grains; asparagus seed, 30 grains; parsley seed, 30 grains; black-haw (bark of root), 60 grains; henbane leaves, 6 grains. The following formula is based upon the foregoing:

Guaiacoz.	1
Asparagus seedoz.	1
Parsley seedoz.	
Black-hawoz.	
Henbanegr.	
Diluted alcoholenous	

Reduce the drugs to powder and percolate with diluted alcohol enough to obtain 16 fluidounces of percolate.

#### Wampole's Tasteless Cod Liver Oil.

A preparation of Cod Liver Oil, combined

cherry, and compound syrup of hypophosphites with iron and manganese. Containing the curative agents from 25 per cent of cod liver oil, and rendered pleasant by aromatics. A preparation with essentially these ingredients and proportions is the following:

morning of the state of the sta	0.1
Fluid extract of wild cherryfl.oz.	2
Fluid extract of licoricefl.oz.	3
Glycerinfl.oz.	1
Simple syrupfl.oz.	1
I'luid extract of maltfl.oz.	6
Compound syrup of hypophos-	
phites, with iron and man-	
ganesefl.oz.	3
Fuller's earth, powdergr.	240
Caramal	

Mix the morrhuol with the glycerin and triturate with the fuller's earth; add the fluid extracts, syrup and malt, shake well, let stand for a day, agitating occasionally; filter, and to filtrate add the syrup of hypophosphites and sufficient caramel to color.

#### Warner's Safe Cure.

The following is said to produce a similar preparation:

Potassium nitrate, powderav.oz.	3/4
Liverwortav.oz.	1
Watersufficie	ent
Alcoholfl.oz.	
Glycerin fl.oz.	
Spirit of wintergreendrops	40

Infuse the liverwort with 16 fluidounces of hot water for 2 hours, strain and filter. Dissolve the potassium nitrate in the liquid; when cold add the alcohol, glycerin and spirit of wintergreen, and make up the measure to 16 fluidounces with water.

#### Warner's Safe Pills.

According to the examination of the Dresden (Germany) Health Department, each pill contains 13/4 gr. of aloes.

## Wayne's Diuretic Elixir.

"Elixir of Buchu, Juniper, Uva Ursi and Potassium Acetate," Part I, contains the essential ingredients of a similar preparation.

### Wei de Meyer's Catarrh Cure.

This consists largely of sodium bicarbonate.-N. I.

#### Weinmann's Dental Anæsthetic.

This contains about 53/2 per cent of cocaine, hydrochlorate, also alcohol, oil of pepperwith extract of malt, fluid extract of wild mint and iodine (indicating possibly aristol).

#### Watt's Anti-Rheumatic Pills.

A similar preparation is made as follows
Aloesgr. 240
Gamboge       gr. 240         Hellebore       gr. 120
Calomel
Yellow sulphide of antimonygr. 15
Oil of clove
Spirit of camphorsufficient

## Weld's Syrup of Chloride of Iron.

Make into 5-gr. pills.

The following makes a non-astringent syrup containing iron chloride:

Solution of chloride of ironfl.dr. 4
Glycerinfl.oz. 13
Citric acidgr. 80
Water,
Ammonia waterof each, sufficient

Mix the solution of iron with the glycerin, dissolve the acid in 1 fluidounce of water, mix the two solutions, add ammonia water until the liquid is only feebly acid, and add the remainder of the water.

## Whiteley's Nipple Wash.

The following is said to be a good wash:
Boraxgr. 40
Tanningr. 10
Glycerin
Rose water

## Winslow's Soothing Syrup.

The following, contributed to the D. C., is said to furnish a similar preparation:

Morphine sulphategr. 1
Sodium carbonategr. 2
Syrup, simplefl.oz. 3
Water
Spirit of fennelfl.dr. 2

## Wistar's Balsam of Wild Cherry.

Fluid extract of wild cherryfl.oz.	1
Fluid extract of ipecacfl.dr.	2
Fluid extract of squillfl.dr.	2
Tincture of opiumfl.dr.	1
Tartar emeticgr.	2
Sugar-house syrupfl.oz.	3
Alcoho'fl.dr.	6
Spirit of anisedrops 1	10
Compound tincture of cudbear,	
Water, of each, enough to make fl.oz.	8
N.	I.

## Wither's Antizymotic Solution.

The following contains the same essential ingredients as were found by Bierbach's analysis in the original:

Corrosive sublimategr. 10	3
Aluminum chloridegr.	3
Zinc chloridegr.	
Potassium chloridegr.	
Sodium chloridegr. 60	
Hydrochloric acid	5
Water, enough to makefl.oz. 16	

## Woolbridge's Treatment of Typhoid Fever.

This method of treatment of typhoid requires the employment of the three formulas:

[.	(Tablets.)	
	Podophyllingr. $\frac{1}{960}$	
	Calomelgr. $\frac{1}{16}$	
	Guaiacol carbonategr. 16	
	Mentholgr. $\frac{1}{16}$	
	Eucalyptolsufficient	
гΥ	(Tableta)	

1. (Tablets.)		
Podophyllin	gr.	000
Calomel	gr.	1
Guaiacol carbonate		
Menthol		
Thymol		
Eucalyptols		
71		

III. (Capsules.)	
Guaiacol carbonate	.gr. 3
Thymol	.gr. 1
Menthol	.gr. ½
Eucalyptol	.m. 5

IV. (Tablets for children.)	
Podophyllingr	. 1
Calomelgr	
Guaiacol carbonategr	
Mentholgr	
Eucalyptolsuffici	

V. (Capsules for children)	
Guaiacol carbonategr.	1/2
Thymolgr	1/8
Mentholgr.	
Eucalyptolm.	
Olive oilsufficien	t

## Wright's Face Powder.

Snow gives the following for a similar preparation:

Talcumav.oz.	30
Starchav.oz.	40
Calcium sulphateav.oz.	
Bismuth oxideav.oz.	<b>J</b> .

#### SUPPLEMENTARY FORMULAS.

The following supplementary formulas are given in response to requests received since the issuance of the first edition:

#### Acacine.

Said to be a mixture of dried casein, sodium bicarbonate and sugar, containing 10 per Barker's Nerve and Bone Liniment. cent of the casein.

## Antinervin.

Salicylio Acetani	nium bromide. c acid	 . part	$\frac{1}{2}$
I.			

This formula is also given:

Acetanilid.....part 1 Sodium salicylate ......part 1 '

## Antipyonin.

Principally a tetraborate or polyborate of sodium .- A. D.

#### Antirheumatin.

This is said to be chiefly a combination of sodium salicylate and methylene blue.-M. B.

## Antiseptin.

Zinc sulphateparts	34
Boric acidparts	4
Zinc iodidepart	1
Thymolpart	1
Arabian Polasam —Goldma	an.

#### abian Balsam

Cottonseed oil	.fl.oz.	15
Origanum oil	fl. 02.	1
Oil of turpentine	.fl.dr.	4
	N.	

#### Armenian Pills.

Balsam of copaibaparts	14
Magnesia, calcined lightparts	2
Cubebs, powderparts	7
Armenian bole, powderparts	7

Heat the copaiba until it acquires the consistence of a plaster, that is, until most of the volatile oil has been dissipated, then mix it with the magnesia and set it aside that the mass may set. Next add the powdered cubebs and Armenian bole, and mix intimately. Make the mass into pills of 8 gr. each and roll them in Armenian bole. - Schacht.

#### Aseptic or Aseptinic Acid.

Said to be an aqueous solution of 5 parts of boric acid in 1000 parts of hydrogen peroxide (5 per cent), with or without 3 parts of salicylic acid. - Thoms.

#### Ayer's Recamier Balm.

A "balm" of this type is made as fo	llows
Zinc oxidegr.	500
Glycerindrops	
Alcoholdrops	
Corrosive sublimategr.	25
Water fl.oz.	30

Camphorav.oz	. 11/4
Oil of tarfl.dr	. 4
Oil of thymefl.oz	
Oil of turpentinefl.oz	
Black or Franklin oil, enough to	
makefl.oz.	16
	J. T
1	V. I.

#### Bate's Salve.

m.	
Beeswaxav.oz.	2
Amber resinav.oz.	3
Linseed oil fl. oz.	10
Red leadav.oz.	4
Boil with constant stirring until the	whole
dark brown B. & C. Drug.	

## Battley's Sedative.

Extract of opiumgr.	600
Boiling waterfl.oz.	
Alcohol fl.oz.	

Dissolve the extract in the boiling water, allow to cool, add the alcohol and the cold water and filter.

#### Bismuthol.

A mixture of bismuth, salicylate and sodium phosphate furnishes an apparently similar product.

#### Borolyptol.

The following formula has been contributed to W. D. as making a similar preparation: Glyceride of boroglycerin....av.oz. 11/4 Formaldehyde solution.....drops 20 Butyric ether.....drops 3 Distilled water..........fl.oz. 16

#### Butyromel.

Fresh butterav.oz.	2
Honeyav.oz	1
Mix until a homegeneous mass is obtai	ned.
Coblent	z.

#### Byrolin.

This appears to be a mixture of boric acid, glycerin and lanolin.

## Calder's Saponaceous Dentifrice.

Powdered castile soapav oz. 4
Precipitated chalkav.oz. 2
Magnesium carbonateav.oz. 1
Sugarav.oz. 1
Oil of wintergreen sufficient to flavor
—N. I.

#### Dean's Cactus Oil.

The following formula has been contributed to W. D. as making a similar preparation:

Neutral oilgal. 1
Oil of mirbanefl.oz. 3
Alkanetsufficient to color

## Duffy's Malt Whiskey.

Diluted and sweetened alcohol has similar physical properties.

#### Formacoll.

A combination of formaldehyde with gelatin.

Garfield Tea. This is a good "senna" tea:
Sennaparts 8
Dog grasspart 1
Liverwortpart 1

#### Glutiform.

A combination of formaldehyde with gelatin.

#### Glutol.

A combination of formaldehyde with gelatin.

## Hall's Catarrh Remedy.

A good catarrh remedy of this type is the following:

Gentian, coarse powderav.oz.	11/2
Bitter orange peel, coarse powder.gr.	300
Cardamom seeds, coarse powder.gr.	100
Potassium iodideav oz.	1
Diluted alcoholsuffice	eient

Macerate the crude drugs in 12 fluidounces of diluted alcohol for 48 hours, then transfer to a percolator and allow to percolate slowly; when the liquid has ceased to percolate pass enough menstruum through the percolator to make the finished product measure 16 fluidounces. In this dissolve the potassium iodide.

In other words, it is potassium iodide dissolved in compound tincture of gentian of the British pharmacopæia.

## Hamburg Tea.

Senna, cutav.oz.	8
Mannaav oz.	3
Coriander av.oz.	1

#### Holman's Liver Pad.

May apple rootoz.	1/2
Leptandra rootoz.	1/2
Bayberry barkdr.	2
Red cinchona barkoz.	2
Fenugreekoz.	1/2
Guiaiac resinoz.	11/2
Oil of eucalyptusfl.dr.	2

Grind the solids to powder, mix with them the oil and make into a pad, using cotton cloth of suitable thickness as the envelope.—Kilner's Form.

#### Kaskine.

This is said to consist largely of sugar.

## Kohler's One-Night Corn Salve.

A good	salve of	this type	is the	follow	ing:
Salicylic	acid			.part	1
Simple	cerate			.parts	3

## Kutnow's Improved Laxative Powder.

This is said to be the name given to an artificial Carlsbad salt.

#### Listol Tablets.

Boric acidgr.	12
Tannic acidgr.	
Salicylic acidgr.	6
Extract of belladonnagr.	11/2
Extract of heloniasgr.	$\frac{1}{3}$
Extract of herbanegr.	9
Extract of opiumgr.	3
Dithymol diiodidegr.	0.0
Alum gr.	
Eucalyptolgr.	11/2
Hydrastine hydrochlorategr.	13

Make into 12 compressed tablets.

The foregoing corresponds with that of the manufacturers, except that they mention 'listol' instead of dithymol diiodide, and colorless extract of hydrastis for hydrastine, and they do not indicate quantities in the three last items.

#### Lloyd's Hydrastis.

See Colorless Solution of Hydrastis, Part I.

#### Maizo-Lithium.

Fluid extract of licorice rootfl.oz.	1
Fluid extract of corn silkfl oz.	2
Lithium citrategr.	120
Simple elixir, enough to makefl.oz.	

#### McGill's Orange Blossom.

The following appears to be similar in physical properties:

2	Zinc sulphate	gr.	60
	Alum	_	15
1	Extract of hyoscyamus	.gr.	1
	White wax		30
	Cacao butter		
1	Make into suppositories of 32 g	r. ea	ch.

#### Merchant's Gargling Oil.

See Gargling Oil.

Miller's Golden Oil.	Phillip's Milk of Magnesia.
Oil of lavender	A Milk of Magnesia is given in Part I.
Oil of sassafrasfl.oz. 1	Phillip's Syrup of Wheat Phosphates.
Oil of turpentine	Said to be similar to:
	Wheat phosphates, Phillip'sfl.oz. 2
Mitchell's Eye Salve.  Petrolatum, whitegr. 350	Simple syrupfl.oz. 14
White waxgr. 130	Phillip's Wheat Phosphates.
Zinc oxidegr. 45 Oxide of mercurygr. 5	This preparation is said to be similar to
Oil of lavenderdrops 10	Pepper's solution of acid phosphate with iron-
Odontodol.	Pray's (Madam) Nail Polish.
Cocaine hydrochloratepart 1 Oil of cherry-laurelpart 1	Paraffin waxgr. 60
Tincture of arnicaparts 10	Chloroform
Solution of ammonium acetateparts 20	Quionin.
Oleoze Co. Oil of lavenderpart 1	A mixture of about 10 per cent of quinine
Oil of clovespart 1	and 90 per cent of the other alkaloids of
Oil of cinnamonpart 1 Oil of thymepart 1	cinchona bark has similar physical properties.
Oil of citronpart 1	Richmond's Samaritan Nervine.
Oil of mace part 1 Oil of neroli part 1	The following is a typical "nervine:"
Peru balsam parts 3 Deodorized alcohol parts 240	Potassium bromideav.oz. 1
Pasta Mack.	Sugar
Sodium bicarbonateav.oz. 5	Waterfl.oz. 5
Tartaric acidav.oz. 4 Powdered starchav.oz. 7	Oil of cassiadrops 10
Sweet almond oil	Roberts' Camphor-Tar Ointment.
Oil of rosedrops 5 Oil of clovesdrops 2	A camphor tar ointment may be made as
Mix the first three ingredients, add the re-	follows:
maining substances, and beat the whole to a	Tarav.oz. 1 Camphorav.oz. 1
stiff paste.—N. I.	Lardav.oz. 8
Patterson's Toothache Wax.	Roche's Embrocation.
A toothache paste may be made by immers- ing absorbent cotton in melted paraffin wax	Oil of amberfl.oz. 1
colored with alkanet and containting 1 per	Oil of clovesfl.oz. 1
cent of carbolic acid.	Olive oilfl.oz. 2
Phalon's Hair Restorative.	Oil of amber, rectifiedfl.oz. 2
The following is recommended as a good preparation of its type:	Tincture of opiumfl.oz. 2 Lardav.oz. 1
Alcoholfl.oz. 8	III.
Castor oil	Asafetidagr. 160
Color this mixture with alkanet root and flavor with oils of bergamot, neroli, verbena	Olive oil
and orange.	Oil of turpentinefl.dr. 2
Phenolid.	Digest the asafetida with the olive oil for
Goldman gives the following:	some hours; decant and mix the solution with
Acetanilidpart 1 Sodium salicylatepart 1	the other oils and add a few drops of oil of gaultheria.
	8

#### Sal Muscatel.

See Tarrant's Aperient or Eno's Fruit Salt.

## Schlotterbeck's Compound Mixture of Helonin.

See Compound Elixir of Helonias, Part I.

## Schultze's Blood-Purifying Powder.

2
14
3
3
4
er.

## Seven Sutherland Sisters Scalp Cleaner.

This is said to be largely borax and salt.

#### Solutol and Solveol.

These are apparently mixtures of crude cresol and solution of soda containing 25 to 50 per cent of cresol.

## Springsteen's Uterine Capsules.

According to a contribution to W. D., these are now sold under the name Viavi, which see.

## Steedman's Soothing Powders.

The following is typical of its class:	
Opium powdergr.	3
Ipecacgr.	1
Milk sugargr.	8
Rice flourgr.	12
Mix and divide into 8 powders.	

## Swift's Syphilitic Specific.

The following is said to yield a satisfactory specific:

Fluid extract of fringe tree bark.fl.oz.	1
Fluid extract of prickly ash bark.fl.dr.	4
Fluid extract of white sumacfl.dr.	2
Fluid extract of red sumacfl.dr.	2
Fluid extract of sarsaparillafl.dr.	6
Copper sulphategr.	8
Pyroligneous aciddrops	20
Alcoholfl.oz.	4
Water, enough to makefl.oz.	16

#### Trask's Magnetic Ointment.

The following are types of "magnetic" ointments.

#### I.

Fine-cut tobacco, Raisins,	
Lardequal parts of each	h
Simmer together and strain.	

#### II.

Cerate of subacetate of Powdered opium	
Mix thoroughly.	

#### Tritica.

An aqueous fluid extract of couch grass, preserved with about 10 per cent of alcohol, has similar physical properties.

#### Trix.

See No. 1, under Cachous, in Part III.

#### Viavi.

According to a contribution to W. D., a mixture of tannin, opium and cacao butter, put up in capsules, resembles this remedy. It is sold as a cure for female diseases.

## Wampole's Laxative Compound.

A teaspoonful of this preparation is said to represent in liquid form an equal amount of compound licorice powder. The following will therefore approximately represent this preparation:

Fluid extract of sennafl.oz.	
Fluid extract of licoricefl.oz. Oil of fennelfl.dr.	
Alcoholfl.dr.	
Syrup, enough to makefl.oz.	

Dissolve the oil in the alcohol, add to the fluid extracts, and then incorporate with the syrup.

## Wistar's Cough Lozenges.

See Troches of Glycyrrhiza and Opium, U. S. P.

#### Wheeler's Elixir.

An elixir of this type consists of equal parts of elixir of cinchona and iron, N. F., and elixir calcium lacto-phosphate, N. F., flavored with bitter almond oil and colored with cochineal.

## Zymoidin.

This is an antiseptic, said to be composed of oxides of zinc, bismuth and aluminum, with iodine, boric, carbolic, gallic and salicylic acids, quinine, etc.

#### PART IV.

## VETERINARY PREPARATIONS.

Many of the formulas here offered, such as those for condition powders, liniments, poultry powders, etc., may be kept on hand in convenient form to be offered for counter sale.

#### SECTION I.—HORSE MEDICINES.

The following doses are intended for grown horses, since foals generally require treatment only for those diseases which are peculiar to foals. As a general rule the quantities ordered may be adjusted to suit the age of the animal, according to the following proportions: For a colt 1 year old, 25 per cent of the full dose; for a 2-year old, 50 per cent, and for a 3- or 4-year old, 75 per cent of the full dose.

#### Anæmia Medicines.

Anæmia is often due to lack of exercise in the open air; this condition is most marked in young animals. It may also be due to improper food. Treatment consists in iron and arsenic, giving green food if possible, and indulging in light exercise in the open air.

Τ.

Sulphurav.oz.	2
Potassium bicarbonateav.oz.	
Sodium chlorideav.oz.	20
Give one tablespoonful with each mea	1.

H.

Sulphurav.oz.	2
Antimony sulphideav.oz.	1
Ferrous sulphateav.oz.	1
Calamusav.oz.	2
Sodium sulphateav.oz.	6
Sodium chlorideav.oz.	8
Reduce all to powder and mix well.	
Give one tablespoonful with each mea	l.

III.

11.	
Saccharated iron carbonateav.oz.	6
Manganese saccharateav.oz.	1
Cinnamonav.oz.	1/2
Clovesav.oz.	1/2
Calamus, in No. 8 powderav.oz.	5
Sodium chloride (common salt). av. oz.	7
Sodium sulphateav.oz.	20
Reduce all to powder and mix well.	
Give one tablespoonful at every meal.	

IV. Fowler's solution.

Give one tablespoonful once daily upon bread.

#### Appetite, for Loss of.

Loss of appetite is often the result of disorders of digestion and may be the accompaniment or precursor of other diseases. The following remedies may be of value:

Γ.

Crude tartar (argols) av.oz. Antimony sulphide	1
Calamus rootav.oz.	4.
Gentianav.oz.	4
Juniper berriesav.oz.	4
Caraway seedav.oz.	2
Mustard seedav.oz.	
Rye flourav.oz.	
Watersufficier	

Mix the drugs in powder form and add enough water to form a mass or paste.

Spread 1 tablespoonful of this on the horse's tongue 3 times a day.

II.

Calamus	av.oz.	1
Gentian	av oz.	1
Ginger	av.oz.	1
Wormwood	av.oz.	1
Sodium chloride	av.oz.	2
Rye flour	av.oz.	2
Tincture of capsicum	.fl.dr.	21/2
Water or simple syrup, sufficier	ıt	,
to form a mass		

This is to be used like the preceding.

III.

Gentian root	:	 .av.oz. 4
Sodium sulphat		
Sodium chlorid		
Sodium bicarbo	nate	 .av.oz. 1

Give 2 tablespoonfuls with each meal.

IV.

Crude tartar (argols)av.oz.	1
Antimony sulphidegr.	
Potassium bicarbonateav.oz.	1 3/4
Gentianav.oz.	134
Caraway seedav.oz.	13/4

Mix and divide into 10 powders.

Give 1 once a day mixed with the horse's eed.

V. See also Condition Powders.

#### Blister.

Cantharides, fine powderav.oz	
Euphorbium, fine powderav.oz	. 3/4
Corrosive sublimateav.oz	. 1/4
Mercurial ointmentav.oz	. 2
Linseed oilfl.da	. 4
Sulphuric acidfl.oz	. 1
Nitric acidfl.oz	
Oil of turpentinefl.oz	. 4
Petroleumfl.oz	

Add the sulphuric acid gradually to the linseed oil, then add the nitric acid very gradually, and after that the turpentine and petroleum. Let stand for several days, decant from residue, and mix with the decanted liquid the mercurial ointment, to which have previously been added the corrosive sublimate, cantharides and euphor-

Extreme care must be taken, in mixing the acids and oil, to add the acids very slowly, with constant stirring, to the oil contained in a broad vessel, such as an evaporating dish.

#### II.

Cantharides, powderav.oz.	21/
Oil of turpentinefl.oz.	11/
Acetic acidfl.oz.	1
Lanolinav.oz.	5
Petrolatumav.oz.	5

Mix the first three and allow to stand for 24 hours, then add the lanolin and petrolatum melted on a water bath and mix, stirring until cold.

#### III.

Cantharidesav.oz.	1
Oil of turpentinefl.oz.	8
Water of ammoniafl.oz.	
Olive oilfl.oz.	2
Oil of sassafrasfl.oz.	1
IV.	
Biniodide of mercurygr. 6	0

Oil of cajuput.....gr. 60 Petrolatum .....gr. 480

#### Catarrh, Remedies for Bronchial.

Bronchial catarrh usually begins with fever and is generally accompanied by coughing, and the discharge of purulent matter from the nose. If the dry cough does not soon loosen, and there is no nasal discharge, the fomentations recommended under "Glanders " should be employed. These fomentations should not be continued longer than to a fine powder, and add them to the oils dis-

necessary to induce the discharge, as otherwise harm may result to the nasal mucous membrane. In addition to using the medicines mentioned below, warm applications should be bound around the throat and warm drinks should be administered.

Ammonium chlorideav.o	z.	1
Fenugreekg		
Fennelg	r.	400
Marshmallowav.o		
Reduce all to powder and mix wel	1	

Give in 2 doses in warm meal or in warm "soft" food.

#### II.

Sodium chloride	.av.oz.	10
Antimony sulphide	.av.oz.	2
Fenugreek	.av.oz.	1
Licorice root	.av.oz.	1

All ingredients should be in powder and should be well mixed.

Give 1 tablespoonful with each meal.

Sodium sulpha	ate.		2V C	7 4
Sodium chlori				
Sodium bicart				
Licorice root.				

All should be in powder and should be well mixed.

Give 1 tablespoonful with each meal.

Mercurial ointmentav.oz.	2
Suetav.oz.	3
Hyoscyamus oilav.oz.	5

To be applied to the throat every morning and evening.

#### Carbolic Composition.

Hager has devised the following for veterinary and farmers' use. It keeps sores clean, cures scab and itch, and kills vermin, and is said to keep flies, mosquitoes and such insects from animals:

Benzoinav.oz.	1
Aloesav.oz.	1/2
Salicylic acidgr.	
Oil of spikefl.dr.	
Oil of anisefl.dr.	
Alcohol fl.oz.	
Oleic acid, crude fl.oz.	1
Caustic sodagr.	250
Boraxgr.	
Waterfl.oz.	
Carbolic acid, crudefl.oz.	
Carbone acid, crude	

Rub the benzoin, aloes and salicylic acid

solved in the alcohol. After a day's maceration, add to the mixture the oleic acid, the soda and the borax dissolved in the water, and, lastly, the carbolic acid. Shake the whole well for half an hour, and after a week's maceration in a cool place decant the clear liquid. For use shake well with twice its bulk of water, and add 100 to 120 times its bulk of water, stirring thoroughly. For scab and itch dilution with only 30 or 40 times its bulk is necessary.

#### Cathartic Medicines.

I.

Cape aloesav.oz.	1
Gingergr.	
Potassium carbonategr.	
Gambogegr.	60
Oil of fenneldrops	20

Make into a mass by means of powdered soap and water, roll into a cylinder about 2 inches long, and cover with gelatin or with thin paper, like tissue paper, suitably oiled or greased to prevent the adhesion of the mass to the paper.

II.

Cape aloesav.oz.	1
Gingergr.	
Potassium carbonategr.	60
Croton oildrops	10
Oil of anisedrops	30
Waterfl.oz.	8
Linseed oilfl.oz.	8
Tincture of opiumfl.dr.	4

Powder the aloes and ginger, add the potassium carbonate, mix with the water, add the oils of anise and croton, then the tincture and finally the linseed oil.

To be given at one dose. In cases of unusual weakness or prostration of the animal, the croton oil can be omitted.

In putting up this formula in large quantity for general sale, it would be better to substitute powdered gamboge for croton oil, using in the prescription above named 60 gr.

III. See also Colic Remedies and Constipation Cure.

#### Colic Remedies.

Colic is usually due to obstinate constipation and retention of urine; sometimes it is caused by diarrhœa. If due to the former, purgative and carminative remedies should

be administered. When the bowels and urinary organs begin to act, the colic generally ceases. During the course of treatment it is advisable to rub the back, abdomen and legs with a turpentine mixture until perspiration ensues.

Τ.

Chlorodyne	fl.oz.	1
Spirit of nitrous ether	fl.oz.	2
Linseed oil	fl.oz.	13

Give at one dose and repeat in 2 hours, if necessary.

II.

Ether		
Castor oil	.fl.oz.	18
Give at one dose.		

III.

Aloes av.oz. 1	1/2
Green soapsufficient to form a mass	_
Give at one dose as soon as colic appear	s.

IV

Oil of turpentinefl.oz.	4
Ammonia waterfl.oz.	1
Alcoholfl.oz.	

To be applied to the abdomen.

V. Hypodermic injections containing eserine may be employed, instead of internal remedies, to relieve colic due to constipation; these act much quicker than internal remedies.

A

Eserine sulphategr. 1 Distilled waterfl.dr.	1 1/2
Di1	12

Dissolve and inject at one dose.

В.

Eserine sulphategr.	11/2
Pilocarpine hydrochlorategr.	5
Distilled waterfl.dr.	

Use at one injection. The latter is particularly valuable when the colic is due to obstinate constipation.

When using these injections, the external treatment should be the same as otherwise.

VI.

Creolin	fl.dr.	4
Oil of turpentine	fl.oz.	2
Aromatic spirit of ammonia.		
Tincture of asafetida		
Linseed oil		

For one dose.

Mix these substances and dispense in heavy

, , , , , , , , , , , , , , , , , , , ,	Mix these substances and dispense in neavy
Tincture of opiumfl.oz. 1	bottles.
Etherfl.oz. 1	The quantity named can be used at one
Alcohol	
Oil of anisedrops 30	dose if a very severe case, and even repeated
Give one tablespoonful every 15 to 30	if the exigency demands it; otherwise, the
minutes in a pint of water.	above recipe can be divided into two doses.
Rub the abdomen and back with straw,	
	Condition Powders.
wet with oil of turpentine.	These preparations are also known as
VIII.	"Horse Powders," "Horse and Cattle
Aloes	E 122 144 Ct 1 E 122 El
Potassium carbonategr. 45	Food" and "Stock Food." Pharmacists
Watersufficient to form a bolus	frequently dispense preparations of this
Give one such bolus every hour.	character made by themselves; as titles, they
IX.	may select those given above, also such as
Tartar emeticgr. 45	"Maud S. Condition Powder," "Prairie
Sodium sulphateav.oz. 3	Condition Powder," "Farmer's Condition
Carawayav.oz. 1	
Make one powder, and give one such pow-	Powder," "Arabian Condition Powder,"
der every half-hour until the bowels move.	"O. K. Condition Powder," etc.
X.	The following list will be of interest, as
	well as of service, in determining what
Magnesium sulphateav.oz. 18	ingredients may enter into the composition
Althæa	of a condition powder:
Rye flourav.oz. $3\frac{1}{2}$	Alteratives.—Sodium hyposulphite, sul-
Add water or simple syrup to form a mass.	
Give one-half of this mass and repeat the	phur.
dose in one-half hour.	Diuretics and Diaphoretics.—Alum, black
XI.	antimony, buchu, cream of tartar, pure and
Sodium sulphateav.oz. 18	crude; juniper berries, lobelia, potassium
Juniper berries, coarse powder.av.oz 31/2	nitrate, resin.
Rye flourav.oz. $3\frac{1}{2}$	
Make into a mass or paste with simple	Expectorants.—Blood root, potassium chlo-
syrup or glucose.	rate, elecampane, licorice root, lobelia, resin.
This is to be administered like the preced-	Tonics. — Iron carbonate, gentian, cin-
ing. It is to be given when the colic is due	chona, poplar bark, iron sulphate.
to both constipation and retention of urine.	-
•	Aromatics and Correctives.—Anise, sodium
XII.	bicarbonate, camphor, cascarilla, capsicum,
Alumgr. 300	cumin seed, fenugreek, ginger, grains of
Althæaav.oz. 1	paradise, mustard, salt, sassafras.
Ginger	Emollients and Laxatives.—Aloes, mag-
White oak barkav.oz. 1½ Juniper berriesav.oz. 1½	nesium sulphate, flaxseed meal, sodium sul-
	•
All of these should be in powder and	phate, oil cake meal.
should be formed into a mass with glucose or	Sedatives.—Asafetida, digitalis, skunk cab-
molasses.	bage, valerian.

Black antimony.....av.oz. 8 Sulphur .....av.oz. 7 Elm bark......av.oz. 31/4 Potassium nitrate.....av.oz. 1½ Anise seed.....av.oz. Reduce all to powder and mix well.

The usual dose of these powders is about

1 tablespoonful 2 or 3 times daily in food.

One-fourth of this mixture is to be given

Tincture of opium.....av.oz.

Spirit of peppermint . . . . . . fl.oz.

Spirit of nitrous ether.....fl.oz.

Ether.....fl.oz.

Diluted alcohol.....fl.oz. Linseed oil.....fl.oz.

Sodium bicarbonate.....gr. 24)

It is of value when the colic is

1

every hour.

XIII.

caused by diarrhœa.

II.	37
Elecampaneav.oz. 8	,
Fenugreekav.oz. 8	
Linseedav.oz. 8	
Juniper berriesav.oz. 8	
Poplar barkav.oz. 8	
Resinav.oz. 8	
Licorice rootav.oz. 6	
omget (title)	
Sodium sulphateav.oz. 6 Sodium chlorideav.oz. 6	
Sulphur	al
Copperas av.oz. 6	aı
Sodium carbonateav.oz. 4	V
Gentianav.oz. 4	
Black antimonyav.oz. 2	
Potassium nitrateav.oz. 2	
Coriander seedav.oz. 2	
Valerian	
Lobelia	
Dried alumav.oz. 1	
III.	
Fenugreekav.oz. 16	
Sulphur	
Cream of tartar	
Licorice rootav.oz. 4	
Black antimonyav.oz. 2	
Gentianav.oz. 1	
Anise	
Common saltav.oz, 1	I.
IV.	
Sodium chlorideav.oz. 1	
Fenugreekav.oz. 4	
Licorice rootav.oz. 4	
Flaxseed, groundav.oz. 7	
Give 1 ounce daily.	
V.	
Black antimonyav.oz. 1	Ι.
Resinav.oz. 1	
Capsicum av.oz. 1	
Gentianav.oz. 2	
Fenugreekav.oz. 2	
Sulphur	C
Saltpeter	
Cream of tartarav.oz. 2	
Ginger	
Licoriceav.oz. 3	
Dose, one tablespoonful once or twice a	
day.	
VI.	
	C
Gentian	Ĭ
Ginger	
Ginger av.oz. 4 Sulphate of iron av.oz. 2	be
Potassium nitrate	uı
Fenugreekav.oz. 6	pΙ
Dose, one dessertspoonful in the feed,	er
morning and night.	
and many	ı

V11.	
Cream of tartarav.oz.	8
Sulphurav.oz.	8
White resinav.oz.	8
Guaiac resinav.oz.	5
Potassium nitrateav.oz.	3
Gentian av.oz.	8

Reduce all to powder and mix well.

This powder is to be recommended if an alterative is desired.

Golden sulphuret of antimony...gr. 240

#### VIII.

Elecampaneav.oz.	4
Licorice rootav.oz.	4
Linseedav.oz.	4
Fenugreekav.oz.	4
Resinav.oz.	4
Aniseav.oz.	2
Capsicumav.oz.	2
Gentianav.oz.	2
Potassium nitrateav.oz.	2
Valerianav.oz.	2
Sulphurav.oz.	2
Copperasav.oz.	2
Juniper berriesav.oz.	2
Black antimonyav.oz.	1
Sodium sulphateav.oz.	1
Sodium chlorideav.oz.	19
	19
37	

#### IX

Exsiccated iron sulphateav.oz.	5
Cantharidesav.oz.	1
Gingerav.oz.	3
Black antimonyav.oz.	6
Potassium nitrateav.oz.	5
Sulphurav.oz.	10
Flaxseedav.oz.	
Gentianav.oz.	7
Cream of tartarav.oz.	3
White resinav.oz.	5
Aniseav.oz.	5

## Condition Powder, Darby's.

Reduce all to powder and mix well.

Sodium sulphateav.oz. 8
Sulphurav.oz. 4
Fenugreekav.oz. 4
Gentianav.oz. 2
Black antimonyav.oz. 2

Reduce all to fine powder and mix well.

#### Constipation Cure.

For constipation, the usual cathartics may be administered. The remedies mentioned under "Colic" as useful in this latter complaint resulting from constipation, may be employed.

See also Cathartics.

#### Cough Remedies.

Coughs should be treated by binding warm applications about the throat or making fomentations as described under "Glanders." If the nasal secretion is too copious, it may be checked by means of a powder containing lead acetate. In addition the following remedies may be employed:

I.

Antimony sulphideav.	oz. 1
Licorice root	oz. 2
Sodium chlorideav.	oz. 5

Mix all in powder form.

Give two tablespoonfuls after each meal.

Π.

Sodium chlorideav.oz.	10
Antimony sulphideav.oz.	
Buckthorn berriesav.oz.	1
Licorice rootav.oz.	1

Mix all in powder form.

Give one tablespoonful after each meal,

III.

Ammonium chlorideav.oz.	31/2
Antimony sulphidegr.	300
Crude tartar (argols)av.oz.	11/2
Linseed Mealav.oz.	7

Divide into 6 powders and give one of these in a mucilaginous or starchy drink twice a day.

IV.

Mustard,	powderav.o	z. 3
Wheat br	ânav.	z. 24

Stir the mixed powders with sufficient water heated to 70 to 75 degs, C, to make a poultice in the usual manner.

The addition of the mustard makes the poultice somewhat of a counter-irritant, but without being so severe as a strong mustard plaster.

V.

Lead acetate				gr.	45
Sugar					
Mix, reduce	to	powder,	and	divide	into
hree parts.					

One of these portions is to be given with food or in drink three times daily.

VI.

Ammoniac	gr. 120
Ipecac	gr. 120
Squill	
Licorice	gr. 60

Make into a mass with syrup or honey.

VII.

Camphor, powder gr.	180
Potassium chlorate, powderav.oz.	11/2
Belladonna leaves, powderav.oz.	2 2
Anise, powderav.oz.	2′
Mix and divide into 6 powders.	

Mix and divide into 6 powders Give one twice a day in food.

VIII

111.	
Althæa	av.oz. 4
Licorice	.av.oz. 4
Elecampane	.av.oz. 2
Kermes' Mineral	.av.oz. 2
Honeysufficient to form	n a mass
Divide into 12 balls.	

IX.

Ipecac. Squill Licorice	gr.	60
-------------------------	-----	----

Mix into a ball with syrup or honey.

X.

Aconite leaves, powdergr.	360
Digitalis, powdergr.	
Arsenicgr.	
Anise, powder gr.	
Mix, and divide into 6 powders.	
The state of the s	

Give one every night in food.

This remedy is useful in chronic cough.

#### Diarrhœa Remedies.

Diarrhœa is often the result of "catching cold," but may also be the precursor or accompaniment of other disorders. Mild cases may be cured by giving dry fodder and warming the drinks. In severer cases, aromatic and bitter substances may be administered, and in some cases astringents may be required. Applications should be made to the entire abdomen; the whole body should be rubbed with a brush of straw or hay, and then covered with blankets so as to retain the perspiration produced by the rubbing. This rubbing of the entire body should be repeated every 3 hours.

 I.
 Alum ∴
 av.oz. 2

 Calamus root.
 av.oz. 5

 Angelica root.
 av.oz. 5

 Wormwood.
 av.oz. 5

 Rye flour.
 av.oz. 5

 Water.
 sufficient

All the drugs should be in powder and enough water should be added to form a paste.

Place a lump, the size of a hen's egg, on

the tongue every $5$ hours, between meals. This is intended for mild cases only.	The following may also be recommended: Fluid extract of buchu fl.oz. 1
II.	Fluid extract of uva ursifl.oz. 1 Sweet spirit of nitrefl.oz. 2
Iron sulphate, powder av.oz. ½ Althæa, powder av.oz. 3	Sweet spirit of nitre
Water sufficient to form a mass	Tincture of opium fl.oz. 1 Glycerin fl.dr. 12
Divide into two doses, and give these	Water
within 3 hours of each other.	Give 1 or 2 fluidounces at a dose.
This is also intended for mild cases.	It sometimes happens that the animal is
III.	simply afflicted with a slight disturbance of
Oak bark (red or white), pow-	the kidneys, producing what is termed by
der av.oz. 13/4	horsemen and veterinary surgeons, yellow
Alum, powdergr. 150	water; for this ailment a simpler remedy is
Althæa, powder av.oz. 134 Rye flour av.oz. 134	demanded, and the following will prove of
Water sufficient to form a mass or paste	service:
Divide into two parts, and give them 5	Fluid extract of buchufl.oz. 2
hours apart.	Sweet spirit of nitrefl.oz. 4
This is also employed in mild cases.	Potassium nitrateav.oz. 1
	Anise waterfl.oz. 9
IV.	Give one or two tablespoonfuls at a dose,
Iron sulphate, powderav.oz. 1	night and morning, for 3 days, and then in
Alum, powder	the morning only, for one week.
Calamus root	Dyspepsia and Indigestion, Reme-
Rye flourav.oz. 4	dies for.
Water, sufficient to form a mass or pas	I.
Place a piece the size of a hen's egg upon	Sodium chlorideav.oz. 5
the tongue every 2 hours.	Sodium bicarbonateav.oz. 2 Calamus root, powderav.oz. 1
This is intended for severer cases.	Calamus root, powder
V.	Give one tablespoonful with each meal.
Alum, powdergr. 300	II.
Althæa, powderav.oz. 1	Arsenious acidgr. 36
Simple syrup or molassessufficient	Potassium bicarbonateav.oz. 2
Make 2 pills, and give them 2 hours apart.	Wormwoodav.oz. 2
This is employed in mild cases.	Sodium sulphateav.oz. 2
VI.	Mix and divide into 12 powders.
	Give one powder once daily on food.
Tannin	Eyes, Inflammation of the.
Simple syrup, molassessufficient	The eye must be protected from bright
Divide into 3 pills and give 1 pill every	light; it should be bathed three times daily
evening.	with water not too cold and covered with
This is used in the more obstinate cases.	cloths moistened with lead water. In severe
	cases, aloes pills should be given sufficient to
VII.	produce purgation, the cheeks should be
Spirit of mustardfl.oz. 2	rubbed with an irritant ointment, and in-
Oil of turpentine	stead of using lead water, apply the follow-
	ing solutions:
Sprinkle half of this on the belly, rub it in	
and cover with a woolen cloth. Repeat in 5 hours.	Zinc sulphategr. 15
	Distilled water
Diuretics.	Miπ and dissolve.
See Remedies for Urinary Diseases.	Fold a cloth so as to form four thick-
1	

nesses, lay over the inflamed eye and moisten IV. with the solution; repeat the moistening every 2 hours.

#### II.

Zinc sulp	ohategr.	15
	tincture of opiumfl.dr.	
Infusion	of elder flowersfl.oz.	16

Mix and dissolve. Tincture of opium may be substituted for the crocated tincture. The infusion may be prepared from 1 av. ounce of drug; plain water may be substituted, if desired, for the infusion.

This preparation is to be applied like the preceding.

#### III.

Silver nitra	ite	.gr. 5
Distilled w	aterf	l.oz. 2

Mix and dissolve.

Drop 2 or 3 drops into the eye once daily after washing with water.

#### Fever Medicines.

Fever in the horse is the result of other diseases. The normal temperature is 37 to 38 degs. C. and may rise in the febrile condition to 40 or even to 41 degs. C.

Potassium nitrateav.oz.	1
Sodium sulphateav.oz.	10
Rye flourav.oz.	4

Make into a mass or paste with simple syrup, glucose, or molasses.

Give one-half in the morning and the other in the evening.

#### II.

Sodium	salicylateav.oz.	3
Licorice	rootav.oz.	2
Rye flo	ırav.oz.	2

Make into a mass with water.

Give one-half of this mixture one morning and the remainder the following morning.

#### III.

Quinine	sulphate	.av.oz. 1
Althæa.		.av.oz. 3

Make into mass with syrup and divide into 4 pills.

Give the four pills during two consecutive days, one each morning and evening.

This medicine is used during malarial or intermittent fever.

Salol										o	r.	300
Althæa										· or	r.	300

Make a mass with simple syrup or glucose and divide into two pills.

Both pills are to be given at one dose.

This remedy is valuable in rheumatic fever.

Acetaniliagr.	300
Althæaav.oz.	

Form into a mass with simple syrup and divide into two boluses.

Give one bolus in the morning and one in the evening.

These pills are of special value against the fever accompanying influenza, glanders, etc. VI.

Tincture of aconite root	.fl.oz. 1
Tincture of veratrum viride	.fl.oz. 1
Sweet spirit of nitre	.fl.oz. 3
Gin	
Water	.fl.oz. 8

Give one-half to one tablespoonful every four or six hours, until the fever abates.

## Founder, Remedies for.

See Rheumatism.

#### Gall, for Saddle.

Blisters or galls are brought about by badly fitting harness or saddles, which produce local sores very difficult to cure. The sores should be washed two or three times a day with water; the ointment should then be applied 2 or 3 times daily on clean soft cloths.

## T

Zinc oxideav.oz.	
Waterav.oz.	1
Salicylic acidav.oz.	1/2
Mutton tallowav.oz.	21/2
Lardav.oz.	5

#### II.

Lead plasterav.oz. 4
Mutton tallowav.oz. 21/2
Lardav.oz. 3
Salicylic acidav.oz. 1/2

#### III.

Tannin					.av.oz.	1
Camphor, powder.					.av.oz.	$^2$
Zinc oxide						

Mix and sift through a fine sieve.

Sprinkle on the raw or injured surfaces,

after having washed them with tepid water II. and carbolic soap.

#### IV.

Tanninav.oz.	1
Camphor, powderav.oz.	1
Golden seal, powderav.oz.	1/2
Compound tincture of benzoinfl.oz.	6
Glycerin	10

V. A most valuable remedy in veterinary practice for all kinds of sores, bruises, cuts, or whenever the skin is broken is the application of "Friar's Balsam," the compound tincture of benzoin of the U.S.P., and may be profitably put up under some appropriate name and sold by pharmacists.

#### Glanders or Strangles, Remedies for.

This is a rather common disease and may result from contracting cold as well as from infection. Two forms are distinguished, a benign and a malignant form; the remedies mentioned apply only to the former.

The diseased horse is languid, perspires easily, has diminished appetite, and coughs. From the inflamed nostrils flows a discharge, watery at first, later becoming thick and mucous. When the secretion assumes the latter character, a swelling appears in the throat, which interferes with mastication. This swelling will become purulent, break open and discharge, and subsequently will heal, after which the horse appears quite well.

Treatment consists in keeping the animal warm by covering with a woolen blanket, also binding a cloth about the throat. Internally give mild cathartics, and assist the process of the swelling by the application of suitable ointments or even by means of poultices.

#### I.

Antimony sulphideav.oz. 1
Sulphurav.oz. 1
Fennel seedav.oz. 1
Calamus rootav.oz. 1
Juniper berriesav.oz. 2
Rye flourav.oz. 2
Oil of turpentine
Watersufficient to make a confection

Give a mass the size of a duck's egg four

Antimony sulphide	av.oz. 2;	12
Ammonium chloride	áv.oz. 3	-
Sulphur		
Juniper berries		
Sodium sulphate	av.oz. 9	
Rye flour.		
Watersufficient to make	a confection	

Give a mass the size of a duck's egg every

#### III.

Antimony sulphideav.oz.	1
Sodium sulphateav.oz.	5
Juniper berriesav.oz.	
Give one tablespoonful with each meal.	

#### IV.

Antimony sulphideav.oz.	1
Sulphurav.oz.	1
Sodium sulphateav.oz. 1	10
Licorice rootav.oz.	4
Buckthorn berries, crushedav.oz.	4
Mix one tablespoonful with each meal	

Buckthorn berries, coarse powder
av.oz. 2
Anise seedav.oz. 2
Sodium chlorideav.oz. 5
Sodium bicarbonateav.oz. 1
Strew two teaspoonfuls on each meal.
This is intended for mild cases.

Linseed meal	av.oz.	2
Chamomile, coarse powdera		
Wheat bran	av.oz.	6

Mix the above with hot soapsuds, and cover the swelling of the neck with the poultice.

#### VII.

Ammonia	linime	ent		fl.oz. 4
Oil of turp	pentin	e		fl.oz. 4
Rub the	neck	with	this	liniment three

times a day.

#### VIII.

I	Mercurial	ointme	nt		av.oz	. 3
(	reen soa	p			av.oz	. 3
(	lycerin.				fl.oz	. 5
Ţ	Sub the	neck tw	ice a	day w	with this	lin

ment.

#### IX.

Ammonium carbonategr.	150
Carbolic acidfl.dr. Oil of turpentinefl.dr.	11/
Oil of turpentinefl.dr.	21/2
Waterfl.oz.	21/2
Put about seven ounces of havseed	into a

This is recommended for obstinate cases. basin, pour hot water into it, and to this add

the above ingredients; cover the horse's head with a cloth, and stir its contents thoroughly, so as to facilitate the liberation of the steam. Keep this up for a quarter of an hour. Repeat the treatment once each day.

This is employed to promote the nasal secretion.

#### Heave Medicines.

[.		
	Sodium sulphateav.oz.	10
	Elecampaneav.oz.	10
	Lobeliaav.oz.	10
	Resin weedav.oz.	10
	Gentianav.oz.	
	Blood rootav.oz.	
	Tartar emeticav.oz.	1
	Alumav.oz.	5
	Fenugreekav.oz.	5
	Linseedav.oz.	15
1	Ţ.	

<del></del>	
Resin weedav.oz.	10
Lobeliaav.oz.	10
Elecampane av.oz.	10
Sodium sulphateav.oz.	10
Linseed av.oz.	
Fenugreekav.oz.	5
Alumav.oz.	
Coniumav.oz.	
Bloodrootav.oz.	3
Gentianav.oz.	3
Tartar emeticav.oz.	1

III.
Lobeliaav.oz. 2
Skunk cabbageav.oz. 4
Elecampaneav.oz. 4
Tartar emeticav.oz. 1
Licorice root av oz 5

This may be diluted if desired with linseed meal, fenugreek, or other drugs.

#### IV.

Balsam of firav.oz. 4
Balsam of copaibaav.oz. 4
Calcined magnesiasufficient

Mix the first two ingredients and add magnesia until the mixture is sufficiently thick to form into balls.

Give a medium sized ball night and morning for a week or ten days.

#### Heels, Ointment for Cracked.

Sulphurav.oz.	1
Lead acetate, powderav.oz.	1/2
Creolin fl.dr.	4
Oil of eucalyptusfl.dr.	4
Petrolatumav.oz.	4
Lanolinav.oz.	
Apply twice daily.	

#### Hoofs, Preparations for the.

Aside from the appearance which it gives a horse's hoofs, the occasional use of a good hoof-dressing really promotes the comfort of the animal and prevents brittleness of the hoof. The following are simple and useful formulas for this purpose:

I.		
	Oil of origanumfl.oz.	1
	Camphorav.oz.	1
	Lardav.oz.	16
	Apply twice every week or two.	

II.	
Tarav.oz	z. 8
Tallowav.ox	z. 8
Use like the preceding.	

111.	
Petrolatumav.oz.	16
Carbolic acid, crystalgr.	180
Camphorgr.	90
Oil of tarfl.dr.	
Oil of origanumfl.dr.	1
IV.	

Should there be any disease of the hoof as hoof-bound, etc., the following ointment will produce satisfactory results:

Ca	amphorav.oz.	1
Ba	alsam of firav.oz.	1
Oi	il of cajuputfl.dr.	21/2
	ompound tincture of iodinefl.dr.	
O	il of turpentinefl.oz.	1
	ardav.oz.	
5.7		,-

The following cement is useful for cracked hoofs:

100131		
Ammoniac	.av.oz.	3
Gum turpentine	.av.oz.	1
Gutta percha	.av.oz.	6

The first two are melted together by means of a water bath, then added to gutta percha, previously melted, with constant agitation. If a black color is desired lampblack may be added.

When used the cement should be softened in hot water and pressed in the hoot crack, which has previously been well cleaned.

#### Indigestion, Remedies for.

See Remedies for Dyspepsia and Indigestion.

#### Influenza Remedies.

The early symptoms of this disease are languor and loss of appetite. After one or

two days there is fever and an increased pulse-rate; the thirst increases while the appetite diminishes. Internal and external remedies are used, the latter being usually of an irritant character.

Τ

Ammonium chlorideav.oz.	3
Potassium nitrateav.oz.	3
Sodium sulphateav.oz.	10
Licorice rootav.oz.	61/2

All should be in fine powder and well mixed.

Give one tablespoonful in warm soft food three times a day.

II.

Camphor, powdergr.	75
Potassium nitrategr.	300
Aloesav.oz.	1
Linseed mealgr.	
Water, sufficient to make two boluse	S

Give one pill every three hours.

III.

Alumgr. 150
Tanningr. 40
Licorice, powdergr. 200
Simple syrup, sufficient to make bolus

Give one pill every five hours.

This is used for the diarrhœa which often manifests itself.

IV.

Aloesgr. 300	
Dried sodium sulphateav.oz. 33	1/2
Linseed mealav.oz. 1	1/2
Soft soap (green soap)gr. 300	
Simple syrup, enough to form an electuar	y

Give in two doses with an interval of two hours.

This is used as a purgative if one be required.

v.

Oil of turpentinefl.oz.	3
Spirit of camphorfl.oz.	

Sprinkle the abdomen, rub with a wisp of straw, and bind a warm wrap about the abdomen.

VI.

Chlorodyne fl.oz.	1
Spirit of nitrous etherfl.oz.	
Solution of ammonium acetatefl. oz.	2
Water	

This dose is to be given every three hours during the first stage when there is much shivering.

#### Liniments.

These preparations may be dispensed under such names as "Horse Liniment," "Veterinary Liniment," "Barbed Wire Liniment," "Stable Liniment," etc.

Many of the liniments in Part II may also be used as veterinary liniments.

I.

Oil of turpentinefl.oz.	15
Carbolic acid, crudefl.oz.	2
Benzine fl.oz.	15
Oil of tarfl.dr.	4
Oil of spikefl.dr.	4
Camphorgr.	120
Capsicumgr.	

H.

L.	
Camphor av.oz.	1
Carbolic acidfl.oz.	1
Oil of origanum fl.oz.	2
Oil of tarfl.oz.	
Crude petroleumfl.oz.	2
Oil of turpentinefl.oz.	
Liquid petrolatumfl.oz.	
Benzinefl.oz.	

III.

Oil of turpentinefl.oz. 1	6
Camphorav.oz.	1
Soap (soft or green)av.oz.	
Water sufficien	ıt

Mix the soap with about 2 fluidounces of water, dissolve the camphor in the turpentine, mix the two and reduce to the desired consistence by the addition of water.

IV.

Oil of turpentine	fl. oz. 4
Stronger water of ammonia	
Olive oil	fl oz 4

This preparation is used as a throat liniment.

## Mange or Scabies Remedies.

Scabies is most apt to affect old or ill-fed horses and generally appears on the side of the neck, on the shoulder, the back, the hips, at the root of the tail and on the feet.

In all cases the first thing to be done is to wash the affected parts with a warm solution of green soap (medicinal soft soap). The animal must also be separated from the others, and special care taken in feeding and grooming it.

I.

Mercurial ointmentav.oz.	3
Salicylic acidav.oz. Lardav.oz.	1/2
Lardav.oz.	$6\frac{1}{2}$

Wash the affected parts with a solution of

green soap; dry, and apply this ointment once a day.

This ointment is recommended for foot scabies.

#### IT.

Creosotefl.oz.	2
Green soapav.oz.	10
Alcoholfl.oz.	

Mix and apply to the affected parts after washing and drying them.

#### III.

Sulphurated potassaav.oz. Green soapav.oz.	3/4
Green soapav.oz.	$1\frac{1}{2}$
Waterfl.oz.	14
Oil of turpentinefl.dr.	$1\frac{1}{2}$

Dissolve the soap and the potassa in the water by the aid of heat, and then add the oil of turpentine.

Wash the affected parts with a weak soda solution, dry with a cloth, and then moisten with the above wash. Repeat this twice each day.

#### IV.

Pine tarav.oz.	10
Green soapav.oz.	5
Alcoholav.oz.	3
Sulphurav.oz.	2
Mix with the aid of a gentle heat.	

Wash the affected parts with warm soap and water, dry with a cloth and spread this salve on with a brush, repeating the application in eight days. As a rule two applications suffice. The salve or paint eventually falls off of itself.

#### Mosquito Oil.

Carbolic acid	.fl.oz. 2
Oil of pennyroyal	.fl.oz. 4
Spirit of camphor	
Oil of tar	
Glycerin	.fl.oz. 4
Lard oil	8.xo.fl.

This is an effective preparation for keeping flies and mosquitoes off horses.

#### Ointments.

Ointments are a class of very useful remedies for the treatment of some of the diseases of stock. Every pharmacist should have on hand, ready for sale, a good veterinary ointment.

I.	
Compound tincture of iodinefl.oz.	2
Camphor av.oz.	
Oil of sassafrasfl.oz.	1
Lard. av oz	15

#### II.

Camphor av.oz.	3
Tanninav.oz.	1/2
Carbolic acid, crystalav.oz.	1
Oil of origanumfl.oz.	1
Zinc oxideav.oz.	3
Lardav.oz.	16

#### III.

Some of the ointments mentioned under "Preparations for the Hoof" and "Remedies for Galls" may also be recommended for general use.

#### Quitter Oil.

Ointment of nitrate of mercury.av.oz.	1
Cottonseed oilfl.oz.	3
Mix together by aid of a gentle heat.	٠

#### Rheumatism and Rheumatic Inflam mation of the Feet (Founder), Remedies for.

This disorder of horses is very common. Usually the soft parts of the hoof are affected; the animal then steps with exceeding caution; it lies down mostly when in the stall, and groans frequently; the pulse-rate is increased, and sometimes there is fever.

Purgative remedies should be given, either by the month or by enema, and care should be taken that there is a normal flow of urine. Stimulant applications should be made externally. Only half rations should be allowed the sick animal, and the food should be of an easily digestible character.

# I. Camphor, powder......av.oz. ½ Potassium nitrate, powder...av.oz. 2 Sodium sulphate, powder...av.oz. 8 Juniper berries, crushed...av.oz. 4 Rye flour...av.oz. 3½

Water......sufficient
Make into an electuary and give one-fourth
of the whole every five hours.

#### TT.

Aloes, powderav.oz.	1
Sodium sulphate, powderav.oz.	8
Rye flourav.oz.	
Watersufficier	

Make into an electuary, divide into two parts and give three hours apart. This is used when there is fever.

#### III.

Ammonia liniment	.fl oz.	10
Oil of turpentine		
Rub in well twice daily		

IV.	
Spirit of camphorfl.oz. 12	,
Tincture of capsicumfl.oz. 1	
Oil of turpentinefl.oz.	3/4
Rub on the lame foot three times a day.	
V.	
Danusia asid	1/

Benzoic	acidav.c	oz. 3½
Sodium	sulphate, powderav.o	z. 13/4
Rye flot	urav.c	oz. 3½
Water		ient

Make into an electuary and give onequarter of the whole every twelve hours. This is to be given when there is fever.

#### Shoulder, for Lame.

Spirit of soapfl.oz.	5
Spirit of camphorfl.oz.	5
Ammonia waterfl.oz.	1

Apply several times daily. See also No. V under "Rheumatism."

If the lameness be of a rheumatic character sodium salicylate should be given.

#### Sinews, for Strained.

T.

Ammonium chloride	.fl.dr.	2
Spirit of camphor		
Diluted acetic acid		
Water		

Mix and apply on a cloth morning and evening for at least eight days.

IT.

Tincture of capsicumfl.oz.	3
Alcoholfl.oz.	4
Spirit of camphorfl.oz.	$\dot{2}$
Spirit of ether	$\tilde{2}$
Oil of turpentinefl.dr.	11/2
Ammonia waterfl.dr.	3
Ammonium chloride av.oz.	1
Sodium chloridegr. I	180
	7

Dissolve the salts in the water and add the remaining ingredients.

Shake up the embrocation well. Dilute half a pint of the fluid with pint and a half of water and with this wash the entire length of the leg and then wrap it up in a woolen bandage.

#### III.

Spirit of ammonia	.fl.oz. 5
Spirit of camphor	.fl.oz. 5
Spirit of ether	fl.oz. 5
Oil of turpentine	
on or imperiment the transfer	

Use like the preceding.

Υ	12	
1	v	٠

Sodium chlorideav.oz.	1
Spirit of camphorfl.oz.	1
Tincture of arnicafl.oz.	2
Water fl.oz.	
Use this like the preceding.	

## Spavin and Ringbone Cures.

Τ.

١.		
	Oil of turpentinefl.oz.	8
	Alcohol	8
	Tincture of iodinefl.oz.	8
	Camphorav.oz.	4
	Crude petroleumfl.oz.	1
	Oil of rosemaryfl.dr.	4
	•	

TT

1.	
Oil of turpentinefl.oz.	141/2
Alcohol fl.oz.	13
Tincture of iodinefl.oz.	13/
Camphor av.oz.	13/
Oil of sassafras fl.oz.	13/
Crude petroleumfl.oz.	1
Corrosive sublimategr.	70

Dissolve the camphor and the corrosive sublimate in the alcohol and add the remaining ingredients.

III.

Venice turpentineav.oz.	2
Cantharides, powderav.oz.	2
Euphorbium, powderav.oz.	1
Red precipitateav.oz.	1
Lardav.oz.	24

Mix all, simmer slowly for an hour, avoiding burning or scorching, and pour off the liquid from the sediment.

For ringbone cut off the hair and rub the ointment well into the lumps once in 48 hours. For spavins, apply once in 24 hours for three mornings. Wash well, previous to each application, with clean soapsuds, rubbing over the places with a smooth stick to remove the thick yellow matter.

TV

• •	
Cantharides, powderav.oz. 4	
Oil of origanumfl.oz. 4	
Oil of amberfl.oz. 4	
Oil of turpentinefl.oz. 4	
Cottonseed oilfl.oz. 4	
Sulphuric acidfl.oz. 1	

Mix all except the acid in a broad, shallow vessel such as an evaporating dish, then very slowly and with constant and rapid stirring add the acid: when the liquid has cooled it may be bottled.

For ringbone or spavin apply with a sponge tied to the end of a stick, until it is

no longer absorbed into the parts; 24 hours afterwards grease well with lard, and in 24 hours more wash off with clean, warm soapsuds. Two or three applications 3 or 4 days apart may be required.

V.

Mercurial ointment	.av.oz. 3
Salicylic acid	.av.oz. 1
Lard	.av.oz. 6

Rub the spavin daily with the ointment for

This is used only for mild cases.

#### VI.

Cantharides	av.oz. 2
Euphorbium	av.oz. 1
Salicylic acid	av.oz. 1
Gum turpentine	.av.02. 3
Lard	av.oz. 2
Cottonseed oil	.fl.oz. 1

Mix, heat for one hour at a temperature of 50 to 70 degs. C., and stir constantly until cool.

Apply once daily for three consecutive days.

## Spavin, for Foot.

Green soap	av.oz.	4
Ammonia water		
Petroleum	fl.dr.	5
Tincture of cantharides	fl.dr.	5

Rub daily for one or two days, then omit for two days and repeat the application. Continue this intermittent application until the ointment is all used.

#### Stomach, for Catarrh of the.

T.

Sodium bicarbonate	av.oz. 4
Sodium chloride	
Sodium sulphate	
Juniper berries	av.oz. 2

Reduce to coarse powder and mix well. Give one tablespoonful with each meal. This is employed only in chronic cases.

II.

Artificial	Carlsbad	salt	av.oz. 8
Oil cake	meal		av.oz. 8

Two tablespoonfuls are given three times daily in about one and a half gallons of warm water.

This, like the preceding, is used only in chronic cases.

#### III.

Precipitated		
Sodium chle	oride	 .av.oz. 3
Gentian		 .av.oz. 1
Fennel		 .av.oz. 1

Reduce all to powder and mix well.

One tablespoonful is to be given with each

This mixture is directed against acute cases.

## Thrush, for Running.

Wash the hoof daily with soap water, then apply the following:

I.

Chlorinated limeav.oz. Waterfl.oz.	11/2
Water	16

First wash the affected parts with a solution of soap, rinse with clear water, and then wash with a solution of chlorinated lime previously warmed. Then press some oakum, first moistened with the solution, into the opening. Repeat the application once daily.

#### II.

Alumav.oz.	1
Copper sulphateav.oz.	
Carbolic acid fl.dr.	11/2
Waterfl.oz.	

Use like the preceding.

#### III.

Alumav oz	. 1
Copper sulphateav.oz	
Pyroligneous acidfl.oz	
Water fl.oz	

Use like the preceding.

IV.

Salicylic	acidgr.	120
Glycerin.	fl.oz.	1
	of aloesfl.oz.	
Tincture	of gallsfl.oz.	5

Wash off the hoof with warm soap and water and paint the thrush with the liquid. Then moisten some oakum with the tincture and press it into the crevices between the swellings. Apply once a day.

#### Urinary Diseases, Remedies for.

Urinary diseases are usually manifested by conditions known as polyuria or the voiding of a large amount of pale urine, and anuria or retention of urine The former is usually caused by feeding with decomposed fodder.

I.	For	po	lyuria:
----	-----	----	---------

Camphor, powderav.oz.	1/2
Gingerav.oz.	11/2
Rye flourav.oz.	
Watersufficient to form a ma	SS

Give one-eighth of this mass mornings and evenings. If the disease still lingers on the fifth day, the following is recommended:

Camphor av.oz.	1/2
Alumav.oz. 1	
Dippel's oilfl.oz. 1	
White oak barkav.oz. 1	
Angelica rootav.oz. 3	,
Rye flourav.oz. 1	1/2
Watersufficient	

Reduce all the drugs to powder and add enough water to form a mass or paste.

Give a piece the size of a hen's egg three times daily spread upon the tongue.

#### II. For anuria:

Juniper berries,	crushedav.oz.	5
	av.oz.	

Make an infusion by pouring on 6 pints of hot water, allow to stand 15 minutes, and strain, expressing the residue in the strainer.

Administer one-third of this infusion and use the remainder as an enema.

#### III. See also Diuretic Remedies.

#### Worms, Treatment for.

Poorly-fed animals are more subject to these parasites than well-fed animals. Treatment consists in giving worm medicines and good food.

I.		
	Oil of tansyfl.dr.	4
	Crude petroleumfl.dr.	
	Wormwood, powderav.oz.	
	Asafetida, powderav.oz.	1/2
	Aloesav.oz.	
	Rye flourav.oz.	2
	Water sufficient to form a mass or n	aste

Every two hours a piece of the size of a hen's egg is to be spread upon the tongue.

L.	
Tartar emeticav.oz.	
Asafetidaav.oz.	5
Gingerav.oz.	4
American wormseedav.oz.	
Saltav.oz.	
Fenugreekav.oz. 1	6
Mustardav.oz. 1	
Poplar barkav.oz. 5	
Corn mealav.oz. 2	

Two tablespoonfuls are to be given each morning before feeding until four doses have been given.

#### III.

Tartar	em	etic		 		 	.av.oz.	1
Aloes.				 		 	.av.oz.	2
Ginger				 	٠.	 	.av.oz.	2
Potassi	um	niti	ate	 ٠.		 	.av.oz.	2

Reduce all to powder and mix well.

Give one teaspoonful twice daily in food. IV.

Crude petroleum (black oil)fl.oz	. 2
Dippel's oilfl.oz	
Levant wormseed, powderav.oz	
Valerianav.oz	. 5

Reduce the drugs to powder and mix with the other ingredients.

A bolus about the size of a hen's egg should be given every two hours.

#### SECTION II—CATTLE MEDICINES.

The doses in the following formulas, when not otherwise specified, are intended for a full grown animal; consequently when the medicines are to be administered to young, small or weak animals a corresponding reduction must be made in the size of the dose, while for unusually large and strong cattle the doses may be increased; for young cattle the following will serve as a general guide to the sedative dose required at different ages:

At one year 25 per cent of the adult dose. At two years 50 per cent of the adult dose. At three to four years 75 per cent of the

adult dose.

## Appetite, Loss of, Remedies for

## I.

Gentian, po	wderav.oz.	1
Magnesium	sulphateav.oz.	10

Mix with one quart of warm water and give for one dose.

#### IT.

Sodium	sulphateav.oz.	3
Sodium	chlorideav.oz.	8
Linseed	mealav.oz.	1

Mix with one quart of lukewarm water and give for one dose.

#### III.

Sodium bio	arbonate.	 		 	.gr.	300
Rhubarb,	powder	 	٠.	 	.gr.	75

Divide into two doses, and give each dose in a cup of chamomile tea or other suitable vehicle.

#### Bowels, Remedies for Inflammation II. of the.

See Remedies for Inflammation of the Stomach and Bowels.

#### Colic Remedies.

Colic is usually the result of too great indulgence in indigestible food; if not properly attended to may cause death. As a rule, the animal is constipated, is very thirsty, groans, etc. If there is no movement of the bowels within three days, the affection is likely to prove fatal.

Treatment consists in evacuating the bowels by means of purgatives and enemas and subsequently giving stomachic medicines.

1	•		
4	•	۰	

Green soapav.oz.	4
Sodium chlorideav.oz.	1
Linseed oilfl.oz.	3
Water	S

Give one injection as above every hour until the bowels act.

#### TT

Magnesium sulphateav.oz.	8
Linseed oilfl.oz.	
Chamomile infusion (1 to 20) flor	

Give one quart every four hours, until all is given, or until the bowels act.

#### III.

Gingergr. 150
Mustardgr. 150
Gentiangr. 150
Reduce all to powder and mix well.

Give one such powder in a pint of warm water when the bowels have acted and the colic has passed off.

#### Constipation Remedies.

Constipation may result from unsuitable food, but may also be the accompaniment of other diseases.

Treatment consists in giving purgatives and enemas.

Ì.

Antimony sulphide gr.	180
Argols	3/4
Aloesav.oz.	1
Sodium sulphateav.oz.	18

All should be in powder and should be well mixed.

One-fourth of this mixture should be administered every 3 hours in warm chamomile tea or other suitable vehicle.

Aloesav.oz.	1
Linseed mealav.oz.	2
Sodium sulphateav.oz.	25

All should be in powder and be well mixed.

Dissolve this powder in one quart hot water and allow to cool somewhat before administering.

#### III.

Aloes, powdergr	. 30	00
Linseed oilfl.oz		16
Mix well, warm and administer	at	one

#### dose. IV.

Rochelle saltav.oz.	2
Aloes, powder gr.	150
Linseed mealgr.	150

Give this mixture, which is suitable for calves only, at one dose in 8 fluidounces of warm water.

Green	soapav.oz.	31/2
Water	soapav.oz.	2

Give as an enema every hour until evacuation of the bowels ensues.

Eserine	sulphategr. waterfl.dr.	21/2
Distilled	waterfl.dr.	11/4
Mix and		

This may be injected subcutaneously to produce an evacuation of the bowels.

#### Diarrhœa, Remedies for.

Diarrhœa may arise from a cold or may be merely an accompaniment of other diseases.

The remedies used are usually efficient, but it is also necessary to cause the animal to perspire by rubbing and then to cover it warmly. It is also essential to put the sick animal in a warm, comfortable and rocmy

The food should be restricted to a little dry feed (good hay, etc.), no green food being permissible and in place of cold water, warm, starchy, or mucilaginous water should be given.

Nutgalls, powder.....av.oz. 11/2 Licorice root, powder.....av.oz. 1½

Give in two doses with an interval of two hours, mixing each dose in a pint of warm water.

II.	
Alum, powderav.oz.	1/2
Oak bark (white or red) powder av oz.	$2^{'}$

Divide into two doses and administer each dose in one pint of warm water with an interval of four hours between the doses.

#### TIT.

Opium, powdergr.	150
Peppermint, powderav.oz.	3,4
Linseed mealav.oz.	1

Give one-half in the morning, the remainder in the evening, in one pint of warm water.

#### Diarrhœa in Sucking Calves, Treatment for.

Wrap up the animal warmly and administer internal medicines as below and also use suppositories to avoid irritating the bowels.

#### I.

Precipitated chalkav.oz.	1/2
Alum, powderav.oz.	1/2
Rye flourav.oz.	
Yolk of eggssufficient	

Make into five pills. Give one pill every five hours.

#### TT.

Alum, powdergr. 20
Salicylic acidgr. 20
Rye flourav.oz. 34
Watersufficient
Make this mixture into five pills.

One pill should be given every five hours.

#### III.

Hydrochloric	acid	fl.dr.	11/4
	fusion (1 to 20)		

In cases of abnormally acid stomach, give the above in two doses with an interval of five hours.

#### IV.

Tanningr.	15
Mutton tallowgr.	45
Cacao buttergr.	135

Mix and form into four suppositories; insert one morning and evening after a passage of the bowels, inserting it as far up as can be done with the oiled finger.

#### Eye, Remedies for Inflammation of.

Inflammation of the eye is usually of a catarrhal character caused by the catching of 'cold.' The eye first appears reddened, then tears begin to flow, and a mucous subday.

stance appears which has a tendency to stick the eyelids together.

Treatment consists in washing with warm milk and applying this solution every hour: Solution of lead subacetate.....fl.dr. 3

Distilled water enough to make..fl.oz. 8

#### Fever Medicines.

	I.
	Potassium nitrateav.oz. 3/4
	Sodium sulphateav.oz. 8
•	Give one-half of the above in one quart of
L	warm bran water at night.

#### II.

Salicylic acidav.oz.	3/1
Sodium bicarbonateav.oz.	3/4 1/2
Magnesium sulphateav.oz.	10
Use like the preceding.	

#### Indigestion Remedies.

The treatment for indigestion is the same as for catarrh of the stomach; hence see Treatment for Catarrh of the Stomach.

#### Jaundice, Treatment for.

Jaundice may emanate from the liver or may result from intestinal catarrh. As in the human subject, it is manifested by yellow discoloration of the mucous membrane of the mouth, of the white of the eye, etc. The urine becomes dark and the feces light colored; the appetite is diminished.

Treatment consists in giving calomel and sodium sulphate and applying irritant ointments in the region of the liver. If no benefit results from this treatment, then the sodium sulphate should be given with aloes, rhubarb and juniper berries.

## I.

٠.					
	Sodium	bicarbor	nate	av.oz.	13/
	Sodium	sulphate		av.oz.	10
	Juniper	berries,	crushed	av.oz.	13/
	Water .			fl.oz.	$64^{'}$

Give half of the above morning and even-

## ing.

5
5
10
10
10

Reduce all to powder and mix well.

Give one heaping tablespoonful in a quart of infusion of juniper berries three times a day.

#### Lice, Treatment for.

Lice are most apt to trouble young animals, only troubling old ones when they are very dirty.

I.

Green soapav.oz.	10
Alcohol	9
Wood alcoholfl.oz.	1
Crude naphthalinav.oz.	2
Water	40

Heat the whole together until dissolved and then stir until cold.

Rub the places infested by lice thoroughly with the above, and wash off the next day with warm soda solution. When the animal is again dry repeat this operation twice. The lice generally die after the second application.

II.

Raw tobacco (not manufactured)
.....av.oz. 5

Pour the boiling water on the tobacco, allow to stand half an hour, strain and add the alcohol.

Moisten the parts infested with lice, and wash off on the following day with warm soda solution. Repeat the operation three or four times as may be necessary.

#### Milk Secretion, Treatment for Diminished.

When diminished milk secretion is not due to age or to disease, the following mixtures will prove useful. They have been dispensed under such names as "Cow Powder," "Milk Powder," and "Pulvis Vaccarum."

т

Carawayav.oz. 4
Calamusav.oz. 4
Sodium chlorideav.oz. 2
Sulphurav.oz. 1
Reduce all to powder and mix well.

Give two heaping tablespoonfuls twice daily in one quart of warm beer.

II.

L,	
Aniseav.oz. 2	
Fennelav.oz. 2	
Antimony sulphideav.oz. 4	
Sodium chlorideav.oz. 4	

All should be in powder and well mixed.

This mixture should be used like the preceding.

III.

Black antimonyav.oz.	2
Sulphurav.oz.	2
Fennelav.oz.	1
Carawayav.oz.	1
Juniper berriesav.oz.	
Sodium chlorideav.oz.	

All should be in powder and well mixed.

This mixture should be used like the pre-

IV.

Potassium nitrateav.oz.	1
Alumav.oz.	1
Sulphurav.oz.	1
Prepared chalkav.oz.	
Aniseav.oz. 1	
Fennelav.oz.	5
Buckbean herbav.oz.	5
Sodium chlorideav.oz. 1	0

All should be in powder and should be well mixed.

Give one or two handfuls with the morning meal.—H.

# Milk, Treatment for Speedy Souring of.

There are conditions of cows when the milk soon becomes sour, in spite of the utmost cleanliness. The following mixture may remove the difficulty:

Т

٦.		
	Sodium bicarbonateav.oz.	31/2
	Precipitated chalkav.oz.	31/2
	Fennelav.oz.	7
	Linseed meal av.oz.	7

All should be in powder and should be well mixed.

Give one-half in one quart of warm water, administering the remainder the next day.

TT

1 1	••		
	Fennel	av.oz. 4	
	Linseed meal		
	Sodium chloride	av.oz. 4	
	Prepared chalk		

All should be in powder and should be well mixed.

Two heaping tablespoonfuls should be given twice daily in one pint of warm water.

This mixture should be employed in obstinate cases.

#### Ringworm, Treatment for

See Treatment for Tetter.

#### Rheumatism Remedies.

I.

Ammonium chlorideav.oz.	2
Potassium nitrateav.oz.	2
Sodium sulphateav.oz.	
boaram sarphate	

Reduce to powder, mix, and divide into four portions.

This is used as a purgative, one powder being given every three hours in a quart of warm water.

II.

When there is fever, sodium salicylate in about 1-ounce doses may be administered every three hours in a pint of warm water.

III.

When there is no fever the following may be recommended:

Arnica flowersav.oz. 3	
Juniper berries, crushedav.oz. 3	1/2
Ammonium chlorideav.oz. 1	
Aloes	
Water, boilingpints 7	
,	

Pour the water upon the arnica and juniper, let stand one-half hour, strain, and in the colature dissolve the remaining ingre-

Warm one quart of this mixture and give every five hours.

IV.

O:1 of	turmontino A du C	
On or	turpentinefl.dr. 6	
Spirit	of camphorfl.oz. 6	

This mixture should be applied to the swollen joints every 6 hours.

#### Shoulder Lameness, Treatment for.

Water of ammonia	.fl.oz.	5
Oil of turpentine		
Spirit of camphor	.fl.oz.	10
Spirit of soap	.fl.oz.	10

Rub well into the lame shoulder three times daily.

## Stomach, Treatment for Catarrh of

Catarrh of the stomach is generally produced by irregular feeding or indigestible food, without taking sufficient exercise, or from eating large quantities of non-nutritious

The treatment consists in the administration of mild laxatives combined with bitter tonics, and of hydrochloric acid.

fed only with easily digestible food, such as flour or bran water.

Antimony sulphidegr.	300
Argols (crude tartar)av.oz.	
Wormwood av.oz.	2
Sodium sulphateav.oz.	15

Give one-fourth of the whole every four hours in a quart of warm water.

Aloes, powderav.oz. 1	
Sodium chlorideav.oz. 3	
Linseed mealav.oz. 4	

Give one-half in the morning and the other half at night, dissolved in one pint of warm water.

III.

Hydrochloric acid	fl.dr.	4
Linseed meal	.av.oz.	3
Water	av.oz.	331/3
Give one-half at night and th		
in the morning.		

This is employed in obstinate cases.

IV.

Aloes, powdergr.	300
Calamus, powderav.oz.	
Argols (crude tartar)gr.	300
Antimony sulphide, powdergr.	150
Linseed mealav.oz.	2
Waterfl.oz.	32

Give one-half at night and the remainder in the morning, first warming the mixture.

This is employed in chronic indigestion.

Calamusav.oz.	1
Linseed mealav.oz.	
Antimony sulphideav.oz.	1/2
Water fl.oz.	32 -

Give two-thirds of the mixture, first warming it, administering the remainder after an interval of four hours.

This is also employed in obstinate cases.

#### Stomach and Bowels, Remedies for Inflammation of the.

It is generally presumed that inflammation of the stomach and bowels is the result either of a cold or of eating some poisonous weed.

The affection is manifested by loss of both appetite and thirst, restlessness, swollen abdomen, constipation, etc.

Warm mashes should be given as food and warm linseed meal water be given in place of During the illness the animal should be cold water to drink. Purgatives containing oil should be given, also enemas, and the II. belly should be rubbed with stimulant applica-

The following may also be administered:

I.

Infusion of chamomile (1 in 10).fl.oz.	32
Sodium sulphateav.oz.	
Linseed oilfl.oz.	26
Salicylic acidgr.	90

Administer one pint every hour until constipation is overcome.

TI.

Infusion of chamomilefl.	oz. 32
Linseed mealav.	oz. 7
Linseed oilfl.	oz. 32

Give one pint every 2 hours after the bowels have acted.

III.

Solution of soapfl.oz.	32
Common saltav.oz.	1 1/2
Linseed oilfl.oz.	3

Give this mixture as an injection every hour until the bowels are relieved.

IV.

Linseed oilfl.oz.	4
Ammonia waterfl.oz.	4
Oil of turpentinefl.oz.	4
Rub the belly with the above every 3 ho	ours

## Tetter or Ringworm, Treatment for.

Tetter usually attacks old, underfed animals where stalls are poorly ventilated and unclean.

The stalls must first of all be thoroughly cleaned, aired and whitewashed with lime, and the affected parts of the animal treated with the following:

Green soapav.oz. 10	)
Water	)
Pine tarav.oz.	,
Heat in a water bath until of a unifo	rm

consistency.

Wash the entire body of the animal every 2 days with warm soft or green soap solution (1 to 20), rinse with warm water and when the animal is dry rub the above ointment well in under the hair.

#### Throat, Inflammation of.

Τ.

Ammonia liniment	fl.oz.	8
Oil of turpentine	fl.oz.	8
Rub the throat three times daily	with	this
mixture and then bind with flannel.		

Alumgr.	375
Salicylic acidgr.	
Honeyav.oz.	1
Diluted acetic acidfl.oz.	2
Waterfl.oz.	32

Mix and dissolve.

Warm the solution and inject into the mouth every half hour. Or a linen cloth moistened with this solution may be employed to wash out the mouth.

# Udder, Treatment for Inflammation

Inflamed udder may result from injury or from cold, or as a result of other diseases. In the first stages, the milk appears normal, subsequently becoming thick and even purulent or bloody.

Treatment consists in gently evacuating the udder mornings and evenings, and limiting the animal to half rations, which should consist of easily digestible food. The udder should be bathed in warm water after milking, and after drying an antiseptic application should be made. If the udder is hot and feverish, the application should contain mercurial ointment. Saline purgatives should be administered internally.

Sa	licylic	acid	l	 	 	.gr.	60
	mpho						
_							

Rubber the udder carefully twice daily.

Salicylic acidgr.	40
Mercurial ointmentav.oz.	1
Liniment of camphorfl.oz.	
Apply like the preceding.	

III.

Potassium	nitrate.					.av.oz.	2
Sodium su	lphate					.av.oz.	20

Give one-third of the above at morning, at noon and at night in one quart of chamomile infusion or other suitable vehicle.

#### Urine, Treatment for Bloody.

This is generally brought about by eating sour food such as oxalis, either fresh or dried, which is generally the cause to be looked for.

Treatment consists in changing the food. If the disease has been contracted while grazing, change to hay food, or if it has been contracted from dry, place the animal at pasture or give it green food. If the change of food alone proves insufficient then use the following:

White leadgr.	45
Sodium acetategr.	150
Camphor, powdergr.	180
Precipitated chalkav.oz.	4

Mix and divide into 6 powders.

Give one powder in a quart of bran mash morning and evening.

#### Worms, Treatment for.

Worms are readily produced by insufficient feeding, and are banished by purgative worm medicines. The important point is to give the animal but little food on the day previous, and to administer the worm medicine and the purgative at the same time.

Wormwood, powderav.oz.	1
Tansy, powderav.oz.	
Aloes, powderav.oz.	1
Dippel's oilfl.dr.	4
Linseed oilfl.oz.	17

Give in two doses, with an interval of five hours.

#### SECTION III—SHEEP MEDICINES.

#### Sheep-Dips.

#### Carbolic Acid Dip.

Soapav.lb.	1
Crude carbolic acidfl.oz.	16
Watergal.	50

Dissolve the soap in a gallon or more of boiling water, add the acid and stir thoroughly. Keep the mixture well thinned, and do not let it get into the mouth, nostrils or eyes of the sheep. Hold each sheep in the bath not less than half a minute.

#### Kerosene Emulsion Dip.

Fresh skimme	d milk	 	gal. 1
Kerosene		 	gal. 2

Churn together till emulsified, or mix and put into the mixture a force pump and direct the stream from the pump back into the mixture. The emulsification will take place more rapidly if the milk be added while boiling hot. Use 1 gallon of this emulsion to each 10 gallons of water required.

#### Kerosene Soap Dip.

Soap	 av.oz.	16
Water	 gal.	1
Kerosene		

Bring the water to a boil and dissolve the soap in it; then add the kerosene and churn until emulsified. Use 1 gallon of this emulsion to 8 of water.

The above are rather prophylactic in their character and are used generally after shearing.

One of the most dangerous of the parasitic diseases of sheep is scab, which may be treated either by rubbing poisonous ointments into the fleece by the hand or by immersing the sheep in aqueous mixtures containing some ingredient which will kill the parasites. When any of the flock are infected, all should be dipped, preferably being first sheared if the season permits it.

Instead of treating the scab by one application, some authorities advise the use of a preliminary dip of alkaline water to soften the scabs, or of oil or glycerin well rubbed in for the same purpose. This is to be followed in two or three days by a poisonous dip. Nearly all advise that the scabs be rubbed with a stiff brush while the sheep is being dipped.

The quantity of dip required for each sheep is variously estimated at from 1 quart to 1 gallon. For small numbers of sheep, say 50 to 100, the larger amount is necessary, while for large flocks, 1 quart for shorn or 2 quarts for unshorn sheep may be allowed. The dip should be kept while in use at a temperature of from 100 to 110 degs. F. The sheep should be dipped again within some 6 or 10 days of the first dipping in order to kill before their maturity any parasites which may have developed from eggs which were left upon the animal at the time of the first treatment, as the dip does not destroy the vitality of the eggs.

The chief poisons used in the dip are tobacco, arsenic and carbolic acid. Of these, tobacco is the favorite, because its use has not been followed by the fatality that has in times past followed the use of arsenic. Carbolic acid is too expensive to be used in large quantities, but is an excellent ingredient when only a few sheep are to be treated. The addition of tar to the dips serves a good purpose, as it is not only healing, but serves an excellent purpose in driving away flies.

#### Texas Tobacco Dip.

Tobacco av	·lbs.	30
Sulphurav	.lbs.	7
Concentrated lyeav	.lbs.	3
Water		

Steep the tobacco in three successive portions of water, expressing each time; then add the other ingredients to the liquor, and stir well while in use.

#### Law's Sheep Dip.

Tobaccoav.lbs. 1	6
Oil of tarfl.oz. 4	8
Soda ash or caustic soda av. lbs. 2	05
Soft soapav.lbs.	
Watergal. 5	

Steep the tobacco as in the previous formula, and add the other ingredients to the liquor.

#### Zundel's Carbolic Dip.

Crude carbolic acidav.lbs.	3
Caustic limeav.lbs.	2
Potash or lyeav.lbs.	6
Soft soapav.lbs.	6
Watergal.	70
Mix and boil.	

#### Dr. Kaiser's Carbolic Dip.

Tobacco av.lbs.	131/2
Soda ash or caustic sodaav.lbs.	8
Freshly slaked limeav.lbs.	4
Soft soapav.lbs.	
Crude carbolic acidav.lbs.	
Watergal.	66

Infuse the tobacco in the water, strain, and to the infusion add the remaining ingredients.

The following arsenical preparations are in favor in England:

Arsenical S	heep-Dip	Paste	(Finlay	Du	nn)
Arsenic			av.	lbs.	2
Pearl ash			av.	lbs.	2
Soft soap.			av.	lbs.	2
Sulphur .			av.	lbs.	2

This is sufficient for 125 gallons of water.

Arsenical Sheep-Dip Pas	te (Prof. Simonds).
Arsenic	av.lb. 1
Soft soap	av.lb. 1
Potassium carbonate	av.oz. 8
Water	gal 4

Boil the arsenic and potash together in half the water, and dissolve the soap in the other half. This is sufficient for 20 sheep. It should be used warm.

#### Colic Remedies.

Colic in sheep may result from exposure to

cold, from constipation, from overfeeding, or from worms.

Treatment consists in making stimulant applications to the body, and giving purgatives and enemas.

#### I.

Capsicum gr.	30
Gingergr.	120
Peppermintgr.	150
Linseed mealgr.	150
Sodium sulphateav.oz.	2

Reduce all to powder, mix well, and divide into four portions.

Give one powder every hour in a cupful of warm coffee or warm beer.

This mixture is useful in the treatment of colic resulting from cold.

#### II.

Castile soap	.gr.	60
Fennel		
Linseed meal		
Chamomile		
Sodium sulphateav		

Reduce all to powder, mix well, and divide into four portions.

Every 2 hours, mix one powder with a cupful of warm water, add a tablespoonful of linseed oil, and give the mixture to the animal.

This preparation is advised when the colic is the result of overfeeding.

#### III.

Soap, castilegr.	75
Soap, castilegi.	
Sodium chlorideav.oz.	11/
Journal Chieffac	- /2

Mix and divide into 5 powders.

One powder is to be mixed with 8 fluidounces of infusion of chamomile and used as an enema, which is to be repeated every hour.

#### Constipation Remedies.

Constipation of the sheep may result from difficultly digestible food, and may also result from a sudden change of food.

Treatment consists in giving sodium sulphate internally as well as enemas.

т

Carawaygr.	150
Sodium bicarbonategr.	150
Linseed mealgr.	
Sodium sulphateav.oz.	2 1/2

Mix all in powder form and divide into three portions.

Stir one into 8 fluidounces of warm water,

add about one-half cupful of linseed oil, and give at one dose; repeat every 3 hours.

#### II.

Soap, powdergr.	75
Rye flourav.oz.	11/2
Mix and divide into 5 portions	

Stir one powder in 8 fluidounces of warm water and give as an enema every hour.

#### Diarrhœa Remedies.

Diarrhœa affects old as well as young animals and may be caused by exposure to cold, by changing from dry to green food, or by feeding with decomposed food.

#### I.

White or red oak barkav.oz.	1
Juniper berriesav.oz.	1/2
Gingergr.	120
Wormwoodgr.	
Sodium chlorideav.oz.	
411 1 11 1 1 1 1 1 1 1 1	11

All should be in powder and be well mixed.

One tablespoonful should be given three times daily in food.

This is best adapted to older animals.

#### II.

Rhubarbav.oz.	
Prepared chalkav.oz.	
Tannic acidgr. Calamusav.oz.	
Rye flourav.oz.	
10 10 110 11 11 11 11 11 11 11 11 11 11	

All should be in powder, be well mixed, and be converted into an electuary by the addition of mucilage or syrup.

A piece the size of a hazel-nut should be given mornings and evenings.

This is best adapted to lambs.

#### Eyes, Inflammation of the.

The eye should be protected from bright light and should be bathed twice daily; then the eye waters mentioned below may be used. It is also advisable to administer mild cathartics.

#### I.

Tincture of opiumdro	ps 40
Lead waterfl.c	
Bathe the eye twice daily.	

#### II.

Zinc sulphategr.	20
Mucilage of quince seed fl.oz.	4
Distilled waterfl.oz.	
Mir and dissolve	

wix and dissolve.

Use like the preceding.

#### III.

Zinc sulphategr.	20
Infusion of chamomile (1 in 20).fl.oz.	8
Mix and dissolve.	

Use like the preceding.

#### Rheumatism Remedies.

Treatment consists in bathing, giving purgatives, and applying stimulant liniments to the limbs.

#### I.

Rosemary herb, cut	 .av.oz.	31/2
Soda ash	 .av.oz.	16
Sodium chloride	 .av.oz.	64

Mix the above with 12 gallons of hot water, allow to stand for 15 minutes, and use for bathing. The liquid may be used 2 or 3 times for a bath if warmed each time.

#### II.

Oil of turpentinefl.dr. 4	Ŀ
Water of ammoniafl.dr. 4	Ł
Spirit of soapfl.oz. 7	
Spirit of camphorfl.oz. 7	1/2
Rub the legs with this mixture twice dai	ly.

#### III.

Sodium sulphateav.oz.	11/2
Aloesgr.	80
Linseed oilfl.dr.	
Infusion of linseed (1 in 20)fl.oz.	7
Mix and dissolve.	

Give one-half at a dose and repeat in 3 hours.

This mixture is intended for full-grown animals.

#### TV.

Sodium salicylategr.	60
Aloesgr.	360
Infusion of linseed (1 in 20)fl.oz.	15
Mix and dissolve.	

Give a teaspoonful 2 or 3 times daily according to the age of the animal.

This mixture is recommended for lambs.

#### Urine, Treatment for Retention of.

Hemp seedav.oz.	2
Magnesium sulphat eav.oz.	1
Bitter almondsgr.	90
Juniper berriesav.oz.	
Rye flourav.oz.	

Reduce all to powder, mix and make an electuary by the addition of water or syrup.

A piece the size of a marble may be given once daily.

#### Urine, Treatment for Bloody.

The animals should be placed in a warm stall and fed with wholesome dry food. The following may also be given:

Lead carbonateav.oz.	1/2
Camphorav.oz.	1/2
Bitter almondav.oz.	1
Linseed mealav.oz.	3
Rye flourav.oz.	5

All should be reduced to powder and be made into a stiff mass with syrup or molasses.

A piece of the size of a hazelnut should be administered once daily.

## Worm, Remedy for Tape.

Aloes	 .gr.	<b>1</b> 50
Oleoresin of male fern	 .gr.	30
Naphthalin		

Make into a mass with spirit of soap and divide into 2 pills.

The dose for a lamb, 4 to 8 months, is 1 pill given in the morning without giving food; the other pill should be given 8 days thereafter. The dose for lambs only is given, as full-grown animals are rarely affected.

#### SECTION IV—SWINE MEDICINES.

#### Appetite, Treatment for Loss of.

Loss of appetite may be indicative of other diseases, but may also be the result of over-feeding or due to the presence of undigested food.

Treatment consists in changing the food, giving an emetic, and subsequently the powder mentioned below.

I.	
Tartar emeticgr. 15	
Ipecac, powdergr. 45	
Althæa rootgr 75	

Mix and make into an electuary by means of syrup, molasses or mucilage.

Give at one dose as an emetic.

TT.

Mix all and reduce to powder.

Give a tablespoonful twice daily.

#### Catarrh Remedies.

ΙI.

Ammonium chloridegr.	150
Antimony sulphidegr.	150
Argols (crude tartar)gr.	300
Licorice rootav.oz.	11/2
Linseed mealav.oz.	11/2
Mix in nowder and add syrup or of	lucose

Mix in powder and add syrup or glucose to form an electuary.

Three times daily give a piece about the size of a walnut.

TT

Potassium nitrategr.	150
Sodium sulphateav.oz.	21/2
Linseed mealav.oz.	34

Mix in powder and add syrup or glucose to form an electuary.

Give a tablespoonful every 2 hours.

#### Colic Remedies.

Colic may result from eating difficultly digestible food or poisonous plants, from exposure to cold, or from worms.

Treatment consists in keeping the animal in a warm stall, administering aromatic, stimulating, and purgative remedies, and giving enemas.

I.

Sodium sulphateav.oz.	11/2
Peppermint	1/2
Sodium chloridegr.	<b>15</b> 0
Reduce all to powder and mix well.	

Give one-half of this mixture in 8 fluidounces of weak coffee and repeat the dose in 3 hours.

II.

Chamomileav	.oz. 1/2
Peppermintav Green soapav	.oz. 1/2
Green soapav	.oz. $2\frac{1}{2}$
Linseed oil	.oz. 2½
Water, boilingfl	

Pour the water upon the chamomile and peppermint, let stand for 15 minutes, strain, and add to the colature the remaining ingredients.

Inject 8 fluidounces of this mixture, previously warmed, into the rectum every half hour.

#### Diarrhœa Remedies.

Diarrheea may result from exposure to cold or from disorders of digestion. If it continues more than 24 hours, treatment is urgently demanded, as it may cause a severer disease. The animal should be kept warm, and astringent and aromatic remedies may be given, and possibly suppositories may be required.

T.

Chamomilegr.	150
Peppermint, cutgr.	150
White or red oak bark gr.	
Tannic acidgr.	

Pour on this mixture 16 fluidounces of boiling water and give one-fourth of the infusion every 2 hours.

II.

Ferrous sulphategr. 40	
Alumgr. 40	
Acaciagr. 375	
Sugar of milkgr. 300	

Mix all in powder and divide into 5 parts. Give one powder in a cupful of warm chamomile tea or other suitable vehicle every 3 hours.

III.

Rhubarbgr.	15
Calcium carbonate gr.	150

Mix in powder and divide into 10 portions. One portion is to be given twice daily in a tablespoonful of warm chamomile tea or other suitable vehicle. This remedy is intended for sucking pigs.

IV.

Tannic acidgr.	8
Cacao buttergr. 30	0
Mix and divide into 5 suppositories.	

After each movement of the bowels, insert one suppository, pushing it as far up into the rectum as possible with the well-oiled finger.

## Eve. Remedies for Inflammation of the.

Wash the eye 3 times daily with lukewarm water and use the following eye waters:

I. Zinc sulphate.....gr. 36 Tincture of opium.....fl.dr. 11/4 Infusion of chamomile (1 in 50).fl.oz. 16

II.

Ammonium	chloridegr.	72
Lead water	fl.oz.	16

#### Hog Cholera Remedies.

Hog cholera, swine fever, or swine plague are the names applied to a group of symptoms produced by three distinct types of disease, quantity as the foregoing.

namely, charbon, contagious pneumo-enteritis and epizootic catarrh, all, however, dependent upon recognized specific germs. The second form, pneumo-enteritis, probably is the one most commonly understood, and it is characterized by an inflamed, ulcerated condition of the alimentary tract, and fetid, bloody discharges-these having determined the popular designation of "cholera."

Prophylaxis consists in keeping the animal in perfect health by insuring proper hygienic food and surroundings. One of the principal, if not the main, cause of the disease is excessive feeding of corn. Corn-fattened animals are not in a physiological condition, as is well understood, and corn, principally consisting of starch, being particularly prone to fermentation, naturally must favor development of fermentative diseases in debilitated organisms. Hence corn should largely be replaced by the grains, bran, peas, beans and other nitrogenous foods, which will produce more muscle and less unhealthy fat. Decaying and fermenting food should not be offered under any condition. Clean, ventilated stalls that will afford protection against inclement weather, but also plenty of exercise in the open air, are prerequisites.

Prophylactic treatment consists in the daily administration of wood or animal charcoal or iron sulphate. Potassium chlorate is also recommended. In fact, experience teaches that the two latter agents are the only ones which proved effective for this purpose or in treating the initial stages of the disturbance. A few formulas are appended.

Arsenicav.oz.	
Antimony sulphideav.oz.	4
Potassium nitrateav.oz.	4
Ferrous sulphateav.oz.	16
Sulphurav.oz.	
Madderav.oz.	

Mix with 12 gallons of slop, and give 1 pint to each hog, the whole being for fifty.

II.

c ·	-1
Capsicumav.lb.	T
Ferrous sulphateav.lb.	2
Madderav.lb.	2
Widuuci	5
Calcium phosphateav.lb.	10
Wood-ashes, siftedav.lb.	10

This may be administered in the same

As soon as an animal develops symptoms of the plague it must be isolated from the herd and surrounded by absolutely antiseptic conditions. Constipation, if present, should be relieved by castor oil or rhubarb. The bowels should be frequently irrigated with warm water; one of the aforementioned prescriptions should be administered; the food should be restricted to well-cooked oat or barley meal gruel, or, if this is not borne, boiled corn starch. Some advise putting a trace of sulphuric acid into the drinking water. For the fever potassium nitrate usually is resorted to. In the convalescent stages tonics must be administered.

When the disease is fully developed there is little hope, and the animal should be killed and buried with quicklime. In England swine fever comes under the contagious diseases act, and treatment is not permitted. The public health authorities must be advised, who destroy the infected animal.

#### III.

Iron carbonate av.oz. Sodium chloride av.oz. Potassium carbonate av.oz. Sulphur av.oz. Lime av.oz. Magnesium carbonate av.oz. Soap av.oz.	5 5 5 10
Soapav.oz. Carbolic acidfl.oz.	10

Mix well and reduce to powder.

One-fourth av. ounce should be given at each meal, mixed with the food.

#### IV.

1.
Wood charcoalav.oz. 4
Sulphurav.oz. 4
Sodium sulphateav.oz. 4
Antimony sulphideav.oz. 4
Sodium chlorideav.oz. 8
Sodium bicarbonateav.oz. 8
Sodium hyposulphiteav.oz. 8
Reduce all to powder and mix well.

A large tablespoonful for each 200 pounds of animal should be given once daily with food.

#### 77

8	•	
	Calcium phosphateav.oz.	16
	Chalkav.oz.	
	Magnesium carbonateav.oz.	4
	Capsicumav.oz.	1
	Reduce all to powder and mix well.	

Give one tablespoonful three times daily with food.

Iron carbonateav.oz.	5
Potassium carbonateav.oz.	2
Sodium chlorideav.oz.	4
Sulphurav.oz.	4
Magnesium carbonateav.oz.	
Wood charcoalav.oz.	
Soapav.oz.	
Carbolic acidav.oz.	2
Chalkav.oz.	50

Reduce all to powder and mix well.

One-half to 1 av.ounce should be given with each meal.

#### VII.

Sodium bicarbonateav.oz.	4
Gentianav.oz.	4
Gingerav.oz.	6
Potassium nitrateav.oz.	2
Chalkav.oz.	16

Reduce all to powder and mix well.

Use like the preceding.

#### Swine Fever Remedies.

This disease, which is exceedingly contagious and prevalent in many localities, is indicated by an increase in the temperature of the hog from 37 degs. C., the normal temperature in health, to 40 degs. C., sometimes rising as high as 41 degs. C. The other symptoms are highly colored urine, cold extremities, loss of appetite, and constipation. The following treatment is recommended:

A cathartic and diuretic should first be given, consisting of:

,	, 8	
	Potassium nitrateav.oz. Sulphurav.oz.	3/4
	Sulphurav.oz.	11/2
	Magnesium sulphateav.oz.	5
	Molassesav.oz.	2
	Water, enough to make fl.oz.	20

Mix the sulphur with the molasses, then add the water gradually, in which the salts have previously been dissolved.

Shake the mixture, and give 1 ounce every morning until relieved. This is the dose for hogs of average size; for larger animals the dose should be increased.

For diarrhœa and weakness usually resulting from subsidence of the fever, the following mixture is administered:

Sodium bicarbonate	.av.oz.	4
Gentian		
Catechu	.av.oz.	4
Cinchona	.av.oz.	4

Reduce all to powder and mix well.

From one-fourth to 1 av. ounce of this pow-

der should be given in food. Condition powders should be administered.

In diarrhoa accompanied with an irritable or relaxed condition, the following is recommended:

Opiumgr.	60
Nutgallgr.	
Pimentogr.	
Sodium bicarbonategr.	
Reduce all to powder and mix well.	

The dose is from 30 to 120 grains made into a bolus with molasses.

#### Vomiting, Remedy for.

Prepared chalkgr.	75
Sodium bicarbonategr.	
Sodium chloridegr.	150
Sodium sulphategr.	
Linseed mealav.oz.	1 1/2
Mills all in manufact and disside into 5	manta

Mix all in powder and divide into 5 parts. Give 1 part every 3 hours in a cupful of warm chamomile tea or other suitable vehicle.

#### Worm Medicine.

Codium aulphoto porudos or or	9
Sodium sulphate, powderav.oz.	
Tansy, powdergr.	
Castor oil fl.dr.	
Naphthalingr.	30
Rye flourav.oz.	3/2
Mir all and add curup glugges of	r mo

Mix all and add syrup, glucose or molasses to form an electuary.

Give one-fourth of this mixture every 2 hours.

#### SECTION V—DOG AND CAT MED-ICINES.

The doses and quantities mentioned in the succeeding formulas are intended for dogs of medium weight—about 50 pounds. Larger animals will require proportionately larger doses and smaller ones smaller doses.

Many of the following remedies are adapted for ailments of cats, the dose being somewhat less than for dogs. Pills and liquids are the best forms of medicines to administer to dogs, while powders and liquids are more suitable for cats.

In giving medicines to dogs, open the mouth of the animal and place in it crosswise a small stick of wood, then thrust the pill, capsule or bolus down the throat with the finger; if a liquid, insert the neck of the bottle on the side of the mouth and hold the head back so as to compel the dog to swallow.

When administering to cats, the powder is best blown through a glass or rubber tube onto the roof of the mouth; the liquid medicine is best poured upon the front paws, which the animal will lick off to clean them.

#### Appetite, Remedies for Loss of.

Dogs are liable to overeating, and this may result in loss of appetite. Treatment consists in giving an emetic and following this with a purgative and stomachic mixture.

I.	
	Tartar emeticgr. ½
	White helleboregr. 2
	Give this powder at one dose.

3	[,	
	Sodium sulphate, driedgr.	90
	Sodium bicarbonategr.	
	Rhubarb gr.	30
	Calamusgr.	

Mix all in powder, make into a mass, and divide into 6 pills.

One pill should be given twice daily.

## Canker of the Ear, Treatment for.

To effect a cure, treatment must be begun in the early stages of the disease. During treatment, the animal must be prevented from scratching the ear.

I.		
	Copper sulphategr.	36
	Alumgr.	
	Waterfl.oz.	
	Mix and dissolve.	

Wash the ear out 3 times daily with this liquid by means of a soft sponge.

#### TT

Lead acetategr.	36
Copper acetategr.	
Glycerin	
Distilled waterfl.oz.	
Mix and dissolve.	- /2
TT 111 41 11 11	

Use like the preceding.

#### III.

Silver nitrategr.	18
Distilled waterfl.oz.	
Wash the diseased portions of the ear	wit

Wash the diseased portions of the ear wit this liquid 3 times daily.

#### IV.

* *	
Tannic acidgr. 6	60
Bismuth subnitrategr. (	30
Simple cerateav.oz.	

Anoint the diseased portions of the ear once daily.

272 THE STANDAR	RD FORMULARY.
V.  Ointment of red oxide of mercury av.oz. ½ Simple cerate	IV.  Yellow sulphide of antimonygr. 10 Ammonium chloridegr. 36 Purified extract of licoricefl.dr. 4 Syrup of althæafl.oz. 3½ Give one tablespoonful every 2 hours.
IV.	ing and evening.
Syrup of buckthornfl.oz. 1	Diarrhœa Remedies.
Give at one dose.	Decomposed or very fatty food, overeating, drinking of too cold water, exposure to cold,
Cough Medicines.	etc., are liable to cause diarrhœa.
I.  Sodium bromidegr. 180 Creosote waterfl.oz. 2 Fennel waterfl.oz. 4  Mix and dissolve. Give one-half tablespoonful 4 times daily.	Treatment consists in keeping the animal warm, rubbing the abdomen with alcoholic liniments, and giving opium, astringents and chalk; suppositories may also be of value. In feeding give meat, cooked rice stirred with yolk of egg, and meat broth.
II.	I.
Tincture of belladonna	Spirit of camphorfl.oz. 2 Spirit of juniperfl.oz. 2 Rub the abdomen with this mixture 3 times daily and then inclose the animal in a warm
Give one teaspoonful 3 times a day.	wrap.
III.  Morphine sulphategr. 2 Bitter almond waterfl.oz. 1 Creosote waterfl.oz. 1	II.  Opium
Mix and dissolve.  Give 20 drops in a little water 3 times daily.	Mix, make into a mass, and divide into 5 pills.  Give one pill mornings and evenings.

III. Tannic acidgr. 50	Eye, Remedies for Inflammation of the.
Bismuth subnitrategr. 30 Licorice rootgr. 50	In simple inflammation of the eye, give
Mix, make into a mass, and divide into 10	mild cathartics and use one of the eye waters mentioned below.
pills.  Give one pill 3 times daily.	I.
	Lead acetategr. 10
IV.  Bismuth subnitrategr. 15 Extract of rhatanygr. 40	Rose waterfl.oz. 2 Drop into the eye every hour.
Sugargr. 100	II.
Mix and divide into 5 powders.	Zinc sulphategr. 5
Give one powder every 4 hours.	Rose water
V	Use like the preceding.
Lead carbonategr. 8 Bismuth subnitrategr. 30	III.
Acaciagr. 40 Sugargr. 80	Magnesium sulphate
Mix and divide into 10 powders.	Mix and dissolve.
Give one powder every 3 hours.	Give one tablespoonful twice daily.
This mixture is intended for obstinate	IV.
cases.	Sodium sulphate, driedgr. 300
VI.	Sodium bicarbonategr. 75 Sodium chloridegr. 75
Rhubarbgr. 75 Cascarillagr. 75	Licorice rootgr. 300
Prepared chalkgr. 150	Buckbeangr. 150
Licorice rootgr. 150	Mix all in powder and form into an electu-
Mix all in powder and form into an electu-	ary by the addition of juice of juniper ber-
ary by the addition of mucilage.	ries, or, if this be not at hand, syrup, molas-
Give a piece the size of a hazelnut three	ses or other suitable agent.
times daily.	Give a piece of the size of a hazelnut twice daily.
VII. Extract of rhatanygr. 18	
Cacao buttergr. 180	Fits, Remedies for.
Make into six suppositories.	See Epilepsy Remedies.
One suppository should be inserted far up	Mange Remedies.
into the rectum after each copious evacua-	In addition to using the remedies men-
tion.	tioned below, the kennel should also be thor-
Epilepsy Remedies.	oughly disinfected.
I.	I.
Zinc oxidegr. 20	Benzinefl.oz. 7
Sulphurgr. 75	Oil of cade
Jalapgr. 75 Extract of green helleboregr. 20	Green soapfl.oz. 2
Extract of gentian or dandelion	Oil of turpentinefl.oz. 2
enough to form a mass	Rub the soap and tar together, add the oil
Divide into 60 pills.	of cade, and then incorporate the other in-
Give one pill three times daily.	gredients.
II.	Rub this in once daily.
Fluid extract of valerian fl.dr. 1	II.
Syrup of buckthornfl.dr. 3  For cats—15 to 20 drops every hour or	Salicylic acidgr. 24 Lardav.oz. 1
two, with little milk or other food, or on	This ointment is recommended for applica-
some herbage like that of catnip.	tion to the head because soap cannot very well
Town was a compression of the co	•

IHE STANDAR	D FORMULARY.
be used in the vicinity of the eyes. It is to	IV.
be applied once daily.	Spirit of camphorfl.oz. 2
III.	Spirit of formic acidfl.oz. 2
Sulphur av.oz. 6	Apply to the painful limbs three times
Ammonium chlorideav.oz. 34 Aloesgr. 90	daily and then wrap with a woollen cloth.
Venice turpentineav.oz. 34 Lard	V.
Lardav.oz. 9	Oil of turpentine
Apply once a day and wash off after four	Spirit of formic acidfl.oz. 2
applications.	Use like the preceding.
IV.	Stomach Catarrh, Treatment for.
Aloesav.oz. 14	Dogs are very liable to overeating or to
Mercurial ointmentav.oz. 1/4 Spirit of turpentinefl.oz. 1/4	eat indigestible or decomposed food, and this
Sulphurav.oz. 2	may result in gastric catarrh.
Lardav.oz. 4	Treatment consists in first giving an em-
Apply twice daily.	etic; then a purgative may be administered
V.	unless there is diarrhea, when medicine must
Sulphurated potassagr. 50 Creosotefl.dr. 34	be given to counteract the latter condition.
Pine targr. 90	Tartar emeticgr. ½
Cocoanut oilfl.oz. 2	Ipecacgr. 15
Cleanse the parts with soft or green soap,	Give at one dose in a spoonful of water.
wash off well, dry, and then apply this oint-	II.
ment.	Aloes gr. 60
VI.	Soap, Waterof each, sufficient
Sperm oil         fl.oz. 7           Kerosene         fl.oz. 7	Mix and make into three pills.
Carbolic acidfl.dr. 3	Give one pill every 5 hours.
Pine tarav.oz. 1 Sulphurav.oz. 1	III.
Apply once daily.	Tannic acidgr. 15
Rheumatism Remedies.	Bismuth subnitrategr. 10 Calamusgr. 150
	Reduce all to powder, mix well, and divide
Treatment consists in giving sodium sali- cylate and mild cathartics and applying stim-	into 5 parts.
ulant liniments.	Two hours after the emetic ceases to act
I.	give one powder in a little water, and repeat
Sodium salicylategr. 192	the dose every 12 hours.
Waterfl.oz. 6	Tonic Medicine.
Give one tablespoonful three times daily.	Gentian gr. 15
II.	Ginger gr. 5 Cascarilla gr. 15
Acetanilidgr. 75	Mix and make into a pill.
Rye flourgr. 75  Mix and make into five pills.	Give one such pill every day.
Give one pill in the morning and one in	Urine, Treatment for Bloody.
the evening.	The usual treatment consists in giving mild
III.	cathartics and enemas.
Sodium nitrategr. 40	I.
Magnesium sulphateav.oz. 1 Solution of ammonium acetatefl.oz. 1	Magnesium sulphateav.oz. 1/2
Waterenough to make fl.oz. 6	Tamarind pulpav.oz. ½ Fennel waterfl.oz. 5
Mix, dissolve and filter.	Mix, dissolve and filter.
Give one tablespoonful every hour.	Give one tablespoonful every 2 hours.

	POULIKY MEDICINES, 215		
II.  Volk of one egg Olive oil	V.  German wormseed, powdereddr. 1 Fluid extract of spigeliafl.dr. 3 Fluid extract of sennafl.dr. 1 Fluid extract of valerianfl.dr. 1 Syrup of buckthornfl.oz. 2 Dose ½ to one teaspoonful night and morning. Suitable for cats.  Flea Powder.  Napthalinav.oz. 4 Starch		
Creosote water	SECTION VI-POULTRY MED-		
Give one-nair and repeat the dose in &			
	ICINES.		
hours.			
	Poultry Powders.  Various mixtures of powdery substances are dispensed under the names "Poultry Powder," "Poultry Food," "Egg Food," and "Egg-Making Food." The latter two names are used because of the alleged property of these mixtures, of increasing the egglaying power of hens.		

III.	Reduce all to moderately coarse powder
Capsicumav.oz. 2	and mix well.
Allspice	Use like the preceding.
Reduce all to powder and mix well.	VIII.
•	Mustard seedav.oz. 10
One tablespoonful to be mixed with every	Fenugreekav.oz. 8
pound of food and fed two or three times a	Sodium sulphateav.oz. 2 Capsicumav.oz. 1
week. The addition of a little dried ants'	Sodium chlorideav.oz. 1
eggs, if not too expensive, would prove bene-	Iron carbonateav.oz. 1
ficial.	Gingerav.oz. 1
IV. Mustardav.oz. 4	Black antimonyav.oz. 1 Bone, groundav.oz. 1
Fenugreek	Corn mealav.oz. 10
Oyster shells, groundav.oz. 21/2	Reduce all to powder and mix well.
Bone, groundav.oz. $1\frac{1}{2}$	Use like the preceding.
Sodium sulphate	IX.
Capsicum	Sunflower seedav.oz. 8
Venetian redav.oz. 2	Mustard seedav.oz. 8
Corn flourav.oz. 4	Ginger
Asafetidagr. 90	Sulphurav.oz. 4
Reduce all to powder and mix well.	Capsicumav.oz. 1/2
A tablespoonful is to be mixed with suffi-	Salt
cient meal or porridge to feed 20 hens.	Bone meal
V. Egg shell, or calcium phosphate av.oz. 4	Oyster shell, groundav.oz. 21/2
Sulphate of ironav.oz. 4	Chalk
Capsicum	Magnesium sulphateav.oz. 1
Fenugreekav.oz. 2	Reduce all to powder and mix well.
Black pepperav.oz. 1	Use like the preceding.
Silver sand	Mustard seedav.oz. 5
Reduce all to powder and mix well.	Fenugreekav.oz. 8
Use like the preceding.	Oyster shell, groundav.oz. 6
VI.	Bone meal
Calcium phosphate, or ground	Sodium sulphateav.oz. 2
bone	Capsicumav.oz. 1
Capsicum	Bayberry barkav.oz. 1
Cantharidesgr. 60	Sodium chlorideav.oz. 1 Gingerav.oz. 1
Sulphurav.oz. 1	Black antimonyav.oz. 1
Potassium nitrateav.oz. 1	Fine sandav.oz. 4
Reduce all to powder and mix well.	Asafetidaav oz. ½
Mix a tablespoonful with a quart of feed,	Wheat flour
and give two or three times a week.	
VII.	Reduce all to powder and mix well.
Oyster shell, groundav.oz. 5 Magnesiaav.oz. 1	Use like the preceding.
Calcium carbonateav.oz. 3	Chicken Lice from Coops, to Remove.
Bone, groundav.oz. 1½	Carbolic acid, crudefl.oz. 1 Waterfl.oz. 15
Mustard branav.oz. 1½	Wash the woodwork with this mixture and
Capsicumav.oz. 1 Saltav.oz. 1	
	sprinkle in nests and on floor.
Sodium carbonate av.oz. 1/2	Chicken Cholera, Remedy for.
Sulphurav.oz. ½ Beef, lean, dried and powdered.av.oz. 10	Sulphuric acidfl.oz. 1
Fine sand av. oz. 10	Iron sulphateav.oz. 16
Corn mealav.oz. 20	Water, enough to makegal. 1
Linseed mealav.oz. 20	Add 1 fluidounce of this mixture to a

pint of water, and supply in place of water, or mix with meal or other food.

#### II.

Reduce all to powder and mix well.

An even teaspoonful is to be given with the feed for a dozen fowl.

#### III.

Give three or four teaspoonfuls of strong alum water once a day. This is recommended by the U. S. Department of Agriculture.

#### Roup Remedy.

Potassium chlorateav.oz. 2
Cubebsav.oz. 2
Anise
Licorice rootav.oz. 3

Reduce all to powder and mix well.

A teaspoonful of this is to be mixed with food for sixty hens.

#### SECTION VII-BIRD MEDICINES.

#### Mixed Bird Seed.

Hemp seedav.oz. 5
Canary seedav.oz. 4
Millet seedav.oz. 1
Maw seedav.oz. 1

Various other mixtures are also used as mixed bird seeds; some of these contain rape seed.

#### Canary Bird Food.

Dried yolk of eggav.oz. 2
Poppy heads, coarse powderav.oz. 1
Cuttlefish, coarse powderav.oz. 1
Sugar, granulatedav.oz. 2
Biscuit from wheat flour, dried
and powderedav.oz. 8

# German Bird Paste or Canary Food. I.

Pass the egg through a fine grater and add to the other ingredients. Beat to a smooth paste with cold water, and granulate the

mass by passing through a coarse grater, then expose the product to the air in a warm place until quite hard and dry.

II.

Beat together the ingredients until a smooth paste is formed, then granulate by pressing through a colander and allow to dry. Some prefer to add to the above the yolks of two eggs, or two or three hard-boiled eggs. Instead of the honey, 2 av. ounces of sugar may be used; then cold water must be added to the mixture to form a paste.

III.

Pea mealav.oz.	8
Sugar, granulatedav.oz.	4
Fine-grated stale breadav.oz.	4
Butter, freshav.oz.	1
Yolks of eggs	1
Poppy seedav.oz.	1
Hemp seed, bruisedav.oz.	16

Mix the first five ingredients, brown slightly in a frying pan and add the remaining ingredients.

#### Food for Larks, Nightingales and Other Insectivorous Birds.

Use either of the three formulas just immediately preceding.

## Mocking Bird Food.

١.		
	Broken crackersav.oz.	8
	Corn av.oz.	9
	Riceav.oz.	2
	Hemp seedav.oz.	
	Capsicumgr.	
	Mix and reduce to coarse nowder	

II.

Hemp seedav.oz.	16
Rape seedav.oz.	8
Crackersav.oz.	8
Rice, unshelledav.oz.	2
Corn mealav.oz.	2
Capsicumav.oz.	2
Lard oilfl.oz.	2

Mix all but the oil, grind to coarse powder, and then incorporate the oil.

#### Food for Redbirds.

Sunflower seed	av.oz.	8
Hemp seed	av.oz.	16
Canary seed		
Wheat, cracked	av.oz.	8
Rice, unshelled	av.oz.	6

Mix and grind to coarse powder.

III.

Oxheartav.oz. 2	
Wheat bread, toastedav.oz. 4	
Maw seedav.oz. 2	
Hemp seedav.oz. 6	

Boil the oxheart well in water, cut small, and place it in a pan in an oven where it must be allowed to become perfectly dry and crisp. All the ingredients must then be mixed and reduced to coarse powder.

#### 

Give a few drops on a lump of sugar in the cage once daily.

This mixture may be dispensed under the name "Pectoral Tonic."

#### Constipation Remedy for Birds.

Fluid extract of sennafl.dr. 2	
Syrup of mannafl.oz. 1	
Fennel water, enough to makefl.oz. 4	
Character to the state of many to	

Give a few drops on a lump of sugar in the cage once daily.

#### Diarrhœa Remedy for Birds.

Tincture of iron chloridefl.dr.	2
Paregoricfl.dr.	
Caraway waterfl.oz.	
Use like the preceding	,-

#### Gapes in Pheasants, Remedy for.

apos III	a moundation, mountains	••
Ferrous	sulphategr.	<b>6</b> 0
Capsicur	mav.oz.	1/2
Fenugre	ekav.oz.	1
Red sau	indersav.oz.	1
Licorice	rootav.oz.	2
Molasses	s suffici	ent

## General Directions for Care of Birds.

When ailing, the first thing is to keep the bird as quiet as possible; this is best accomplished by covering the cage closely. Most ailments of birds are due to a cold, the cause of which is generally owing to exposure to a draught of air or keeping the bird in an overheated room.

To cure a cold, add to the regular food equal parts of a paste consisting of hardboiled eggs, corn meal and grated apple, the whole well spiced with Cayenne pepper. Sometimes the bird seems hoarse and has apparently lost his voice; this is frequently occasioned by over-singing, and is remedied by dissolving a little pure rock candy in the Birds troubled with diardrinking water. rhæa can usually be relieved by placing a rusty nail in their drinking water; if this is not effective, recommend "Diarrhoa Remedy." Should the bird be troubled with costiveness, mix an additional quantity of green fruit to the regular food; if this does not relieve, reccommend "Constipation Remedy."

This is made into a paste, of which a piece the size of a hazelnut is dissolved in a gallon of water and given the birds to drink.

#### Ointment, Healing and Soothing.

Peru balsamgr.	60
Cold creamav.oz.	1

#### Tonic Medicine.

[.			
	Tincture	of cinchonafl.dr.	1/2
		of chloride of irondrops	
	Glycerin		1
		waterenough to make fl.oz.	
	Put a fev	y drops on a lump of sugar in	the

Put a few drops on a lump of sugar in the cage daily.

II.

Iron sulphategr.	120
Diluted sulphuric aciddrops	15
Waterfl.oz.	
D' 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1

Dissolve the sulphate in the water and add the acid.

A teaspoonful of this mixture is to be added to each quart of the drinking water of the birds. This is especially recommended for moulting birds.

This preparation is known as "Douglass" Mixture."

III.

111.			
	Compound tincture of cinchonafl.dr.	2	
	Compound tincture of gentianfl.dr.	2	
	Syrup of orangefl.oz.		
	Simple elixirfl.oz.	$2\frac{1}{2}$	

Put a few drops on a lump of sugar in the cage once daily.

These preparations may be dispensed under the names "Bird Tonic," or "Tonic Elixir."

When moulting, special attention should be paid to the birds, as any disease to which they are predisposed will show itself then. Give warming, nourishing food, keep the bird and cage clean, place in the warm sunlight and keep them out of draughts. Should this not be effective to recuperate them, recommend "Tonic Elixir."

The feet are often the seat of disease. To remove accumulations of dirt from the feet, hold them in lukewarm water for three or four minutes each day till the dirt drops off. If the feet are warty and seem to be sore, bathe them as above and grease them with fresh cream, or, still better, with an "Ointment of Cold Cream."

Pimples, or obstruction of the rump gland, is caused where the bird is not using the gland. Open the gland with a fine needle and apply the "Ointment of Cold Cream."

For lice, supply the cage daily with clean, fresh sand mixed with some "Insect Powder."

#### PART V.

## TOILET PREPARATIONS.

This division naturally subdivides into the entirely; a residuary odor remains in the case following sections:

SECTION 1. Distinctively Odor Preparations.

SECTION 2. Skin Preparations.

Hair Preparations. SECTION 3.

SECTION Mouth (Tooth, Lip, Breath) Preparations.

SECTION 5. Bath Preparations.

#### SECTION I OF PART V.

Preparations used mainly or largely for their odor, including Handkerchief Extracts, Sachet Powders, Solid Perfumes, Pot Pourris, Fumigating Pastilles, Powders, etc.; Cologne Waters, Toilet Waters, Bay Rum, Toilet Vinegars, and Smelling Salts.

#### Perfumes.

Perfumes, as druggists employ the term, embrace Handkerchief Extracts, Cologne and Toilet Waters, and Sachet Powders.

Fine perfumes can be prepared only by the use of good material; in fact, the best obtainable is none too good for a discriminating taste. In the following paragraphs will be outlined the requisites of the best materials.

One of the first requisites in the manufacture of fine perfumes is a first-class quality alcohol, free from fusel oil or other odorous contamination that would modify or destroy the delicate floral odors which are the main ingredients of fine perfumes. best kind of alcohol is what is known as "deodorized alcohol." (See process of manufacture in Part I.) Another grade of alcohol which is fairly good is what is sold as "cologne spirit," which is the "middle run" obtained in the manufacture of ordinary alcohol, and is fairly free from fusel oil.

The difference between ordinary alcohol and purified alcohol can be determined very easily by rubbing a few drops between the bands and allowing the alcohol to evaporate

of ordinary alcohol, which is quite marked and very pungent and objectionable.

It is in the case of volatile oils probably that the greatest difficulty is experienced in securing first-class material. These substances should always be of the very best quality and of the best grade, free from adulteration, and should be preserved in small well-stoppered bottles in a cool place. Such oils as those of the citrine variety-orange, lemon, etc.—should be preserved by the addition of some alcohol, this being added as soon as the oils are received.

Oil of Bitter Almond is now obtainable deprived of hydrocyanic acid, and owing to the excessively poisonous character of the latter, such an oil should be preferred for perfumery purposes. The artificial oil should not be substituted for the natural product.

Oil of Neroli, or Orange Flowers, is of several varieties. The most highly esteemed and the one which should always be used for fine perfumes is what is known as Oil of Neroli Petale, distilled from the flowers of the sweet orange. The other varieties are Oil of Neroli Bigarade, obtained from the flowers of the bitter orange, and Oil of Petit Grain, which is produced by distilling the leaves and unripe fruit.

Much confusion exists regarding Oil of Rose Geranium. Only the best French or Spanish oil should be used. In the case of Oil of Cloves, what is known as the Bourbon variety is most highly esteemed. In the case of Oil of Orange, the Oil of Bitter Orange peel is considered the best. Of the oils of Lavender, the English, or so-called "Mitcham," is the best.

Oil of Orris occurs in two forms, concrete and liquid, but the latter only is to be employed in the formulas given below. Oil of Sandalwood is largely adulterated, and only the best English or East Indian oil should be used.

Volatile oils are frequently employed in the

form of spirits, or dilutions with alcohol, in the manufacture of liquid perfumes.

What is known as "pomades" are also of very great importance in the manufacture of perfumes. These are known by numbers which indicate the number of times purified lard has been impregnated with the flower odors by the method of enfleurage.

These pomades include the following odors: Rose, orange flowers, jasmine, tuberose, jonquil, cassie, reseda or mignonette, violet, lily of the valley, mimosa, and helictrope. Before these pomades can be used they must be extracted by means of strong alcohol (to be described below); this alcoholic extract is known in this work by the title of "Essence."

Various animal substances are used in the manufacture of perfumes. These include musk, civet, and ambergris. Inasmuch as these are very expensive, they are subjected to the grossest adulterations, and the utmost care must be exercised to obtain the very purest of product.

These animal substances are not altogether used for their own odor but for their peculiar property of "fixing" the fugitive flower odors and making them permanent.

The best musk is the variety known as

Tonquin in "grains." This substance is employed usually in the form of a tincture or alcoholic extract. The extractive action of the alcohol is assisted by a small amount of alkaline water and placing the vessel containing the mixture in a warm place for at least 30 days.

Ambergris occurs in several varieties, the most esteemed of which is that kind known as the gray variety. Both it and civet are used in the form of alcoholic tinctures. To facilitate extraction, they are first triturated with about an equal amount of finely granulated orris root.

Among various crude vegetables employed in perfumery are orris root, vanilla, tonka, benzoin, tolu, storax, and Peru balsam. The best orris root is the kind known as Florentine; it is generally employed in the form of a concentrated alcoholic tincture. The other substances mentioned occur also in different grades or varieties, but only the very best should be selected for perfumery purposes. Like orris, all are generally employed in the form of alcoholic tinctures.

Among the other substances employed in the manufacture of perfumes are various chemicals such as heliotropin, hyacinthin, terpineol (also known commercially as

The following table is an interesting one as showing the relationship between the different odors:

Classification of Odors.

CLASSES.	TYPES.	ORDERS BELONGING TO THE SAME CLASS.		
Rose	Rose	Geranium, sweet brier, rosewood. Lily of the valley, white pond lily, ylang ylang.		
Orange flower	Orange Flower	Acacia, syringa, orange leaves.		
Tuberose Violet	Tuberose	Lily, jonquil, hyacinth. Orrisroot, mignonette.		
Balsamic	Benzoin	Balsam of Peru, tolu, benzoin, storax.		
Vanilla	Vanilla			
Spice	Cinnamon			
Camphor	Camphor			
Sandal	Sandalwood			
Citrine	Lemon Lavender			
Mint	Peppermint	Spearmint, balm, rue, sage.		
		Caraway, dill, coriander, fennel, star anise.		
		Peach kernels, mirbane, cherry-laurel. Civet, musk-seed, sumbul, ambergris.		
Fruit	Pear	Apple, pineapple, quince.		

lilacin or muguet), coumarin, vanillin, benzoic acid (so-called true or English should be employed in perfumery, this being prepared from benzoin and containing, therefore, some oily substance derived from the resin),

Another chemical employed in making perfumes is ionone, which appears in the market in the form of a 10 per cent alcoholic solution; this is used in imitating the odor of violets. In late years these chemicals really a most important part in perfumery play. manufacture.

Rose and orange flower waters are employed in some perfumes. In making handkerchief extracts, only the best imported "triple" water should be used.

### Handkerchief Extracts.

Handkerchief extracts, or "extracts," as they are more commonly termed, are produced by incorporating with alcohol one or more essences (see definition of term above), one or more spirits (alcoholic solutions of volatile oil) or possibly the oils themselves, and a tincture of some of the animal or resinous substances mentioned above. Occasionally other ingredients are added, such as one of the chemicals, which really take the place of the volatile oil or spirit. This mixture should be set aside for some time (several weeks at least) to allow it to "ripen" or blend, after which it may be filtered.

If the ingredients of handkerchief extracts be classified according to their function, the classification would be as follows: (1) Odorous agents; (2) fixing agents, and (3) vehicle.

The essential odorous agents are the essences, spirits or oils, and chemical substances. The "fixing" agents consist of the musk, ambergris, civet, resins, orris, etc. The distinction between the first and second classes is not sharply drawn, for the reason that some of the "fixing" agents are used for their odor as well as for their "fixing" prop-The third class practically embraces erties. but one substance, viz., alcohol, although rose and orange flower water are sometimes added after the alcohol, and may therefore also be considered as vehicles.

to hold the perfume to, or fix, or fasten it upon, the handkerchief or other fabric to which it is applied, it being understood that the odoriferous matters are all very volatile and therefore evanescent when exposed to the atmosphere. The "fixing" agents also serve to hold or secure the delicate flower odors to the vehicle while the moisture is still present in the container as "extract."

The "fixing" agents are of three varieties: (1) Fatty matter derived from pomades during extraction with alcohol; (2) resinous substances, such as Peru balsam, tolu, storax, benzoin, orris, etc., and even myrrh, asafetida, aloes, etc., may be employed; and (3) animal substances, which include musk, ambergris and civet.

The objection to the first class is that they soon decompose and impart to the perfume a rancid odor, and consequently pains must be taken to exclude or remove all fat from the

The objection to the second class is but an æsthetic one and hence is less serious than in the preceding. The resins impart a high color to the perfume and when the latter is dropped on a white handkerchief the fabric at once presents a soiled appearance. This is one reason why benzoin is often replaced by benzoic acid obtained by sublimation of the resin. Orris does not have the objection of imparting a high color, but the amount of resinous matter is so small that a very concentrated alcoholic tincture must be employed. Of course, all resins are objectionable because of their own odor, which may perceptibly modify the other odors, and this is sufficient reason why different resins are selected for different perfumes.

A suggestion may be thrown out here for using resins as "fixing" agents without obtaining any great amount of color, and that is to heat rather strongly a resinous substance, or suitable mixture of resinous substances, in a test tube plugged with a large wad of cot-The latter becomes impregnated with odorous substances naturally contained in the resins, also some volatile empyreumatic substances formed by the destructive distillation, and will possess a certain amount of color. The office of "fixing" agents is chiefly Upon placing in alcohol, these various principles will be dissolved, and the alcoholic liquid may be used as before for "fixing" purposes.

The third class of "fixing" agents is remarkable in that but trifling amounts will serve the purpose, and hence they are the most largely employed. The one objection to their use is their pronounced and very persistent odor, which is likely to remain after the other odorous substances have dissipated. These substances include, as stated, musk, civet, and ambergris.

In all perfumes, the appearance, next to the odor, plays the most important part. The rule is that they should be either colorless or, at the most, of a very pale tint, pale green, for example, being a popular tint for "extracts." This color should be of such a character as to disappear on drying. Essence of cassie is sometimes added to "extracts" with a view of imparting this green tint. Sometimes a very trifling amount of green aniline is employed for the same purpose; the latter is more objectionable owing to the fact that it is liable to produce stains, and also owing to the presence of a poisonous substance, arsenic, in almost all anilines. An acceptable green coloring agent for white rose extract is a tincture of patchouly made in the proportion of 4 av. ounces of the leaves to 16 fluidounces of alcohol. The resins mentioned as "fixing" agents also impart color, this being brown, but is objectionable, as stated above, owing to its staining properties.

Care should be taken to preserve all perfumes, but particularly the "extracts," with proper care. They should never be exposed to strong light; exposure to air is also prejudicial and so is exposure to heat. Cold may precipitate or separate out some of the substances present in solution in the alcoholic liquid, and hence extracts should be preserved in locations of a moderate temperature, and this temperature should also be tolerably uniform. It may therefore be stated that "extracts" should be preserved in rather small (about 8 ounce) glass-stoppered bottles, out of contact with strong light, and at a uniform, medium temperature.

Some extracts named after flowers are prepared from essences or oils derived from the

flowers, to which are added suitable diluents and fixing agents. A great many flower odors are, however, so weak that they cannot be captured, or, if successfully and economically captured, cannot be properly "fixed"; in such instances it is customary to blend other odors so as to imitate the natural product. Heliotrope is, for example, a mixture of vanilla and rose, slightly modified by means of bitter almond. The odors of lily of the valley and lilac are imitated in a similar manner. Frequently the reputed imitations are but very poor imitations indeed.

Other odors besides flower odors are imitated, such, for example, as that of new mown hay. Then, again, there are other "extracts" which are pleasing mixtures or combinations of odors, which have received arbitrary titles such as "Upper Ten," "Marie Stuart," etc., and frequently also are known as "bouquets."

A convenient and popular, though indefinite, classification of "extracts" is into "delicate odors" and "heavy odors," the former including such as violet, heliotrope, lilac, etc., the latter musk, jockey club, etc.

A few remarks on how to sell perfumes will not be amiss at this juncture. mon practice among pharmacists in showing these goods to prospective customers is to remove the stopper from the stock bottle, and allow the customer to smell directly from the latter. The vehicle alcohol being more volatile than the volatile oils which form the real perfume, obtrudes itself upon the olfactory nerves and crowds away, as it were, the more delicate odors. The proper method of showing the "extracts" and indicating the differences between them is to wet the stopper by inverting the bottle, then to moisten a strip of blotting paper or a piece of rice (cigarette) paper, allowing the latter to become almost dry before handing to the patron. Another good suggestion is to have a small sample rack of perfumes containing a number of glass-stoppered half or one ounce bottles; this will enable the pharmacist to keep the stock bottles in a dark place to be opened only when an "extract" is to be dispensed. The sample rack may also be kept in a dark place when not in use. If the glass stoppers of

the bottles of the rack are of the elongated kind, so as to reach to the bottom of the bottle, the tissue or rice paper may be dispensed with.

In showing "extracts," the order in which they are shown is also a matter of consequence, especially when the prospective customer does not have any preferences. The finest, most delicate odors should always be shown first; if the heavy odors be presented first, the olfactory nerves will have become so impressed with the latter that the delicate odors which are shown subsequently will appear to be decidedly lacking in quality, and thus a sale may be lost.

The sale of "mixed odors," that is, a mixture of two or more "extracts," should be discouraged because each "extract" is a harmonized blending of odors, and mixing "extracts" is liable to destroy this harmony. However, this matter may not be of such great consequence, because those that ask for "mixed odors" do not properly appreciate the value of a well-made "extract."

In preparing the "extracts" which follow, the subjoined essences, spirits and tinctures should be employed

It is quite essential that the amateur manufacturer of "extracts" carefully peruse the preceding pages to inform himself upon the varieties of materials used in perfumes, the method of preservation, etc.

It may be stated that the "extracts" can be cheapened by the use of weaker pomades or inferior oils, or of a tincture of musk of one-half the strength given, by the substitution of civet for the ambergris, except in the case of ambergris extract, by the addition of larger proportions of alcohol and of some water, etc.

# Essences.

Essences (extracts, extraits and esprits, they are also termed) are prepared, as already stated, by extraction of the odorous substances from pomades by means of alcohol, this process being known generally as "washing." The best method of washing is as follows:

Sixteen av.ounces of the pomade are cut into small pieces and placed in a bottle of sufficient capacity, such as a fruit jar, in

which is put 16 fluidounces of pure alcohol. Place the bottle, suitably stoppered, in a water bath, and apply heat sufficient to barely melt the pomade, shake well together, and repeat the shaking frequently until the fatty matter solidifies. In this way the pomade will be reduced to a finely divided or granular state, permeated thoroughly by the alcohol. Allow this to stand for a week-a month or even longer would be better-giving it an occasional shake, then drain off the liquid into another bottle; if this fall short of 16 fluidounces, repeat the operation with a sufficient quantity of alcohol to make up to this measure.

By subsequent and similar treatment, a second and even a third quantity of essence may be made, which, although much weaker, will be found useful in the preparation of colognes and toilet waters. Some operators use the second essence for washing a new quantity of original pomade, thus making a stronger essence or enabling one to use a weaker pomade. The residual pomade may be utilized for making hair pomades or as a diluent for mercurial or other ointments, or for making fine soaps (by manufacturers).

Essences prepared by the above method contain some fatty matter and will soon become rancid. This retained fatty matter can be separated by the application of cold, and in order that it may all be separated, the following process should be adopted in preference to the preceding one: Into an ice cream freezer of a size commensurate with the quantity of essence to be prepared, pour the requisite quantity of alcohol, then heat the pomade cautiously upon a water bath until melted, pour this into the freezer, put on the cover, set the apparatus in motion and continue the beating for 15 or 20 minutes, by which time probably all the odorous matter will have been extracted. Now surround the can with a freezing mixture, composed of ground rock salt and cracked ice, resume the beating until the liquid is thoroughly chilled and the fat has all adhered to the paddle and the sides of the can. open the freezer, pour off the liquid, and pass through a well-covered filter to separate the particles of fat still suspended in the liquid.

fluidounces of alcohol.

Spirit of Almond.	Spirit of Orange.
Oil of bitter almonds m. 80	Prepare according to the formula in Part
Alcohol, enough to makefl.oz. 16	I, from 1 fluidounce of oil of orange and 19 fluidounces of alcohol.
Spirit of Ambrette.	As stated in the introductory remarks on
Oil of ambrettefl.dr. 2 Alcoholfl.oz. 8	perfumes, the oil of bitter orange peel is to be
Spirit of Bergamot.	preferred.   Spirit of Orris.
Oil of bergamot	Oil of orris, liquidfl.dr. 1 Alcoholfl.oz. 8
Spirit of Cedarwood.	Spirit of Patchouly.
Oil of cedar (Lebanon)	Oil of patchoulyfl.dr. 4 Alcoholfl.oz. 15½
Spirit of Cinnamon.	Spirit of Pimento.
Make according to formula in Part I, from 1 fluidounce of oil of cinnamon (Ceylon) and	Oil of pimento
9 fluidounces of alcohol.	Spirit of Rose.
Spirit of Cloves.	Oil of rose         fl.dr. 4           Alcohol         fl.oz. 7½
Oil of cloves	Spirit of Rose, Compound.
Spirit of Hyacinth.	Spirit of rose goronium :fl.oz. 2
Hyacinthingr. 60	Spirit of rose geraniumfl.oz. 2 Alcoholfl.oz. 4
Alcoholfl.oz. 16 Spirit of Lavender.	Or mix
Prepare according to the formula in Part I, from 1 fluidounce of oil of lavender flowers	Oil of rose
and 19 fluidounces of alcohol.	The oil of rose geranium is added to give
Spirit of Lemon.	permanence to the spirit.
Prepare according to the formula in Part I,	Spirit of Rose Geranium.
from 1 fluidounce of oil of lemon and 19 fluid- ounces of alcohol; the lemon peel may be	Oil of rose geranium
omitted.	Spirit of Sandal.
Spirit of Lemon Grass.	Oil of sandalwoodfl.dr. 2 Alcohol, enough to makefl.oz. 16
Oil of lemon grass	Spirit of Vetivert.
Spirit of Lilac.	Oil of vetivert
Terpineol or lilacingr. 120 Alcoholfl.oz. 8	Spirit of Ylang Ylang.
Spirit of Linaloe.	Oil of ylang ylangfl.dr. 3
Oil of linaloefl.dr. 1½ Alcoholfl.oz. 8	Alcoholfl.oz. 16  Tincture of Ambergris.
Spirit of Neroli.	Ambergrisgr. 120
Oil of neroli	Orris root, powdergr. 120 Alcoholfl.oz. 16
Spirit of Nutmeg.	Rub the ambergris and orris in a mortar until reduced to a fine powder; transfer to a
Make according to the formula in Part I,	bottle, and add the alcohol. Macerate for 30
from 1 fluidounce of oil of nutmeg and 19	days, agitating occasionally, and filter through

paper

# Tincture of Ambrette.

Musk	seed.									av.oz. 4
Alcoho	1									.sufficient

Reduce the drug to fine powder, and extract by slow percolation so as to obtain 16 fluidounces of product.

# Tincture of Benzoin.

Siam benzoin,	fine	powder	av.oz.	2
Alcohol			fl.oz.	16

Mix, macerate for 30 days, agitating occasionally, and filter.

# Tincture of Civet.

Civetgr.	<b>6</b> 0
Orris root, powdergr.	
Alcoholfl.oz.	16

Proceed as with tincture of ambergris.

### Tincture of Musk.

Muskgr.	120
Lime water, warmfl.oz.	
Alcoholfl.oz.	12

Rub the musk to a fine paste with the water, macerate in a covered mortar for 2 hours, add the alcohol, and transfer to a tightly corked bottle, macerate for 30 days or longer, preferably in a rather warm place; agitate frequently.

# Stronger Tincture of Orris.

Orris root, powderav.oz. 8	
Alcoholsufficient	

Extract the drug by slow percolation so as to obtain 16 fluidounces of product.

# Weaker Tincture of Orris.

Orris root, powdergr.	600
Alcohol, enough to makefl.oz.	

Prepare like the preceding.

# Tincture of Peru Balsam.

Peru balsamav.oz.	1
Alcoholfl.oz.	15

Mix, macerate for 14 days, agitating occasionally, and filter.

# Tincture of Storax.

Storaxa	v.oz. 1
Alcohol, enough to makef	1.0z. 10

Mix, macerate for 14 days with occasional agitation, and filter.

# Tincture of Tolu.

Prepare according to formula in Part I, from 13/4 av. ounces of tolu and enough alcohol to make 16 fluidounces.

### Tincture of Tonka.

Tonka												.av.oz. 3
Alcohol	Ι.											sufficient

Reduce the bean to moderately fine powder, add 16 fluidounces of alcohol, macerate for 14 days, agitating occasionally; filter, and add enough alcohol through the filter to make 16 fluidounces of product.

# Tincture of Vanilla.

Vanillagr.	480
Sugar (granulated or rock candy).gr.	480
Alcoholfl.oz.	16

Cut the drug into small pieces, beat with the sugar in a mortar until reduced to coarse powder, macerate with the alcohol for 30 days, and filter.

# Acacia Extract.

Essence of cassie	fl.oz.	4
Stronger tincture of orris	fl.oz.	2
Tincture of vanilla	fl.dr.	12
Tincture of ambergris	. fl.dr.	4
Alcohol		

# Ambergris Extract.

Compound spirit of rosefl.oz,	3
Tincture of ambergrisfl.oz.	8
Tincture of muskfl.oz.	4
Tincture of vanillafl.oz.	1

# Clove Pink Extract.

I.		
Essence of rose	fl.oz.	(
Essence of cassie	fl.oz.	4
Essence of orange flowers	fl.oz.	4
Tincture of vanilla		

# Oil of cloves......drops 10

Essence of rose	fl.oz.	9
Essence of orange flowers	fl.oz.	41/
Tincture of vanilla	fl.oz.	31/2
Oil of cloves	m	100

# Crab Apple Blossom Extract.

Essence of violet	4
	2
	1
Essence of cassie fl.oz.	1
Spirit of ylang ylangfl.dr.	63/
Spirit of linaloefl.oz.	11/2
Spirit of neroli fl.dr.	24
Compound spirit of rosefl.dr.	4
	2
Tincture of muskfl.dr.	1
	4
Alcoholfl.oz.	1
Essence of jasmine, second	
washing, enough to make fl.oz. 1	6

Cosmos Bouquet Extract.	IV.
Essence of jasmine	Oil of bergamot       fl.oz. 1         Oil of lemon       fl.dr. 1         Oil of neroli       fl.dr. 1         Essence of reseda       fl.oz. 14
Oil of lavender         drops         9           Oil of cloves         drops         3           Coumarin         gr.         2½	Tincture of ambergris fl.dr. 4 Alcohol
Heliotropingr. 3/8 Tincture of civetfl.dr. 1/2 Weaker tincture of orrisfl.oz. 15	Essence of rose
Mix and dissolve.  Elder Flower Extract.	Oil of rose
Essence of jasmine	Oil of bergamot
Essence of tuberose	As much of the latter may be added as is possible without causing precipitation.
Spirit of ylang ylang	Esterhazy Bouquet Extract.
Tincture of ambergrisfl.dr. 2 Terpineolgr. 60	Essence of orange flowers fl.dr. 2½  Spirit of vetivert fl.oz. 2  Spirit of rose fl.dr. 1
Mix and dissolve.	Spirit of rose geraniumfl.dr. 3 Spirit of nerolifl.dr. 3
Egyptian Lotus Extract.	Spirit of sandalfl.dr. 4 Spirit of clovesfl.dr. 2
Essence of jasmine	Spirit of clovesfl.dr. 2 Tincture of tonkafl.oz. 3½
Tincture of vanilla	Tincture of vanillafl.oz. 2
Tincture of civet	Stronger tincture of orrisfl.oz. 1
Tincture of benzoinfl.dr. 5	Tincture of ambergrisfl.dr. 2
Spirit of nerolifl.dr. 7	Alcoholfl.oz. 5
Spirit of clovesfl.dr. 3½	Evening Primrose Extract.
Spirit of patchoulyfl.dr. 1	Essence of orange flowersfl.oz. 3
Oil of rose	Essence of rosefl.oz. 2
111conor	Essence of jasminefl.oz. 2
Ess. Bouquet Extract.	Compound spirit of rosefl.oz. 5
I.	Spirit of rose geranium fl.dr. 10
Compound spirit of rosefl.oz. 8	Spirit of ambrettefl.dr. 4
Spirit of lemon	Spirit of patchoulyfl.dr. 11/4
Spirit of bergamot	Tincture of benzoinfl.dr. 4 Tincture of muskfl.dr. 1
Spirit of nerolifl.oz. 1	Tincture of muskfl.dr. 1 Alcoholfl.dr. 12
Tincture of ambergrisfl.oz. 1	Alcohol
Stronger tincture of orris fl. oz. 1	Fashion Bouquet Extract.
Essence_of cassie	Oil of rose
Alcoholfl.dr. 4	Oil ot neroli
II.	Spirit of bergamot
Compound spirit of rosefl.oz. 6	Tincture of muskfl.oz. 6
Spirit of lemonfl.dr. 18	Stronger tincture of orrisfl.oz. 1
Spirit of rose geraniumfl.dr. 12	Essence of jasminefl.oz. 8
Stronger tincture of orrisfl.oz. 4½	Benzoic acidgr. 30
Tincture of ambergris	Alcohol, enough to makefl.oz. 16
Oil of bergamot	Frangipanni Extract.
	I.
III.	Essence of orange flowersfl.oz. 23/4
Stronger tincture of orrisfl.dr. 21/2	Spirit of sandalfl.oz. 4
Tincture of muskfl.dr. 1	Spirit of neroli
Essence of jasminefl.oz. 12	Spirit of rose
Oil of rosedrops 16	Tincture of muskfl.dr. 3
Oil of neroli	Tincture of tonkafl.dr. 11
Alcohol, enough to makefl.oz. 16	Alcoholfl.oz. 3½

II.	III.
Tincture of musk	Essence of jasmine
Stronger tincture of orrisfl.oz. 3	Oil of rosedrops 6 Oil of bitter almondsdrops 3
Essence of orange flowersfl.oz. 3	Tincture of muskdrops 6
Essence of tuberose	Ess. bouquet extractfl.oz. 4½ Heliotropingr. 22
Oil of rosefl.dr. 2	
Oil of rose geraniumfl.dr. 1	Mix and dissolve.
Oil of sandal	IV.
III.	Spirit of bergamotfl.oz. 6
Essence of orange flowersfl.oz. 4	Tincture of benzoin
Essence of cassie	Heliotropingr. 10
Spirit of sandalfl.dr. 17	Alcohol
Compound spirit of rosefl.dr. 22	Mix and dissolve.
Spirit of vetivert	Heliotrope (White) Extract.
Tincture of muskfl.dr. 1½	Essence of jasminefl.oz. 4
Tincture of tolu	Essence of rose
Oil of neroli bigaradedrops 30	Essence of tuberosefl.oz. 2
Frangipanni (Roman) Extract.  Musk	Spirit of ylang ylangfl.dr. 634 Tincture of civetfl.dr. 6
Musk	Tincture of muskfl.dr. 1
Vanilla, cut fine and trituratedgr. 60	Heliotropingr. 50
Tonka, bruised gr. 120 Essence of cassie fl. dr. 2	Coumarin
Essence of rosefl.dr. 2	Mix and dissolve.
Essence of orange flowers, fl.dr. 2	
Essence of tuberosefl.dr. 2 Stronger tincture of orrisfl.oz. 2½	Hesperis Extract.
Oil of rosedrops 22	Essence of cassie
Oil of cedar, Lebanondrops 22 Oil of rose geraniumdrops 10	Tincture of muskfl.dr. 4
Oil of rose geraniumdrops 10 Oil of neroli petaledrops 12	Tincture of benzoinfl.oz. 2
Oil of orange (from bitter	Tincture of tonkafl.dr. 5½ Oil of bergamotfl.dr. 2
orange) drops 3 Alcohol	Oil of clovesfl.dr. 1
Mix and macerate for several months, agi-	Spirit of lavender fl.oz. 2½ Spirit of rose
tating occasionally.	Spirit of rose
Flowers of Ireland Extract.	Alcohol, enough to makefl.oz. 16
White rose extractfl.oz. 15	Honeysuckle Extract.
Tincture of vanillafl.dr. 12	Essence of rosefl.oz. 4
Heliotrope Extract.	Essence of violet
I.	Essence of tuberosefl.oz. 4
Tincture of ambarrais flor 1	Tincture of vanilla
Tincture of ambergrisfl.oz. 1 Tincture of civetfl.oz. 1	Tincture of musk
Compound spirit of rosefl.oz. 3	Spirit of bitter almondfl.dr. 3¼ Spirit of nerolifl.dr. 1½
Oil of bitter almonddrops 5 Essence of rosefl.oz. 3	Alcohol
II.	
Essence of jasminefl.oz. 1	Hyacinth Extract.
Tincture of vanillafl.oz. 4 Tincture of muskfl.oz. 1	Hyacinthin
Tincture of storaxfl.dr. 1	Tincture of muskfl.dr. 2½
Spirit of nerolifl.dr. 5½	Tincture of benzoinfl.dr. 5
Spirit of almond	Essence of jasminefl.oz. 3 Alcoholfl.oz. 12
Alcohol	Orange flower water, triplefl.oz. 1½

Jockey Club Extract.	Kiss-Me-Quick Extract.
I.  Compound spirit of rose. fl.oz. 4 Essence of rose. fl.oz. 1 Essence of tuberose. fl.oz. 4 Essence of cassie. fl.oz. 2 Essence of jasmine. fl.oz. 1 Essence of orange flowers fl.oz. 1 Tincture of civet fl.oz. 2 Tincture of musk. fl.oz. 1 II.	Essence of cassie       fl.oz.       2½         Essence of jasmine       fl.dr. 10         Essence of tuberose       fl.dr. 10         Tincture of vanilla       fl.dr. 2½         Tincture of benzoin       fl.dr. 1¾         Stronger tincture of orris       fl.dr. 1¾         Tincture of musk       m. 50         Spirit of bergamot       fl.dr. 7         Spirit of neroli       fl.dr. 5½         Alcohol       fl.oz. 9
Essence of tuberose. fl.oz. 5½ Essence of cassie. fl.oz. 2½ Essence of jasmine. fl.oz. 1½ Compound spirit of rose. fl.oz. 3½ Spirit of ambrette. fl.oz. 1 Spirit of neroli fl.dr. 2¼ Tincture of civet. fl.dr. 4 Alcohol, enough to make fl.oz. 16	Lilac Extract.  I.  Essence of tuberose
III.  Essence of orange flowersfl.oz. 4 Essence of cassiefl.oz. 2 Essence of jasminefl.oz. 2 Compound spirit of rosefl.oz. 4 Spirit of rose geraniumfl.dr. 15 Spirit of ambrettefl.dr. 4 Spirit of bergamotfl.dr. 1½ Spirit of clovesfl.dr. 1 Tincture of muskfl.dr. 1 Alcohol, enough to make .fl.oz. 16	H. Essence of jasmine fl.oz. 3 Essence of tuberose fl.oz. 2 Essence of rose fl.oz. 2 Spirit of lilac fl.oz. 4 Spirit of hyacinth fl.dr. 10½ Spirit of ylang ylang fl.dr. 6½ Tincture of civet fl.dr. 6 Tincture of musk fl.dr. 1 Heliotropin gr. 20 Alcohol, enough to make fl.oz. 16
*IV.  Essence of jasmine	Essence of tuberose. fl.oz. 2½ Essence of orange flowers fl.oz. 3½ Essence of violet fl. dr. 10 Essence of cassie fl.oz. 1 Spirit of ylang ylang fl.dr. 27 Tincture of civet fl.oz. 7  IV. Essence of tuberose fl.oz. 4 Essence of orange flowers. fl.oz. 1
Essence of jasmine fl.dr. 10 Essence of violet fl.dr. 4 Tincture of musk fl.dr. 2 Tincture of vanilla fl.dr. 4½ Spirit of rose fl.dr. 6 Spirit of sandal fl.oz 3 Spirit of neroli fl.dr. 5½ Benzoic acid gr. 30 Alcohol fl.oz 14 Mix and dissolve.	Tincture of civet
VI. Oil of rose	Essence of tuberose. fl.oz. 8 Essence of jasmine fl.oz. 1 Essence of orange flowers fl.oz. 1 Essence of cassie fl.oz. 2 Essence of rose fl.oz. 2 Compound spirit of rose fl.oz. 1 Oil of bitter almonds drops 2 Tincture of vanilla fl.oz. 1

II.	Locust Blossom Extract.
Essence of rose	Essence of jasminafl.oz. 3
Essence of violet	Essence of cassie
Essence of jasminefl.oz. 1	Tincture of vanilla
Essence of cassiefl.oz. 1	Tincture of civet
Spirit of linaloe	Alcohol
Spirit of ylang ylangfl.dr. 6½	
Compound spirit of rosefl.dr. 4 Spirit of nerolifl.dr. 2½	May Bells Extract. (Mabel Bouquet
Spirit of neroli	Extract.)
Oil of mace, volatiledrops 6	I.
Alcohol	Essence of jasminefl.oz. 31/4
Essence of rose, second wash-	Ylang ylang extract No. 1fl.oz. 3½
ing, enough to makefl.oz. 16	Stronger tincture of orrisfl.oz. 23/4
III.	Fluid extract of cardamomm. 30
Essence of tuberosefl.dr. 21	Alcohol
Essence of tuberose	
Essence of jasminefl.dr. 2½	The fluid extract may be replaced by 3
Essence of orange flowersfl.dr. $5\frac{1}{2}$	drops of oil of cardamom.
Spirit of rose	II.
Spirit of rose geraniumfl.dr. 11	Essence of jasminefl.oz. 12
Spirit of almondfl.dr. 2	Ylang ylang extractfl.oz. 4
Alcohol fl.oz. 9	Orris root, granulatedav.oz. 2
IV.	Cardamom seed, powdergr. 60
Essence of jasminefl.oz. 6	
Cardamom, powdered without	Mix, let stand 7 days, agitating occasion-
membranes	ally, and filter.
Spirit of ylang ylangfl.dr. 21	Marie Stuart Extract.
Alcohol	I.
, -	Essence of cassiefl.oz. 2
Mix, macerate for several weeks or a month,	Essence of orange flowersfl.oz. 2
agitating frequently, and filter.	Essence of rosefl.oz. 2
V.	Compound spirit of rosefl.oz. 4
Essence of jasminefl.oz. 20	Spirit of bergamotfl.dr. 12
Cardamom, powdered without	Spirit of rose geraniumfl.dr. 10
membranesgr. 60	Spirit of patchoulyfl.dr. 21/4
Oil of ylang ylangfl.dr. 1	Tincture of musk
Weaker tincture of orrisfl.oz. 4	Tincture of tonkafl.dr 5½
Prepare like the preceding.	Tincture of tolufl.dr. 5 Tincture of benzoinfl.dr. 4
Lavender Extract.	Coumarin
	Oil of verbenam. 15
Essence of rosefl.oz. 2	Alcohol fl.oz. 1
Spirit of lavenderfl.oz. 10	
Alcoholfl.oz. 4	Mix and dissolve.
Lily (White) Extract.	II.
Essence of rosefl.dr. 19	Essence of rose
Essence of orange flowersfl.dr. 9½	Essence of jasmine
Essence of cassie	Tincture of muskfl.dr. 3½
Tincture of vanillafl.dr. 14	Tincture of ambergrisfl.dr. 7
Spirit of clovesfl.dr. 4½	Tincture of vanillafl.dr. 7
Alcoholfl.oz. 9	Stronger tincture of orrisfl.dr. 7
Liriodendron Extract.	Alcohol, enough to makefl.oz. 16
Essence of cassiefl.dr. 5½	Mousseline Extract.
Essence of tuberosefl.dr. $5\frac{1}{2}$	
Essence of orange flowers fl.dr. 5½  Essence of orange flowers fl.dr. 5½	Esterhazy bouquet extractfl.oz. 5 Essence of cassiefl.dr. 7
Tincture of benzoinfl.oz. 1	
Tincture of civet	Essence of jasmine
Tincture of orris	Spirit of sandal
Spirit of bergamotfl.oz. 2	Compound spirit of rosefl.dr. 13
Spirit of rosefl.dr. 5	Spirit of rose geraniumfl.dr. 3½
Alconol	Alcohol

Millefleurs Extract.	II.
Compound spirit of rosefl.oz. 3	Tincture of tonkafl.oz. 12
Essence of rose	Essence of rose 2
Essence of jasmine	Essence of jasmine fl.oz. 2
Essence of orange flowersfl.oz. 2	Oil of rose geraniumfl.dr. 1
	Oil of rosedrops 15
Essence of cassie	Oil of nerolidrops 10
	III.
Tincture of tonkafl.oz. 1	
Tincture of ambergrisfl.dr. 4	
Tincture of musk	
Oil of bergamot	J
Oil of bitter almonddrops 3	Compound spirit of rosefl.oz. 3
Oil of nerolidrops 3	Spirit of ambrettefl.oz. 2
Oil of clovesdrops 3	Spirit of lavenderfl.dr. 10
Musk Extract.	Tincture of vanillafl.oz. 2
	Tincture of tonkafl.dr. 5½
I.	Tincture of civetfl.dr. 4
Tincture of muskfl.oz. 11	Tincture of benzoinfl.dr. 1½
Tincture of civet	Oil of limettedrops 30
Compound spirit of rosefl.oz. 4	IV.
This is a rather high-priced article, but the	Essence of orange flowersfl.oz. 3½
	Essence of tuberosefl.dr. 7
tincture of musk may be reduced one-half	Spirit of rosefl.dr. 5
with alcohol and still yield a satisfactory	Spirit of nerolifl.dr. 5
product.	Tincture of tonkafl.dr. 7
•	Alcohol
II.	
Essence of orange flowersfl.oz. 2	Night-Blooming Cereus Extract.
Essence of cassiefl.oz. 2	Essence of jasminefl.oz. 4
Spirit of rose	Tincture of tonka
Tincture of muskfl.oz. 1½	Tincture of civetfl.oz. 2
Tincture of civet	Tincture of benzoinfl.oz. 4
Tincture of vanillafl.dr. 4	Spirit of rosefl.oz. 1
Tincture of tonkafl.dr. 6	Spirit of rose geraniumfl.oz. 1
Tincture of tolufl.dr. 5	
Tincture of benzoinfl.dr. 4	Ocean Spray Extract. ("Sea Breeze"
Alcoholfl.oz. 5	Extract.)
III.	Forence of oronge flowers flow 6
Muskgr. 288	Essence of orange flowersfl.oz. 6
Civetgr. 54	Essence of jasminefl.oz. 3
Ambergrisgr. 36	Compound spirit of rosefl.oz. 5
Musk seedav.oz. 6	Spirit of ambrettefl.dr. 4
Alcohol	Spirit of clovesfl.dr. 414
Triturate the muck seed to powder add	Tincture of muskfl.dr. 1½
Triturate the musk-seed to powder, add	Tincture of vanillafl.dr. 6
the alcohol, macerate in a warm place for 2	Orange Flower Extract.
or 3 weeks, agitating frequently; place into	-
a percolator, and percolate until 15 fluid-	I.
ounces of liquid have been obtained. To	Essence of orange flowersfl.oz. 12
	Escence of cassie flor 2
this liquid, add the musk, civet and amber-	Tincture of muskfl.oz. 2
gris, making an intimate mixture; place the	
whole away in a well-stoppered bottle in a	II.
warm place for 2 months, agitating fre-	Essence of orange flowersfl.oz. 434
	Stronger tincture of orris fl.dr. 14
quently, and filter.	Tincture of muskfl.dr. 1/2
This is reputed to produce a very fine	Spirit of nerolifl.dr. 1
musk extract.	Alcohol, enough to makefl.oz. 16
	Patchouly Extract.
New Mown Hay Extract.	
I.	I.
Coumaringr. 8	Spirit of patchoulyfl.oz. 5
Vanillingr. 4	Spirit of rosefl.dr. 4
Weaker tincture of orrisfl.oz. 16	Alcohol
canci difference of offis,	

II.	Rose (Musk) Extract.
Spirit of patchouly fl.dr. 8½ Spirit of bergamot fl.oz. 2 Essence of jasmine fl.oz. 4 Essence of rose fl.oz. 2 Tincture of benzoin fl.dr. 2 Alcohol, enough to make fl.oz. 16  III. Patchouly leaves, cut av.oz. 5 Alcohol	Essence of rose.fl.dr. $9\frac{1}{2}$ Essence of tuberose.fl.dr. $4\frac{1}{2}$ Essence of jasmine.fl.dr. $4\frac{1}{2}$ Stronger tincture of orris.fl.dr. $14$ Tincture of musk.fl.dr. $7$ Spirit of orange.fl.dr. $1\frac{1}{2}$ Alcohol, enough to make.fl.oz. $16$
Alcohol fl. oz. 18 Water fl. oz. 16	Rose (Tea) Extract.
Salt	Essence of rose
fluidounces of distillate are obtained, and to	II. Essence of rosefl.dr. 10½
Pooch Blossom Extract	Compound spirit of rosefl.dr. 22
Essence of orange flowers         fl.oz.         3½           Essence of tuberose         fl.dr.         7           Spirit of lemon         fl.dr.         13           Spirit of almond         fl.oz.         2½           Tincture of Peru balsam         fl.dr.         5½           Alcohol         fl.oz.         6	Spirit of rose geranium
Rondeletia Extract.	I. Compound spirit of rose flor 8
I. Tincture of musk. fl.dr. 4 Tincture of ambergris. fl.dr. 4 Tincture of vanilla. fl.dr. 4 Spirit of lavender fl.oz. 5 Spirit of cloves fl.oz. 4 Spirit of bergamot fl.oz. 2 Spirit of rose fl.oz. 1 Alcohol fl.oz. 2	Compound spirit of rose. fl.oz. 8 Essence of rose. fl.oz. 3 Essence of jasmine. fl.oz. 4 Patchouly extract, No. 1 fl.oz. 1  II. Oil of rose. fl.dr. 1½ Spirit of rose geranium fl.oz. 1½ Essence of rose fl.oz. 3 Essence of jasmine fl.oz. 1½
II.	Tincture of musk
Spirit of lavender. fl.oz. 3 Spirit of cloves. fl.oz. 3 Spirit of bergamot. fl.cz. 1½ Compound spirit of rose. fl.dr. 10 Spirit of rose geranium. fl.dr. 5 Tincture of musk. fl.dr. 6 Tincture of ambergris. fl.dr. 1½ Tincture of vanilla. fl.dr. 2¼ Alcohol, enough to make. fl.oz. 15	Alcohol
III. Spirit of hercamet A or 2	IV.
Spirit of bergamot         fl.oz. 2           Spirit of lemon         fl.oz. 2½           Spirit of cloves         fl.oz. 4           Spirit of lavender         fl.oz. 2½           Spirit of neroli         fl.oz. 1           Spirit of rose         fl.dr. 2½           Alcohol         fl.oz. 4	Essence of violet fl. oz. 4 Essence of jasmine fl. oz. 2 Compound spirit of rose fl. oz. 8 Spirit of patchouly fl. dr. 2½ Tincture of ambergris fl. oz. ½ Alcohol fl. oz. 1
Rose (Moss) Extract.	V. Essence of rose
Compound spirit of rose. fl.oz. 9 Essence of orange flowers. fl.oz. 3 Essence of rose. fl.oz. 2 Tincture of civet fl.oz. 1 Tincture of musk fl.oz. 1	Essence of violet

VI.	Sweet Pea Extract.
Spirit of rosefl.oz. 1	Essence of tuberosefl.oz. 5
Spirit of cedar	Essence of orange flowersfl.oz. 5
Spirit of patchoulyfl.dr. 4 Spirit of orangefl.dr. 2½	Essence of rosefl.oz. 5 Tincture of tonkafl.oz. 2
Essence of tuberosefl.dr. 4	
Essence of violet	Upper Ten Extract.
Tincture of muskfl.dr. 4	Tincture of vanillafl.oz. 4 Tincture of ambergrisfl.oz. 3
Benzoic acidgr. 15	Stronger tincture of orrisfl.oz. 3
Alcohol	Compound spirit of rosefl.oz. 3
VII.	Essence of orange flowersfl.oz. 3 Oil of bergamotfl.dr. 1½
Spirit of rose	Oil of lemondrops 15
Spirit of patchoulyfl.dr. 4	Verbena Extract.
Spirit of pimento	I.
Essence of jasmine	Essence of orange flowersfl.oz. 3 Essence of tuberosefl.oz. 3
Rose water, triplefl.dr. 10	Compound spirit of rosefl.oz. 3
Sweet Brier Extract.	Spirit of lemon grassfl.dr. 13½
Oil of bergamotfl.dr. 6	Spirit of neroli
Oil of lemon	Oil of orangefl.dr. 23/4
Oil of lavenderfl.dr. 4 Oil of verbenadrops 8	Alcoholfl.oz. 3
Spirit of rosefl.oz. 2	II. Oil of orangedrops 30
Spirit of almondfl.dr. 12½ Tincture of muskfl.oz. 2	Spirit of lemonfl.oz. 10
Alcohol	Spirit of lemon grassfl.oz. 6
Rose (Yellow) Extract.	III.
Essence of rosefl.oz. 2	Spirit of lemon
Essence of tuberosefl.oz. 2	Stronger tincture of orrisfl.oz. 1
Tincture of tonkafl.dr. 10 Verbena extractfl.dr. 2¼	Alcohol
Alcohol	IV.
Rose Geranium Extract.	Spirit of lemon
Oil of rose geraniumfl.oz. 1	Oil of orangefl.dr. 10
Alcohol	Violet Extract.
Spring Flowers Extract.	I.
Essence of rosefl.oz. 7	Essence of violet
Essence of violet	Tincture of muskfl.oz. 1
Compound spirit of rosefl.oz. 1	Stronger tincture of orrisfl.oz. 2
Tincture of ambergris	Essence of cassiefl.oz. 6
	Essence of rosefl.oz. 3
Stephanotis Extract.  Essence of cassiefl.dr. 5¼	Essence of tuberosefl.oz. 3
Essence of tuberosefl.dr. 5¼	Stronger tincture of orrisfl.oz. 3 Spirit of almondfl.oz. 1
Essence of jasminefl.dr. 23/4	III.
Stronger tincture of orrisfl.oz. 4 Tincture of tonkafl.oz. 1	Essence of violetfl.oz. 434
Tincture of muskfl.oz. 1	Essence of cassiefl.dr. 24 Stronger tincture of orrisfl.dr. 34
Spirit of rose	Stronger tincture of orrisfl.dr. $3\frac{1}{2}$ Spirit of almondfl.dr. $1$
Benzoic acidgr. 30	Alcohol
Alcohol, enough to makefl.oz. 16	IV.
Tuberose Extract.	Essence of cassie
Essence of tuberosefl.oz. 15	Essence of violet
Tincture of ambergrisfl.oz. 1	Essence of tuberosefl.oz. $2\frac{1}{2}$

Violet de Parme Extract. (Parmese	West End Extract.
Violet Extract.)	I.
I.	Essence of jasminefl.oz. 1
Essence of violetfl.oz. 7	Essence of cassiefl.oz. 1
Essence of cassiefl.oz. 4	Stronger tincture of orrisfl.oz. 3
Spirit of orris	Tincture of muskfl.oz. 2
Compound spirit of rosefl.oz. 1	Tincture of storaxfl.dr. 2
Spirit of almondfl.dr. 3/4	Spirit of rosefl.dr. 4
Tincture of civet	Spirit of cedarfl.dr. 5
Alcohol	Spirit of nerolifl.dr. 4
	Oil of verbenadrops 4
II.	Benzoic acidgr. 15
Essence of cassie	Alcohol, enough to makefl.oz. 16
Stronger tincture of orrisfl.oz. 3	II.
Spirit of rosefl.dr. 4	Essence of jasminefl.oz. 1
Spirit of almond	Essence of cassiefl.oz. 1
Alcohol	Stronger tincture of orrisfl.oz. 3
	Tincture of musk
Violet (Wood) Extract.	Tincture of storaxfl.dr. 2
	Spirit of rosefl.dr. 4
Violet extract, No. IIfl.oz. 16	Spirit of cedarfl.dr. 5
Oil of bitter almonddrops 15	Spirit of nerolifl.dr. 4
I.	Oil of verbenadrops 4
Essence of violetfl.dr. 13	Benzoic acidgr. 15
Essence of cassiefl.dr. 10	Alcohol, enough to makefl.oz. 16
Essence of rosefl.dr. $6\frac{1}{2}$	View or View or Finders of
Essence of tuberosefl.dr. $6\frac{1}{2}$	Ylang Ylang Extract.
Stronger tincture of orrisfl.dr. 10	I.
Spirit of almondfl.dr. $1\frac{1}{2}$	Spirit of ylang ylangfl.oz. 8
Alcohol fl.oz. 9	Compound spirit of rosefl.oz. 4
II.	Essence of jasminefl.oz. 2
Essence of violetfl.oz. 8	Tincture of civetfl.oz. 2
Essence of cassiefl.oz. 1	II.
Essence of tuberosefl.oz. 1	
Tincture of vanillafl.dr. 9	Essence of jasminefl.dr. 21
Tincture of tonkafl.dr. $2\frac{1}{2}$	Essence of tuberosefl.dr. 14 Essence of orange flowersfl.dr. 7
Tincture of muskfl.dr. 6	
Spirit of rose	Spirit of ylang ylangfl.dr. 11 Alcoholfl.oz. $9\frac{1}{2}$
Spirit of bergamotfl.dr. 4	
Spirit of neroli	III.
Alcohol	Oil of ylang ylangdrops 15
Widow McPhelan Extract.	Oil of nerolidrops 2
Essence of cassiefl.oz. 2	Oil of rosedrops 5
Essence of violetfl.oz. 1	Oil of lemondrops 2
Stronger tincture of orrisfl.oz. 3	Tincture of muskfl.dr.
Spirit of nutmegfl.dr. $5\frac{1}{2}$	Alcoholfl.oz. 16
Spirit of pimentofl.dr. 4½	IV.
Spirit of rosefl.dr. 2	Essence of jasminefl.oz. 2
Spirit of cinnamonm. 30 Ylang ylang extract No. IIfl.oz. 3	Essence of violetfl.oz. 2
Ylang ylang extract No. IIfl.oz. 3	Spirit of rosefl.dr. 2
Alcoholfl.oz. 6	Spirit of ylang ylang fl.oz. 51/4
Vild Olive Extract.	Tincture of civetfl.oz. 2
	Alcohol, enough to makefl.oz. 16
Essence of rosefl.oz. 4	57
Essence of violet	V.
Essence of jasminefl.oz. 2	Essence of jasminefl.oz. 2
Essence of cassiefl.oz. 1	Essence of rosefl.oz. 4
Spirit of bergamotfl.oz. 1	Weaker tincture of orrisfl.oz. 6
Spirit of rose	Tincture of civetfl.oz. 1
Spirit of lavenderfl.dr. 2½	Oil of ylang ylangfl.dr. 1
Alcohol, enough to makefl.oz. 16	Alcoholfl.oz. 4

# Sachet Powders.

As is well known sachet powders are mixtures, in the form of moderately fine powder, which are to be inclosed in little sacks of cloth and placed with linen or wearing apparel, or stationery, etc. Sachets are preferred by some to "extracts," because there is no fear of using too much and thus making the user appear "loud" or vulgar. The objection to sachet powders is the want of permanency; they are liable to lose their odor even if carefully preserved, and the purchaser may therefore receive a sachet powder which can not be compared, in strength or delicacy, with a good "extract." For this reason sachet powders are frequently "freshened" by the addition of the corresponding "extract," viz., violet sachet by violet extract,

Sachet powders are composed of two kinds of ingredients, viz., the "body" or vehicle, and the odorous agents. The first almost invariably contains orris root; this may be the only "body," or it may be combined with rose petals, orange peel, or lavender flowers, or a mixture of several of these. The odorous agents are the same as those employed in making "extracts." Remarks made with reference to the production of the latter will in a large measure apply to the sachet powders. For example, the best ingredients are required to make good sachet powders, whereas indifferent ingredients cannot but produce poor products. All such substances as orris, orange peel, etc., must be perfectly fresh and odorous, and the oils, etc., must be of superior quality.

In preparing sachet powders, the orris, rose petals, lavender flowers, and similar solid ingredients should be mixed and ground in a mill; the musk, civet, ambergris, vanilla, and tonka should be triturated to an intimate mixture with a portion of this powder; the solid resins or gum resins, like benzoin, should be contused in a mortar until reduced to moderately fine powder; all these ingredients should now be mixed, placed in a large mortar, the oils, tinctures, and other liquids added, and the whole mixed intimately by trituration. When the quantity of volatile oil is very small, it may be

advantageous first to dissolve in a small amount of alcohol.

Sachet powders should be preserved in rather small, well-stoppered bottles in a location of moderate temperature and be protected from light.

# Cassie or Acacia Sachet.

Cassie	flowers.	 	 				.av.oz.	8
Orris r	oot	 	 				.av.oz.	8

Pass each separately through a mill to reduce to tolerably fine powder, then mix, and pass through mill again, to reduce to finer powder.

# Clove Pink Sachet.

Orris rootav.oz.	8
	4
ratementy reares	3
Clovesav.oz.	1
Deer tongueav.oz.	1
Pimentoav.oz.	$\frac{1}{2}$
Muskgr.	3
Oil of rosedrops 4	)
Oil of nerolidrops 4	3
Oil of sandalwooddrops 8	)
Oil of lavenderdrops 4	0

Mix the first six ingredients, grind to a moderately fine powder, triturate the musk to an intimate mixture with a portion of this powder, add the remainder of the powder and the oils, and mix the whole thoroughly.

The deer tongue may be replaced by tonka if desired.

# Essence Bouquet Sachet.

1.		
	Orange peel, recently driedav.oz.	4
	Sandalwood av.oz.	4
	Rose petalsav.oz.	4
	Orris rootav.oz.	4
	Muskgr.	2
	Coumaringr.	4
	Vanillingr.	4
	Oil of rosedrops	12
	Oil of bergamotdrops	12
	Oil of nerolidrops	5
	Oil of ylang ylangdrops	5
	Oil of rose geraniumdrops	4
	Oil of cassiadrops	5
	Oil of bitter almonddrops	3
	Essence of jasminefl.oz.	1

Mix the first four ingredients, grind to powder in a mill, triturate the musk, coumarin, and vanillin with a portion of the ground material, add to the remainder of the powder, now add the oils and essence, and again mix well.

PEX
II.
Orris powder
of the orris and the oils, and triturate to a
intimate mixture.
Frangipanni Sachet.
I. Orris root. av.oz. 8 Rose petals av.oz. 8 Vanilla av.oz. ½ Benzoin av.oz. ½ Oil of lavender fl.dr. 1 Oil of bergamot fl.dr. 1 Oil of cassia drops 24 Oil of pimento drops 40 Oil of sandalwood fl.dr. 2 Oil of neroli fl.dr. 1 Oil of rose drops 32 Grind the orris and rose separately to coarse powder, mix, pass through mill again to reduce to somewhat finer powder, triturate the vanilla to powder with a portion of this
mixture, also reduce the benzoin to powder
mix the whole, add the oils, and triturate to an intimate mixture.  II.  Orris root
Orris, powder         av.oz. 12           Vetivert         av.oz. 1           Sandalwood         av.oz. 1           Vanilla         av.oz. 1           Tonka beans         av.oz. ½           Oil of neroli         drops 15           Oil of sandalwood         drops 10           Oil of bergamot         drops 15           Oil of rose geranium         drops 15           Oil of rose         drops 8           Tincture of musk         fl.dr. 2           Tincture of civet         fl.dr. 1           Prepare like the preceding.

_		~00
	Heliotrope Sachet.	
	I.	
	Rose petalsav.oz.	74
ì	Orris rootav.oz.	$3\frac{2}{3}$
	Lavender flowersav.oz.	334
	Tonka beansav.oz.	1
Į	Benzoin av.oz.	1
Į	Musk	$90^{\frac{7}{2}}$
Į	Oil of bitter almonddrops	10
Į	Oli of sandalwoodfl.dr.	2
ļ	Oil of nerolidrops	
ļ		40
l	Prepare like the preceding.	
Į	II.	
۱	Orrisav.oz.	8
l	Red rose petalsav.oz.	4
١	Tonka	2
l	Vanillaav.oz.	1
l	Muskgr.	
١	Oil of bitter almonddrops	2
	Mir the free true in medicate and i	1.
l	Mix the first two ingredients and grin	d to
	powder in a mill; contuse the vanilla, to	
	and sufficient of the orris root together i	ıntil
	the first two are reduced to a fine pow	
	triturate the musk with a portion of	
	powder, add the remaining ingredients,	and

III.	
Orrisav.oz.	4
	4
Sandalwood av.oz.	-
Orange peel, recently driedav.oz.	7
	1
Vanillin gr.	2
Coumaringr.	į
Tincture of ambergrisdrops	6
Oil of rosedrops 1	2
	2
Oil of nerolidrops	3

mix the whole intimately in a mortar.

Reduce the first four ingredients to powder by grinding in a mill, add the other ingredients and mix well.—D.

Oil of ylang ylang drop 1
Spirit of almond fl.dr. 1
Essence of jasmine fl.dr. 4

# Jockey Club Sachet.

I.		
	Lavender flowersav.oz.	2
	Rose petalsav.oz.	6
	Orrisav.oz.	8
	Vanillagr.	120
	Muskgr.	16
	Essence of jasminefl.oz.	1
	Oil of sandalwooddrops	
	Oil of nerolidrops	
	Oil of rosedrops	
	Doduge the first three insmediane	

Reduce the first three ingredients to powder by grinding in a mill, contuse the vanilla with a portion of the ground mixture

to powder; also triturate the musk with a portion of the powder intimately, add the oils and essence, and mix the whole thoroughly by trituration in a mortar.

### II.

Orris, powderav.oz.	12
Sandalwood, groundav.oz.	2
Oil of bergamotfl.dr.	
Oil of rosedrops	
Tincture of muskfl.dr.	4
Tincture of civetfl.dr.	2

Mix the orris and sandal, add the other ingredients, and triturate until well mixed.

### III.

~~*	
Orrisav.oz.	4
Rose petalsav.oz.	4
Orange peel, recently driedav.oz.	63
Sandalwood gr.	140
Cloves	35
Sumatra benzoingr.	375
Tincture of civetdrops	10
Tincture of muskdrops	5
Coumarin gr.	1/2
Oil of rosedrops	$7^{\frac{1}{2}}$
Oil of bergamotdrops	10
Oil of rose geraniumdrops	3
Oil of nerolidrops	2
Oil of cassiadrop	1
Oil of corianderdrop	1
Oil of bitter almonddrop	1
Oil of ylang ylangdrop	ī
Essence of jasminefl.dr.	$\bar{6}$
and the second of the second o	U

Mix the first four ingredients, reduce to powder by grinding in a mill, contuse the benzoin to powder, and then mix all the ingredients thoroughly by trituration in a mortar.—D.

# Lavender Sachet.

I.	
Lavender flowersav.oz.	13
Benzoin	
Oil of lavenderfl.dr.	$1\frac{1}{2}$

Reduce the lavender and benzoin each to powder, mix, add the oil, and triturate until well mixed.

# II.

A •	
Lavender flowersav.oz.	13
Benzoinav.oz.	
Oil of bergamotfl.dr.	11/
Oil of lavenderfl.dr.	24
Prenare like the preceding —H	-

# III.

1.	
Lavender flowersav.oz.	16
Benzoinav.oz.	1
Oil of lavenderfl.dr.	4
Tincture of muskfl.dr.	4
Prepare lile the preceding.	

# Marechale Sachet.

Sandalwoodav.oz.	41
Orrisav.oz.	33/
Rose petalsav.oz.	3
Clovesav.oz.	$2\frac{1}{4}$
Cassia barkav.oz.	
Tincture of muskdrop	1

Reduce the first five ingredients to moderately fine powder by grinding in a mill, add the tincture of musk and mix well by trituration.—H.

# Millefleurs Sachet.

Lavender flowersav.oz	. 3
Clovesav.oz.	1
Cassia budsav.oz.	
Coriander	
Benzoingr.	
Nutmeggr.	
Orrisav.oz.	8
Vanillaav.oz.	$\frac{1}{2}$
Musk	20
Oil of rosedrops	
Oil of nerolidrops	16
Oil of patchoulydrops	8
Oil of lavender (English)drops	16
Oil of verbenadrops	. 8
Oil of sandalwooddrops	40

Contuse the orris and nutmeg, add the lavender, cloves, cassia, and coriander, grind all together in a mill to moderately fine powder, triturate the vanilla and musk each with a portion of this powder until well mixed, contuse the benzoin to powder, mix the whole, add the oils and mix all thoroughly by trituration.

# H.

Lavender flowersav.oz.	21
Orrisav.oz.	
Rose petalsav.oz.	$2\frac{7}{2}$
Benzoinav.oz.	$2\frac{1}{2}$
Tonkaav.oz.	$2\frac{1}{2}$
Vanillaav.oz.	
Sandalwoodav.oz.	
Clovesav.oz.	
Cardamonav.oz.	
Cassia barkav.oz.	
Muskgr.	7

Grind the lavender, orris, rose petals, sandal, cloves, cardamom, and cassia together in a mill to moderately fine powder, triturate the vanilla and tonka with a portion of this mixture until reduced to powder, also triturate the musk with another portion until well mixed, contuse the benzoin to fine powder, and mix the whole together thoroughly.—II.

II.	
Lavender flowersav.oz.	$2\frac{1}{2}$
Cassia flowersav.oz.	$2\frac{1}{2}$
Rose petalsav.oz.	$2\frac{1}{2}$
Orrisav.oz.	5
Sandalwoodav.oz.	1 1/4
Cloves gr.	135
Cinnamon gr.	135
Benzoin gr.	270
Tonka gr.	270
Vanilla gr.	200
Tincture of civet	40
Tincture of musk	40
Oil of bergamotm.	40
Oil of rose geraniumdrops	5
Oil of patchoulydrops	2

Mix the first seven ingredients, grind together in a mill to moderately fine powder, contuse the benzoin to fine powder, triturate the vanilla and tonka with some of the ground material to fine powder, mix all three, add tinctures and oils, and mix the whole intimately by trituration in a mortar.

# Musk Sachet.

Oil of rosedrops	2
Ammonium carbonategr.	6
Muskgr.	
Orris, powderav.oz.	16
Mix intimately. The proportions ma	
altered if desired.	

# New Mown Hay Sachet.

I.	
Orrisav	.oz. 4
Rose petalsav	.oz. 4
Orange flowersav	.oz. 2
Musk seedav	.oz. 2
Tonkaav	.oz. 2
Benzoin	gr. 290
Oil of verbenadr	
Oil of bitter almonddr	ops 15

Grind the orris, rose petals, orange flowers, and musk seed together in a mill to moderately fine powder, triturate the tonka with a portion of this to fine powder, also contuse the benzoin to powder, mix all, add the oils, and mix the whole intimately by trituration. 11

1.	
Orris, powderav.oz.	13
Tonkaav.oz.	11
Vanillaav.oz.	11
Oil of bitter almonddrops	2
Oil of rose geraniumdrops	24
Oil of rosedrops	6
Oil of bergamotdrops	12
Tincture of muskfl.dr.	21

Triturate the tonka and vanilla with the orris to fine powder, add the oils and tinc- moderately fine powder in a mill, add the ture, and mix well by trituration.

IJ	11.	
	Deer tongue leavesav.oz. 8	3
	Orrisav.oz. 4	Ł
	Orange flowersav.oz. 4	Į
	Rose petalsav.oz.	$\frac{1}{2}$
	Mix and grind to moderately fine power	

in a mill.

# Opoponax Sachet.

Orrisav.oz.	8
Rose petalsav.oz.	24
Cassie flowersav.oz.	$2\frac{1}{2}$
Tonkagr.	290
Vanilla av.oz.	1
Musk	75
Oil of citronelladrops	3
Oil of lemondrops	5
Oil of bergamotdrops	20
Oil of patchoulydrops	5
Oil of rose geraniumdrops	10
Oil of rosedrop	1
Tincture of civetm.	40

Mix the first three ingredients, grind to moderately fine powder in a mill, triturate the vanilla and the tonka with a portion of this powder until a fine mixture is produced, add the remainder of the ground mixture, the oils and the tincture, and mix well by trituration.

# Oriental Sachet.

Orris:av.oz.	24
Calamusav.oz.	$2\frac{1}{2}$
Sandalwood av.oz.	11/4
Rosewood av.oz.	
Clovesgr.	270
Cassia gr.	
Orange peel, recently driedav.oz.	$2\frac{1}{2}$
Rose petalsav.oz.	$2\frac{1}{2}$
Musk seedav.oz.	11/
Benzoingr.	
Myrrhgr.	270
Tincture of ambergrism.	20

Mix all but the myrrh, benzoin, and tincture; grind to moderately fine powder, contuse the benzoin and myrrh to powder, mix all, add the tincture of ambergris, and triturate until well mixed.

# Patchouly Sachet.

	I. Patchouly herbav.oz.	8
	Lavender flowersav.oz.	3
	Orrisav.oz.	2
	Clovesav.oz.	
i	Oil of bergamotfl.dr.	1
i	Oil of patchoulydrops	$^2$
l	Tincture of ambergrisfl.dr.	1
ŀ	Tincture of muskfl.dr.	1/2

Mix the first four ingredients, grind to other ingredients and mix well.

THE STANDAR	D TORM OLZIK I.
II.	Sweet Brier Sachet.
Patchouly leaves	Orris, ground.         av.oz. 18           Sandalwood, ground.         av.oz. 3           Oil of rose geranium.         drops 6           Oil of neroli.         drops 12           Oil of verbena.         drops 24           Oil of bergamot.         drops 8           Oil of lemon.         drops 12           Oil of rose.         drops 6           Tincture of ambergris.         m. 45           Tincture of musk.         m. 45           Mix the whole thoroughly.           Verbena Sachet.           Orris, ground.         av.oz. 16           Oil of bergamot.         m. 40           Oil of verbena.         fl.dr. 1           Oil of rose geranium.         drops 10           Tincture of musk.         m. 80
contuse the benzoin to powder, add to the previous mixture, to the whole add the oils,	Mix the whole thoroughly.
and mix intimately.  Rondeletia Sachet.  Orris	Violet Sachet.         av.oz. 16           Orris, ground.         av.oz. 16           Oil of bergamot.         drops 10           Oil of bitter almonds.         drops 7           Oil of rose         drops 7           Tincture of musk.         fl.dr. 2½           Mix the whole thoroughly.           Wild Flowers Sachet.           Canada snake root.         av.oz. 4           Coriander.         av.oz. 3           Deer tongue         av.oz. 2           Lavender flowers         av.oz. 2           Sweet flag root.         av.oz. 1           Patchouly leaves         av.oz. 1           Nutmeg         av.oz. 1           Oil of bergamot         fl.dr. 2           Oil of neroli         drops 40           Oil of sandalwood         drops 80
powder, then add the oils and tincture, and mix the whole thoroughly.  Rose Sachet.  I. Rose petals	Oil of verbena
powder, add the other ingredients, and mix thoroughly.  II.  Orris	I.       Orris       av.oz. 5         Rose petals       av.oz. 5         Orange peel, recently dried       av.oz. 6½         Coumarin       gr. 1         Vanillin       gr. 2         Tincture of civet       drops 48         Tincture of musk       drops 24         Oil of ylang ylang       drops 12         Oil of rose       drops 8         Oil of bergamot       drops 4         Essence of jasmine       fl.dr. 3

patchouly, grind to moderately fine powder in a mill, add the other ingredients and mix well. Mix the first three ingredients, reduce to moderately fine powder in a mill, dissolve the

coumarin and vanillin in the remaining ingredients, and mix the whole thoroughly.

### II.

Orris av.oz.	8
Rose petalsav.oz.	21/2
Cassie flowersav.oz.	21/2
Pimentogr.	
Tonka gr.	140
	140
Benzoin gr.	70
Oil of pimentodrops	10
Oil of bergamotdrops	20
	10
Oil of rose geraniumdrops	20
Oil of ylang ylangdrops	30
Oil of rose drops	
Tincture of muskfl.dr.	11/4
Tincture of civetdrops	40

Mix the first four ingredients, grind to moderately fine powder in a mill, triturate the tonka and vanilla with a solution of this mixture to powder, contuse the benzoin to powder, mix all, add the oils and tinctures, and mix the whole thoroughly.

### III.

Orris, groundav.oz.	15
Benzoinav.oz.	
Musk	110
Oil of ylang ylang drops	30
Oil of rosedrops	15

Contuse the benzoin to powder, triturate the musk intimately with a portion of the orris, add the remainder of the orris, the benzoin, and the oils, and mix the whole thoroughly.

# Solid Perfumes.

The novelty sold under this name is prepared by melting paraffin in a water bath, adding odorous substances when nearly cool, and molding into small tablets. The following formulas may be employed in preparing them, each of the mixtures given being sufficient for 4 avounces of paraffin.

### T.

Oil of lavender fl.dr. 2
Oil of cloves
Oil of rose geranium
Oil of bergamotfl.dr. 2
Vanillingr. 10
· · · · · · · · · · · · · · · · · · ·
Oil of nerolifl.dr. 1/2
Oil of rose geranium fl.dr. 1/2
Oil of lavenderfl.dr.
Oil of bergamotfl.dr. 1
Oil of clovesdrops 2
Heliotropingr. 10

т	т	т		
T	т	1	٠	

Oil of linaloefl.c	lr. 2
Oil of bergamot	
Oil of lemon	
Heliotropin	

# IV.

(	Oil of ylang ylangfl.dr.	2
(	Oil of nerolifl.dr.	1
(	Oil of sandalwoodfl.dr.	1/2
(	Coumaringr.	20
	Tincture of muskm.	

# V.

Oil of bergamotfl.dr.	4
Oil of rose geranium	0
Oil of neroli	0
Oil of lemonfl.dr.	
Oil of orangefl.dr.	
Oil of rosemary	0
Oil of lavender 2	0

These tablets are intended to replace sachet powders.

# Pot Pourris.

These are mixtures of odorous substances, in rather coarse condition, to be placed in open jars and intended for scenting rooms. The individual particles should be of about the size of a split pea, and such substances as orris, benzoin, etc., should be reduced to this size by appropriate means. In making good pot pourri mixtures, the best materials are required.

Extra perfume, such as an "extract," may be added to these pot pourris if desired.

Ι.

1.		
	Lavender flowersav.oz.	4
	Orrisav.oz.	4
	Rose petalsav.oz.	4
	Clovesav.oz.	1
	Cinnamonav.oz.	1
	Siam benzoinav.oz.	1
	Pimentoav.oz.	1
	Table saltav.oz.	2
	Vanillaav.oz.	3/4
	Musk gr.	1003/4
	Oil of bergamotdrops	30
	Oil of lemondrops	30
	Oil of lavenderflowersdrops	15
	Oil of sandalwooddrops	15
	Oil of rose geraniumdrops	
	Oil of rosedrops	
	Tincture of ambergrisfl.dr.	1/2
	D. t the sumin many metals alarmas	.:

Reduce the orris, rose petals, cloves, cinnamon, benzoin, pimento and vanilla to particles of suitable size, add the lavender, salt and musk, and then the oils, and tincture and mix

II. To 1 pint of rose petals, add.	
Orrisav.oz.	2
Pimentogr. 2	20
Clovesgr. 2	20
Cascarilla gr. 1	10
Musel-	2
Musk	$\tilde{\tilde{2}}$
Prepare like the preceding.	
•	
III.	
Sandalwoodav.oz.	6
Orris av.oz.	6
Benzoinav.oz.	1
Clovesav.oz.	1
Tonkaav.oz.	1
Maceav.oz.	1/
Musk	20
Oil of rosedrops	20
Oil of lavenderdrops	
Oil of bergamot	1
Oil of lemonfl.dr.	1
Prepare like the preceding.	
1V.	
Rose petalsoz.	4
Lavender flowersoz.	8
Vanillagr.	60
Cloves gr.	60
Storaxgr.	60
Benzoin gr.	60
Ambergrisgr.	20
Oil of rose drops	
Prepare like the preceding.	
r repare like the preceding.	

# Fumigating Pastilles.

I.

These are cone-shaped bodies produced by mixing red saunders or wood charcoal with potassium nitrate, odorous substances, and mucilage, and forming a mass. By the use of charcoal, black pastilles are obtained, while saunders produces the red variety.

When heated, these pastilles emit a pleasant odor; they are employed for this odor and for preventing and removing disease germs or foul odors caused by disease. That they are of any use except to produce a pleasant odor is doubtful.

The pastille mass may be formed into cones by means of a pastille machine, or by means of the hand similarly to the handmade suppositories.

Benzoinav.oz. 10
Charcoal
Sassafras

Mix the first four in fine powder, add the mucilage, form a mass, and make into conical pastilles.

L	.,	
	Potassium nitrategr.	375
	Waterfl.oz.	25
	Charcoal wood, powderav.oz.	30
	Tragacanth, powdergr.	375
	Storaxgr.	300
	Benzoingr.	300
	Vanillingr.	8
	Coumaringr.	3
	Muskgr.	3
	Civetgr.	11/2
	Oil of rosedrops	20
	Oil of bergamotdrops	15
	Oil of ylang ylangdrops	10
	Oil of rhodiumdrops	10
	Oil of sandalwooddrops	5
	Oil of cinnamondrops	5
	Oil of orrisdrop	1
	Oil of cascarilladrop	1

Saturate the charcoal with the potassium nitrate dissolved in the water, dry the mass, powder, add the other ingredients and mix thoroughly. Beat the mixture to a plastic mass with the addition of sufficient mucilage of tragacanth containing 2 per cent of saltpeter in solution, and form into cone-shaped pastilles. In order to evenly distribute the storax throughout the mass, it may be previously dissolved in a small amount of acetic ether.—D.

H	II.		
	Benzoin	.av.oz.	2
	Cascarilla		
	Myrrh		1
	Potassium nitrate		1/2
	Potassium chlorate		
	Charcoal, wood		
	Oil of cloves		
	Oil of cinnamon	fl.dr.	1
	Oil of lavender		
	Mucilage of tragacanth	.suffici	ent

Mix the first six ingredients previously reduced to fine powder, add the oils, and then incorporate enough mucilage to form a mass. Divide this into pastilles weighing about 60 gr. and dry.

Charcoal, powder	.av.oz.	30
Potassium nitrate		
Water		
Tragacanth, powder		
Tincture of benzoin		
Peru balsam		
Storax, crude		
Tolu balsam		
Oleo-balsamic mixture		
Coumarin		
	0	

Saturate the charcoal with the potassium nitrate dissolved in the water, then dry,

2

reduce to powder, and incorporate the tragacanth and then the remaining ingredients. Now form a mass by the addition of sufficient mucilage of tragacanth containing 2 per cent of potassium nitrate in solution, and divide into pastilles .- D.

7.		
	Red saunders, powderav.oz.	24
	Potassium nitrateav.oz.	21
	Waterfl.oz.	33
	Fragacanth, powderav.oz.	1
	Tincture of benzoinfl.dr.	12
	Peru balsamgr.	300
	Storax, crudeav.oz.	11
	Tolu balsam av.oz.	11
	Oleo-balsamic mixturefl.dr	24
	Mucilage of tragacanth, con-	,
	taining 2 per cent of potas-	
	sium nitratesuffic	ient

Prepare like the preceding.—D.

VI.	
С	ascarilla av.oz. 5
В	Senzoinav.oz. 4
0	Olibanumav.oz. 4
A	mberav.oz. 1
	andaracav.oz. 1
	Charcoal, wood, or red saunders.av.oz. 6
	otassium nitrateav.oz. 2
	toraxav.oz. 6
P	'eru balsamav.oz. 1

Mix the first seven ingredients previously reduced to powder, add the remaining ingredients, form a mass with mucilage of acacia or tragacanth, divide into pastilles and dry.-H.

### Fumigating Powder. (Fumigating Species.)

Fumigating powder is of similar composition to the pastilles and is employed for the same purposes. It is in the form of coarse powder, free from any fine powder as well as from large, coarse pieces, and is of variegated brilliant colors, which are often produced by the use of aniline colors dissolved in alcohol and different portions separately tinctured, or sawdust is thus colored and added to the aromatics.

Ι.		
	Benzoingr.	240
	Tolu balsamgr.	240
	Storaxgr.	
	Alcoholfl.oz.	
	Peru balsamgr.	60
	Oil of cinnamondrops	4
	Oil of lavender flowersdrops	4
	•	

alcohol, agitate occasionally for several days, unsized paper with the liquid.

filter and add the other ingredients. Moisten clean pine sawdust with this liquid.

I.	
Sandalwood, yellowav.oz	51/2
Sassafrasav.oz	z. 3¾
Cinnamonav.oz	. 11/4
Clovesav.oz	. 11/4
Cascarillagr	:. 275
Potassium nitrategr	:. 275
Waterfl.oz	. 10
Storax:av.oz	
Tolu balsamav.oz	
Etherfl.oz	
Benzoinav.oz	z. 3
Olibanumav.oz	21/2
Juniper berriesgr	r. 275

Reduce the first five ingredients to a coarse powder, free from dust, then thoroughly impregnate it with a solution of the potassium nitrate in the water, and dry the mass thoroughly in the air. Dissolve the storax and balsam of tolu in the ether, and evenly and thoroughly saturate the above, after which dry quickly in the air and immediately add the last three ingredients, previously reduced to coarse powder. Finally preserve in a well-When used, a quantity of stoppered bottle. the powder should be thrown upon live coals.

III.

Corianderav.oz.	3
Cassia barkav.oz.	3
Cascarillaav.oz.	3
Lavender flowersav.oz.	3
Peony flowers av. oz.	3
Orris rootav.oz.	3
Juniper berriesav.oz.	_
Rose petalsav.oz.	3
	.3
Blue flowersav.oz.	3
Calendula flowersav.oz.	3
Clovesav.oz.	
Peru balsamgr.	300
Storaxgr.	300
Oil of bergamotfl.dr.	5
Oil of lemonfl.dr.	5
Tincture of benzoinfl.dr.	5
Oil of lavender flowersfl.dr.	1/3
Oil of cloves fl.dr.	1/2
Oil of cassia	1/2
Tincture of ambergrisdrops	
Tincture of muskdrops	12
	-

Mix the drugs, reduce to coarse powder, and thoroughly incorporate with the other ingredients which have previously been well mixed.—H.

# Fumigating Paper.

I. Make a tincture as in No. I. under Mix the benzoin, tolu and storax with the the preceding heading and saturate pieces of II.

Storaxav.oz.	1
Benzoinav.oz.	
Fumigating essencefl.oz.	
Alcoholfl.oz.	2
Ether fl.oz.	1
Acetic acid, glacial drops	

Dissolve the benzoin and storax in a mixture of the alcohol and ether, filter, and add the fumigating essence and the acetic acid. Spread the mixture upon bibulous paper by means of a broad soft brush and allow it to dry. To prevent sticking, dust the surface with talcum and preserve in waxed paper. When used the paper is simply warmed on a hot plate or over a lamp.—D.

# Fumigating Essence or Tincture.

Benzoinav.oz.	1
Storaxgr.	300 -
Peru balsamgr.	75
Oil of bergamotdrops	30
Oil of rosedrops	15
Oil of ylang ylangdrops	8
Oil of rhodiumdrops	8
Oil of rose geraniumdrops	5
Oil of sandalwooddrops	5
Oil of sassafrasdrops	5
Oil of cassiadrops	5
Oil of clovesdrops	5
Oil of bitter almondsdrops	2
Oil of orrisdrop	1
Coumaringr.	1
Vanillin gr.	3/4
Tincture of muskfl.dr.	11/2
Acetic etherfl.dr.	21/2
Essence of jasminefl.oz.	1
Alcoholsuffici	-
	CIIL

Mix all of the above, using 5 fluidounces of alcohol, macerate for several days, agitating occasionally; filter, and add through the filter enough alcohol to make the product measure 8 fluidounces.

This preparation may be cheapened by using four times as much alcohol and twice as much benzoin, storax, Peru balsam and oil of bergamot.—D.

# Fumigating Vinegar.

Fumigating tincture	.fl.oz.	31/4
Acetic ether	.fl.dr.	11/2
Acetic acid		

Mix, and after standing in a cool place a few days, filter.

In fumigating sick rooms the vinegar is vaporized either by heating it in a spoon or by pouring it upon a hot iron.—D.

# Cologne Waters.

The cologne waters resemble the handkerchief extracts, but are much weaker; in fact, they may be produced by diluting extracts to any desired extent with alcohol, adding distilled water gradually with frequent agitation until the mixture becomes milky, and then filtering. A white rose extract would thus make a White Rose Cologne; a violet extract, Violet Cologne, etc.

Instead of diluting extracts, colognes may be prepared from second washings of pomades and adding suitable oils, tinctures, etc., as directed under "Handkerchief Extracts." They may also be prepared by mixing the ingredients specified in the formulas below. The oils, tinctures, essences, etc., should be mixed with the alcohol, and then the water should be added gradually with agitation. The mixture should then be set aside for several weeks until the mixture has blended and 'ripened,' when it may be filtered.

As in the case of other odorous mixtures, superior ingredients will produce a good product, inferior ingredients a poor product.

The essences, tinctures and spirits mentioned in the formulas should be prepared according to the systems enumerated under "Handkerchief Extracts."

Suitable names for cologne waters are Newport Cologne, Opera Cologne, German Cologne, Lafayette Cologne, Farina Cologne, Victoria Cologne, Superior Cologne, Imperial Cologne, Oriental Cologne, etc.

Cologne waters should be preserved in well-stoppered bottles, away from strong light and heat.

Τ.

Musk	gr. 31/2
Vanilla, the meat or interior	r. 7
Oil of rose	m. 30
Oil of bergamotfl.c	ir. 51/2
Essence of jasminefl.c	lr. 5
Essence of violetfl.c	lr. 5
Oil of neroli	n. 40
Asafetida, powder	gr. 1
Alcoholfl.c	z. 59
Waterfl.c	z. 5

Mix the musk, vanilla, water and 11 fluidounces of alcohol, macerate for 30 days, add the remaining ingredients, macerate another 30 days, agitating frequently, and filter.

PERF	UMES. 303
II.  Oil of bergamot	VII.  Oil of lemon fl.dr. 4 Oil of bergamot fl.dr. 6 Oil of melissa fl.dr. 1/2 Oil of rosemary fl.dr. 1/3 Oil of neroli fl.dr. 1/2 Tincture of musk fl.dr. 1/2 Alcohol fl.oz. 78 Oil of rose m. 30 Acetic ether m. 45 Mix the first eight ingredients, distill over 64 fluidounces, and to the distillate add the oil of rose and acetic ether.  VIII. Oil of bergamot fl.oz. 1 Oil of lemon fl.oz. 1 Oil of rosemary fl.dr. 2 Oil of neroli fl.dr. 2
Essence of jockey club	(ltl of lemon m 50)
Alcohol	Oil of lemon         fl.dr.         2           Oil of neroli         fl.dr.         2           Oil of orange         fl.dr.         2           Oil of lavender         m.         20           Oil of rosemary         m.         10           Oil of bergamot         fl.dr.         7½           Alcobol         fl.dr.         6½           XI.         Oil of bergamot         fl.dr.         4½           Oil of lemon         fl.dr.         6           Oil of lavender         m.         45           Oil of neroli         m.         45           Oil of rosemary         m.         25           Tincture of musk         fl.dr.         3           Tincture orris         fl.oz.         3           Orange flower water         fl.oz.         13           Alcohol         fl.oz.         48
VI.  Oil of bergamot fl.dr. 6 Oil of lemon fl.dr. 3 Oil of neroli drops 60 Oil of origanum drops 18 Oil of rosemary drops 60 Orange flower water fl.oz. 3 Alcohol fl.oz. 60 Mix all but the water, then add latter gradually with agitation, set aside for some time and filter.	Ull Of Tose

1112 511112111	_
XIII.	r
Oil of lemon fl.dr. 1	
Oil of rosedrops 80	t1
Oil of neroli drops 30 Stronger tincture of orris fl.dr. 6	n
Tincture of tonkafl.dr. 11	12
Tincture of muskfl.dr. 4	p
Alcoholfl.oz. 47	e
XIV.	a
Oil of bergamotfl.dr. 4	b
Oil of lemon grass	d
Oil of lavender	1
Oil of nerolidrops 15	1
Oil of bitter almondsdrops 15	
Oil of cassia	
Oil of wintergreendrops 12	
Waterfl.oz. 12	
Alcohol'fl.oz. 52	
Mix all but the water, add the latter grad-	1
ually with agitation, set aside for some time,	
and filter.	
XV.	
Oil of lemon	
Oil of bergamot	t
Oil of rose	f
Oil of neroli m. 40	1
Tincture of musk	I
Orange flower water	
Mix all but the water, add the latter grad-	
ually, set aside for some time, and filter.	
XVI.	
Oil of jasminefl.oz. 1	
Oil of bergamotfl.dr. 4	
Oil of lemonfl.dr. 1	
Oil of lavender	
Oil of neroli	ŀ
Oil of clovesdrops 2	f
Oil of rosedrops 5	1
Tincture of muskfl.dr. 2 Tincture of vanillafl.dr. $6\frac{1}{2}$	- 1
Stronger tincture of orrisfl.dr. 4	
Alcohol	
Waterfl.oz. 16	
Mix all but the water, add the latter grad-	-
ually with agitation, set aside for some time,	
and filter.	
XVII.	
Oil of nerolifl.dr. 2	]
Oil of rose geraniumfl.dr. 1	
Stronger tincture of orris fl.oz. 4	
Tincture of civetfl.dr. 2	
Alcoholfl.oz. 56	
Mix and filter.	1

# Coilet Waters.

Toilet waters are similar in composition to the cologne waters. They are usually named after the principal ingredient, viz., lavender water, because oil of lavender is the principal; violet water, because containing essence of violet, etc. Florida water is another toilet water, the principal ingredients being oils of lavender and bergamot or lavender, bergamot and lemon.

# Bretfeld Cosmetic or Toilet Water.

(Eau de Bretfeld.)

Oil of bergamotfl.oz.	1
Oil of lemon fl.dr.	2
Oil of cloves	50
Oil of lavender	50
Oil of neroli	30
Oil of rosedrops	10
Vanillingr.	
Tincture of muskfl.dr.	
Alcoholfl.oz.	
Waterfl.oz.	5

Mix all ingredients except the water, agitate well, add the water, shake again, set aside for 8 days, and filter.—D.

# Florida Water.

I.		
	Oil of lavender flowersfl.oz.	1
	Oil of bergamotfl.dr.	4
	Oil of clovesfl.dr.	2
	Oil of rose geraniumm.	30
	Oil of cinnamondrops	15
	Oil of spearmintdrops	4
	Benzoic acidgr.	120
	Waterfl.oz.	4
	Alcoholfl.oz.	60

Mix the oils, acid and storax with the alcohol, shake thoroughly, add the water, and filter.

II		
	Oil of lavenderfl.dr.	
	Oil of bergamotfl.dr.	4
	Oil of lemonfl.dr.	4
	Oil of nerolifl.dr.	2
	Oil of melissa fl.dr.	1
	Oil of rosedrops	20
٠	Tincture of curcumafl.dr.	
	Alcohol fl. oz.	64
	3.51	

Mix and filter.

III.	
Oil of lavender flowersfl.c	)z. 1
Oil of lemonfl.c	lr. 4
Oil of orangefl.c	lr. 4
Oil of cloves	ir. 2
Alcoholfl.c	

Mix and filter.

IV.	II.
Oil of lemon       fl.dr.       4         'Oil of bergamot       fl.dr.       3         Oil of neroli       fl.dr.       2         Oil of lavender flowers       fl.dr.       1         Oil of white thyme       m.       50         Oil of mace       m.       50         Oil of cassia       m.       50	Oil of lavender       fl.dr. 6         Oil of bergamot       fl.dr. 2½         Oil of rose       m. 25         Tincture of tonka       fl.oz. 6         Tincture of musk       fl.dr. 5½         Alcohol       fl.oz. 59
Oil of rosemary	Oil of bergamot       fl.oz. 1         Oil of lavender       fl.dr. 2         Oil of cloves       fl.dr. 1         Oil of sandalwood       m. 40         Oil of rose       m. 40         Tincture of ambergris       fl.dr. 2
sionally, and filter.	Tincture of muskfl.oz. 1 Alcoholfl.oz. 63
V.	IV.
Oil of lavender flowers fl.dr. 2 Oil of bergamot fl.dr. 1 Oil of cloves m. 40 Oil of rosemary flowers m. 40 Oil of cinnamon drops 20 Oil of neroli drops 6 Tincture of musk drops 8 Tincture of ambrette fl.dr. 2 Alcohol, enough to make fl.oz. 64	Oil of lavender fl.dr. 10 Oil of bergamot fl.dr. 1½ Oil of neroli drops 4 Oil of rose geranium drops 6 Oil of sandalwood drops 7 Tincture of musk drops 8 Alcohol fl.oz. 30 Water fl.oz. 30
A portion of the alcohol may be replaced by distilled water, using only enough of the latter to produce opalescence.	V.   Oil of lavender
VI.  Oil of lavender flowers fl.oz. 1 Oil of bergamot fl.dr. 4 Oil of cloves fl.dr. 2 Oil of rose drops 20 Oil of cinnamon drops 10 Benzoic acid gr. 30 Alcohol fl.oz. 62	Tincture of musk.
Geranium Water.	Lilac Water.
Oil of rose geranium	, Essence of tuberose
add the water, and filter.	Add the essences, oil and tincture to the
Lavender Water.	alcohol, add water gradually with agitation until the liquid becomes very slightly milky,
I.	and filter.
Oil of lavenderfl.dr. 6 Alcoholfl.oz. 48	Violet Water.

Essence of violet .....fl.oz.

Essence of rose. fl. oz. 2½
Essence of cassie. fl. oz. 2½

Water may be added if a cheaper prepara-

Alcohol ......fl.oz. 56

tion is wanted. Tincture of orris may also

Triturate the oil with the magnesium carbonate, add the alcohol, and then the water, and filter. It may be colored a light brown tint with caramel if desired.

Magnesium carbonate . . . . . . av.oz. 1

be a desirable addition. The preparation may also be tinted with chloropyhll.

Essence of violet.....fl.oz. Essence of cassie.....fl.oz. Spirit of rose.....fl.oz. Alcohol ......fl.oz. 48

This may be diluted, modified or tinted like the preceding.

III. Violet water may also be prepared by diluting violet extract with alcohol, adding some water, and filtering.

Other toilet waters may be prepared having odors different from any of the above. Such preparations may be made as "White Rose Water," " Heliotrope Water," etc.

# Bay Rum. (Spirit of Myrcia.)

Ī.		
	Oil of bayfl.dr.	4
	Oil of orange	15
	Oil of pimentom.	15
	Alcoholfl.oz.	39
	Waterfl.oz.	25

Mix the oils and alcohol, add the water, set aside in a well-stoppered bottle for 8 days and filter.-U. S. P.

TT Oil of bay.....fl.dr. Acetic ether.....fl.dr. 21/2 Stronger tincture of orris . . . . . fl.dr. New England rum.....fl.oz. Alcohol ......fl.oz. 32 Water, enough to make.....fl.oz. 64

Mix the oils and tincture with the alcohol, add the rum, then the water, and filter.

III.

Oil of bay	.fl.dr.	1 1/2
Oil of pimento		
Oil of lemon grass		
Oil of orange		
Oil of cloves		
Oil of rose geranium		
Acetic ether		
Alcohol		
Water		

Mix the oils and ether with the alcohol, add the water, and filter.

IV.	
Oil of bayfl.dr.	1
Oil of pimentodrops	8
Oil of rose geraniumdrops	4
New England rumfl.oz.	8
Alcoholfl.oz.	32
Water, enough to makefl.oz. 6	34

incorporate the water previously slightly warmed, and filter.

V.

Oil of bayfl.dr.	2
Jamaica rumfl.oz.	4
Alcoholfl.oz.	24
Waterfl.oz.	

This can be made clear and bright by filtering through magnesia and charcoal. It makes a cheap article for barbers' use.

VI.

- (	Oil of bayfl.dr.	11/2
-	Oil of pimentom.	45
	Acetic etherfl.dr.	11
	Alcoholfl.oz.	32
	Waterfl.oz.	32

Mix the oils and ether with the alcohol, add the water, and filter.

VII.

Oil of bay	.fl.dr.	4
Oil of orange		
Alcohol	.fl.oz.	40
Water	.fl.oz.	40
Calcium phosphate	gr.	120
Jamaica rum		

Mix the oil and calcium phosphate, add the alcohol, and stir; then add the water and filter clear. Finally add enough rum to give the desired color.

VIII.

Oil of bayfl.dr.	2
Alcoholfl.oz,	40
Waterfl.oz.	24
Bay leavesgr.	120

Mix, macerate for 14 days, and filter.

# Toilet or Aromatic Vinegars.

Such a preparation may be known simply as toilet or aromatic vinegar, or as lavender, rosemary, eucalyptus, witch hazel, cosmetic, antiseptic. hygienic, preventive, vinegar according to fancy or to the main constituent, or it may be known by the French title "vinaigre de toilette."

These preparations consist of aromatic oils and other odorous substances combined with alcohol and acetic acid.

Lavender flowers...... ..av.oz. Alcohol ......fl.oz. Diluted acetic acid.....fl.oz. 56

Mix, macerate for 8 days, agitating frequently, express in a straining cloth, heat the Mix the oils and alcohol, add the rum, then colature nearly to boiling, place for several

days in a cool place,	, then fil	ter, place	the	fil-
trate in small bott	les, and	preserve	in	the
dark.—D.				

I.	
Oil of lavenderdrops	4
Oil of rosemarydrops	4
Oil of juniperdrops	4
Oil of peppermintdrops	4
Oil of cassiadrops	4
Oil of lemondrops	8
Oil of clovesdrops	8
Alcoholfl.oz.	23/
Acetic acidfl.oz.	23/
Distilled water, enough to make.fl.oz.	16

Dissolve the oils in the alcohol, add the acid, and lastly the water. Warm the mixture for several hours to a temperature not exceeding 76 degs. C., taking care there is no loss by evaporation; set aside for a few days and filter.—N. F.

III. The aromatic vinegar of the German pharmacopæia differs from the preceding only in containing 50 per cent more of each of the oils.

# IV.

Oil of lavenderdrops 15	ĭ
Oil of rosemarydrops 15	
Oil of nerolidrops 18	3
Oil of lemonfl.dr.	3
Oil of bergamotfl.dr.	3
Oil of orangefl.dr.	31
Tincture of muskdrops 15	5
Tincture of benzoin, U. S. P fl. oz.	L
Tincture of tolufl.oz.	l
Glacial acetic acidfl.oz. 16	3
Alcoholfl.oz. 48	5

Mix the oils and tinctures with the alcohol, add the acid, let the mixture stand for several days and filter.

# V.

Rosemary leavesav.oz.	-7
Alcoholfl.oz.	8
Diluted acetic acidfl.oz.	56
Prepare like No. ID.	

# VI.

Tincture of benzoin, U. S. P fl.dr.	6
Tincture of tolufl.dr.	6
Oil of orangefl.dr.	2
Oil of lemon	2
Oil of bergamotdrops	50
Oil of nerolidrops	12
Oil of rosemarydrops	8
Acetic acidfl.oz.	$1\frac{1}{2}$
Diluted alcohol, enough to make.fl.oz.	64
711	

### VII.

Glacial	acetic	acid	l			fl.dr.	10
Cologne							
Mix, let	stand	for	seve	ral	days	and	filter
D.							

|--|

Oil of rosedroj	os 16
Oil of bergamotdroj	os 16
Oil of ylang ylangdroj	os 2
Oil of orris, liquiddro	os 4
Oil of cassiadroj	os 2
Tincture of muskdroj	
Tincture of ambergrisfl.d	
. Coumaring	r. 1
Heliotroping	
Essence of jasminefl.d	r. 11
Glacial acetic acidfl.d	r. 12
Alcoholfl.o	
Distilled water	

Mix, macerate for at least 8 days, and filter.—D.

### IX.

Rose petalsav.oz.	14
Essence of rosefl.oz.	
Diluted acetic acidfl.oz.	54

Mix, macerate for 1 day, strain, set the liquid aside for several days, and filter.

### X

_			
	Essence of cassie	.fl.dr.	4
	Essence of violet	.fl.dr.	4
	Essence of rose	.fl.dr.	4
	Stronger tincture of orris	.fl.oz.	4
	Diluted acetic acid	.fl.qz.	16
	Distilled extract of witch hazel,		
	enough to make	.fl.oz.	32

Mix and filter.

### Smelling Salts.

Under the name of "smelling salts," "inexhaustible salts," "Preston salts," and "pungent" there have been prepared moist granular preparations which are aromatized and give off a fragrant odor in connection with the pungent odor of ammonia. They consist of either coarsely powdered ammonium carbonate with or without the further addition of ammonia water, or of a coarsely powdered mixture, which slowly evolves the odor of ammonia, a mixture of ammonium chloride and potassium carbonate or lime, the whole being aromatized by the addition of volatile oils, or pomade essences, or hand-kerchief extracts.

These preparations are usually dispensed in small ornamental glass containers which can be securely closed when not in use.

These are used by inhalation in the fainting spells or headaches of ladies.

These preparations may be made stronger in ammonia strength by adding a fresh portion of ammonia water.

I. Ammonium carbonateav.oz.	11
Caustic potassaav.oz.	$\frac{1}{2}$
Orris rootav.oz.	1
Ammonium chlcridegr. 12	0
Lemon peelgr. 12	0
Rosemarygr. 12	
Lavender flowersgr. 12	
	0
	0
	0
	1
Concentrated water of ammonia. fl. dr.	1
Mix the first nine ingredients previous	11517

Mix the first nine ingredients previously reduced to a granular condition, then add the other ingredients, mix quickly, and bottle.

### TT.

Fill the bottle with small crystals of potassium sulphate and add enough of the following to moisten:

Water of ammoniafl.oz.	4
Oil of lemondrops 1	5
Oil of bergamotdrops	3
Oil of lavender flowersdrops	4

Or mix equal parts of ammonium chloride and potassium carbonate, and perfume, or fill the vial with coarsely powdered ammonium carbonate, the fragments of which have been brought to a uniform size by sifting out both the fine and coarse pieces, and perfume as above.

### III.

Ammonium chloride, granular.av.oz.	1
Quicklime, granularav.oz.	1
Mix and moisten with the following:	
Stronger water of ammoniafl.oz.	2
Oil of bergamotdrops	12
Oil of lavender flowersdrops	6
Oil of rosedrops	
Oil of cassiadrops	
Oil of clovesdrops	5
<u> </u>	H.

, V •	
Ammonium chlorideav.oz.	1 1/2
Potassium carbonateav.oz.	1 3/4
Ammonium carbonategr.	180
Camphor gr.	60
Oil of clovesdrops	10
Oil of bergamotdrops	

Reduce the ammonium carbonate to coarse powder, add the ammonium chloride and potassium carbonate in a granular condition and the camphor in powder form, and then the two oils.

Menthol smelling salts may be prepared by substituting 60 grains of menthol for the camphor and oils in the above.

V.	
Oil of clovesdrops	6
Oil of lavender flowersdrops	
Oil of bergamotdrops	30
Stronger water of ammoniafl.oz.	

Fill the smelling bottles with coarsely powdered ammonium carbonate, and add to the salt as much of the aromatic solution as it will absorb.

VI

4.	
Oil of lemondrops	12
Oil of lavenderdrops	6
Oil of clovesdrop	1
Stronger water of ammoniafl.oz.	3

Fill the smelling bottles with crystallized potassium sulphate and pour into each bottle as much of the aromatic ammonia as the salt can retain without spilling. This makes a much prettier looking smelling salt than ammonium carbonate, and as it does not cake together like it, the bottle need not be emptied when the ammonia has evaporated. All that is necessary is to fill it up again with the aromatic ammonia.

VII. Moisten coarsely powdered ammonium carbonate contained in a suitable bottle with the following mixture:

Oil of lavender flowersfl.dr.	2
Spirit of ammoniafl.dr.	. 1
Absolute alcoholfl.dr	17

This preparation should be dispensed as lavender salts.—D.

VIII. Violet salts may be prepared in a manner similar to the preceding, substituting for the lavender oil and alcohol a concentrated tincture of orris root, in proper quantity, spirit of orris, or, if preferred, violet extract. It may be tinted green by the addition of green aniline or tincture of grass.

Other odors like heliotrope, etc., may be produced by substituting the corresponding spirit, oil, extract or other substance for the lavender or violet.

IX. The following pungent is quite different from any of the preceding and is given here for the sake of variety:

Sodium acetate, small crystals. av. oz.	0
Glacial acetic acidfl.dr.	$2\frac{1}{2}$
Acetic etherfl.dr.	$2\frac{1}{2}$
Oil of rosedrops	20
Oil of bergamotdrops	
Oil of nerolidrops	
Oil of ylang ylangdrops	2
Oil of orris, liquiddrops	
Coumarin: ar	1

Mix all but the sodium acetate and add to the latter contained in a suitable vial.

This is to be used like the preceding pungents. If desired of a red tint, it may be colored by the addition of a trifling amount of fuchsin to the solution.—D.

# SECTION II—SKIN PREPARA-TIONS.

Preparations intended mainly or entirely for the skin, viz.: Ointments and Creams, Skin Food, Toilet Milks, Toilet Lotions, Toilet Jellies, Camphor Ice, Toilet Lanolin, Almond Paste, Almond Meal, Toilet Powders, Liquid Cosmetics, Face Bleach, Remedies for Blackheads, Freckles, Pimples, Tan, Sunburn, etc.; Rouge, Grease Paints, Depilatories, Liquid Soaps, Shaving Creams and Powders, and Manicure Cosmetics.

# Cold Creams.

Preparations known as "Cold Creams" are mixtures of solid fats, like wax, spermaceti, petrolatum, lanolin, etc.; oils, like almond oil, water, rose water, glycerin, distilled extract of witch hazel, or other similar fluid, and some flavoring substances. In preparing them the solid fats must be melted, the oil (castor oil is preferable to other oils) then mixed with the fluid mentioned, the whole stirred vigorously and constantly in a large mortar until well mixed and quite solid, finally adding the flavoring substances. The mixing may also be by means of stirring with a large spatula or an egg beater.

Such a preparation receives a name depending on its constitution or flavor. may be called simply "cold cream; if flavored with bitter almond oil, "almond cold cream"; if it contain cucumber juice, "cucumber cream, pomade, or ointment"; if witch hazel extract, "witch hazel cold cream"; if glycerin, "glycerin cold cream"; if camphor, "camphor cold cream"; lanolin, "lanolin cold cream"; salicylic acid, "salicylic or salicylated cold cream"; petrolatum, "petrolatum (or vaselin) cold cream"; violet extract or orris, "violet cold cream"; if orange flower water and Peru balsam, "sultana cold cream"; boric acid and glycerin, "boroglycerin cold cream," etc.

1.	
Spermacetiav.oz.	2
White waxav.oz.	2
Sweet almond oilfl.oz. 1	0
Stronger rose waterfl.oz.	3
Borax, powdergr. 4	

Melt the wax and spermaceti at a moder ate heat, add the oil, place the mixture in a warmed mortar, carefully add the rose water in which the borax has previously been dissolved, and stir rapidly and continuously until the mixture becomes uniformly soft and creamy.—U. S. P.

•	
II.	
Sweet almond oilfl	.oz. $6\frac{1}{2}$
Castor oil (odorless)fl	.oz. $1\frac{1}{2}$
Lard, benzoinatedav	.oz. 2
White waxav	
Rose water (in winter less, in	
summer more, than quantity	
named)fl	.oz. 3
Orange flower waterfl	
Oil of rosedr	
Essence of jasminefl	
Essence of cassiefl	
Borax, powder	.gr. 240
Glycerinfl	.oz. 1

Melt the oil of sweet almond, wax and lard together and stir in the castor oil, make a solution of the borax in the glycerin and rose and orange flower waters, add this solution, a little at a time, to the melted fat, stirring constantly to insure thorough incorporation; finally, add the oil of rose dissolved in the essences, and beat the ointment until cold.

# III.

11.	
Castor oilav.oz. 12	
Spermacetiav.oz. 4	
White waxav.oz. 1	
Oil of rosedrops 10	
Oil of bitter almonddrops 3	

Melt the wax and spermaceti, add the castor oil; when cooling beat to a cream, add the remaining oils, and mix well.

### TV

White waxav.oz.	1
Lardav.oz.	4
Sweet almond oilfl.oz.	$4\frac{1}{2}$
Water, warnıfl.oz.	31
Borax, powdergr.	15
Zinc oxidegr.	120
Oil of rosedrops	8
Oil of bergamotdrops	5
Oil of rose geraniumdrops	5
Alcoholfl.dr.	1

Melt the wax and lard together, add the oil, then the warm water with which the

borax and zinc oxide have previously been solved, add the water in which the borax has mixed, stir the whole thoroughly in a mortar until well mixed and nearly cold, add the oils previously dissolved in the alcohol, and mix again.

### V.

Spermacetiav.oz.	2
White waxav.oz.	11/2
Castor oilav.oz.	$4\frac{1}{2}$
Cottonseed oil, bleachedfl.oz.	$6\frac{1}{2}$
Rose waterfl.oz.	43/4
Borax, powder gr.	$120^{\circ}$
Oil of rosesufficient to fla	vor

Melt the spermaceti and wax, add the castor and cottonseed oils, then incorporate the water in which the borax has previously been dissolved, and finally add the oil of rose.

# VI.

Spermacetiav.oz. 3	13
White waxav.oz. 3	į
Castor oilav.oz. 3	
Cottonseed oilfl.oz. 12	
Rose waterav.oz. 7	
Oil of lavender flowersdrops 12	,
Prepare like the preceding.	

# Almond Cold Cream. (Almond Cream.)

•	
Spermacetiav.oz.	2
White waxav.oz,	2
Sweet almond oilfl.oz.	14
Water, distilledfl.oz.	7
Borax, powdergr. 6	60
	1
Oil of bergamotdrops	$24^{-}$
Oil of rosedrops	6
Oil of bitter almonddrops	8
Tincture of ambergrisdrops	
Prepare like the preceding.—D.	

### Camphor Cold Cream. (Camphorated Cold Cream.)

•		
Spermaceti	.av.oz.	2
White wax		2
Camphor	.av.oz.	1 1/4
Sweet almond oil	fl.oz.	13
Water, distilled		$6\frac{1}{2}$
Borax, powder		60
Coumarin	gr.	$\frac{1}{2}$
Oil of rose		18
Oil of rose geranium	.drops	4
Oil of ylang ylang		
Oil of bitter almond	.drops	2
Oil of orris, liquid		1
Tincture of musk		10
Tincture of civet		

Melt the wax and spermaceti, add the sweet almond oil, add the camphor, first broken into pieces, stir until the latter is dis- lanolin may be used, the amount of water

previously been dissolved, stir until well mixed and nearly cold, add the other ingredients and again mix well .- D.

# Boroglycerin Cold Cream.

Boric acid, powdergr.	90
Glycerinfl.oz.	2
Water, distilledfl.oz.	
Lanolin, anhydrousav.oz.	
Parafin whiteav.oz.	
Oil of rosedrops	6
Oil of bergamotdrops	6

Heat together the first three ingredients until solution is effected (one hour). Melt together the lanolin and petrolatum, incorporate the boroglycerin solution by rapid and constant stirring, then add the oils, and mix again.—D. modified.

# Glycerin Cold Cream.

Spermacetiav.oz.	3
White waxav.oz.	1
Sweet almond oilfl.oz.	8
Borax, powdergr.	240
Glycerinfl.oz.	3
Orange flower waterfl.oz.	1
Oil of nerolidrops	5
Oil of rosedrops	3
-	

Melt the wax, spermaceti, and almond oil together; dissolve the borax in the orange flower water and glycerin previously mixed; pour the solution, a little at a time, into the melted mixture, stirring the preparation without ceasing until all the solution has been fully incorporated, and a homogeneous product results; finally add the essential oils.

# Lanolin Cold Cream. (Lanolin Cream.)

٠.		
	Sweet almond oilav.oz.	$8\frac{1}{2}$
	Lanolin, anhydrousav.oz.	33/4
	White waxav.oz.	11/4
	Spermacetiav.oz.	11/4
	Borax, powdergr.	40
	Rose waterfl.oz.	6

Melt together the first four ingredients, then incorporate the solution of borax in the rose water.

An equally good preparation is obtained by melting together 3 av. ounces of anhydrous lanolin and 6 av. ounces of benzoinated lard, and adding 9 fluidounces of rose water.

Instead of anhydrous lanolin, ordinary

being reduced if necessary. Any desired perfume or combination may be added to these mixtures.

### II.

White waxgr. 4	180
Spermacetigr. 4	
Sweet almond oilfl.oz.	7
Lanolinav.oz.	$3\frac{1}{2}$
Borax, powdergr.	
Waterfl.oz.	
Perfumeto s	uit

Melt the wax and spermaceti, add the lanolin and oil, and when melted add the water containing the borax in solution; stir until cool and add suitable perfume.

### III.

White petrolatumav.oz.	14
Paraffin waxav.oz.	1
Lanolin av.oz.	4
Waterfl.oz.	
Alcoholfl.dr.	
Vanillingr.	4
Oil of rosedrops	5

Melt the petrolatum and paraffin on a water bath, pour into a warm mortar, add the lanolin and with constant stirring incorporate the water. When of the consistence of a thick cream add the vanillin and oil dissolved in the alcohol.

### IV.

Petrolatumav.oz.	10
Lanolinav.oz.	
Soap, powdered, white gr.	140
Waterfl.oz.	5
Oil of rosedrops	
Oil of nerolidrops	6
Oil of bergainotdrops	6
771 4 4 4 4 4	

Mix the petrolatum, lanolin, and soap, incorporate the water with this mixture and then add the oils. - D.

# Petrolatum Cold Cream.

Ţ.

White waxav.oz.	$1\frac{1}{2}$
Spermacetiav.oz.	$1\frac{1}{2}$
Petrolatumav.oz.	$4^{-}$
Sweet almond oil fl.oz.	9
Borax, powdergr.	90
Waterfl.oz.	
Coumaringr.	1
Oil of rhodiumdrops	10
Oil of bergamotdrops	
Oil of rose geraniumdrops	3
Oil of rosewooddrops	2
Oil of orris, liquiddrop	
Tincture of civetdrops	8

almond oil and petrolatum, and when melted with glycerin. - D.

incorporate the water in which the borax has previously been dissolved; finally add the coumarin, oils, and tincture and mix well.

White petrolatum is preferred in the above, as in all cold cream preparations.

Petrolatumav.oz.	12
Soap, powdered, whitegr.	108
Glycerinfl.dr.	
Waterfl.oz.	
Oil of rosedrops	5
Oil of nerolidrops	5
Oil of begamotdrops	5

Mix the petrolatum and soap, then gradually incorporate the glycerin and water previously mixed, and finally add the oils.-D.

Petrolatum, whiteav.oz.	6
Spermacetiav.oz.	3
White waxav.oz.	3
Rose waterfl.oz.	$6\frac{1}{2}$
Borax, powdergr.	40
Oil of rosedrops	

Melt the spermaceti and wax, add the petrolatum, then incorporate the water in which the borax has previously been dissolved, and finally add the oil of rose.

# IV.

White petrolatum	av.oz.	10
White wax		
Cottonseed oil, bleached	fl.oz.	1
Rose water	fl.oz.	1
T) 111 .1 11	4 41	

Prepare like the preceding, adding any desired perfume.

V. Some of the lanolin cold creams contain petrolatum and may be dispensed as a petrolatum cold cream.

# Salicylated Cold Cream. (Salicylic Cold Cream.)

White wor	2
White waxav.oz.	
Spermacetiav.oz.	2
Sweet almond oilfl.oz.	12
Distilled waterfl.oz.	2
Glycerinfl.oz.	2
Salicylic acidgr.	90
Coumarin	1/2
Oil of rosedrops	5
Oil of nerolidrops	5
Oil of bergamotdrops	5
Oil of wintergreendrops	3
Oil of ylang ylangdrop	1
Tincture of muskdrops	2
•	

Prepare in the usual manner for making cold cream, but add the salicylic acid last, Melt the wax and spermaceti, add the previously triturating it to a smooth paste

### Rose Cold Cream.

Rose cold cream may be prepared from any of the "cold creams" by tinting a rose color by means of solution of carmine.

# Sultana Cold Cream.

Spermacetiav.oz.	$\frac{1}{2}$
White waxav.oz.	$\frac{1}{2}$
Sweet almond oilfl.oz.	8
Cacao butterav.oz.	8
Orange flower waterfl.oz.	$^2$
Peru balsamav.oz.	

Melt the wax and spermaceti, add the cacao butter, melt again, add the oil, and then incorporate the orange flower water and balsam.

# Witch Hazel Cold Cream. (Witch Hazel Cream.)

White petrolatumav.oz.	9
White waxav.oz.	
Spermacetiav.oz.	
Distilled extract of witch hazelfl.oz.	3

Melt the first three ingredients together; allow to cool to some extent, then add the witch hazel extract and stir. When nearly cold add any desired perfume and stir vigorously.

# Perfumes for Cold Cream.

I. Coumaringr. 1
Oil of rosedrops 30
Oil of nerolidrops 10
Oil of rose geraniumdrops 7
Oil of ylang ylangdrops 3
Oil of orrisdrop 1
Tincture of ambergrisdrops 10
II.
Coumaringr. 2
Oil of rosedrops 20
Oil of nerolidropś 20
Oil of bergamotdrops 20
Oil of wintergreendrops 12
Oil of ylang ylangdrops 2
Tincture of muskdrops 10
III.
Coumaringr. 1
Oil of bergamotdrops 40
Oil of rosedrops 10
Oil of bitter almonddrops 12
Tincture of ambergrisdrops 7
These mixtures may be employed for per-

## Cucumber Cream or Ointment.

fuming cold cream.-D.

White waxav.oz.	3
Spermacetiav.oz.	3
Benzoinated lardav.oz.	
Cucumbersav.oz.	3
Borax, powdergr. 1	150

Melt the wax and spermaceti, and add the lard. Peel the cucumbers, chop into small pieces, mix with the foregoing and add 90 grains of borax. After this mixture has stood for 12 hours, melt on a water bath, and strain through cheese cloth. Place the mixture on ice; when the fat has solidified, drain off the fluid, add the remainder of the borax to the fat, and thoroughly incorporate the fat with the borax and the residual fluid it contains. This may be colored green with chlorophyll if desired.

II.

Melt together 11 av. ounces of lard and 7 av. ounces of veal suet, the suet first having been rendered, in a water bath. To the fat, strained into a jar of the capacity of one gallon, when it begins to thicken add onethird of the juice obtained by grating and expressing 3 av. pounds of green cucumbers and beat the mixture with a spatula until all of the odor of the cucumber has been absorbed by the fat, which will require several hours. The watery fluid should then be rejected and the remaining juice added in two consecutive portions, incorporated in the previous manner and the watery liquid again rejected. The fat should then be heated in a closed vessel on a water bath for one hour. the albuminous coagulum should be skimmed off, the watery matter should be removed on cooling, the ointment should again be melted and then strained. Before use, this fat should be triturated with a little rose water until it is of a creamy character.

### III.

Lardav.oz. 1	0
Lanolin, anhydrousav.oz.	5
Tincture of benzoinfl.oz.	
Cucumber juice, freshly ex-	
pressed and strained through	
absorbent cottonfl.oz. 1	0
Min the land and langlin incomposate	+1

Mix the lard and lanolin, incorporate the tincture, and lastly the juice.—H. modified.

V•	
Lardav.oz.	10
Beef tallowav.oz.	6
Benzoic acidgr.	15
Rose waterfl.dr.	1
Cucumber juice, freshly ex-	
pressed and strained through	
absorbent cottonfl.oz.	10

Melt the lard and tallow, add the acid,

previously triturated to fine powder, add the cucumber juice, and incorporate thoroughly, as described under "cold cream." Perfume may be added if desired..

V.

Cucumbers, cut fine or grated av.oz. 3	0
Alcohol fl.dr. 1	
Benzoinated lardav.oz. 1	0
Beef tallowav.oz.	5

Mix the cucumbers and alcohol, macerate for one hour, and then distill off 9 fluid-drams. This distillate may be known as spirit of cucumber.

Melt the lard and tallow on a water bath and incorporate with it the above distillate.—H.

The tallow above may be replaced by 3 av. ounces of spermaceti.

# Skin Food.

Preparations sailing under this name are usually cold creams or other bland ointments suitably perfumed and sometimes tinted. Fat forms the basis and gives them their hygienic effect as it imparts fullness and softness to the skin. When applied, these preparations are to be rubbed into the skin abundantly, and the friction asssits the absorbed fat in developing the muscles. Any of the cold creams may be dispensed under the name of "skin food" or the following may be used:

Petrolatum, whiteav.oz.	14
Paraffin waxav.oz.	
Lanolin av.oz.	4
Waterfl.oz.	6
Oil of rosedrops	5
Vanillingr.	4
Alcohol fl.dr.	2

Melt the wax, add the petrolatum and lanolin, pour into a warm mortar, and, with constant stirring, incorporate the water. When nearly cold, add the oil and vanillin dissolved in the alcohol.

These preparations may be tinted red by means of alkanet root.

# Face or Toilet Creams.

These preparations are mucilaginous in character, consisting of tragacanth or quince seed mucilage combined with water, alcohol, glycerin, borax, or boric acid, and sometimes other ingredients. Their purpose is for application to face, neck and hands to assist in through muslin.

healing cracks and all roughness of the skin. They are readily absorbed and dry quickly. The formulas do not specify coloring matter, but the preparations may be tinted any desired color; red, for example, by means of solution of carmine; violet by means of violet aniline, etc.

Such preparations may be dispensed under the names "face cream," "toilet cream," "skin cream," "cosmetic cream," "glycerin cream," "fragrant cream," or similar titles. If containing menthol, the title "menthol cream" may be used; if containing camphor, "camphor cream"; if containing calendula, "calendula cream," etc. Other titles which are used are "cream of roses, "cream of lilacs," "calendula and glycerin lotion," etc. I.

Tragacanth, wholegr.	50
Waterfl.oz.	
Glycerinfl.oz.	2
Alcohol	1
Boric acidgr.	80

Macerate the gum in the water until perfectly soft, strain by expression through muslin, dissolve the acid in the glycerin by the aid of heat, add to the mucilage, then incorporate the alcohol, perfume to suit, and add if necessary enough water to make 16 fluidounces.

II.

Gum tragacanth, whole av.oz.	3/4
Glycerinfl.oz.	8
Salicylic acidgr.	60
Alcoholfl.oz.	8
Water fl.oz.	48

Soften the gum in about 32 fluidounces of water, strain forcibly through muslin, add the remainder of the water and the other ingredients, first dissolving the acid in the alcohol. Shake well and perfume to suit with some "extract."

III.

Gum tragacanth, whole	gr. 120
Waterfl	.oz. 14
Glycerinfl	.oz. 1
Tincture of benzoinfl.	.dr. 2
Borax, powder	gr. 120
White rose extractfl	.dr. 4

Macerate the tragacanth in the water until perfectly soft, and dissolve the borax in the glycerin. Mix the two solutions, add the tincture and strain with expression through muslin.

IV.	
Tragacanth, powdergr.	240
Glycerinfl.oz.	
Borax, powdergr.	
Water, enough to makefl.oz.	

Dissolve the tragacanth and borax in the water and strain through cheese cloth, add the glycerin, and perfume to suit.

Quince seed	3/
DOLIC acidgi. ov	
Carbolic acid	
Tincture of benzoinfl.dr. 12	
Glycerinfl.oz. 6	
Lilac extractfl.dr. 2	
Water, enough to makefl.oz. 64	

Pour 32 fluidounces of boiling water upon the seed, add the two acids, macerate for several hours, stirring occasionally, decant the liquid, upon the residue pour 16 fluidounces of boiling water, macerate again as before, decant as before, mix and strain the two liquids, add the glycerin, incorporate the tincture gradually with vigorous agitation, add the perfume, and finally the remainder of the water.

# Calendula Toilet Cream.

Tragacanth, powdergr.	120
Boraxgr.	
Ammonium chloridegr.	
Tincture of calendulafl.oz.	
Glycerinfl.oz.	
Waterfl.oz.	

Triturate the gum with glycerin to a smooth paste; dissolve the borax and ammonium chloride in a portion of the water, filter this solution, mix the two liquids, add the remainder of the water, and incorporate the tincture (and some "extract" to perfume) with this mixture by agitation.

# Glycerin Toilet Cream.

Quince seedgr. 9	90
Boric acidgr.	
Glycerinfl.oz.	4
Alcoholfl.oz.	
Carbolic acidfl.dr.	$\frac{1}{2}$
Cologne waterfl.dr.	4
Oil of lavender flowersdrops 4	
Glycerite of starchav.oz.	4
Waterenough to make fl.oz. &	

Dissolve the boric acid in 16 fluidounces of water, macerate the quince seed in the solution for three hours, press through a straining cloth, add the glycerin, carbolic acid and glycerite, and mix well. Mix the

alcohol, cologne water and lavender oil, add the mixture to the mucilage and mix the whole thoroughly.

# Camphor Toilet Cream.

Quince seedgr. 6	0
Water, hotfl.oz. 1	4
Borax, powdergr. 6	0
Glycerinfl.oz.	
Spirit of camphorfl.oz.	1
Oil of bitter almonddrops 1	0

Macerate the quince seed for several hours with the water, strain, add the glycerin and in the mixture dissolve the borax. Dissolve the oil in the spirit of camphor and gradually add to it the mucilage and mix thoroughly.

# Menthol Toilet Cream.

Tragacanth, powderg	r. 80
Glycerinfl.d	
Menthol	
Alcoholfl.d	
Water, enough to makefl.o	
Solution of carminesuffi	

Triturate the tragacanth with the glycerin, dissolve the menthol in the alcohol, mix the gummy and alcoholic liquids by trituration, color a light pink by the addition of the carmine solution, and then gradually and thoroughly incorporate the water.

# Creams Containing Almond.

Some toilet creams differ from the preceding kind in that they contain almond in some form, either an emulsion of sweet or bitter almonds, sweet almond oil in emulsion or saponaceous form, or a flavor of bitter almond. These receive such names as "almond cream," "honey and almond cream" (usually these contain no honey), and "amandine"; also "cream of roses," if flavored strongly with rose or if tinted a rose color (for example, Nos. III, V and VI).

These preparations are used for cleaning, softening and whitening the skin.

ı.		
	Ointment of rose waterav.oz.	1
	Oil of sweet almondsfl.oz.	
	Glycerinfl.oz.	1
	Boric acidgr.	100
	Solution of sodafl.oz.	21/4
	Mucilage of quince seedfl.oz.	4
	Water, enough to makefl.oz.	40
	Oil of rose,	
	Oil of bitter almonds. of each,	

Heat the ointment, oil and solution of soda

together, stirring constantly until an emulsion or saponaceous mixture is formed. warm together the glycerin, acid, mucilage and about 30 fluidounces of water; mix this with the emulsion, stir until cold and add the remainder of the water. Lastly, add the volatile oils.

The rose water ointment used should be the "cold cream" of the U.S. P.

### II.

Sweet almonds, blanched av. oz.	5
Castile soap, whitegr.	120
White waxgr.	
Spermacetigr.	
Oil of bitter almondsdrops	
Oil of bergamotdrops	
Alcoholfl. oz.	6
Water suffic	ient

Make an emulsion of the almonds with water so as to obtain 16 fluidounces of product, straining through cotton, which has previously been washed to remove stiffening. Dissolve the soap with the aid of heat in the necessary amount of water to form a liquid, add the wax and spermaceti, continue the heat until the latter is melted, transfer to a mortar, and incorporate the almond emulsion slowly with constant stirring until all has been added and a smooth cream has been formed. Finally add the two volatile oils.

### III.

Sweet almonds, blanched av. oz.	8
Rose waterfl. oz.	32
Alcoholfl oz.	4
Oil of rosesfl. dr.	1
White waxgr.	240
Spermaceti gr.	240
White castile soapgr.	240

Shave the soap, place it in a vessel, add several ounces of rose water and heat on a water bath until dissolved. When the soap is dissolved, add the wax and spermaceti, continue the heat and stir occasionally. While this is going on, blanch the almonds, carefully excluding every damaged particle. Then beat them up in a scrupulously clean mortar, and allow the rose water to trickle into the mass by degrees. When the emulsion of almonds is finished, strain it without pressure, through clean-washed muslin. The previously prepared saponaceous mixture is now put in the mortar, and the emulsion carefully and gradually blended with it. As the last of the tion to impart a rose tint.

emulsion is run into the mortar, the alcohol, in which the oil of rose has been dissolved, is made to follow it and mixed very gradually with the other ingredients. A too sudden addition of the alcohol frequently coagulates the milk and causes it to be curdled. As it is, the temperature of the mixture rises, and every means must be taken to keep it down. Finally strain the product.

The almond residue may be washed with a few ounces of fresh rose water, to prevent any loss in bulk in the whole quantity. The newly formed milk should be allowed to stand at rest for 24 hours, when the clear portion may be drawn off the sediment, and is ready for bottling.

### IV.

Spermacetigr.	240
White waxgr.	
Castile soap, whitegr.	
Bitter almonds, blanched av. oz.	
Alcoholfl.oz.	6
Oil of bitter almondsdrops	10
Oil of bergamotfl.dr.	1
Water, distilledfl.oz.	16

Triturate the almonds with the water to a smooth paste, melt the spermaceti and wax by means of a water bath, mix the two intimately, slowly add the alcohol in which the oils have previously been dissolved, and finally strain through cheese cloth.

V. The bitter almonds in the preceding may be replaced by sweet almonds and oil of rose substituted for the oils of bitter almond and bergamot.

## VI.

Honey	av.oz.	2
Castile soap, white powder.		
Oil of sweet almond		
Oil of bitter almonds	fl.dr.	1
Oil of bergamot	fl.dr.	1/2
Oil of cloves		
Peru balsam		
Liquor potassa,		
Civilia	.1 .00 .	,

Solution of carmine...of each, sufficient

Mix the honey with the soap in a mortar, and add enough liquor potassa (about 1 fluidram) to produce a nice cream. volatile oils and balsam with the sweet almond oil, mix this with the cream, and continue the trituration until thoroughly mixed. Finally add, if desired, enough carmine solu-

VII.	
Bitter almondsav.oz.	3
Yolk of eggav.oz.	3
Honeyav.oz.	6
Sweet almond oilfl.oz.	6
Oil of bergamotdrops	45
Oil of lemondrops	35
Oil of clovesdrops	35

Bruise the almonds, previously macerated in cold water, and deprived of its coat, and rub through a fine sieve; add the essential oils and the mixed yolk of egg, honey and almond oil, and beat all together until they are incorporated.

# Milk of Roses.

Some toilet preparations are known by this title. As examples may be mentioned Nos. III, V and VI, under the preceding title. The following formulæ may also be used for making a preparation of this name:

# I. English milk of roses.

Sweet almonds, blanched av.oz.	11/2
Sweet almond oilfl.dr.	1
Soft soap (as white as possible)gr.	60
Rose waterfl.oz.	
Oil of rosedrops	
Alcoholfl.oz.	

Make an emulsion of the sweet almond oil, soap and water, strain and incorporate the oil of rose previously dissolved in the alcohol. B.

Sweet almond oil fl.oz. 1	
Soft soap (as white as possible).av.oz. 1	L
Potassium carbonategr. 30	)
Water, boilingfl.oz. 4	Ł
Alcoholfl.oz. 2	3
Oil of rosedrop 1	L
Rose water, enough to make fl.oz. 16	;
The state of the s	

Mix the first four ingredients intimately, allow to cool, add the alcohol containing the oil of rose, and finally add the rose water.

# II. French milk of roses.

١.	
Tincture of benzoinfl.dr. 4	Ŀ
Tincture of storaxfl.dr. 2	,
Spirit of rosefl.dr. 2	
Alcohol fl.oz. 2	
Rose waterfl.oz. 16	1/2

Mix the first four ingredients, and add the rose water gradually with frequent and vigorous agitation.

В.		
Tincture of benzoin	fl.dr.	2
Tincture of Peru balsam	drops	40
Alcohol		
Rose water	fl.oz.	14

Mix the tinctures and alcohol and then add adhere more readily.

rose water gradually and with vigorous agitation.

# III. German milk of roses.

Diluted solution of lead subace-	
tatefl.oz.	1
Spirit of lavenderfl.oz.	1
Rose waterfl.oz.	
Water fl.oz.	16
Waterfl.oz.	16

While this formula is given here for the sake of completeness, its use is not recommended owing to the presence of the poisonous lead compound.

### IV.

* •	
Salicylic acidgr.	15
Benzoic acidgr.	61
Rose waterfl.oz.	28
Glycerin	10
Alcoholfl.oz.	2
Tincture of benzoinfl.dr.	5
Oil of rosedrops	5
Oil of bergamotdrops	2
Oil of nerolidrop	1
Weaker tincture of orrisdrops	3
Essence of jasminedrops	10
Tincture of muskdrops	6
Coumarin sugargr.	30
Solution of carminesufficie	ent
Solution of Carmine	

Triturate the two acids with the rose until an intimate mixture is formed, then add the remaining ingredients, and mix well, using enough of the solution of carmine to impart a pale rose tint.—D.

### Toilet Milks.

Some toilet washes are known by the appellation of "milks." These consist of an oily substance combined with borax, powdered soap, and a large quantity of water, and are therefore in a sense an imperfect and thin soap, the whole being suitably flavored. The only substance may consist of lanolin, cacao butter, cocoanut oil or a combination of these. According to this fatty constituent, the preparation receives the name of "cocoa milk," "cacao milk" or "lanolin milk." Another "milk" which is also used is "cucumber milk," which also contains borax, some oil or soap, and cucumber juice.

These preparations are used as a substitute for cold cream in cleansing the skin; it may also be applied to the skin at night instead of cold cream and may be applied to the face before using powder, to enable the latter to adhere more readily.

110
1/2
34
134
20
8
8
4
1
1

Γriturate the first four ingredients together until well mixed, then gradually add the rose water previously warmed to 40 degs. C., triturating constantly during this addition, and add the oils. - D.

II.

White castile soap, powdergr.	22
Lanolinav.oz.	
Tincture of benzoinfl.dr.	$1\frac{1}{2}$
Distilled watersufficie	ent

Dissolve the soap in 2 fluidounces of warm water, also mix the lanolin with 2 fluidounces of water, then incorporate the two with each other, finally adding the tincture. The latter may be replaced by 90 gr. of powdered borax. III.

Borax, powdergr.	140
White castile soap, powderav.oz.	
Cacao butter, gratedav.oz.	11/2
Cocoanut oilav.oz.	
Waterfl.oz.	. 2
Rose waterfl.oz.	28
Oil of bergamotdrops	20
Oil of nerolidrops	5
Oil of orris drop	) 1
Vanilla sugargr.	150

Triturate the first five ingredients together in a warm mortar until well mixed, then gradually incorporate the rose water previously warmed to 40 degs. C., and add the remaining ingredients previously triturated together.-D.

IV.

v •	
Borax, powdergr.	90
White castile soap, powdergr.	180
Cocoanut oilav.oz.	$1\frac{1}{2}$
Waterfl.oz.	1
Rose waterfl.oz.	17
Oil of bergamotdrops	8
Oil of nerolidrops	4
Oil of wintergreendrops	2
Oil of ylang ylangdrop	1
Oil of bitter almonddrop	1

Triturate the first four ingredients together until well mixed, gradually add the rose water, previously heated to 40 degs. C., until or cochineal if desired. 21

thoroughly incorporated and finally add the oils.—D.

V. Take fresh strained cucumber juice, bring to a boil as quickly as possible, cool down rapidly, and to every 5 fluidounces of this juice, add:

Borax, powdergr.	175
Sodium acetategr.	
Tincture of quillajafl.oz.	
Tincture of benzoinfl.dr.	4
Rose or orange flower waterfl.oz.	16

Mix the whole thoroughly. It may be tinted a pale green with chlorophyll and perfumed with essence of cassie.

VI.

Cucumber	juice,	boiled	and		
cooled				.fl.oz.	4
Spirit of so.					
Rose or ora					

This may be colored and perfumed like the preceding.

VII. The last two are known as 'cucumber milk" or "milk of cucumbers"; this preparation has received the same title, but is devoid of cucumber in any form:

Borax, powdergr.	225
Sodium acetategr.	
Spirit of soapfl.dr.	5
Tincture of benzoinfl.dr.	5
Glycerinfl.dr.	10
Rose waterfl.oz.	21
Oil of bergamotdrops	4
Oil of rosedrops	2
Spirit of orrisfl.dr.	11/2
Tincture of muskdrops	3
Coumarin sugargr.	12
•	

Dissolve the borax and sodium acetate in the rose water, add the remaining ingredients, and mix well.-D.

#### Toilet Lotions.

Under this title are included transparent preparations, which are employed instead of some of the preceding "creams" and "milks," for roughnesses of the skin, cracked hands, These may be denominated "face lotion," "glycerin lotion," "cosmetic lotion."

Glycerin.....fl.oz. 3 Rose or orange flower water....fl.oz. 13

This may be tinted with solution of carmine

II.	
Castile soap, whiteav.oz.	1/2
Honeygr. 120	)
Boraxgr. 120	)
	3
Glycerinfl.oz.	3
Alcoholfl.oz.	3
Solution of cochineal, or car-	
minesufficient to colo	r
Water, enough to makefl.oz. 10	3

Dissolve the soap and borax in boiling water, allow to cool, add the other ingredients, macerate for 24 hours, and filter.

#### TIT.

Tincture of arnicafl.oz. 4
Glycerinfl.oz. 8
Rose waterfl.oz. 4
Mix and filter.

#### $\mathbf{T}\mathbf{V}$

V.	
Boraxgr.	300
Glycerinfl.dr.	10
Rose waterfl.oz.	-31
Coumarin sugargr.	15
Tincture of ambergrisdrops	3
,	
Oil of rosedrops	6
Oil of neroli drop	1
Solution of carminesuffic	ient

Dissolve the borax and sugar in the water, add the glycerin and other ingredients, shake well, and filter. Sufficient of the carmine solution is to be used to impart a pale rose tint.-D.

#### Toilet or Cosmetic Jellies.

These are preparations of stiff or thick consistence intended as emollient toilet applications for the skin. The body consists either of gelatin, starch, tragacanth, or similar substance. Other common ingredients are glycerin, water, and perfumed ("extract," essence, or volatile oil). Other additions are boric acid, salicylic acid, fluid extract or tincture of arnica, fluid extract of calendula and distilled extract of witch hazel. Sometimes they are tinted a pale rose color with cochineal or carmine.

The common appellation for these jellies is "glycerin jelly." If containing arnica, they may be known as "arnica jelly"; if containing calendula, "calendula jelly"; witch hazel, "witch hazel jelly"; if tinted a rose color and flavored with oil of rose, "rose jelly" or "jelly of roses," etc.

tions must be dispensed in wide-mouthed bottles, or jars, or in collapsible tubes.

Gelatingr. 160 to 24	0
Boric acidgr. 24	10
Glycerinfl.oz.	
Waterfl.oz. 1	.0

Perfume to suit.

Dissolve the gelatin in the water by the aid of heat, also the acid in the glycerin, mix, allow to cool somewhat and incorporate the perfume.

The amount of gelatin may be varied to suit the thickness desired.

The perfume must be one which mixes without opalescence, or otherwise it mars the beauty of the preparation. Orange flower water or rose water may be substituted for the water if desired, or another perfume consisting of:

Vanillingr. 4	
Coumaringr. 4	
Spirit of bitter almondfl.dr. 1	1/2
Alcoholfl.dr. 3	

added to the quantities given above would prove agreeable.

II.

Gelatingr.	240
White of eggav.oz.	
Salicylic acidgr.	25
Rose waterfl.oz.	
Glycerin, enough to makefl.oz.	

Dissolve the gelatin in the rose water by the aid of the water bath, using a gentle heat. Allow to cool, and before it jellifies, add the albumen and stir together. Mix the salicylic acid with the glycerin, and after again applying heat to the gelatin solution, add it to the latter, stirring constantly. When the mixture is quite homogeneous, remove from the fire and filter, by means of a hot filtration apparatus, directly into receptacles in which it solidifies. Instead of rose water, any other distilled perfumed water, such as orange flower water, may be used.

III.

Gelatinav.oz.	1
Glycerinfl.oz.	16
Waterfl.oz.	
Oil of rosedrops	2
Oil of lavender flowersdrops	10

Soak the gelatin in the mixed glycerin and Owing to their thickness, these prepara- water for 12 hours, then heat on a water bath

until dissolved, and finally add the oils. Other flavors may be used. Also other additions may be made; carbolic acid, for example, would make it a carbolated glycerin jelly.

IV.

Fluid extract of arnicafl.oz. 1
Glycerinfl.oz. 6 to 8
Gelatin av.oz. 1
Watersufficient

Cover the gelatin, contained in a suitable vessel, with cold water; allow it to macerate several hours or until soft and pliable; drain off the excess of water, dissolve by heat in the residual water and the glycerin, the quantity of the latter varying with the season, using more in the winter than in hot weather. When dissolved, add the arnica, perfume to suit and color with solution of carmine.

Suitably perfume glycerite of starch and color it with solution of cochineal, and add extract of arnica. This and the preceding are known as "Arnica Jelly."

VI.

Glycerinfl.oz.	61/4
Waterfl.oz.	61/4
Starchav.oz.	
Fluid extract of arnicafl.oz.	11/2
Spirit of bitter almondfl.dr.	21/2
Carbolic acidfl.dr.	1

Mix the glycerin and water, add the starch, rub to a smooth mixture, and heat over a direct flame with constant stirring until a perfectly smooth jelly is formed; allow to cool, and when nearly cold, add the fluid extract, spirit and acid.

VII.

Glycerite of starchav.oz. 14
Fluid extract of calendulafl.oz. 134
Solution of cochineal or car-
minesufficient to color a rose tint
Oil of rosesufficient to perfume
VIII.
Glycerinfl.oz. 32
Tragacanth, powdered, enough

to thicken, or about.....av.oz. Borax . . . . . . . . . . . . . . . . . av.oz. Orris root, powder . . . . . . . . av.oz. Essence of cassie, . . . . . . . . . fl.oz. 1 Essence of jasmine.....fl.dr.

mix intimately with the tragacanth, adding the essences and orris root with trituration.

Owing to the odor, this should be denominated "violet jelly" or "jelly of white violets."

IX.

Tragacanth, powdergr.	
Glycerin	51/2
Waterfl.oz.	101/2

Triturate the gum with the glycerin and water to a smooth paste, and then perfume as desired.

X.

Mucilage of Irish mossav.oz.	4
Glycerinfl.oz.	6
Distilled extract of witch hazelfl.oz.	4
Cologne waterfl.oz.	
Boraxgr.	30

Dissolve the borax in the witch hazel extract, mix with 3 fluidounces of glycerin and with the cologne, add slowly to the mucilage previously mixed with the remainder of the After standing a few hours strain glycerin. the mixture.

Υĭ

¥1.	
Russian isinglassgr.	108
Clarified honeyav.oz.	1
Glycerin fl.oz.	5 1/2
Distilled extract of witch hazel fl. oz.	4
Distilled water	6
Oil of nerolidrops	30

Dissolve the isinglass in the water by aid of a gentle heat, add the witch hazel extract, strain and finally add the oil.

XII.

Spermacetiav.oz.	3
White waxav.oz.	11/2
Sweet almond oilfl.oz.	6
Glycerinfl.oz.	3
Tragacanth, powdergr.	45
Rose waterfl.oz.	

Dissolve the tragacanth in the glycerin, with gentle heat, and add the rose water; melt the other ingredients by heat, add to the glycerin mixture, and beat with an egg-beater until nearly cold, or triturate vigorously in a wide and capacious mortar.

XIII.

Glycerite of starch av.oz.	12
Lanolinav.oz.	4

Triturate the lanolin with a small portion Mix the essences with the powdered orris of glycerite until thoroughly mixed, then add root; dissolve the borax in the glycerin and the remainder of the glycerite gradually.

rubbing thoroughly after each addition. Any suitable perfume may be added.

This preparation may be known as lanolin jelly or lanolin glycerite.

#### Lime Juice and Glycerin.

This is a cosmetic lotion of indefinite character, usually containing no lime juice and frequently even no glycerin.

This preparation is not to be confounded with the preparation of the same title intended for internal use.

The following formulas may be employed: I.

Boraxdr.	2
Sweet almond oilfl.oz.	
Castile soap, whitedr.	2
Waterfl.oz.	
Liquor potassa fl.dr.	3
Perfume to s	uit

Dissolve the soap, finely shredded, and the borax in the water over a water bath, place in a large bottle, and gradually add the oil, shaking well after every addition; then add the liquor potassa, and shake well till cold; lastly add the perfume and give an occasional shake for 12 hours.

II.	
Sweet almond oil fl.oz	6
Castor oilfl.oz	. 2
Lime waterfl.oz	. 4
Glycerinfl.oz	. 2
Mix well by agitation.	

#### III.

Olive oil, best	.fl.oz. 8
Lime water	.fl.oz. 8
Oil of lemon	.fl.dr. 2

#### IV.

Lime juicefl.oz.	8
Rose waterfl.oz.	
Alcoholfl.oz.	2
Oil of lemondrops	24
Oil of lavender flowersdrops	
Glycerinfl.oz.	

Mix the lime juice and rose water and add about two-thirds of the alcohol; shake well together, let stand about 24 hours, strain, add the other ingredients, first dissolving the oils in the alcohol.

#### Camphor Ice.

These are solid preparations containing fatty bodies like wax and spermaceti in combination with camphor, and are intended for inunction of the hands and face where there

is roughness or cracking of the skin. They are prepared by melting the fatty substances, allowing to cool somewhat, stirring in the camphor, allowing to cool, adding flavoring oil, if the latter be used, and pouring into molds. The best material for the latter is block tin. It may be chilled before casting the mixture as this renders adhesion less likely.

Much cheaper, though less elegant, molds may be made of tinned iron.

The usual way of putting up camphor ice for sale is to wrap it first in thin smooth paper, then in an outer covering of tin foil, and lastly to inclose it in a paper box.

Camphor ice may be known by this title; if it contain glycerin, it should be known as glycerin camphor ice (sometimes also known as "compound glycerin cream"), and if containing petrolatum as petrolatum camphor ice.

#### Τ.

Spermacetiav.oz.	2
White waxav.oz.	
Sweet almond oil fl.oz.	8
Camphorav.oz.	2
Oil of bitter almondfl.dr.	1
Expressed oil of macegr.	60

Melt the wax and spermaceti, add the sweet almond oil, then the camphor in small pieces, stir constantly until dissolved, allow to cool, stirring frequently; when quite cool, add the remaining oils, and finally pour the mixture into molds.

#### II.

Mutton suetav.oz.	6
Spermacetiav.oz.	4
White waxav.oz.	
Camphorav.oz.	11/2

Melt together by a gentle heat, reserving the addition of the camphor until the other ingredients are liquefied, on account of its volatility. Stir well as the mixture begins to cool, continuing until ready to set, then pour into molds.

#### III.

Mutton suet, strained or fil-	
tered clearav.oz.	12
Spermacetigr.	
White waxgr.	320
Camphorav.oz.	

bination with camphor, and are intended for Melt the suet, spermaceti, and wax, add inunction of the hands and face where there the camphor in small pieces, stir until dis-

solved, stir occasionally until quite cool, and pour into molds.

# Glycerin Camphor Ice. (Compound Glycerin Cream.

Stearin (stearic acid) av.oz.	
Lardav.oz.	5
White waxav.oz.	21/2
Spermacetiav.oz.	21/2
Borax, powder gr.	
Glycerinfl.dr.	
Camphorav.oz.	
1	

Melt the first four ingredients on a water bath. Dissolve the borax in the glycerin. Add the latter gradually to the former; when at the point of cooling, stir well, add the camphor in pieces, stir again until dissolved and pour into molds.

H.

L.	
Spermacetiav.oz.	2
White waxav.oz.	
Olive, castor or cottonseed or	, -
other pure bland, fixed oil fl.oz.	10
Camphorav.oz.	11/2
Glycerinfl dr.	
Rose waterfl.oz.	11/2
Boraxgr.	40
9	

Melt together the two fats, add the oil, then the camphor and stir constantly until dissolved. Now, add the glycerin mixed with the rose water in which the borax has previously been dissolved, stir the whole until nearly cold, and pour into molds.

#### Petrolatum Camphor Ice

Paraffin waxav.oz. 5	1/2
White petrolatumav.oz. 8	
White waxav.oz. 3	3/1
Camphorav.oz. 1	

Melt the two waxes together, add the petrolatum, and then the camphor in pieces, stir until the latter is dissolved, allow to cool, and pour into molds.

#### Toilet Lanolin.

Under this heading are included fatty combinations consisting mainly of lanolin, which may be put up in stick form by the usual method of casting in molds. (See "Camphor Ice.")

I.

Benzoinated suetav.oz.	3
Lanolinav.oz.	6
Boric acid, powderav.oz.	1

Melt the suet, add the lanolin, stir in the acid, and form into sticks.—D.

11	•	
	Benzoinated suetav.oz.	2
	Yellow waxav.oz.	
	Lanolinav.oz.	51/2
	Carbolic acid, crystalav.oz.	1/2
	· ·	,

Melt the wax and suet, add the lanolin and acid, and form into sticks.—D.

TTT

Benzoinated suetav.oz.	21/2
Yellow waxav.oz. Lanolinav.oz.	3.4
Lanolinav.oz.	$6\frac{1}{2}$
Salicylic acidgr.	90

Melt the suet and wax, stir in the acid, add the lanolin, and form into sticks.—D.

#### Almond Paste. (Amandine.)

L.		
î	Bitter almondsav.oz.	7
	Orris root, powderav.oz.	13/
	White castile soap, powderav.oz.	134
	Glycerite of starch av.oz.	31/2
	Clarified honeyav.oz.	2
	Oil of lavender flowersfl.dr.	1
	Oil of bergamot fl.dr.	2
	Oil of bitter almonds drops	8
	Solution of cochineal, to color sufficient	nt

Blanch the almonds, heat them with a small quantity of water to a smooth paste, add the other ingredients, and mix intimately.

TT

Π		
	Sweet almonds, blanchedav.oz.	71/2
	Bitter almonds, blanchedav.oz.	5′~
	Borax, powdergr.	320
	Liniment of camphorfl.dr.	10
	Spermacetiav.oz.	11/4
	Starchav.oz.	$5^{'}$
	Talcum, powderav.oz.	21/2
	Rose waterfl.oz.	51/2
	Oil of bergamotdrops	12
	Oil of rosedrops	6
	Oil of cassiadrops	4
	Oil of clovesdrops	2
	Oil of sassafras drops	2
	Oil of ylang ylangdrop	1
	Oil of orris, liquiddrop	1
	Tincture of civetdrops	10
	Tincture of muskdrops	5
	Coumaringr.	1

Mix the first three ingredients and 4 fluidrams of rose water intimately so as to form a perfectly smooth paste. To this mixture add the liniment and spermaceti previously melted together, now incorporate the talcum and starch previously made into a smooth paste with the remaining rose water, and finally add the other ingredients, also previously mixed so as to dissolve the coumarin.

The whole may be colored a pale rose tint, if	
desired, by means of alkannin.—D.	

#### III.

Bitter almonds, blanched	av.oz. 4
Honey	av.oz. 7½
Yolk of egg	
Sweet almond oil	fl.oz. 7
Oil of bergamot	fl.dr. 1
Oil of cloves	fl.dr. 1

Beat the almond to a fine paste, then gradually and thoroughly add the remaining ingredients, which have previously been well mixed.

#### IV.

Sweet almonds, blanchedav.oz.	8
	-
Bitter almonds, blanchedav.oz.	8
Rose waterfl.dr.	3
Eggs, white and yolk	3
	1
Borax, powderav.oz.	.1.
Potassium carbonate, fine pow-	
dergr.	150
Glycerinfl.oz.	$2\frac{1}{2}$
Corn meal flourav.oz.	8
Tincture of curcumasufficient to c	olor
Oil of rosedrops	20
Oil of bergamotdrops	8
Oil of nerolidrops	5
Oil of rose geraniumdrops	2
Oil of sassafrasdrops	2
Oil of orris, liquid drop	1
On or orns, nquid urop	
Tincture of muskdrops	6
Coumaringr.	11/2
	$\tilde{8}^{\prime 2}$
Vanillingr.	0

Reduce the sweet and bitter almonds with the rose water to a smooth paste, mix the eggs, borax, potassium carbonate, and glycerin, add to the preceding mixture, then incorporate the flour, the curcuma tincture and the perfumes, the latter being previously mixed so as to dissolve the coumarin and vanillin.—D.

#### Almond Meal. (Mandelklei.)

Sometimes this is prepared from ordinary bitter or sweet almonds and sometimes from almonds from which the oil has been expressed. This is ground fine, sifted, mixed with powdered orris, soap or other ingredients, and perfumed.

#### I.

Almonds, blanched and powderedav.oz. 8
White castile soap, dried and
powderedav.oz. 4
Orris root, powderav.oz. 1
Pumice stone, fine powderav.oz. 3
Oil of bitter almondsfl.dr. 1

-	
	II.
	Sweet almonds, blanched and
	powderedav.oz. 8
	Wheat flourav.oz. 8 Orris root, powderav.oz. 2
	Oil of lemon
	Oil of bitter almonddrops 8
	III.
ĺ	Almond meal, prepared from
	blanched bitter almonds from
	which the oil has been ex-
	pressed, in very fine powder.av.oz. 6 Orris root, fine powderav.oz. 4
	Wheat flourav.oz. 4
	White castile soap, powderav.oz. 1
	Borax, powderav.oz. 1
	Oil of bitter almonds drops 10 Oil of bergamot
	Oil of bergamot
	Mix well and pass through a fine sieve.
	IV.
	Sweet almonds, blanched and in
	fine powderav.oz. 9
	Bean flour (or starch)av.oz. 9
	Orris, fine powder
	White castile soap, powderav.oz. 3 Spermacetiav.oz. 34
	Sodium carbonate, driedav.oz. ½
	Oil of bergamotfl.dr. 1
	Oil of lemonfl.dr. 1
	Oil of lavender flowersfl.dr. 1
	Mix all intimately to form a fine powder
	and sift.
	V
	Sweet almonds, blanched and
	powderedav.oz. 2 Bitter almonds, blanched and
	powderedav.oz. 1
•	Orris root, powderav.oz. 7

To reduce the cost the almonds may be replaced by almond press cake (deprived of oil), orris root by wheat flour, and the volatile oils reduced or changed.

#### VI.

1.
Cacao butterav.oz. 2
Talcum, powderav.oz. 4
Bean flour (or starch)av.oz. 20
Sweet almonds, blanched and
powdered av.oz. 10
Glycerin
Cologne waterfl.oz. 2
Coumaringr. 2
Oil of bitter almondsdrops 25
Tincture of ambergrisdrops 6
Melt the cacao butter, thoroughly incor-

porate the talcum, then add the bean flour (or starch) and the almonds. Dissolve the coumarin and oils in the cologne water, add the glycerin and tincture, and incorporate this mixture with the preceding combination—D.

#### Complexion or Face Powders.

Complexion powders contain such constituents as talcum, starch, precipitated chalk, bismuth subnitrate, oxide, hydrate, subcarbonate or oxychloride, zinc oxide, magnesium carbonate or oxide, and orris root, together with suitable perfume and with coloring matter if a colored powder be desired. These solids must be in the form of fine powder, must be intimately mixed, and then sifted through a fine bolting cloth sieve; whatever fails to pass through the sieve must be returned to the mortar and be still further triturated until all will pass through. If colored powder is to be prepared, the coloring matter should be added to the powder before sifting. After sifting, the perfume may be added; the whole should be again passed through the sieve to break up any lumps which may have formed by the addition of the moist perfume to the dry powder.

If a flesh-colored powder is desired, carmine is employed as the coloring agent. Sometimes a brunette powder is desired and then burnt umber is used as the coloring agent; cream powder is prepared by the use of cadmium yellow or chrome yellow (lead chromate), a trace of carmine being added sometimes. In the formulas given below, no coloring agents are mentioned, it being intended that these should be added only as desired.

In preparing face powders, the best materials should be employed. For example, Hubbuck's zinc oxide only should be used. The best talcum is what is known as Venetian chalk; the best precipitated chalk is the kind known as the English; the preferred bismuth subnitrate should be a very light variety, etc.

Some face powders are made with a lead salt; the latter should, owing to its poisonous character, never be employed.

Mix well and perfume with a mixture of oils of bergamot, ylang ylang and nerch.

This preparation is known as "Sarah Bernhardt's Face Powder," also as "la Diaphane."

Any other perfume may be used if desired.

I.
Talcum, powder.....av.oz. 15

Starchav.oz.	
Orris rootav.oz.	$1 \nu$
Oil of bergamotdrops	12
III.	
Zinc oxideav.oz.	4
Precipitated chalkav.oz.	24
Talc, powderav.oz.	4
Starchav.oz.	8
Essence of rosefl.dr.	2
Essence of jasminefl.dr.	2
Essence of orange flowersfl.dr.	2
Essence of cassie fl.dr.	2
Tincture of muskfl.dr.	1
If this powder be too light, a porti-	on o
the precipitated chalk may be replaced	witl
prepared chalk.	
IV.	
Talcum, powderav.oz.	5
Zinc oxideav.oz.	5
Chalk, preparedav.oz.	21
Starchav.oz.	15
Essence of jasminefl.dr.	4
Oil of bergamotdrops	15
Oil of rosedrops	8
Oil of ylang ylangdrops	2
Oil of nerolidrops	8
Oil of orrisdrop	1
Tincture of muskdrops	5
V.	
Talcum, powderav.oz.	16
Bismuth oxideav.oz.	1
Zinc oxideav.oz.	1
Perfume to suit.	
VI.	
Zinc oxideav.oz.	4
Rice powderav.oz.	14
Precipitated chalkav.oz.	4
Talcum powderav.oz.	2
Orris root, powderav.oz.	2
Perfumesufficie	ent
VII.	
Zinc oxideav.oz.	2
Orris root, powderav.oz.	2
Pice flour	16

Rice flour .....av.oz. 16

Oil of rose geranium......drops

Oil of ylang ylang......drop Coumarin .....gr.

Acetic ether......drops 10

Mix the first three ingredients, mix the

other ingredients so as to dissolve the coumarin, and incorporate this mixture with the powder.—D.

#### VIII.

Zinc oxideav.oz.	2
Orris root, powderav.oz.	$^4$
Talcum, powderav.oz.	
Starchav.oz.	
Essence of jasminefl.dr.	21/2
Oil of rosedrops	5
Oil of bergamotdrops	5
Oil of ylang ylangdrop	1
Tincture of musk drops	5
Coumaringr.	1/2

Mix the first four ingredients intimately, dissolve the coumarin in the essence, add the oils and tincture, and incorporate this mixture with the powder.—D.

#### IX.

Zinc oxide	77)
Coumaringr. Acetic ethersufficien	

Mix the first four ingredients intimately, add the oils, tincture and coumarin, first dissolving the latter in a small amount of acetic ether.—D.

## Х.,

Zinc oxideav.oz.	4
Starchav.oz.	6
Talcum, powderav.oz.	4
Oil of bergamotdrops	
Oil of rosedrops	
Oil of lemondrops	
Oil of orris, liquiddrop	
Tincture of ambergrisdrops	
Coumaringr	1/2
Acetic etherm.	50

Mix the first three ingredients intimately to smooth powder, and add the remaining ingredients previously mixed, so as to dissolve the coumarin. One-half of the zinc oxide may be replaced by bismuth subnitrate.

—D.

#### XI.

Magnesium carbonate av.oz.	8
Talcum, powderav.oz.	8
Oil of rosedrops	8
Oil of nerolidrops 2	
Essence of jasminefl.dr.	
Tincture of muskfl.dr.	

#### XII.

Bismuth subcarbonate	.av.oz	1/2
Zinc oxide	.av.oz. 4	Į.
Talcum, powder	.av.oz.	5
Precipitated chalk	av.oz.	5
Starch		
Oil of rose geranium	.drops 40	)

#### XIII.

Lanolin, anhydrousav.oz.	1
Starchav.oz.	
Talcum, powderav.oz.	20
Coumaringr.	24
Oil of rosedrops	
-	

The lanolin and the perfume are gradually mixed, the talcum, and then the starch is added.

Lanolin may also be incorporated in face powders by dissolving some volatile solvent like ether, chloroform, or benzine, incorporating this solution quickly with magnesia, chalk or other powder, allowing the solvent to vaporize, and incorporating other suitable ingredients with the residue.

Lanolin is introduced into some face powders owing to the dryness of the skin or to prevent the latter from becoming dry and scaly. The fat imparts to the powder a desirable smoothness, increases the power to adhere to the skin, and preserves the latter in a smooth and supple condition.

#### Complexion Tablets.

Any of the preceding powders may be converted into the tablet or cake form by adding a small amount of powdered tragacanth, beating into a stiff paste with water, pressing into suitable molds, and then drying.

#### Toilet and Nursery Powders.

I.

1.	
	Talcum
	Salicylic acidgr. 165
	Boric acidav.oz. 11/4
	All should be in fine powder and should
be	mixed intimately.—N. F.

#### II.

Talcum	<i>.</i>	av.	oz. 11
Starch		av.	oz. $1\frac{1}{4}$
Salicylic acid	.1		gr. 165

Prepare like the preceding.—Germ. Pharm.
This preparation as well as the preceding may be perfumed as desired; or they may also be carbolated.

. SAIN I REI
III.
Starch         av.oz. 15           Orris root.         av.oz. 1           Talcum.         av.oz. ½           Oil of lemon         fl.dr. ½           Oil of bergamot         fl.dr. ½           Oil of neroli         drops 15           Oil of bitter almond         drops 3           Oil of verbena         drops 3           The first three must be in very fine powder
and the whole should be well mixed, after
which it should be sifted through a very fine
sieve.
IV.       Salicylic acid       .gr. 60         Starch       av.oz. 12         Talcum       av.oz. 4         Orris root       .av.oz. 1         Oil of bergamot       .drops 8         Oil of neroli       .drops 4         Tincture of musk       .drops 5         Mix and reduce to impalpable powder.         V.         Orris, powder       .av.oz. 16         Oil of rose       .drops 2         Oil of lavender flowers       .drops 4         Oil of bergamot       .drops 15         Tincture of musk       .drops 15         Essence of violet       .drops 15
VI.       Camphor       .av.oz.       3/4         Zinc oxide       .av.oz.       1         Starch       .av.oz.       4         Mix well, reduce to powder, and sift
through bolting cloth.
This is known as "McCall Anderson's
Dusting Powder."
VII.       av.oz. 1         Boric acid
VIII.
Fuller's earth       av.oz. 8         Talcum       av.oz. 8         Oil of lavender       .drops 5

Mix the first two ingredients in fine

Wheat flour......av.oz. 24
Orris root, powder.....av.oz. 2

Oil of lemon......drops 15

Oil of bergamot......drops 15

Oil of neroli......drops 8

Oil of bitter almond......drop 1 Tincture of musk......drops 8

powder and incorporate the oil.

IX.

X.		
Starch	av.oz.	24
Orris root	.av.oz.	2
Oil of lemon		
Oil of bergamot	drops	20
Oil of neroli	drops	10
Oil of bitter almond	drop	1
Tincture of musk	fl.dr.	$\frac{1}{2}$
Mix the starch and orris	in very	fine
powder and add the other ingre	edients.	

I.	
Fuller's earthav.oz.	9
Boric acidav.oz.	13
Talcumav.oz.	3
Starchav.oz.	9
Orris rootav.oz.	15
Oil of bergamotfl.dr.	2
	Fuller's earth. av.oz. Boric acid. av.oz. Talcum av.oz. Starch av.oz. Orris root av.oz.

XII.	
Fuller's earthav.oz.	2
Talc, powderedav.oz.	9
Starch, powderedav.oz.	9

The two preceding, being in very fine powder and perfumed, are known as "Medicated Fuller's Earth" Powder.

#### Liquid Cosmetics.

Any of the various mineral powders used as cosmetics may be converted into lotions by mixing with water, which may or may not be perfumed or colored; a small amount of glycerin is frequently added. The powders do not dissolve, but are simply suspended by shaking.

Substances suitable for the purpose indicated are: Prepared chalk, magnesium carbonate, zinc oxide, the oxychloride or subnitrate of bismuth. The last two resemble chalk in density, but much excel it in whiteness.

Unscrupulous or careless persons sometimes use white lead in the preparation of cosmetics, and calomel is said to be the chief ingredient of a proprietary cosmetic that has had some reputation. The character of both these substances renders their use for such purposes dangerous, as even external application may produce the constitutional effects of lead or mercury.

In making these preparations a perfectly smooth mixture must be made by long trituration, preferably triturating the powder first with the glycerin. If a white preparation is desired, no coloring agent is added; but if a flesh-colored preparation, then solu-

tion of carmine or an alcoholic solution of eosin must be added. No coloring agent or perfume is mentioned in the formulas below, as these may be added as desired.

These liquid cosmetics are usually known by such titles as "enamel," "oriental cream," "pearl cream," "liquid pearl," "balm," "liquid face paint," etc. They are usually dispensed in white (opaque) bottles.

1 1 1
I.
Calomelav.oz. 2
Bismuth subnitrateav.oz. 1
Zinc oxideav.oz. 1
Glycerin
0
II.
Zinc oxideav.oz. 2
Glycerinfl.oz. 6
Waterfl.oz. 8
Oil of bergamotdrops 4
Oil of lemondrops 4
III.
Precipitated chalkav.oz. 3
Glycerinfl.dr. 6
Waterfl.oz. 15
IV.
Bismuth subnitrateav.oz. 2
Glycerinfl.oz. 1
Water, enough to makefl.oz. 16
V.
Zinc oxideav.oz. 21
Precipitated chalkav.oz. 1½ Alcoholfl.oz. 3
Waterfl.oz. 12
VI.
Bismuth oxychlorideav.oz. $3\frac{1}{2}$
Glycerin
Alcohol
Waterfl.oz. 10
77 21 3

#### Face Bleach.

Under this name are marketed a number of preparations of which the principal or active constituent is corrosive sublimate. The use of such preparations is not countenanced, and the following formula is given only because the preparation is in occasional demand:

Mercury bichloridegr.	1
Emulsion of bitter almondfl.oz. 1	2
Tincture of benzoinfl.dr.	1

The mercury salt is dissolved in the emulsion and the mixture gradually added to the tincture. This preparation does not keep long; should be kept in small opaque vials and be dispensed only with a "shake well" label.—D.

The emulsion may be prepared from one av. ounce of almond.

Other "bleaches" contain from one-half to one grain corrosive sublimate to the fluidounce.

The above is known as Hebra's Oriental Cosmetic Water.

#### Applications for Blackheads or Comedones.

Before retiring bathe the parts affected with very hot water, dry well, then rub thoroughly with a lotion consisting of:

Ether			fl.oz. 1
Alcohol		. <b></b>	fl.dr. 6
Aromatic	spirit of	ammonia	fl.dr. 2

The object of this is to dissolve the sebaceous deposits. To further soften and loosen the same, a paste is applied, composed of:

Acetic aciddr. 2
Glycerindr. 3
China claydr. 4
Perfumesufficient

This paste is removed by washing on the following morning. After a few days the comedones can be easily expressed, if necessary, with the aid of a watch key.

Friction with a Turkish towel, the use of soap containing tar, resorcin or ichthyol, rubbing with resorcin solution in spirit of ether, etc., constitute a good after treatment.

11.	
Resorcin	60
Zinc oxidegr.	
Starchgr.	
Petrolatumgr.	240
TIT	

111		
	mmonium carbonategr. 20	)
]	ther	Ĺ
7	ater fl.oz.	L
	4 . 1 . 1 . 1	

IV.	
Lanolin	)()
Petrolatumgr. 20	00
Hydrogen peroxidefl.dr.	4

Apply at night.

V.		
Petrolatum	av.oz.	1
Lanolin (anhydrous)		
Hydrogen peroxide	fl. oz.	1
Acetic acid		

VI. The preparation employed for the removal of pimples may usually also be used against blackheads.

Freckle Remedies.  I.  Borax	Spermaceti. gr. 120 White wax gr. 120 Oil of sweet almonds fl.dr. 4 Salicylic acid gr. 16 White precipitate gr. 4 Rub this on, night and morning, for five minutes, with a clean finger. VI.
salts and filter.  Label: Apply with a soft sponge several times a day.  This forms a mild harmless lotion and frequently answers completely in mild cases.  The same remarks apply to the next preparation.	Copper oleate         .gr. 30           Petrolatum         av.oz. 1           Apply twice a day.           VII.         Zinc sulphocarbolate         .gr. 15           Glycerin         .fl.dr. 4           Rose water         .fl.oz. 2½           Alcohol         .fl.dr. 2
II.       Sodium sulphocarbolate	Cologne water
III.  Salicylic acid	allow the latter to dry upon the skin.—D.  This is intended for the removal of summer freckles.  VIII.
ness of the skin, which should be subdued by the use of glycerite of starch. Care must be taken not to get any of it too near the eyes, nostrils and lips. Should the skin be- come red and irritated, the lotion must be	Zinc sulphocarbolategr. 60 Oil of lemonfl.dr. 1 Alcoholfl.oz. 1 Collodion flexiblefl.oz. 5 After bathing the face with hot water and drying, apply with a brush.
suspended for a few days to allow the inflammation to subside.  IV.	Lactic acid
Mercuric chloridegr. 6 Alcoholfl.oz. 1 Green soapav.oz. 2 Oil of lavenderdrops 10 Label: Apply at night and wash off in the	X. Citrin ointmentgr. 60 Cold creamgr. 420 Apply night and morning. XI.
morning. Some kind of glycerin cream should be applied during the day.  This is to be used only in obstinate cases.  The same applies to the next formula.	Zinc sulphategr. 20 Cold creamav.oz. 1 Dissolve the salt in a small amount of water and incorporate with the ointment. Use like the preceding.
V. Corrosive sublimate	XII. Some of the preparations intended to remove tan or sunburn may frequently also be employed to remove freckles.  Applications for Pimples or Acne.
inen, cut to fit, at night. Remove when dry. After a few nights redness of the epidermis is induced, when it peels off in fine scales; then may be used an ointment composed of:	Mercuric iodide

V. When there is much pustulation the following ointment may be used:

 Bismuth subnitrate.
 gr. 30

 Ammoniated mercury.
 gr. 30

 Ichthyol.
 gr. 30

 Petrolatum
 gr. 360

For external use. Every evening at bedtime, the pustules should be smeared with this ointment.

VI. Most of the preparations intended for the removal of blackheads may also be used against pimples.

In the treatment of acne it may be found advisable to assist the action of external remedies by the exhibition of internal medicines.

# Applications for Tan and Sunburn.

Hydrochinonegr.	48
Glacial phosphoric acidgr.	30
Glycerinfl.dr.	
Water, enough to makefl.oz.	
Mix and dissolve.	

This is to be applied twice daily, first washing and drying the skin carefully.

II. Under the name albadermine has been devised a foreign process for the removal of tan, sunburn, and summer freckles which requires the use of two solutions:

#### Solution "A."

Potassium iodidegr.	120
Iodine gr.	
Glycerinfl.dr.	3
Infusion of rosefl.oz	4

Dissolve the potassium iodide in a small quantity of the infusion and one fluidram of the glycerin; with this fluid moisten the iodine in a glass mortar and rub it down, gradually adding more liquid until complete solution has been obtained; then stir in the remainder of the ingredients, and bottle the mixture.

#### Solution "B."

Sodium hyposulphite.....gr. 240
Rose water......fl.oz. 16
Dissolve and filter.

With a small camel's hair pencil or piece of fine sponge apply a little of "A" to the tanned or freckled surface, until a slight but tolerably uniform brownish-yellow skin has been produced. At the expiration of fifteen or twenty minutes moisten a piece of cambric, lint or soft rag with "B" and lay it upon the affected part, removing, squeezing away the liquid, soaking it afresh, and again applying until the iodine stain has disappeared. Repeat the entire process thrice daily, but diminish the frequency of the application if tenderness be produced. In the course of from three or four days to as many weeks the discoloration will either have disappeared entirely, or its intensity will be very greatly diminished.

#### TTT

Sweet	almonds,	blanched	.av.oz.	1
Bitter	almonds,	blanched	.av.oz.	$\frac{1}{2}$
Corros	ive sublim	ate	gr.	15
Alcoho	ol		fl.dr.	2
Water			. sufficie	ent

Mix and crush the almonds and add enough water so as to obtain 16 fluidounces of emulsion. To the latter add the corrosive sublimate dissolved in the alcohol.

This preparation is well known under the name "Gowland's Lotion." It should be applied at night and be washed off in the morning, after which an emollient preparation like a mixture of glycerin and rose water, or cold cream, should be applied.

IV.

Boraxgr.	320
Zinc oxidegr.	320
Glycerinfl.oz.	$2\frac{1}{2}$
Bay rumfl.oz.	$2\frac{1}{2}$
Distilled waterfl.oz.	10
Mix and apply freely 5 or 6 times da	ily.

V. Most of the preparations employed in the

treatment of freckles may also be used for the removal of tan and sunburn.

#### Miscellaneous Cosmetic Preparations.

Under this heading are grouped a number of preparations which are demanded occasionally and cannot be appropriately classed under any other heading.

#### I.—A. Kummerfeld's Cosmetic or Toilet Water.

Sulphur, precipitatedgr.	150
Glycerinfl.dr.	
Camphorgr.	
Cologne waterfl.oz.	
Boraxgr.	
Distilled waterfl.oz.	144
Tincture of muskdrops	$2^{-}$

Triturate the sulphur to a smooth paste with the glycerin, dissolve the camphor in the cologne water and the borax in the distilled water, mix all, and add the tincture.

The sulphur will be found at both the bottom and the top of the mixture. This may be overcome, in a great measure at least, by adding to the above mixture 1 fluidounce of ether.—D.

Acadea, powder gr. 180
Sulphur, precipitated av.oz. 2½
Rose water fl.oz. 9
Lime water fl.oz. 9

Mix the first three ingredients intimately, gradually add the rose water with constant trituration, and then add the lime water.—D.

Another formula uses the same amount of solids for 30 fluidounces of each of the waters.—H.

### II. Eau de Lys de Lohse.

•	
Zinc oxidegr.	75
Talcum, powdergr.	
Glycerinfl.dr.	
Rose waterfl.oz.	
Tincture of benzoinfl.dr.	
Essence of jasminem.	40
Coumarin sugargr.	8
Tincture of muskdrops	
Spirit of orrisdrops	
Spirit of ylang ylangdrops	
•	

Triturate the first two ingredients with the glycerin, add the tincture of benzoin, incorporating thoroughly and then add the remaining ingredients with trituration.—D.

#### III. Hebra's Cosmetic Liniment or Sulphur Paste.

Potassium carbonateav.oz.	Э
Glycerinfl.oz.	4
Sulphur, precipitatedav.oz.	5
Waterfl.oz.	
Alcoholfl.oz.	4
Etherfl.oz.	$6\frac{1}{2}$
Dissolve the potassium carbonate in	the

Dissolve the potassium carbonate in the glycerin, incorporate the sulphur and add the remaining ingredients previously mixed.

Used against flesh worms. Apply at night and wash off in the morning.—D.

# IV. Lilionese (or Lilionese Cosmetic Water).

Boraxgr.	110
Potassium carbonategr.	
Rose waterfl.oz.	
Cologne waterfl.dr.	4
Talcum, powdergr.	720
Glycerinfl.dr.	5
Tincture of benzoinfl.dr.	

Dissolve the borax and potassium carbonate in the rose water and add the cologne water previously mixed with the tincture; rub the talcum to a smooth paste with the glycerin and gradually add to it the previous mixture.—D.

## V. May Dew Water.

Boraxgr.	40
Sodium hyposulphitegr.	375
Glycerinfl.dr.	5
Distilled waterfl.oz.	14
Cologne waterfl.oz.	1
Oil of nerolidrops	5
Oil of ylang ylangdrop	1
Essence of jasminedrops	10
Tincture of ambergrisdrops	2
Tincture of muskdrop	1

Mix the first four ingredients and add the remaining ingredients previously mixed.—D.

# VI.—A. Lait Virginal (Virgin's Milk).

Tincture of tolufl.dr.	. 25
Tincture of benzoinfl.dr.	$2\frac{1}{2}$
Spirit of soapfl.dr.	21
Glycerinfl.dr.	5
Boraxgr.	110
Orange flower waterfl.oz.	$3\frac{1}{2}$
Rose waterfl.oz.	5
Distilled waterfl.oz.	81
Essence of jasmine	40
Oil of rosedrops	2
Spirit of orrisdrops	30
Tincture of civetdrops	2
Coumarin sugargr.	4

Dissolve the borax in the water and add the solution to the tinctures by trituration,

"bloom of roses."

then add the spirit of soap and the other	I.
waters and finally the perfumes.—D.	Carminegr. 30
В.	Ammonia waterfl.dr. 1
Tincture of benzoinfl.dr. $3\frac{1}{2}$	Rose water
Tincture of tolufl.dr. 5	Mix, set aside 24 hours, agitating fre-
Rose or orange flower waterfl.oz. 15	
Mix the two tinctures and add the water	quently, and filter.
very slowly in a thin stream with vigorous	II.
and constant stirring.	Eosingr. 16
C.	Water
Tincture of benzoinfl.dr. 4	Glycerin fl.dr. $\frac{1}{2}$ Alcohol fl.oz. $3$
Glycerin	Cologne waterfl.oz. $2\frac{1}{2}$
Rose waterfl.oz. 16	Mix and dissolve.
Mix the first two ingredients and add the	III.
water gradually with vigorous stirring.	
D.	Rosanilingr. 5 White waxgr. 50
Tragacanth, powdergr. 10	Spermacetigr. 50
Glycerin	White petrolatumgr. 380
Rose water	Alcohol
Make a smooth mixture of the glycerin,	Perfumeto suit.
	Dissolve the dye in alcohol, add this solu-
gum and water, and then thoroughly incor-	tion to the fats previously melted, and incor-
porate the tincture.	porate the whole together, continuing the
VII. Eau d'Hebe.	stirring until the mixture has cooled.
Oil of lavender flowersdrops 25	IV.
Oil of lemondrops 10	Eosingr. 3
Oil of bergamotdrops 5 Oil of rosedrops 3	Rosanilin
Alcoholfl.oz. 5	White waxgr. 50 Spermacetigr. 50
Diluted acetic acidfl.oz. 10	White petrolatumgr. 380
Mix the oils with the alcohol, add the acid,	Oil of lavender flowersdrops 2
and filter.—H.	Oil of bergamotdrop 1
VIII. Copeland's Cosmetic Water.	Alcohol
Emulsion of bitter almondsfl.oz. 3\frac{1}{2}	Prepare like the preceding.
Rose waterfl.oz. 4	V
Orange flower waterfl.oz. 4	Eosingr. 5
Boraxgr. 60	Waxgr. 50 Spermacetigr. 50
Tincture of benzoinfl.dr. 2	Spermacetigr. 50 Petrolatum, whitegr, 380
The emulsion may be prepared from one-	Oil of rosedrops 2
half av. ounce of bitter almonds.	Alcoholfl.dr. ½
IX. Goddard's Cosmetic Lotion.	Prepare like the preceding.
Corrosive sublimategr. 6	VI.
Tincture of benzoinfl.dr. 2	Carminegr. 90
Rose waterfl.oz. 6	Ammonia waterfl.dr. 3
Pougos	Talcum, powderav.oz. 3
Rouges.	Dextrin
These are preparations of deep red tint	Oil of bergamotdrops 6 Oil of rosedrops 2
employed usually for heightening the color	Oil of sassafrasdrop 1
of the cheeks. They may be in liquid, oint-	Simple syrupsufficient
ment, or tablet forms. The coloring agent	Mix the talcum and dextrin, add the oils,
is either carmine, eosin, or rosanilin. The	then incorporate the carmine previously dis-
liquid preparations receive a fanciful name	solved in the ammonia, now add enough
like "vinegar rouge," "maiden's blush," or	syrup to make a stiff mass and form into
"hloom of roses"	cakes —D

cakes.—D.

#### Theatrical Grease Paints.

In preparing these the solid fats should be melted first, the fixed oils then added, incorporate the coloring agent, add the perfumes, and form into sticks if desired

#### I. White:

A

2X•
Prepared chalkav.oz. 4
Zinc oxideav.oz, 4
Bismuth subnitrateav.oz. 4
Asbestos, powderav.oz. 4
Sweet almond oilabout fl.oz. 21/2
Camphorgr. 40
Oil of peppermintfl.dr. 3
Ess. bouquet extractfl.dr. 3
Sufficient almond oil should be used t
form a mass of proper consistence.
В.
White waxav.oz. 2
Olive or sweet almond oilfl.oz. 3
Talc, powderfl.oz. 1
Zinc oxide $fl.oz$ . $\frac{1}{2}$
C.
Bismuth oxychlorideav.oz. 5
White waxav.oz. 2
Sweet almond oilfl.oz. 5
D.
Zinc oxideav.oz. 1
Olive oilfl.oz. 1½
Paraffin waxav.oz. 31
Cacao butterav.oz. 4
Oil of bergamotdrops 5

#### II.-Red:

A. Same as any of above, coloring with an ammoniacal solution of carmine.

Oil of rose......drops 5

Oil of lemon......drops 3

Tincture of civet......drops 5

Cacao butterav.oz.	4
White waxav.oz.	
Olive oilfl.oz.	2
Oil of rosedrops	8
Oil of bergamotdrops	3
Oil of nerolidrops	2
Tincture of muskdrops	2
Carminegr.	
Ammonia waterfl.dr.	3
No. 11 at a second seco	

Melt the cacao butter and wax, add olive oil, stir in the carmine first dissolved in the ammonia, add the volatile oils and tincture, and form into sticks.—D.

## C.

Carm	ine		orr	112
Wate	r of ammoni	a, suffi	cient to	
dis	solve about.		fl.oz. 1	l or 2
Rub	intimately	with	powdered	talcun

(av. oz.  $1\frac{1}{2}$ ) till dry; rub with this vehicle.

The latter two should be first melted together.

#### III.-Pink:

Zinc oxideav.oz.	21
Bismuth subnitrateav.oz.	$2\frac{1}{2}$
Asbestos, powderav.oz.	21
Sweet almond oil, aboutfl.oz.	1
Camphor av.oz.	
Oil of peppermintfl.dr.	
Ess. bouquet extractfl.dr.	
Eosingr.	
8	

#### IV .- Black:

Α						
	Soot			 	av.oz.	2
	Sweet	almond	oil	 	fl.oz.	2
	Cacao	butter.		 	av.oz.	6
	Perfur	ne		 	sufficie	nt.

The soot should be derived from burning camphor and repeatedly washed with alcohol.

The soot should be triturated to a smooth mixture with the oil; then add to the melted cacao butter, add the perfume, and form into sticks.

#### B.

Lampl	black, bestav.oz.	1
Cacao	butterav.oz.	3
Oil of	nerolifl.dr.	21/2

Melt the cacao butter, add the lampblack and while cooling make an intimate mixture, adding the perfume toward the last.

#### C. La

Lampblackav.oz.	1/2
Olive oilfl.oz.	
Cacao butterav.oz.	$3\frac{1}{2}$
White waxav.oz.	4
Oil of rosedrops	6
Oil of bergamotdrops	2
Tincture of muskdrops	

Triturate the lampblack to a smooth mixture with the olive oil, melt the wax and cacao butter, add the oily mixture, then the volatile oils and tincture, and form into sticks.

#### V.-Brown:

A brown paint may be prepared according to either of the formulas of the preceding, substituting finely levigated burnt umber, sienna or similar earth for the lampblack.

### Depilatories.

Depilatories are preparations for removing hair. They are usually in the form of powder, which should be in impalpable con-

dition. The main or active ingredient is usually a sulphide or sulphhydrate of one of the alkalies or alkaline earths, although the older depilatories were made with caustic alkalies. In using these depilatories they should be made into thin paste with water, applied in a thin layer to the skin, allowed to remain a few minutes and then scraped off with a blunt instrument, when the hair will have softened sufficiently to remove without pain.

Too long contact of depilatories with the skin should be avoided, as they are liable to cause erosions and even ugly sores. To avoid any bad after-effect, the skin should be thoroughly cleansed and then anointed with a bland oil.

Ţ

Prepare sulphuretted baryta (barium sulphide) by making heavy spar (natural barium sulphate) and charcoal into a stiff paste by means of linseed oil, forming this mass into cylindrical rolls and subjecting to the heat of a coal fire. The dark gray coke, after pulverizing, is then made up as follows:

Crude sulphuretted barytagr.	120
Zinc oxidegr.	
Starch	60

With the aid of water this powder is converted into a soft paste, and applied to a hairy skin in a layer as thick as a straw. After drying (about ten minutes), the pellicle is scraped off with a paper knife, or similar blunt instrument, and with it the hair. The face should be washed clean and anointed with some bland oil.

Ordinary barium sulphide may be used in place of the sulphuretted baryta.

II.

Sodium sulphhydrategr.	100
Slaked limegr.	
Starch	20
Lime waterfl.dr.	4

When using apply this like the paste formed in the preceding formula.

Sodium sulphhydrate is prepared by supersaturating at ordinary temperature a solution of sodium hydrate of spec. grav. 1.35 (made from 1 av. ounce of caustic soda and 2 fluidounces of water), and then setting aside the well-closed jar for several days in a cold, dark place, when the crystals formed may be

removed and preserved in a well-closed vial, protected from the light.

#### III.

Sodium sulph	ihydrate	 gr. 100
Chalk		
Make into a		_
No. I.		
V		

Sodium sulphhydrate...gr. 130 Slaked lime...av.oz. 1 Starch...av.oz. 1

Use like the preceding.

Barium sulphide......gr. 80
Precipitated chalk.....gr. 400
Use like the preceding, removing in 3 or 4 minutes.

The barium sulphide should be absolutely dry.

#### VI.

Orpime	ent															.gr.	120
Slaked	lime.	•	•		•	•	•	•	•		•	•	•	•	•	.gr.	360

Use like the preceding.

#### VII.

Orpimentgr.	60
Slaked limegr.	300
Starchgr.  Use like the preceding.	100

#### VIII.

Quicklimegr.	120
Sodium sulphidegr.	240
Starchgr.	80
Orris root, powdergr.	40
Orris root, powdergr.	40

Rub the necessary portion of this powder into a thin paste with water, and apply as directed in formula 1.

#### IX.

Calcium sulphide	gr. 120
Glycerin	
Orris root powder	
Oil of lavender	
Oil of ylang ylang	
Camphor water	
Mala a smooth posts and	

Make a smooth paste and use as directed in formula 1.

 Strontium sulphide
 g. 300

 Zinc oxide
 gr. 100

 Starch
 gr. 100

 Menthol
 gr. 5

Mix intimately, reducing to very fine powder.

In using this preparation, make a paste with water and apply as in No. 1.

Strontium sulphide has the advantage over

barium sulphide of being non-poisonous and Shaving Cream. over almost all other depilatory agents of not evolving hydrogen sulphide.

#### XI.

Barium sulphidegr.	120
Starchgr.	200
Orris root, powdergr.	200

Make a paste with water and use as in formula I.

#### XII.

Charcoalgr.	8
Quicklime gr.	30
Sodium carbonategr.	
Glycerinfl.dr.	
Lardav.oz.	

This is applied to the skin for ten or twelve days, when the latter assumes a rose tint and the hairs may be extracted.

#### XIII.

Iodine collodion, Part I, may be employed In using apply once daily as a depilatory. for about 4 days, when the film may be removed and the hair will come off with it.

#### Liquid Soaps.

Liquid soaps are solutions of soap in alcohol, water, glycerin, etc., put up in attractive forms for cosmetic or medicinal purposes. Most preparations of this kind are put up under the title "liquid glycerin soap."

Potassa soap, whiteav.oz.	11
Glycerin fl. oz.	8
Simple syrupfl.oz.	71
Alcoholfl.oz.	4
Oil of bergamotfl.dr.	1
Oil of sassafrasdrops	40
Oil of cassiadrops	20
Oil of rose geraniumdrops	
Oil of wintergreendrops	
Oil of clovesdrops	
Oil of citronelladrops	
Oil of mirbane or bitter almonds.drops	
Tincture of muskfl.dr.	1

Mix, allow to stand a few days, and filter. -D.

### II.

Potassa soap, whiteav.oz.	11
Glycerin	
Alcoholfl.oz.	

Mix, add the perfumes of the preceding mixture, allow to stand a few days, and filter. -D.

I.	Lard	
	Caustic potassaav.oz. 2	,
	Waterfl.oz. 6	i
	Perfumeto suit	

Met the lard in a porcelain vessel over a salt water bath; dissolve the potassa in the water, and run the lye, thus formed, very slowly into the melted grease, stirring thoroughly all the time, until saponification is complete.

A pearly appearance can be given to the "cream," which is simply a soft soap, by long trituration in a mortar with a little alcohol, say 2 fluidrams to each pound of the soap.

Bitter almond oil may be used as a perfume for the "cream." Only a very minute proportion is required. A few drops dissolved in the alcohol used as above will

Glycerin should be added to this cream to retain it in a permanently soft condition.

11,	
Castile soap, whiteav	oz. 3
Rose waterfl.	oz. 12
Sweet almond oilfl	.oz. 11/2
Cacao butterav	oz. 1 ½
Tincture of benzoinfl	.dr. 3
Oil of rose geraniumdi	rops 3
Oil of bitter almond di	
Glycerinsu	
•	

Dissolve the soap in the water by the aid of water-bath heat, add the sweet almond oil and cacao butter previously melted together, then, while still warm, incorporate the tincture, oils and sufficient glycerin to give the proper creamy consistence.

#### Shaving Powder.

# Coop powder

Soap, powderav.oz.	10
Orris root, powdergr	375
Starchgr.	375
Sodium carbonate, puregr.	
Oil of bergamotdrops	36
Oil of lemondrops	
Peru balsamdrops	
Tincture of muskdrops	
Mix well reducing to fine powder —	

.1.		
Soap, powderav.		
Sodium carbonateav.	oz.	11/4
Starchav.	oz.	2
Orris root, powder	gr.	288
Oil of bergamotdro	ops	25
Instead of the orris root the san	ne v	veight

of powdered quillaja and a very little oil of orris may be used. An addition of glycerin will render the powder milder in use.

#### Manicure or Finger-Nail Cosmetics.

Different preparations of tin have been used successfully for cleansing and polishing finger nails, probably on account of their detergent and astringent qualities. The use of tin oleate has been highly recommended as imparting a splendid luster to the nails, and when colored with a little carmine, giving to them a fine roseate tint. The oleate is prepared as follows: To a solution of white castile soap in warm water 1 av.ounce to the pint, gradually add a 10 per cent solution of tin chloride until it ceases to produce a pre-The insoluble substance formed, after being washed and dried, is tin oleate. It is a soft solid, and is used without further preparation, unless, as stated, it be tinted with carmine. It may be perfumed if desired.

Another substance which is used as a nail powder is pure tin oxide, perfumed with oil of lavender, and tinted with carmine. It is applied either by rubbing it in on to the nail with the finger, or with a nail polisher covered with leather.

Still another polish for finger nails is tin stearate. It is superior to the oleate, being stiffer and thus nicer to use. It may be prepared like the oleate by precipitation, using solutions of sodium or potassium stearate and tin chloride. This may be tinted with carmine if desired. If a cheaper preparation is wanted, it can be mixed with equal parts of zinc oxide.

Another suitable powder is the following: Pumice stone, very fine powder.av.oz. 2 Talcum, powder.....av.oz. ½ Mix well. Add 15 gr. of carmine if desired, as well as some suitable perfume.

Cuttle-fish bone in very fine powder is also employed for polishing the nails.

The following is employed as a varnish or polish after the application of powder:

Paraffin waxgr 60	
Chloroform	
Oil of rosedrops 3	

Polish the nails, apply the varnish, then rub with chamois skin.

This preparation is employed as a nail wash.

Oxalic acid	 gr. 30
Rose water	 fl.oz. 1

Mix and make solution.

Apply to the discolored nails by means of soft leather or flannel with friction.

Citric acid or acetic acid may be substituted for oxalic acid in the above.

The following ointment is employed for softening the nails, curing hang nails, etc.:

Petrolatum	av.oz. 1
Castile soap, white, power	dergr. 60
Oil of bergamot or other	
ing oil	sufficient

White petrolatum should be preferred as it will make a nice appearing preparation.

This should be applied at night and the fingers covered with gloves.

#### SECTION III—PREPARATIONS FOR THE HAIR, SCALP, AND MUSTACHE.

#### Shampoo Liquids.

These are applications to the head for cleansing purposes, to remove dirt, dandruff, etc., from the scalp and hair. They should be applied freely and rubbed in thoroughly until considerable lather is formed, which latter should be removed by means of a large quantity of water. The "Sea Foam Liquids," which follow are very similar:

1.					
Potassiı	ım carb	onate.	 av	.oz.	1
Borax .			 av	.oz.	1
Distille	d water		 fl	.oz.	32
Mix and	d dissol	ve.			

II.	
Soft or green soapav.oz.	11/4
Potassium carbonateav.oz.	21/2
Alcoholfl.oz.	3
Waterfl.oz.	25

Dissolve the potassium carbonate in the water and add the remaining ingredients.

TTT

111.		
Potassium carbonate	av.oz.	1
Ammonia water	.fl.oz.	11/2
Tincture of cantharides	.fl.dr.	6
Bay rum	.fl.oz.	4
Alcohol	.fl.oz.	4
Water	.fl.oz.	6

Dissolve the potassium carbonate in the water, and add the remaining ingredients.

HAIR PRE	PARATIONS, 335
IV. Potassium carbonate	Dissolve the quinine in the cologne and tincture of quillaja with the aid of heat; then add the remaining ingredients and filter if necessary.  This may be put up under the title "tonic shampoo."  Sea Foam Liquids.  The difference between these and the preceding class of preparations is not a well defined one, and in fact, the difference is largely in the matter of application or use, the shampoos being employed in liberal quantities and rubbed in vigorously to produce a copious lather, which will then require a large amount of water for removal. In the case of sea foams, a more or less volatile alkali, ammonia, for example, usually forms the principal ingredient, which foams but slightly, and the hair may be cleansed by rubbing with a wet towel.  I.  Ammonia water
Mix and dissolve.  VIII.  Borax	II.

V.     Tincture of arnica	these are known but some of any egg, but Examples herewith, so without it.  Castile so Curd soa
VII.       Bay rum       fl.oz.       16         Alcohol       fl.oz.       8         Glycerin       fl.oz.       8         Tincture of cantharides       fl.oz.       1         Ammonium carbonate       gr. 360	Potassium Honey Perfume Make a h a sufficient
Mix and dissolve.  VIII.  Alcohol	Ammonia Cologne Alcohol Water Whites of The whit oughly beat with the wa remaining in and the who III. White cas Ammonia Bay rum, Glycerin. Water Dissolve of heat; wh
Rose water	ingredients.  IV.  Borax Glycerin. Rum Bay rum. Whites of Incorpora the glycerin gradually a

Beat the eggs thoroughly, mix with the rose water, add the other ingredients, and strain.—D.

Oil of rose.....drops

Oil of bergamot......drops Oil of rose geranium.....drop

Oil of bitter almonds.....drop

### Shampoo Paste, Cream or Jelly.

Many shampoo preparations are now put up in the form of pastes or jellies. Many of

own by the name "egg shampoo," these preparations do not contain t are merely a perfumed soft soap. of shampoo pastes are given ome containing egg, some being

oap, white.....av.oz. 4 p, powder.....av.oz. 2 n carbonate . . . . . . . . av.oz. 1 .....av.oz. 1

to suit.

omogeneous paste by heating with quantity of water.

Ammonia water	fl.dr. 3
Cologne water	fl.dr. 3
Alcohol	fl.oz. 5
Water	fl.oz. 5
Whites of eggas many as	desired

tes of egg (about two) are thorten up previous to being mixed ater and water of ammonia; the ngredients are added in their order ole stirred briskly.

White castile soap, in shavings av.oz. Ammonia water fl.oz.	
Bay rum, or cologne waterfl.oz.	1
Glycerinfl.oz. Waterfl.oz.	

the soap in the water, by means nen nearly cold, stir in the other

Boraxav.oz.	2
Glycerinfl.oz.	
Rumfl.oz.	
Bay rumfl.oz.	10
Whites of egg	. 2

ate the borax in fine powder with n and add the bay rum and rum and with constant stirring to the The previously well-beaten white of egg is added lastly, and the whole stirred thoroughly until an even mixture results.

2

Castile soap, whiteav.oz.	4
Potassium carbonateav.oz.	1
Waterfl.oz.	6
Glycerinfl.oz.	2
Oil of lavender flowersdrops	
Oil of bergamotdrops	10

To the water, add the soap, in shavings,

and the potassium carbonate, and heat on a water bath until thoroughly softened; add the glycerin and oils. If necessary to reduce to proper consistency, more water may be added.

#### Shampoo Powders.

Examples of powdery mixtures are herewith given, which may be dispensed in small boxes or in envelopes, each package being sufficient for about one shampoo.

*
I.
Borax, powderav.oz. 1
Sodium carbonate, powderav.oz. 1
Camphor, powder
Oil of rosemarydrops 10
This is sufficient for one quart of water.
II.
Borax, powderav.oz. 3
Sodium carbonate, driedav.oz. 6
Quillaja, fine powderav.oz. 3
Perfume to suit.
III

III

Borax, powderav.oz.	3
Camphor, powdergr. 86	0
Cochineal, powdergr. 49	0
Oil of rosemarydrops 2	
Mix well.	

This may be put up in half-ounce packages, each of which is sufficient for one pint of water.

# **Hair Oils.** (Huile Philocome.—Huile Antique.)

Hair oil may be prepared from any of the bland, fixed, non-drying oils such as olive, mustard, rapeseed, peanut, or benne oil, also from liquid petrolatum, and from a mixture of castor oil and alcohol. These mixtures must be rendered pleasantly odorous by the addition of suitable perfume. Frequently hair oil is colored red by the use of alkanet root or of its coloring constituent, alkannin.

Hair oil in its usual yellow condition is frequently dispensed under the name "bear's oil"; if colored red, it is often known as "rose oil."

Ι.		
	Castor oilfl.oz.	4
	Alcoholfl.oz.	26
	Tincture of cantharidesfl.oz.	2
	Oil of lavenderfl dr.	1
	Oil of rosemaryfl.dr.	1
	Oil of clovesfl.dr.	2
	Oil of bergamotfl.dr.	4
	Alkanet root, powdergr.	120

Mix the oils in a bottle; put the alkanet root on a filter or pack in a funnel and percolate the alcohol through it; mix this percolate with the oily mi..ture, and add the cantharides tincture.

II.	
Castor oilfl.oz.	-8
Alcoholfl.oz.	24
Oil of bergamotfl.dr.	3
Mix well.—H.	

#### III.

-			
	Olive or benne oil	fl.oz.	32
	Burdock root, fresh	.av.oz.	4
	Castor oil		
	Oil of bergamot		
	Oil of rose geranium		
	Alkannin or alkanet root		

Digest the olive or benne oil with the burdock root for about one-half hour at a moderate heat, then decant the clear liquid, add the other oils, and color, if desired, with alkannin. Alkanet may be employed instead of the latter and may then be added to the burdock root during digestion.

This preparation, as well as the following, is known as "Burdock Root Hair Oil."

IV.		
Olive oil	fl.oz.	27
Benzoinated oil	fl.oz.	3
Alkannin	gr.	-8
Chlorophyll	gr.	40
Oil of bergamot		
Oil of lavender flowers		

Dissolve the alkannin and chlorophyll in the first two oils by gentle warming and add the other ingredients. If not clear, filter.

—D.

Oil of rose.....drops 8

V. This preparation and the following two are known as "Macassar Oil."

Peanut or olive oil	.fl.oz.	32
Alkannin	gr.	15
Oil of bergamot	drops	45
Oil of lemon		
Coumarin	gr.	1
Mix and dissolve.—D.		
VI.		
Cocceput oil	fl oz	4

VI.	
Cocoanut oilfl.	
Castor oilfl.	oz. 3
Alcoholfl.	oz. 7
Oil of lavender flowersfl.	dr. 1
Oil of bergamotdro	ps 30
Oil of rose geraniumdro	
	-

Melt the cocoanut oil, and add it to the castor oil dissolved in the alcohol. Shake

well together and add the volatile oils. On cooling, the mixture acquires a crystalline appearance, characteristic of cocoanut oil.

This preparation is known as "cocoanut oil" hair dressing.

#### VII.

Castor oil
Alcoholfl.oz. 3
Oil of nutmeg, essentialdrops 30
Oil of rosemarydrops 10
Oil of sweet marjoramdrops 10
Oil of nerolidrops 10
Oil of rosedrops 20
Tincture of muskfl.dr. 1
Alkanetsufficient to color

#### VIII.

Cassia buds, bruisedav.oz.	1 1
Alkanet, bruisedav.oz.	11
Olive oilfl.oz.	$32^{\circ}$
Oil of cinnamonfl.dr.	2
Oil of rosedrops	10
Oil of bergamotdrops	20
Oil of bitter almonddrops	

Mix the olive oil with the cassia buds and alkanet, macerate in a warm place for 3 days, agitating occasionally, filter, and add the other oils.

#### Perfumes for Hair Oils and Pomades.

The following mixtures may be employed for perfuming oily mixtures which are intended to be used as hair oils:

I.

Oil of lavender flowers	fl. oz.	1
Oil of rosemary	.fl.oz.	1
Oil of cloves		
Oil of cassia,	.fl.dr.	1
I.		
Oil of bergamot	fl 07	21

 Oil of lemon.
 fl.oz.
 1

 Oil of cloves.
 fl.dr.

 Oil of orange.
 fl.dr.

 Oil of cinnamon.
 drops 20

 Oil of bitter almonds.
 drops 10

TI	т	

Oil of	bergamot	.fl.dr.	7
Oil of	rose	.drops	50
	rose geranium		
	cloves		
7.7			

#### IV.

Oil of	bergamot	. fl dr.	4
	sandalwood		
	orris, liquid		
	cloves		
	rose		

V.	
Oil of bergamot	fl.dr. 6
Oil of bitter orange	drops 30
Oil of neroli, petals	drops 80
Oil of petit grain	drops 50
Oil of rose geranium	drops 30
VI.	
Oil of rose geranium	fl.dr. 3
Oil of cloves	drops 50
Peru balsam	
Heliotropin	

Dissolve the latter in a little warm olive oil before adding the volatile oils and balsam.

#### VII.

· 11:	
Heliotropingr. 8	
Coumaringr. 1	
Oil of orris, liquiddrop 1	
Oil of rosedrops 16	
Oil of bergamotdrops 32	
VIII.	
Coumaringr. 1	
Oil of lemondrops 16	

## Oil of bergamot......drops 48

	Coumaringr.	1
	Oil of bitter almondsdrops	
	Oil of cassiadrops	
	Oil of lavender flowersdrops	
	Oil of lemondrops	
	Oil of bergamotdrops	
٧,	0	

#### X.

••	
Coumaringr.	2
Oil of wintergreendrops	
Oil of clovesdrops	
Oil of cassiadrops	
Oil of lavender flowersdrops	16
Oil of lemon	48
Oil of bergamotdrops	72

#### XI.

Oil of lemon	.fl.oz.	1
Oil of bergamot		
Oil of cloves	.fl.dr.	1/2
Oil of orange	.fl.dr.	1/2
Oil of cinnamon	.drops	20
Oil of bitter almond	drops	10

#### Hair Pomades or Pomatums.

These preparations are made with various substances such as lard, beef marrow, cacao butter, wax, cocoanut oil, petrolatum, ceresin, lanolin, castor oil, olive, cottonseed oil etc. To obtain a good product the various fatty ingredients must be in perfectly fresh condition.

These fatty bodies should be rendered pleasantly odorous by the addition of perfumes or mixtures of various oils, tinctures, etc. In some of the formulas given below, the perfume is mentioned; in such as specify no perfume, the scents or perfumes under the preceding title may be employed.

Pomades may be colored red or pink by means of alkanet root or alkannin.

Some formulas for hair pomades mention water as an ingredient. This is employed for cheapening purposes.

In combining the ingredients of pomades, the less fusible fats, like wax or spermaceti, should be melted first, the lard, petrolatum, or fixed oil should then be added, and the whole allowed to become nearly cold before incorporating the perfume.

I. Yellow waxav.oz.	
Cottonseed oilfl.oz.	
Lardav.oz.	5

Mix by fusion, stir until cold, incorporating perfume with the mixture during cooling.

11.

Yellow wax	
Castor oilfl.oz. 6	
Sweet almond oilfl.oz. 1	
Olive oilfl.oz. 4	
Benzoinated lardfl.oz, 3	

Melt the wax at a gentle heat, add the oils and lard, stir till all is melted, allow to cool, stirring constantly till hard. While it is cooling any desirable perfume may be incorporated.

III.

Lardav.oz.	24
White waxav.oz.	21/2
Boraxgr.	150
Distilled water, warm fl.oz.	6 1/2

Melt the wax, add the lard, allow to cool, and then stir with a broad spatula or wooden paddle until the mixture has thickened. Now add the borax dissolved in the water, and continue stirring until the fatty mixture becomes brilliantly white and foam-like. This may be perfumed as desired.—D. IV.

Cacao butter......av.oz. 4
Lard, benzoinated.....av.oz. 12

Mix by fusion and stir occasionally until cool.—D, modified.

This may be perfumed as desired.

Melt the lard and lanolin, add the oil, stir until cool, and add suitable perfume.—D.

VI.

Lard, benzoinatedav.oz. 18
Beef tallowav.oz. 2
Cacao butterav.oz. 2
Orange flower waterfl.dr. 12
Oil of bergamotfl.dr. 3
Oil of lemonfl.dr. 1
Oil of citronelladrops 39
Oil of rose geraniumdrops 30
Tincture of musk drops 5

Melt lard, tallow, and cacao butter together, incorporate the water, and when nearly cold add the remaining ingredients.

—H.

VII.

Expressed oil of nutmegav.oz.	1/2
Lardav.oz.	
Beef tallowav.oz.	134
Yellow waxav.oz.	
Cinnamonav.oz.	34
Clovesav.oz.	1/4
Lemon peel, freshav.oz.	11/2
Benzoinav.oz.	1/2
Alkanet rootav.oz.	1/2
Oil of bergamotfl.dr.	4
Oil of citronellafl.dr.	1

Mix the cinnamon, cloves, lemon peel, benzoin, and alkanet, reduce to coarse powder, add to the nutmeg oil, lard, wax, and tallow previously melted together, maintain the whole at a water bath temperature for three hours, strain, allow the fine particles present to subside, decant the clear liquid, and add the volatile oils.—H.

#### Castor Oil Hair Pomade.

Castor oil. av.oz. 16
White wax av.oz. 4
Oil of bergamot fl.dr. 3
Oil of lavender flowers drops 30
Benzoic acid gr. 10

Melt the wax, add the castor oil, mix well and when nearly cold add the remaining ingredients.

TT

4.4.	
Castor oilav.oz. 16	
Petrolatumav.oz. 4	1/2
Yellow waxav.oz. 2	1/2
Melt the wax, add the petrolatum and o	
and when nearly cold add suitable perfum	e.
TIT	

Melt the wax, add the other ingredients, and stir until cool. Suitable perfume may now be added.

IV.	
Spermacetigr.	300
White waxgr.	480
Castor oilfl.oz.	16
Oil of rosemaryfl.dr.	1
Oil of verbenafl.dr.	1
Oil of bergamotfl.dr.	1
Tincture of curcumafl.dr.	

Melt the wax and spermaceti, add the castor oil, and stir in the perfume and tincture.

#### Cocoanut Oil Hair Pomade.

Lardav.oz.	3
Cocoanut oilav.oz.	12
Ceresin, white and odorlessav.oz.	3
Boraxgr.	150
Distilled water, warmfl.oz.	

Prepare like the preceding.—D.

#### Crystal Hair Pomades.

Under this heading are grouped hair pomades which are made to assume a crystalline appearance.

Ι.

Castor oilav.oz.	17
Olive or peanut oilav.oz.	12
Spermacetiav.oz.	4
Oil of ylang ylangfl.dr.	
Oil of rosedrops	8
Oil of bergamotdrops	8
Oil of nerolidrops	5
Oil of rose geraniumdrops	2
Oil of bitter almond drop	1
Heliotropingr.	1
72 11 0 1 11 11 1	. 4

Fuse the first three ingredients together and add the perfume before it has congealed.

The pomade appears to best advantage in transparent glass jars and in order that it may appear as coarsely crystalline as possible, allow the congealing to go on as slowly as possible by placing the filled jars in warm water, and allow the mixture to stand undisturbed for 6 hours.—D.

Castor oil	.av.oz.	16
Spermaceti	.av.oz.	3
Oil of bergamot	fl.dr.	3
Oil of verbena	fl.dr.	1/2
Oil of lavender flowers	fl.dr.	1/2
Oil of rosemary	fl.dr.	1/2
Benzoic acid		10

Melt the spermaceti, add the castor oil, then the other oils and acid, and allow the whole to cool very slowly and undisturbed.

#### Marrow Hair Pomades.

Under this heading are included a number of hair pomatums known by this name and allow to cool somewhat, add the other ingrewhich usually contain beef marrow.

1.	
	Beef marrowav.oz. 8
	Beef suetav.oz. 4
	Yellow waxav.cz. 1
	Castor oilav.oz. 4
	Oil of bergamotfl.dr. 2
	Oil of lemonfl.dr. 1
	Oil of orangefl.dr. 1

Melt the wax, suet, and marrow, strain, allow to cool somewhat, and add the volatile oils.

II.

Beef	marrow	av.oz.	4
Lard		av.oz.	12

Melt together, strain, and perfume as desired.—D.

III.

Beef marrow	av.oz.	19
Peru balsam	gr.	270
Oil of cinnamon	.fl.dr.	21/2
Oil of bergamot	.fl.dr.	11/2
Tincture of cantharides	.fl.dr.	1
Alcohol	.fl.dr.	6

Melt the marrow, add the other ingredients, strain, and stir until solidified .- H.

This has been known as "Dupuytren's pomade."

The following has also been known as "marrow pomatum:"

Lardav.oz.	16
Suetav.oz.	8
Oil of lemonfl.dr.	2
Oil of bergamotfl.dr.	1
Oil of clovesdrops	40

Mix the lard and suet by fusion, allow to cool, incorporate the oils, and stir until solid.

#### Bear's Grease.

A number of hair pomades are known by this title. It is almost needless to state that none of the preparations bearing this title are made with any ingredients from the animal which furnishes the name,

I.		
	Beef marrowav.oz.	8
	Lardav.oz.	24
	Oil of lemon fl.dr.	6
	Oil of rosedrops	15
	Oil of bergamotdrops	15
	Oil of cinnamon drop	1
	Coumaringr.	1

Mix the marrow and lard by fusion, strain, dients and stir frequently until solid.—D.

II.
Beef marrowav.oz. 15
Spermacetiav.oz. 1
Cacao butterav.oz. 1
Mix by fusion and perfume to suit.—H.
III.
Lardav.oz. 8
Veal suetav.oz. 8
Olive oil
Compound tincture of benzoinfl.dr. 4
Melt the lard and suet, add the olive oi

and thoroughly incorporate the tincture.

#### Anti-Kink Hair Pomade.

Beef suetav.oz. 16
Yellow waxav.oz. 2
Castor oilav.oz. 2
Benzoic acidgr. 10
Oil of lemonfl.dr. 1
Oil of cassiadrops 15
Mix the suet and wax, add the castor o

il and acid, allow to cool somewhat, and incorporate the other oils.

This is used for taking the kinks out of and straightening the hair.

#### Bandolines.

These are mucilaginous preparations used by ladies for application to the hair before the process of "frizzing." The object in employing them is to cause the hair to remain in curl for a longer period of time.

Usually these preparations are perfumed; if containing oil of rose, they are known as "rose bandoline"; if oil of bitter almonds, "almond bandoline," etc.

L.	
Gum tragacanth	.gr. 120
Waterfl	.oz. 12
Alcoholfl	.oz. 2
Oil of rosed	rops 5

Leave the gum tragacanth in the water over night. In winter the mixture should be kept in a warm place. When the gum is thoroughly disintegrated, strain the mucilage with pressure through a cloth and add it to the oil of rose dissolved in the alcohol. other essential oil may be substituted for that of rose. The preparation may be tinted pink or red by the addition of solution of carmine.

If a thicker preparation is desired, more tragacanth may be added.

Quince seed, bruised.....gr. 180 Water.....fl.oz. 16

Macerate at a temperature just short of boiling, with frequent agitation, until a thick mucilage is formed; strain and add 2 fluidounces of good cologne water in which 30 grains of salicylic acid has been dissolved, or 60 grains of borax may be substituted for the salicylic acid. The borax if used should be dissolved in the mucilage. Instead of cologne as a perfume oil of rose dissolved in alcohol may be used.

III.

Irish	moss.	 	 	 	 .av.oz.	4
Wate	r	 	 	 	 fl.oz.	16

Boil and strain; when cold add 2 ounces of cologne and preservative agent as in the preceding.

IV.

Tragacanthav.oz. Glycerinfl.oz.	1/2
Glycerinfl.oz.	1
Rose waterfl.oz.	12
Diluted alcoholfl.oz.	11/2

Macerate the gum in the glycerin and water until thoroughly softened, strain forcibly through muslin, and add the diluted alcohol containing some perfuming agent in solution.

V.	
Tragacanth	gr. 150
Rose waterfl.	
Fuchsin	gr. 8
Oil of rosedro	
Alcohol fl.o	

Macerate the tragacanth in rose water until softened, and strain forcibly through muslin; dissolve the oil and fuchsin in the alcohol and add this to the preceding mixture.-H. modified.

VI. Bandoline in powder is usually powdered tragacanth perfumed, by trituration, with a suitable flavoring volatile oil.

# Curlique or Hair Curling Liquid or

#### I. For keeping hair in curl:

Borax powderav.oz.	1
Gum arabicgr.	30
Spirit of camphorfl.dr.	6
Water, warmfl.oz.	16

Dissolve solids in warm water, and when cool, add the camphor.

Wet the hair with above and roll on papers as usual, let dry, unroll, and form into ringlets.

II. For curls without papers:
Potassium carbonate
In using, moisten hair, adjust them loosely,
and they will curl upon drying.
III.
Potassium carbonate
hair; adjust it loosely, when it curls upon
drying.
IV.
Gum arabic       dr. 1         Sugar       dr. 1         Rose water       oz. 2
Mix and dissolve. Moisten the hair with
the solution at bedtime; roll in twists on
paper.
V.
Potassium carbonate         gr. 120           Cochineal, powder         .gr. 30           Water of ammonia         fl.dr. 1           Glycerin         fl.dr. 2           Alcohol         fl.oz. 1½           Distilled water         fl.oz. 18

Digest with agitation for a week, then decant or filter. The hair is moistened with it and then loosely adjusted. The effect occurs as it dries.

Oil of rose ......drops 5

37	Т
v	1.

Resin												gr.	16	
Alcohol											fl.	OZ.	16	

Mix, dissolve and add suitable perfume.

#### Remedies for Dandruff.

Dandruff requires treatment mainly for the reason that it may be either the precursor or the cause of baldness.

Most of the remedies given below require that the scalp be first cleansed with a shampoo, although some dandruff remedies combine detergent, or cleansing, and curative properties.

The remedies mentioned are to be applied once, or possibly twice, daily.

The various hair tonics, hair oils, and hair pomades, as well as the remedies for baldness, are also useful for the cure of dandruff. I months.

	I.	
	Salicylic acidgr.	25
	Glycerinfl.dr.	1
	Diluted alcoholfl.oz.	2
į	Oil of wintergreendrops	3
	Oil of rosedrop	1
	Oil of nerolidrop	1
ı	Waterfl.oz.	
	3.61 .1 11 1 11 1 1 1 1 1	

Mix the acid and oils with the glycerin and alcohol, add the water and filter.

II.	
Pilocarpine hydrochlorategr.	3
Quinine muriategr.	60
Sulphur, precipitatedgr.	150
Peru balsamgr.	
Beef marrowav.oz.	3
III.	
Salicylic acidgr	. 30
Borax, powdergr	. 15
Peru balsamgr	
Oil of anisedrop	s 5
Oil of bergamotdrop	

Petrolatum .......av.oz.

Mix well, making a smooth ointment.

#### IV.

Resorcingr.	240
Castor oilfl.oz.	
Alcoholfl.oz.	
Peru balsamgr.	
8	

Rub in daily with a piece of flannel.

• •	
V.	
Peru balsamgr.	
Betanaphthol gr.	60
Lanolingr.	
Lard, benzoinatedgr.	120
***	

VI.				
Resorcin	 	 	 gr.	60
Ether	 	 	 .fl.dr.	1
Olive oil.	 	 	 .fl.dr.	1
Alcohol.	 	 	 .fl.oz.	6

To be well shaken and applied to the scalp by a bristle brush about twice as large as the ordinary mucilage brush, by insinuating it between the locks of hair. The head to be well washed with soap and warm water twice a week.

#### VII.

Potassium hydrategr.	18
Carbolic aciddrops	
Cocoanut oilgr.	240
Lanolingr.	240

Dissolve the potassium hydrate in a small amount of water and add the other ingredients.

Label: Rub into the scalp twice a day. A cure is usually effected in from one to three months.

VIII.	
Borax, powdergr.	60
Sulphurgr.	120
Glycerinfl.dr.	
Spirit of camphorfl.oz.	1
Soap linimentfl.oz.	2
Waterfl.oz.	12

Dissolve the borax in the water; triturate the sulphur with the glycerin, mix the two, and add the remaining ingredients.

Directions: Apply to the scalp two or three times a day.

#### IX.

Betanaphtholgr.	360 -
Glycerinfl.oz.	
Oil of wintergreendrops	30
Oil of rosedrops	15
Oil of nerolidrops	15
Terpineoldrops	15
Oil of orangedrops	5
Heliotropingr.	11/3
Tincture of quillajafl.oz.	
* *	

Wash the hair, dry it, apply the above lightly with a sponge, tie a cloth over the head, and allow it to remain for one-half hour.—D.

#### Χ.

Carbolic	acid.	 	 .H.dr.	1
Bay rum		 	 .fl.oz.	32
ΧI.				

#### \_\_\_\_

Chloral hydrategr.	120
Glycerinfl oz.	
Bay rum, enough to makefl.oz.	
KII.	

Chloral hydrategr.	30
Bay rumfl.oz.	
Glycerin	2
Tincture of cantharidesfl.dr.	1
Rose waterfl.oz.	
·	

#### Remedies for Baldness and Falling Out of Hair.

While the formulas given below are intended for alopecia or baldness and loss of hair, it should also be understood that the formulas immediately preceding ("Remedies for Dandruff"), and those that follow this heading, may usually also be employed to strengthen or tone the hair follicles and thus cure or prevent alopecia.

I. Baldness is frequently an infectious disease, caused by the promiscuous use of combs and brushes, more especially in tonsorial establishments. Cure is possible when treatment is commenced reasonably early, and

carried out with systematic energy. The scalp should be, daily for 6 to 8 weeks, later on less frequently, thoroughly rubbed for 10 minutes with a lather of good tar soap, then washed with lukewarm water, and finally with cold water, and partially dried; then with a mixture of:

Mercuric chloridegr. 7	
Waterfl.oz. 5	
Cologne waterfl.oz. 2	
Glycerin	

The scalp should then be rubbed with alcohol containing ½ per cent of betanaphthol; completely dried, and then thoroughly saturated with a mixture of purified lard or lard oil having 2 per cent of salicylic acid and 3 per cent of tincture of benzoin. After a short time the hair will begin to appear, and will, in most cases, soon resume its former vigor and growth. Mercuric chloride (corrosive sublimate) appears to promote the growth of the hair, as has been observed where it has been used as an antiseptic dressing.

With reference to the use of pomade on the hair, it is recommended that all pomade be carefully removed by a washing before it becomes rancid, but that the scalp be not washed without applying pomade afterward. For pomade, animal is preferable to vegetable or mineral fats; the following is recommended:

#### II.

Pilocarpine hydrochlorategr.	1
Petrolatumgr.	100
Lanolingr.	
Oil of lavender flowersdrops	8

#### III.

Carbolic acid	drops	30
Tincture of nux vomica		
Tincture of cantharides		
Tincture of cinchona compour	d.fl.dr.	8
Cocoanut oil, enough to make		

To be rubbed in the scalp with a small sponge twice daily.

#### IV.

Canthariding	r. 1
Acetic etherfl.d	
Alcohol	
Castor oilfl.o.	
Oil of lavender flowersdrop	

establishments. Cure is possible when treatment is commenced reasonably early, and small sponge every third day. After apply-

ing a few times the head should be washed, as an accumulation of the liniment may cause too much irritation.

This preparation is known as linimentum crinole.

crimore.
V.
Oleate mercury av. oz. 1/2
Oil of almond, sweetfl.oz. $2\frac{1}{2}$
Oil of rosedrops 2
Oil of bergamotdrops 2
VI.
Tincture of cantharidesfl.dr. 2
Acetic acid fl.dr. 2
Bay rumfl.oz. 4
Infusion of tea (1:10), enough
to makefl.oz. 32
Apply once every one or two days.

#### Hair Tonics.

The preparations known by the general title of hair tonics are intended as tonic to hair and scalp to prevent and cure dandruff, baldness, dryness of the hair, etc.

Some of these preparations contain cinchona or quinine; some contain sulphur and lead salts; others contain fixed oils like benne or cocoanut oil, and others contain still other ingredients.

They are known by such titles as "hair restorer," "hair restorative," "hair invigorator," "hair vigor," "hair promoter," "hair grower," "hair lotion," "hair renewer," "hair wash," "hair renovator," "hair balsam," etc.

#### Quinine and Cinchona Hair Tonics.

These preparations are usually dispensed under the titles "quinine hair wash," "quinine hair tonic," and "eau de quinine."

They may be colored red, if desired, by means of red saunders.

I.	
Quinine sulphategr.	20
Bay rumfl.dr.	4
Glycerin	4
Tincture of cantharidesfl.dr.	2
Tincture of capsicumfl.dr.	2
Water, enough to makefl.oz.	16
Mix and dissolve.	
II.	
Quinine sulphategr.	20
Glycerinfl.oz.	1
Cologne waterfl.oz.	2
Bay rumfl.oz.	2
Rose waterfl.oz.	11

Rub the quinine with the glycerin and add the other ingredients in order named. The addition of fluid extract of jaborandi is recommended to stimulate the growth.

#### HI.

Quinine sulphategr.	30
Acetic acidfl.dr.	
Resorcin	120
Waterfl.oz.	4
Oil of eucalyptusfl.dr.	2
Tincture of cantharidesfl.dr.	3
Alcoholfl.oz.	12

Mix all, dissolve by agitation, and filter.

#### TV

V <b>.</b>	
Quinine sulphategr.	20
Tincture of cantharidesfl.dr.	2
Fluid extract of jaborandifl.dr.	2
Alcoholfl.oz.	2
Glycerinfl.oz.	2
Bay rumfl.oz.	6
Rose water, enough to makefl.oz.	
. 3	

The quinine should be dissolved in the alcoholic liquids by warming slightly, then the other ingredients added, and the whole filtered.

#### V.

Tincture of red cinchonafl.oz.	1
Tincture of cantharidesfl.dr.	2
Glycerinfl.oz.	
Compound spirit of lavenderfl.dr.	2
Alcohol, enough to make fl.oz.	16
171	

#### VI.

Ouinine sulphategr.	90
Diluted sulphuric acidm.	
Alcohol	2
Glycerinfl.oz.	1/2
Rose waterfl.oz.	14

Triturate the quinine with the acid, gradually adding the water until solution is effected; then add the remaining ingredients and filter.

#### VII.

0 1 11	A
Orris rootav.oz.	
Clovesgr.	10
Nutmeggr.	5
Red saundersgr.	20
Quinine sulphategr.	40
Cologne waterfl.oz.	1
Glycerinfl.oz.	11/2
Oil of lavender flowersdrops	4
Oil of rose geraniumdrops	8
Oil of nerolidrops	4
Water,	
Alcoholof each, sufficient	ent

Mix the orris, cloves, nutmeg and saunders, reduce to moderately fine powder, and

percolate with a mixture of one volume of water and three of alcohol, so as to obtain 29 fluidounces of percolate. To the latter add the remaining ingredients and dissolve by agitation.

#### VIII.

Fluid extract of pale cinchonafl.dr.	
Tincture of cantharidesfl.dr	-2
Glycerinfl.oz.	1
Bay rumfl.dr.	4
Rose water, enough to makefl.oz.	

#### TX

record water, enough to make,, moz	. ~0
X.	
Chloral hydrate gr.	240
Tincture of cantharidesfl.dr.	
Tincture of cinchonafl.dr.	4
Glycerinfl.oz.	1
Orange flower water,	
Rose water of each, equal parts,	
to makefl.oz.	16

Brush into the roots of the hair every morning, and rub in a little lanolin at night.

#### Х.

Tincture of rhatanyfl.dr.	1
Tincture of cantharidesfl.dr.	4
Spirit of lavenderfl.dr.	2
Glycerinfl.oz.	2
Quinine sulphategr.	15
Alcohol, enough to makefl.oz.	16
Mix, dissolve and filter.	

#### ΧŢ

_	.l. ,	
	Carbolic acidfl.dr.	11/2
	Tincture of nux vomicafl.oz.	1
	Tincture of cinchonafl.oz.	31/2
	Tincture of cantharidesfl.dr.	2
	Cologne waterfl.oz.	7
	Castor oilfl.oz.	4
	Shake well before using.	

#### XII

11.	
Alcohol fl.oz.	9
Spirit of soapfl.oz.	31/
Tincture of cinchonafl.oz.	2
Tincture of cantharidesfl.dr.	1
Peru balsamfl.dr.	5
Oil of bergamotfl.dr.	2
Oil of orangefl.dr.	2
Oil of rose geraniumfl.dr.	1
Mix and filter.	

#### XIII.

Quinine sulphategr. 15
Cologne waterfl.dr. 3
Rum
Alcoholfl.oz. 4
Glycerin, fl oz. 3
Rose waterfl.oz. 21
Alkanninsufficient to color
Mix, dissolve and filter.—D.

#### Lead and Sulphur Hair Promoters.

Hair preparations containing sulphur and lead salt not only act as alleged curatives, but also act as dyes or coloring agents, deepening the color of the hair. The sulphur and lead combine upon exposure to light to form black lead sulphide. For this reason such preparations should be kept darkened by means of an adherent wrapper. The public should be told lead preparations are not harmless.

I.		
	Lead acetateav.oz.	
	Sulphur, precipitatedav.oz.	1
	Tincture of cantharidesfl.dr.	4
	Glycerin fl.oz.	8
	Alcoholfl.oz.	4
	Oil of citronellafl.dr.	1
	Oil of bergamotdrops	30
	Water, enough to make fl.oz.	

Dissolve the oils in the alcohol, add the glycerin and tincture of cantharides, and mix with the water, then add the sulphur and the acetate of lead.

II. Here is a formula which makes a preparation without sediment. It should be kept from the light.

Lead acetateav.oz.	3/4
Sodium hyposulphiteav.oz.	21/2
Glycerinfl.oz.	8
Alcohol,fl.oz.	
Oil of lemonfl.dr.	1
Oil of bitter almondsdrops	15
Oil of clovesdrops	15
Rose waterfl.oz.	
Water, enough to makefl.oz.	64

Dissolve the lead acetate and sodium hyposulphite, each separately in two pints of hot water, and mix the solutions. Dissolve the oils in the alcohol, adding 16 fluidounces of water, and rub with 120 grains of magnesium carbonate in a mortar; filter and add the filtrate to the other mixture, now incorporate the glycerin, and the remainder of the water.

#### III.

Sulphur, precipitated av.oz.	1
Lead acetateav.oz.	1
Tincture of cantharides fl.dr.	4
Glycerinfl.oz.	2
Water, enough to makefl.oz.	

Dissolve the lead salt in a portion of the water, and add the remaining ingredients.

This preparation is known as Gen. Twigg's Hair Restorer.

IV.	V.
Precipitated sulphur         .gr. 60           Lead acetate         .gr. 60           Bay rum         fl.oz. 4           Jamaica rum         fl.oz. 2           Sodium chloride         .gr. 60           Rose water         fl.oz. 4           Glycerin         fl.oz. 2	Tincture of cantharides
Sulphur, precipitated       av.oz.       1         Lead acetate       av.oz.       1         Glycerin       fl.oz.       6         Rose water, enough to make       fl.oz.       32	twice daily.  VI.  Sweet almond oil
VI.       Precipitated sulphur.       av.oz.       1         Lead nitrate       av.oz.       34         Tincture of cantharides       fl.oz.       1         Glycerin       fl.oz.       8	Chloroform m. 50 Oil of rosemary fl.dr. 1½ Oil of lemon drops 12 Alcohol, enough to make fl.oz. 16 This preparation is well known under the
Alcohol fl.oz. 4 Oil of lavender flowers fl.dr. 2 Oil of bitter almonds drops 15 Water, enough to make fl.oz. 64	name Erasmus Wilson's Hair Lotion.  VII.  Castor oil
Dissolve the lead salt in the water and the oils in the alcohol, mix and add the remaining ingredients. Lead nitrate has the advantage over the acetate of being odorless.	Tincture of cantharides       .fl.dr. 3         Benzoic acid       .gr. 120         Tannic acid       .gr. 90         Alcohol       .fl.oz
Hair Tonics Containing Oil.	Mix and filter.
Castor oil.   fl.oz. 16     Alcohol.   fl.oz. 24     Tincture of cantharides   fl.dr. 5     Oil of bergamot   fl.oz. 1     Oil of rose   drops 10	Miscellaneous Hair Tonics.   I.   Oleo-balsamic mixture.   fl.oz. 4   Glycerin   fl.oz. 5   Rose water.   fl.oz. 20   Tincture of cantharides   fl.dr. 4
Red saundersenough to color Mix, macerate for several days, agitating occasionally and strain.	Ammonium carbonateav.oz. 1  Mix, shake thoroughly, let stand for one hour, and filter.
II.	II.
Carbolic acid	Tincture of cantharidesfl.dr. 3 Tincture of capsicumfl.dr. 1 Ammonia waterfl.dr. 2 Glycerinfl.oz. 1 Cologne water, enough to make.fl.oz. 16 III.
III.       Castor oil.       fl.oz.       16         Tincture of cantharides       fl.oz.       2         Tannin.       gr. 120         Oil of bergamot       fl.dr.       2	Tincture of cantharidesfl.dr. 4 Glycerinfl.oz. 4 Bay rumfl.oz. 11½  IV. Salicylic Hair Tonic:
Oil of cloves       fl.dr.       2         Oil of lavender       fl.dr.       ½         Oil of rosemary       fl.dr.       ½         Alcohol       fl.oz.       48	Salicylic acid.       .gr. 50         Borax       .gr. 150         Tincture of cantharides.       .fl.oz. 1         Bay rum.       .fl.oz. 6
IV.  Castor oil	Rose water

V.	'Pilocarpine Hair Wash.''
	carpine nitrate or muriategr. 4
Tine	ture of cantharidesfl.dr. 4
Colo	gne water fl.oz. 4
Soap	liniment, enough to make fl.oz. 16
App	ly to the scalp once daily.
See :	also next formula.

#### VI.

Oil of bergamot. fl.dr. 19 Oil of lavender flowers fl.dr. 19 Stronger water of ammonia fl.dr. 3 Tincture of cantharides fl.dr. 3 Castor oil fl.oz. 2 Alcohol, enough to make, fl.oz. 20
Alcohol, enough to makefl.oz. 20

The pilocarpine and oils should be dissolved in some of the alcohol, then the tincture and the ammonia added, and enough alcohol to make up the required volume.

The scalp should be thoroughly cleansed by washing with warm soap and water containing a little borax, and dried; the tonic is then to be applied with a small sponge.

#### VII.

Benne oil	.fl.oz.	16	
Lime water	.fl.oz.	24	
Oil of bergamot	.fl.oz.	2	
Oil of lemon	.fl.dr.	1	,
Oil of lavender flowers	.fl.dr.	I	2
Oil of cloves	. drops	15	

Mix the benne oil and water thoroughly and add the remaining ingredients.

This hair preparation is known by the misapplied appellation of "Lime Juice and Glycerin."

#### Hair Dyes.

These preparations are generally made by the use of silver, lead and bismuth salts, the first mentioned being usually the most effective and most expeditious. Some of these preparations require the use of a second or mordanting agent, others are made without the use of the latter. In the case of the twobottle preparations, the two vials are put up together in a neat pasteboard box.

The usual color desired is black, although brown is also largely in demand. The dyes that produce a black color will produce a brown by appropriate weakening or dilution.

Before using any dye, the hair should be freed from grease by washing with an alka-

by the abundant use of water, subsequently drying thoroughly.

The dye should be distributed among the hair by means of a tooth brush and occasional combing with a new comb: contact with the scalp should be avoided.

#### $I. - \Lambda.$

Pyrogallic acid	 	 		 dr.	1
Alcohol					
Water. distilled	 	 	• •	 fl.oz.	4

27		
Silver nitrate	 	dr. 1
Water, distilled		
Ammonia water.	 	enough

After dissolving the silver nitrate in the water, gradually add water of ammonia, stirring constantly, until the brown turbidity produced has vanished and the liquid appears colorless. Then add

Water, distilled, enough to make.fl.oz. 1

A large excess of ammonia tends to produce a brownish dye. Various shades of brown may be produced by increasing the amount of water in the silver solution. should be remembered that the hair must. previous to treatment, be washed with warm water containing sodium carbonate, well rinsed with clear water, and dried.

#### II.

Silver nit	rate	 		 		.gr.	120
Copper 1	nitrate	 	٠.			.gr.	12

Dissolve the two salts in 2 fluidounces of water and add ammonia water, constantly stirring, until the precipitate at first formed is dissolved and the liquid becomes clear and transparent. Make another solution of

Pyrogallic acidgr.	40
Acetic acidfl.dr.	
Alcohol fl.dr.	

Apply the pyrogallic solution to the hair with a stiff brush (a tooth brush will answer), taking care not to wet the scalp. When partially dry, apply the silver solution in the same manner, using another brush. For a brown dye, decrease the amount of pyrogallic acid; as little as one-half grain to the fluidounce is sometimes used. Sometimes the acetic acid is omitted, and in most of the commercial hair dyes diluted alcohol is used as a solvent of the acid. Copper sulphate line carbonate or soap, removing the latter might possibly be substituted for the nitrate. The use of the copper salt is to avoid the dull reddish tint imparted to hair by the use of silver nitrate alone.

#### III. Make two solutions as follows:

Bismuth subnitrate gr. 200
Waterfl.oz. 2
Nitric acid.gr. 420 or sufficient to dissolve
Use heat to effect solution also

Tartaric acidgr.	150
Sodium bicarbonategr.	168
Waterfl.oz.	

When effervescence of the latter has ceased, mix the cold liquids by pouring the latter into the former with constant stirring. Allow the precipitate to subside, transfer it to a filter or a strainer, and wash with water until free from the sodium nitrate formed, as this salt would be an unnecessary impediment to the operation of the dye. Now allow the magma to drain until its weight is reduced to at least 42 av. ounces. This can be readily determined without removing it from the filter and funnel, if both have been previously weighed. Transfer the magma, which consists of bismuth tartrate, to a dish, and dissolve it by the addition of sufficient stronger water of ammonia.

Next dissolve 100 grains of sodium hyposulphite in 3 fluidounces of water, and mix the two liquids. The total volume of the product should be about 7 or 8 fluidounces, which would make the solution contain about 10 per cent of bismuth tartrate, the product from above quantities being nearly 300 grains. The addition of 1 fluidounce of glycerin is calculated to make it more effective in coloring the hair, as this ingredient prevents entire drying up of the constituents, and thus favors a continuation of the decomposition.

Should it be desired to produce a jet black, this may be accomplished (after the dye given above has first been applied and allowed to dry) by the application of a solution of an alkaline sulphide or sulphuret. It is not necessary that the latter salt should be absolutely pure, as the commercial sulphuret of potassium answers well if fresh or undecomposed. The application of these dyes and mordants is usually made by means of a toothbrush and comb, so as to avoid staining the scalp.

IV.	
Silver nitrateav.oz.	1
Ammonia waterfl.oz.	2
Sodium carbonate, puregr. 1	
Distilled watersufficie	

Dissolve the silver salt in 8 fluidounces of water and ammouia water until the precipitate first formed is redissolved; dissolve the sodium salt in this solution, add enough water to make 12 fluidounces, and after standing a few days, decant the clear liquid.

As a mordant use:

Pyrogallic acidgr.	144
Alcohol fl.oz.	4
Waterfl.oz.	8

Mix and dissolve.

This is used as a brown dye.

<b>7.</b>	
Silver nitrategr. 48	0
Copper nitrate	0
Distilled waterfl.oz.	8
Ammonia watersufficien	

Dissolve the two salts in the distilled water and add the ammonia water until the liquid becomes a clear fluid.

In using, apply to the hair carefully with a tooth brush, after thoroughly cleansing the hair, and expose the latter to the rays of the sun.

#### VI.

Silver, metallicgr.	150
Iron, filings or reducedgr.	300
Nitric acidfl.dr.	10
Watersuffici	ent

Mix the first three, allow to stand until dissolved or nearly so, dilute with water to 4 fluidounces and filter.

Apply by means of a tooth brush to the hair previously cleaned with soap and water and dried.

#### VII.

Litharge	 	.av.oz.	1
Lime, freshly slaked	 	av.oz.	2
Starch	 	av.oz.	2

Mix all in fine powder and perfectly dry, and keep in well-stoppered bottles.

This powder is to be made into a thin paste or cream with water for use as a black dye or a thin milk for a brown dye. Before using, the hair should be freed from grease by means of soap and water, subsequently drying. Now apply the liquid by means of a sponge, brush or the fingers, observing to rub it well

into the roots of the hairs, and to pass a comb occasionally through the latter to insure uniform distribution and contact with every part. The hair should be kept in a moistened state during 3 or 4 hours, which can be effected by wrapping about the head a moist cloth or wearing a cap of oiled silk. When the coloring has been effected, remove the powder by washing it out with warm soap and water.

The danger in using this preparation is the possibility of lead poisoning.

VIII. The following is known as "walnut hair oil or dve:"

Green walnut	shellsa	v.oz. 2
Alum	a	v. oz. 1/4
Olive oil		fl. oz. 4

Heat together in a water bath until the water has been completely expelled; then express, filter and perfume.

#### Brilliantines.

These are preparations intended for application to the mustache and consist of mixture of alcohol, fixed oils and glycerin, with some perfuming agent. They are of two varieties, separable and non-separable, the former separating on standing into two layers. The cause of the separation is the presence of fixed oils other than castor and alcohol, and possibly glycerin or water. The non-separating kind consists of castor oil in alcoholic solution. If a colored preparation is desired, it may be tinted with saffron tincture, alkanet root, or alkannin.

Į
31
ĺ
3
3
3
į
ĺ
9
Ł
9
2
3
Ŀ
3
)
3

V.
Castor oil
Alcoholfl.oz. 8
Water
Oil of verbenadrops 10
Oil of rose geraniumdrops 15
Oil of lemondrops 18
Mix and color with tincture of saffron.
VI.
Castor oilfl.oz. 3
Glycerin
Alcohol
Perfume to suit.
VII.
Glycerin
Castor oil
Mix and perfume.
VIII.
Glycerin
Oleic acid         fl.oz. 1           Alcohol         fl.oz. 15
Mix and perfume.
_
IX.
Glycerin
Alcohol
Mix and perfume.
X.
Veal or beef suetav oz. 4
Spermacetiav.oz. 2
Castor oilav.oz. 12
Oil of bitter almondsdrop 1 Oil of clovesdrops 10
Oil of housement drops 10

Melt the suet and spermaceti, add the castor oil, and incorporate the flavoring oils, allowing to cool somewhat before adding the latter.

Oil of bergamot......drops 20

This preparation is more of the nature of a pomatum.

# Stick Mustache Pomades. (Stick Pomatum.—Cosmetique.)

T.	•	•		
I. White wax			av.oz.	. 8
Lard			av.oz.	8
Oil of bergamot.			fl.dr.	. 4

Melt the wax and lard together with a gentle heat, stir well as the mixture begins to cool, add the oil just before it is ready to set, and then pour into molds.

Hard pomade is usually cast in little bars or rolls varying in size according to fancy.

These bars or rolls are wrapped in well sized paper, and then enveloped in tin foil.

The molds should be of heavy metal, fashioned after the style of bullet molds. They should be chilled by immersion in very cold water just before casting; this will greatly lessen the chances of the pomade adhering to the molds. When only small quantities are made so that it is desirable to avoid the expense of the regular apparatus, paper molds may be substituted. They can be easily made by folding stout manilla or similar paper over a suitable model and securing it with a little mucilage at the edges. One end is turned in; the molds are then secured in an upright position and filled from the end remaining open. When cast in this way, the mold itself usually answers as the inner wrapper.

Stick pomade made as above is, as will be noticed from the recipe, white. It is frequently required to be brown or black. For the former it is colored by the admixture of a sufficient quantity of burnt umber. The umber must first be rubbed to perfect smoothness on a slab or tile with a little of the melted pomade. When black is desired, lampblack is substituted for the umber.

The perfume is, of course, varied at will. When strong smelling oils are used, yellow wax will usually answer as a substitute for the white when the pomade is to be colored. An article much superior in point of perfume to the foregoing may be made by the substitution of flower-scented pomades for the lard.

#### II.

White	wax.									av.oz.	6
Suet		 								av.oz.	3

Melt, mix thoroughly, add suitable perfume, and pour into molds. Color as described in No. I.

#### III.

White waxav.oz.	4
Beef tallowav.oz.	
Oil of bergamotfl.dr.	4
Oil of cassiadrops	40
Oil of thyme (white)drops	20

A yellow color is produced by tincture of safron or tincture of turmeric, a brown color by burnt umber in oil, and a black color by animal charcoal ground or triturated in oil.

#### IV.

White wax	.av.oz.	8
Suet	.av.oz.	8
Tuberose pomade	.av.oz.	4
Jasmine pomade	.av.oz.	4
Oil of roses		

First melt the wax, then add the suet, afterward the pomade. When beginning to cool, incorporate the oil of roses and cast into molds.

V. For other mustache pomades, see the next heading.

Hungarian Mustache Pomade. (Hungarian Mustache Wax.—Pomade Hongroise.)

This is a mixture of gum, soap, wax or spermaceti, and water, with perfume and coloring matter. It is in the form of a sticky or tenacious paste which, when applied to the ends of the mustache, retains the latter in extended fashion.

Owing to the fact that it will become hard and dry when exposed to the atmosphere, it must be preserved and dispensed in wellstoppered wide-mouth bottles.

#### I.

Waxav.oz.	41/2
White castile soapav.oz.	
Oil of bergamotfl.oz.	
Gum arabic, powderav.oz.	
Waterfl.oz.	

Dissolve the gum in the water; melt the wax and soap together on a water bath, stir in the solution of gum, and lastly, just before cooling, add the bergamot oil. Color, as required, with burnt umber or lampblack rubbed to perfect smoothness on a slab with a little of the melted wax.

#### II.

Spermacetiav.oz.	1
Wax av.oz.	4
Gum arabic, powderav.oz.	-3
White castile soapav.oz.	2
Glycerinav.oz.	1
Waterfl.oz.	

The soap should be finely shaved, and it and the gum stirred up with 4 fluidounces of water to a homogeneous paste. The spermaceti and wax should then be heated with the remainder of the water on a water bath, and stirred carefully into the gum and soap paste. Lastly, the glycerin should be added drop by drop. Perfumery may be added to

suit the taste; if a brown color is desired, umber should be mixed with the glycerin, and for black, lampblack.

#### III.

Gum arabic, powderav.oz.	4
White castile soap, powderav.oz.	4
White waxav.oz.	4
Rose waterfl.oz.	4
Oil of bergamot m.	80
Oil of sandalwoodm.	40

Triturate the gum with the water to a smooth paste; melt the wax, add the soap, stir in the gummy mixture before cooling, add the oils and color, if desired, as in the preceding formulas.

#### IV.

White castile soap, powderav.o	oz. 2
White waxav.d	oz. 5
Mucilage of acaciafl.	oz. 6
Distilled water	oz. 5
Glycerinfl.	oz. 11/2
Oil of bergamotdro	
Oil of lemondro	
Oil of rose dro	

Triturate the soap with the mucilage previously mixed with the water to a smooth mixture. To this add the wax and glycerin, heat the whole on a water bath, stirring constantly, until the wax is melted, and the mixture is homogeneous. Now, incorporate the volatile oils, also coloring matter, if a colored pomade is desired. For blond pomade, use yellow ocher; brown, burnt umber; and black, lampblack, each previously triturated to a smooth paste with the glycerin.—D.

#### SECTION IV—MOUTH PREPARA-TIONS.

#### Tooth Preparations. (Dentifrices.)

Tooth Powders.—These preparations must be reduced to fine powder by trituration and sifting, and the ingredients must also be mixed intimately. The customary ingredients are chalk, orris root, myrrh, cuttlefish bone, soap, pumice, etc.

The name given varies according to composition and according to the fancy of the maker. It may be called "myrrh tooth powder," "saponaceous tooth powder," "charcoal tooth powder," "thymol tooth powder," "salicylated tooth powder," "quinine tooth

powder," "coral tooth powder," "pearl tooth powder," "rose tooth powder," "antiseptic tooth powder," "salol tooth powder," "violet tooth powder," "aromatic tooth powder," "imperial tooth powder," "crown tooth powder," etc. In each case, the word "dentifrice" may be substituted for "tooth powder."

Tooth powders are usually flavored, the oils of peppermint and wintergreen being the most popular flavors.

Frequently they are also colored with carmine. This must be triturated to a smooth and intimate powder with a portion of the mixture before adding the remaining ingredients.

Tooth Creams and Pastes. - These preparations differ in consistence, the former being rather thin, the latter being a rather hard mass. The former are dispensed in collapsible, the latter in white or opal, jars. Both creams and pastes may be produced from tooth powders by the addition of sufficient glycerin, honey or simple syrup, the first mentioned being preferred on account of its antiseptic property which prevents the preparation from spoiling. The creams are usually made by the use of glycerin, these requiring more of the diluting agent than the pastes. Creams also differ from pastes in almost invariably containing soap, which is usually incorporated in the powdered condition.

Coloring matters and perfumes are added to tooth creams and pastes in the same manner as to tooth powders.

Suitable names for tooth pastes and creams are the following: "Menthol glycerin tooth cream (or paste)," "Castilian tooth cream (or paste)," "Persian tooth cream (or paste)," "cherry tooth cream (or paste)," "creta cream (or paste)," "Oriental tooth cream (or paste)," "saponaceous tooth cream (or paste)," "damask rose tooth cream (or paste)," "eucalyptus tooth cream (or paste), "coca tooth cream (or paste)," " coral tooth cream (or paste)," " salicylated tooth cream (or paste)," "odontine," "rose tooth cream (or paste)," "kalodont," "salol tooth cream (or paste)," "thymol tooth cream (or paste)," "violet tooth cream (or paste)," "dentine," "dentalba," "dental cream (or paste)," "antiseptic tooth cream (or paste)," "carbolated tooth cream (or paste)," "camphorated tooth cream (or paste)," "charcoal tooth paste," "ruby tooth cream (or paste)," "myrrhine tooth cream (or paste)," "Vienna tooth cream (or paste)," etc. The word "dentifrice" may in each instance be substituted, if desired, for the word "tooth" or for the phrase "tooth cream" or "tooth paste."

Tooth Washes or Liquid Dentifrices.— These are preparations made from quillaja or soap, and are colored and flavored. They replace the tooth powders, pastes, creams and soaps.

They may be known by such titles as "eudonto," "dentine," "quillaja tooth wash," "kalliodont," "odontine," "saponaceous tooth wash," "aromatic dentifrice," "dentoline," "antiseptic liquid dentifrice," "almond tooth cream," "Oriental tooth wash," "carbolated tooth wash," etc.

See also under heading "Mouth Washes." Tooth Soaps.—These differ from tooth pastes in being of still firmer consistence and in always containing soap. The other ingredients are practically the same as are employed in the production of the pastes.

#### Tooth Powder.

II.	
Precipitated chalkav.oz.	15
Sugarav.oz.	5
Boraxav.oz.	5
Orris rootav.oz.	5
Cardamomgr. 2	70
•	

Mix all of these ingredients, previously reduced to fine powder, flavor and color, if desired, with carmine.

#### III.

Precipitated chalkav.oz.	20
Orris root, powderav.oz.	2
Tanningr.	80
Oil of rosedrops	18
Oil of clovesdrops	15
Oil of pimentodrops	2
Tincture of muskdrops	30
Carminegr.	8
Alcoholfl.dr.	1

Mix the chalk, orris and tannin, thoroughly incorporate the carmine, and then add the oils and tincture previously mixed with the alcohol.

#### IV.

Cuttle fish, powderav.oz.	
Orris root, powderav.oz.	8
Chalk, precipitatedav.oz.	16
Oil of lemonfl dr.	4
Oil of nerolidrops	30
-	

#### V.

Precipitated chalk	av.oz.	12
Pumice stone		
Cuttlefish bone	av.oz.	2
Magnesium carbonate	av.oz.	1
Armenian bole	av.oz.	1
Oil of rose		
Oil of geranium		
Oil of cloves	.drops	20

Reduce the pumice stone and fish bone to fine powder, add the other ingredients and mix well.

#### VI.

Prepared chalkav.oz.	12
Orris rootav.oz.	
White castile soapav.oz.	1
Sugarav.oz.	
Oil of wintergreenfl.dr.	1

Reduce the chalk, orris root and sugar to fine powder, add the oil and mix well.

#### VII

Y 1.1.	
Chalk, precipitated	av.oz. 16
White castile soap, power	lerav.oz. 1
Licorice root, powder	av.oz. 1
Magnesium carbonate	av.oz. 2

Mix, flavor as desired, and color, if desired, with carmine.

#### VIII.

Talcumav.oz.	12
Cream of tartarav.oz.	1
Burnt alumav.oz.	1
Cochinealav.oz.	
Oil of peppermintfl.dr.	2

Powder carefully and finely.

#### Tooth Cream or Paste.

### 

II.	oils, then the glycerin and water previously
Precipitated chalkav.oz. 8	mixed, and triturate until well mixed.
White castile soap, powderav.oz. 4 Orris root, powderav.oz. 4	VIII.
Oil of sassafras drops 40	White castile soap, powder av.oz. 8½
Oil of baydrops 80	Precipitated chalk
Honey sufficient to form paste	Carminegr. 40
III.	Oil of peppermintfl.dr. 1½
Orris root, powderav.oz. 8	With the aid of glycerin make a paste.
Myrrh, powderav.oz. 2	
Pumice, powder	IX.
Oil of cloves. fl.dr. 2 Oil of lemon fl.dr. 2	Salicylic acidav.oz. 1 Precipitated chalkav.oz. 10
Oil of rosedrops 30	Talcum, powder
Solution of carminesufficient to color	White castile soap, powderav.oz. 2½
Honeyenough to form paste	Pumice stone, powderav.oz. $2\frac{1}{2}$
IV.	Sugar
Pumice, powderav.oz. 1	Oil of peppermint
Orris, powderav.oz. 4	
White castile soap, powderav.oz. 1/2 Tragacanth, powdergr. 70	With the aid of glycerin make a paste.
Chalk, precipitatedav.oz. 8	X. Honeyav.oz. 8
Solution of potassafl.dr. 1	Chalk, precipitatedav.oz. 8
Glycerin	Orrisav.oz. 8
Oil of cloves drops 20 Oil of rose drops 15	Carminegr. 60
Oil of rose geraniumdrops 15	Oil of cloves drops 30 Oil of nutmeg drops 30
V.	Oil of rose drops 30
	Simple syrupenough to form a paste
Precipitated chalkav.oz. 15 White castile soap, powderav.oz. 15	XI.
Orris, powderav.oz. 5	Pumice, powderav.oz. 4
Oil of peppermintfl.dr. 1	Orris, powderav.oz. 4
Oil of cinnamonfl.dr. ½	Myrrh, powderav.oz. 1
Carmine	Honey
VI.	Spirit of lemonfl.dr. 3
	Oil of clovesfl.dr. 1
Precipitated chalk	Oil of rose drops 16 Cochineal coloring sufficient to color
Orris, powderav.oz. 5	
Cinnamon, powderav.oz. 3	XII.
Cloves, powder	Cream of tartarav.oz. 10
Oil of cloves	Sugar of milk
	Essence of peppermint fl. dr. 1
Mix the ingredients and add sufficient	Mix well and make into a paste with a mix-
honey to form a mass. If a colored prepara-	ture of 3 parts honey and 1 part glycerin by
tion is desired, carmine or solution of carmine	weight.
may be added. VII.	XIII.
Chalk, precipitatedoz. 3	Calcium carbonate, precipitated. av. oz. 3
Magnesium carbonatedr. 2	Sugarav.oz. 2
Castile soap, white, powdergr. 40	Cream of tartargr. 60
Oil of cloves	Glycerin fl.dr. 4 Rose water fl.dr. 12
Oil of cassia	Castile soap, whiteav.oz. 8
Oil of lavender flowers	Alcoholfl.oz. 2
Oil of rose geranium	Oil of peppermintfl.dr. 2
Glycerinfl.oz. 1	Carmine, dissolved in ammoniagr. 60
Waterfl.dr. 6	Mix the calcium carbonate, sugar and
Mix the first three ingredients, add the	cream of tartar, and make into a paste with a

mixture of the glycerin and 4 fluidrams of the rose water. Dissolve the soap in the alcohol and the remainder of the water by the aid of heat, add to the previous mixture, and lastly incorporate the oil and carmine.

### XIV.

Precipitated chalkav.oz	. 16
White castile soap, powderav.oz	. 4
Oil of clovesdrops	s 40
Oil of nutmegdrops	s 40
Oil of rosedrops	s 20

Form into a paste with a mixture of equal parts of glycerin and water.

### XV.

Precipitated chalkav.oz.	8
White castile soap, powder av.oz.	4
Orris, powderav.oz.	4
Oil of sassafras drops	40
Oil of baydrops	8
Honey, enough or aboutav.oz.	13

### XVI.

Orris, powderav.oz. 1	12
Alum, powderav.oz.	$^2$
Prepared chalkav.oz. 1	10
Cochinealav.oz.	11/2
Potassium bitartrateav.oz.	1
Oil of clovesdrops 4	10
Oil of rosedrops 4	10
Rose waterfl.dr. 1	
Glycerin sufficient to form a mass	

# Tooth Wash.

### T.

Quillaja, coarse powderav.oz.	11/2
Cochineal, powdergr.	15
Glycerin	3
Oil of wintergreendrops	25
Alcoholfl.oz.	10
Peppermint water, sufficient to	
makefl.oz.	32

Mix the quillaja with the alcohol and 12 fluidounces of peppermint water, macerate for a few days, add the cochineal, glycerin and oil, macerate for another day, agitating occasionally and filter, adding the remainder of the water through the filter.

### H.

Castile soap, shavingsav.oz. 3
Glycerinfl.oz. 8
Alcoholfl.oz. 12
Water, hotfl.oz. 12
Oil of peppermintdrops 40
Oil of wintergreendrops 60
Oil of cloves
Incture of vanilla
Cochineal solutionsufficient

add the glycerin and vanilla tincture. Dis- and filter.

solve the oils in the alcohol. Mix both solutions, add sufficient coloring to produce the desired shade, and after having allowed it to stand for 24 hours, filter through paper.

# III.

Quillaja, powder	.av.oz. 4
Cudbear	gr. 60
Diluted alcohol	sufficient
Heliotropin	gr. 2
Oil of peppermint	drops 20
Oil of anise	drops 10
Alcohol	fl.oz. 1
Glycerin	fl.oz. 2

Macerate the quillaja and cudbear with diluted alcohol or extract by percolation, in each case obtaining 30 fluidounces of prod-To this add the heliotropin and oils dissolved in the alcohol, macerate for several days in a warm place, filter if necessary, and add the glycerin.

### $\mathbf{IV}$

T	v .	
	Quillaja, coarse powderav.oz.	2
	Orris, coarse powderav.oz.	1/2
	Boraxav.oz.	1
	Saccharingr.	10
	Oil of peppermintfl.dr.	1
	Oil of wintergreenfl.dr.	1/2
	Waterfl.oz.	18
	Glycerinfl.oz.	2
	Alcoholfl.oz.	12
	Solution of carminefl.dr.	2

Mix, macerate for 7 days, agitating occasionally, and filter.

Quillajaav.oz.	2
Orris rootav.oz.	1
Canada snake rootav.oz.	
Clovesav.oz.	
Alcoholfl.oz.	
Waterfl.oz.	
Honeyfl.oz.	2

Extract the drugs in powder form by percolation, using the above mixture of alcohol and water as a menstruum; to the percolate, add the honey, and filter.

### VI.

Thymolgr.	30
Oil of wintergreenfl.dr.	1
Oil of peppermintfl.dr.	1
Compound tincture of cardamom.fl.oz.	$-2\frac{1}{2}$
Glycerinfl.oz.	2
Sandalwoodav.oz.	1
Tincture of quillajafl.dr.	2
Alcoholfl.oz.	13
Water, enough to make fl.oz.	32

Dissolve thymol and oils in alcohol, add Dissolve the soap in the hot water, and other ingredients, let stand at least 24 hours

VII.
White castile soapav.oz. 2
Oil of orange peeldrops 15
Oil of cinnamondrops 10
Waterfl.oz. 8
Alcohol fl.oz. 8
Liquor coccineus, N. F. sufficient to color
Dissolve, mix and filter.
VIII

Alcohol	.fl.oz.	10
Water	.fl.oz.	121/2
White castile soap		
Oil of wintergreen		
Red saunders		
red bullingers, it is it is it is it.		

Dissolve the soap in a mixture of the alcohol and water, add the saunders, flavor with the oil, add enough water to make the liquid measure 30 fluidounces, and filter.

### IX.

Quillajaav.oz. 3
Star aniseav.oz. 1
Cloves gr. 120
(innamon gr. 120
Cudbeargr. 60
Oil of peppermintdrops 12
Diluted alcoholsufficient

Mix the drugs, reduce to powder, and extract by percolation with diluted alcohol so as to obtain 32 fluidounces of product in which the oil is to be dissolved.

# X

_
3/4
1/2
_

Dissolve the soap in one-half the alcohol and water mixed; macerate the myrrh with the remainder of the alcohol and water for several days, filter each, mix and add the remainder of the ingredients. If the mixture is to be colored with alkannin, it may be added now; if alkanet is used, it may be added to the myrrh during maceration.

### XI.

Soap bark, powderav.oz. 4
Glycerin
Diluted alcoholfl.oz. 30
Oil of wintergreendrops 20
Oil of peppermintdrops 20
Mississian and the state of the

glycerin and dilute alcohol for three or four days, and filter through a little magnesia previously triturated with the volatile oils.

### XII.

Castile soap, whitegr. 270
Glycerin
Simple syrupfl.oz. 2
Waterfl.oz. 13
Alcoholfl.oz. 13
Tincture of cardamomfl.dr. 2
Tincture of Canada snakerootfl.dr. 2
Oil of peppermintdrops 25
Oil of wintergreendrops 25
Oil of clovesdrops 6
Oil of cassiadrops 6
Solution of carminesufficient to color

Mix the soap, glycerin, syrup and water, stir well and add the alcohol. Add the remainder of the ingredients, let stand a few days and filter at a low temperature, so that no soap will afterward precipitate.

# XIII.

Quillajaav.	oz. 2
Orris root	gr. 120
Cinnamon	gr. 120
Cochineal	gr. 60
Benzoic acid	
Tannin	gr. 60
Borax	
Oil of wintergreen	m. 30
Oil of peppermint	m. 30
Sugarav.	oz. 4
Glycerinfl.c	
Alcoholfl.c	
Waterfl.c	oz. 20

Reduce the drugs to powder, add to the remaining ingredients, macerate for at least and filter.

# XIV.

White castile soapav.oz.	2
Oil of peppermintdrops	20
Oil of wintergreendrops	50
Glycerin fl. oz.	2
Waterfl.oz.	4
Alcoholfl.oz.	8
Solution of cochineal, sufficient to co-	lor
Dissolve the soap in the alcohol and w	ater.

add the other ingredients, and filter.

### XV.

Quillaja barkgr.	280
Cochineal gr.	10
Oil of wintergreenm.	40
Alcoholfl.oz.	41/2
Peppermint waterfl.oz.	5
Glycerinfl oz.	2
Water, enough to makefl.oz.	30
Mix the cochineal and quillaja with 4	fluid-

Macerate the soap bark in the mixture of ounces of alcohol and 6 of water, macerate

for 7 days, agitating occasionally; filter, and add the remaining ingredients.

This may be pleasantly modified by the addition of a few drops of either oil of cinnamon or oil of cloves, or both.

# Tooth Soaps.

Precipitated chalk	.av.oz.	4
Carmine	gr.	25
White castile soap, powder	.av.oz.	10
Oil of peppermint	.fl.dr.	21/2
Alcohol	fl.oz.	11/2
Ammonia water	.sufficie	ent

Triturate the carmine with a few drops of ammonia water and add the precipitated chalk, mixing intimately. Dissolve the oil of peppermint in the alcohol; add the solution to the soap contained in a mortar and thoroughly incorporate; then add the precipitated chalk, and when the whole is homogeneous, transfer to suitable molds and dry.

TT.

White castile soap, powderav.oz.	10
Talcum, powderav.oz.	4
Pumice stone, powderav.oz.	2
Cuttle-fish bone, powderav.oz.	2
Cochineal, powdergr.	90
Sodium carbonate, driedgr.	180
Diluted alcoholfl.oz.	1
Glycerinfl.dr.	6
Oil of peppermintfl.dr.	2
Rose watersuffic	ient

Mix the soap, talcum, pumice and cuttlefish bone, then add the cochineal previously triturated to a fine powder. Having mixed these ingredients thoroughly, add the soda, alcohol and glycerin, stirring well, incorporate the oil, and then add enough rose water to form a mass. Divide into pieces or press in the boxes, and allow to dry.

# III.

White castile soap, powderav.oz.	10
Tincture of rhatanyfl.oz.	31/
Precipitated chalkav.oz.	334
Benzoic acidav.oz.	1/2
Potassium chlorate, powderav.oz.	3/1
Borax, powderav.oz.	3/4
Saccharingr.	10
Oil of cinnamon sufficient to flav	or

Make into a hard mass by the addition of glycerin and water, press into tin boxes, and dry,

IV.

Talcum, powderav.oz.	10
Pumice, powderav.oz.	1/2
Orris root, powderav.oz.	2

Mix well and color with carmine if a pink or red color is desired, and with chlorophyll, if a green color is desired, and flavor with a mixture consisting of:

Oil of peppermint Oil of sage. Oil of calamus. Oil of thyme, white. Coumarin	.fl.dr. .fl.dr. .fl.dr.	1 1/4 1 1/2
Now mix		
White castile soap, powder Alcohol, Glycerin	.fl.oz.	5

Beat together to form a soft paste, and then gradually incorporate the previous mixture of powders. Press the mass into molds, and, after removing the cakes, brush the latter over with tincture of benzoin containing a little oil of peppermint. When dry, cover with tin foil. The mass may also be pressed into tin boxes and allowed to dry in the latter.

# Camphorated Tooth Powder.

1.
Precipitated chalkav.oz. 10
Orris root, powderav.oz. 5
Camphorav.oz. 1
II.
Prepared chalkav.oz. 8
Orris rootav.oz. 4
Camphor
Cinnamonav.oz. 1
Reduce all to powder, and mix well.
ÍII.
Prepared challs over 16

Prepared chalkav.oz.	16
Camphor av.oz.	8
Cuttle-fish boneav.oz.	
Myrrhav.oz.	
Boraxav.oz.	2
Rose pinkav.oz.	1
Reduce all to powder, and mix well.	

# Charcoal Tooth Powder.

I.		
	Orris rootav.oz.	5
	Myrrhav.oz.	5
	Charcoal	11
	Cinchonaav.oz.	11

All should be in very fine powder and the whole should be well mixed and finally passed through a fine sieve.

II.

Charcoalav.oz.	12
Myrrhav.oz.	2
Pale cinchonaav.oz.	2

Mix well, having first reduced each to fine powder.

### Harlan's Tooth Paste.

Precipitated chalkav.oz. 8
Orris root, powderav.oz. 8
White castile soap, powder av.oz. 2
Borax, powderav.oz. 2
Myrrh, powderav.oz. 1
Honey,
Glycerin

of each sufficient to form a soft paste Carmine.....sufficient to color

Perfume to suit.

# Hunter's (John) Tooth Powder.

Cream of tartarav.oz.	
Alumav.oz.	
Cochinealav.oz.	2
Cinnamonav.oz.	1
Sugarav.oz.	4

Mix all, reducing to fine powder. Color, if desired, with carmine.

### Marshall's or Hudson's Dentifrice.

Chalk, preparedav.oz.	15
Myrrh, powderav.oz.	5
Orris, powderav.oz.	5
Rose pinkgr.	125

Mix well, reduce to fine powder, and sift.

### Menthol Glycerin Tooth Cream.

Precipitated chalkav.oz. 8
White castile soap, powderav.oz. 4
Magnesium carbonateav.oz. 2
Menthol (dissolved in alcohol),
Solution of carmine,
Glycerinof each, sufficient

Rub the first three ingredients into a paste with glycerin, then flavor and color to suit with the menthol and carmine solutions.

# Rose Tooth Powder.

Prepared chalkav.oz.	221/
Sugar of milkav.oz.	8
Orris root, powderav.oz.	11/2
Carminegr.	
Oil of rosedrops	16

Rub the chalk, orris root and 5 av.ounces of the sugar of milk together in a capacious mortar, and pass the mixture through a No. 80 sieve. Then rub the carmine in the mortar, and gradually add to it, while rubbing, the remaining 3 av.ounces of sugar of milk. To this mixture add the oil of rose, and, after rubbing all well together, add to it about 3 av.ounces of the sifted mixture. Stir this well together and also pass it through the sieve. Finally return all the sifted powder

into the mortar and thoroughly mix the whole of it.

# Quinine Tooth Powder.

Precipitated chalkav.oz.	29
Orris rootav.oz.	31/2
Sugar of milkav.oz.	31/2
Saccharingr.	4
Pumice stonegr.	390
Magnesium carbonategr.	390 -
Tannic acidgr.	300
Quinine hydrochlorategr.	80
Oil of rosedrops	16
Oil of peppermintdrops	80
Oil of ylang ylangdrops	5
Oil of bitter almondsdrops	5
-	

Mix all, and reduce to a fine, uniform powder.—D.

# Salicylated Tooth Powder.

Sodium salicylategr.	120
Sodium bicarbonateav.oz.	4
Precipitated chalkav.oz.	16
Myrrĥav.oz.	3,1
White castile soapav.oz.	11/2
Orris rootav.oz.	3′~
Licoriceav.oz.	2
Oil of wintergreendrops	3
Oil of rose geraniumdrops	30
o i	

Reduce the myrrh, soap, orris and licorice to fine powder, mix all the ingredients, and color, if desired, with carmine or solution of carmine.

# Salicylated Tooth Paste.

Precipitated chalkav.oz.	16
White castile soap, powder av.oz.	4
Sugar, powderav.oz.	4
Orris, powderav.oz.	
Pumice, powderav.oz.	
Sodium salicylategr.	80
Glycerinfl.oz.	
Carmine or solution of car- minesufficient to co	
Water sufficient to form a m	ass

Mix well and perfume with oil of peppermint, wintergreen or other oil.

### Saponaceous Tooth Powder.

I.

White castile soap, powder av.oz. 8
Precipitated chalkav.oz. 4
Magnesium carbonate av.oz. 3
Sugar, powderav.oz. 2
Oil of wintergreen sufficient to flavor

II.

Precipitated chalkav.oz.	14
White castile soap, powderav.oz.	2
Saccharingr.	
Oil of wintergreendrops	

# Thymol Tooth Paste.

Calcium carbonateav.oz.	16
Magnesium carbonate av.oz.	3/4
Orris root, powderav.oz.	3
Thymolgr.	60

Mix well and make a mass with sufficient of the following mixture:

Gelatin, puregr.	70
Glycerinfl.oz.	3
Waterfl.oz.	1

Dissolve by the application of a gentle heat.

# Thymol Tooth Powder.

Precipitated chalkav.oz.	15
White castile soap, powder av.oz.	1
Saccharingr.	
Thymolgr.	
Camphorgr.	
Vanillingr.	
Oil of rosedrops	6

Rub the camphor and thymol together in a mortar, and warm gently so as to render the mixture liquid; then add the chalk in small portions at a time, reserving about 1 av. ounce; next add the other ingredients, the perfumes being first separately rubbed with the remainder of the chalk.

# Mouth Washes.

These are preparations intended for cleansing, purifying and deodorizing the mouth, and frequently also for cleansing the teeth; before use, they are usually diluted with water, about one teaspoonful being added to a cupful of the latter. They always contain antiseptic and flavoring constituents, usually also an astringent substance like tannic acid, kino, rhatany, oak bark, etc., and frequently also a coloring constituent.

They are usually dispensed under such names as "prophylactic tooth and mouth wash," "dentifrice elixir," "thymol dentifrice," "salol mouth wash," "aromatic mouth wash," "astringent mouth wash," "chinoline mouth wash," "tooth tincture," "mouth essence," "elixir of roses," "violet mouth wash," "rubicreme," "favorite tooth and mouth wash," "mentholated dentifrice," "mentholine tooth wash," "eau dentifrice," "Imperial mouth wash," "salicylated mouth wash," "eau angelique," "carbolated tooth wash," "arnica tooth wash," "mouth water," etc.

ı	1.
ļ	Salolgr. 75
	Alcohol fl oz. 16
	Solution of cochinealfl.dr. 4
	Oil of rosedrops 8
	Oil of peppermintdrops 15
	1 1 1 1

Dissolve the salol in the alcohol, add the remaining ingredients, and filter.

II.		
White oak bark	av.oz.	21/2
Rhatany root		1/2
Sassafras	gr.	60
Red cinchona	gr.	180
Cardamom seeds	gr.	30
Cloves	gr.	30
Ceylon cinnamon	gr.	20
Oil of wintergreen.	fl. dr.	1
Oil of anise		1/2
Alcohol,		, -
Water	of each, suffic	ient

Reduce the solid substances to a coarsepowder, and extract by percolation so as to obtain 32 fluidounces of product, using as menstruum a mixture of 3 volumes of water and 5 of alcohol; to the percolate add the two oils.

III. The preparation known by the name "Eau de Botot," is dispensed according to different formulas, as follows:

Α.		
	Star anisegr.	720
	Cassia gr.	288
	Cloves gr.	288
	Cochineal gr.	72
	Waterfl.oz.	8
	Alcoholfl.oz.	24
	Oil of peppermintfl.dr.	11/
	Oil of rosedrops	

Mix the drugs, reduce to coarse powder, add the alcohol and water, macerate for 7 days, agitating occasionally, filter, and add the oils. The drugs may also be extracted by percolation as in the preceding formula.

В.	
Clovesav.oz.	1/2
Cinnamonav.oz.	1/2
Star aniseav.oz.	1/2
Rose waterfl.oz.	31/2
Alcoholfl.oz	
Cochineal, powdergr.	48
Cream of tartargr.	
Oil of peppermintfl.dr.	11/4

Reduce the cloves, cinnamon and anise to coarse powder, macerate in the rose water and alcohol for 24 hours, add the remaining ingredients, macerate for another 24 hours, agitating frequently, and filter.

C.		
	Star anisegr.	360
	Cloves gr.	360
	Galangagr.	360 -
	Ceylon cinnamongr.	360 -
		144
	Tannic acidgr.	72
	Peru balsamgr.	72
	Oil of peppermintfl.dr.	21/2
	Oil of rosedrops	15
	Oil of nerolidrops	8
	Oil of orrisdrop	1
	Coumarin sugargr.	15
	Waterfl.oz.	8
	Alcoholfl.oz.	24

Reduce the drugs to powder, mix all the ingredients, macerate for 8 days, agitating occasionally, and filter.—D.

This preparation may also be made by extracting the mixed and ground drugs by percolation, by means of a mixture of one volume of water and 3 of alcohol, so as to obtain 32 fluidounces of percolate; to the latter, add the acid, balsam, oils and sugar, and dissolve.

### IV.

Kino, powderav.oz.	5
Tincture benzoinfl dr.	1
Tincture tolufl.dr.	1
Tincture vanillafl.dr.	1/2
Oil of peppermintfl.dr.	1
Oil of anise	1/2
Oil of cinnamonfl.dr.	1/2
Alcohol fl.oz.	32´¯

Mix, macerate for 7 days, agitating occasionally, and filter.

This preparation is known as Eau de Mialhe.

### v.

Myrrh, powderav.oz.	1
Borax, powderav.oz.	1
Red saundersav.oz.	1
Sugarav.oz.	1
Cologne waterfl.oz.	6
Alcoholfl.oz.	
Waterfl.oz.	9

Mix, macerate for several days, agitating occasionally, and filter.

### VI.

Chinoline tartrategr.	150
Oil of peppermintfl.dr.	
Alcoholfl.oz.	
Waterfl.oz.	28

Add a teaspoonful to a tumblerful of water and use as a mouth wash.

3.7	1	т	

Oil of sagefl.dr.	23/4
Oil of lemonfl.dr.	114
Alcoholfl.oz.	61/2
Waterfl oz.	251/2

This preparation has been known as Eau de Salvia.

#### VIII.

Oil of sassafras (preferably
" Safrol ")fl.dr. 2
Oil of pinus pumiliofl.dr. ½
Oil of curacaofl.dr. 1/2
Oil of wintergreendrops 8
Oil of calamusdrops 10
Oil of anisedrops 2
Oil of rose geraniumdrops 2
Oil of vetivertdrops 2
Betanaphtholgr. 20
Solution of saccharinfl.dr. 11/2
Chloroformfl.dr. ½
Glycerin
Alcoholfl.oz. 8
Calcium phosphateav.oz. 34
Water, enough to makefl.oz. 32

In the alcohol dissolve the oils, naphthol, and chloroform. Add to the solution the glycerin and solution of saccharin, and gradually add water until the product measures 32 fluidounces. Allow the mixture to stand some time, frequently shaking; then mix it intimately with the calcium phosphate, and filter.

### IX.

Saccharingr.	2
Sodium bicarbonategr.	60
Salicylic acidav.oz.	1/2
Alcoholfl.oz.	32

Mix, dissolve, and filter.

A few drops in a glass of water make a gargle or collutory which is to be used frequently for bad breath.

# X.

Potassium	permanganategr.	30
	aterfl.oz.	

Use a teaspoonful to a tumblerful of water for a mouth wash.

# Lip Salves or Pomades.

These are preparations for anointing the lips to cure or prevent "cracking." They consist usually of white or yellow wax, spermaceti, paraffin, cacao butter, petrolatum, or lard mixed with an oil like olive or sweet almond, the whole being flavored, usually with oil of rose. Frequently they are tinted

a rose color by means of carmine, alkannin, or alkanet root. The first mentioned must first be triturated to very fine powder before adding oil or fat, and during cooling the fatty mixture must be frequently stirred to prevent it from subsiding. Alkannin is easier to use because it dissolves in the fats. colors the fatty mixture during a rather prolonged maceration assisted by heat. If a colored preparation is to be made, yellow wax should be substituted for white wax, wherever the latter is mentioned, owing to the better keeping qualities of the former. Camphor may be added to these preparations, also salicylic acid.

The preparations are usually known by such titles as "rose lip salve," "tulip salve," "salicylated lip salve," "lip pomade," "rose cerate," "lip ointment" and "coral lip salve."

Ι.

White waxav.oz. 1
Sweet almond oilfl.oz. 2
Carminegr. 1
Oil of rosedrop 1

Melt the wax, add the sweet almond oil, triturate the carmine to very fine powder, mix intimately with the fats, and then incorporate the oil of rose. - Codex.

TT.

White waxav.oz.	1
Spermacetiav.oz.	1
Liquid petrolatumfl.dr.	6
Tincture of benzoinfl.dr.	2
Alkanet root, bruisedgr.	120
Oil of rose drop	

Melt the wax and spermaceti, add the liquid petrolatum and the root, macerate the latter with the melted fats for about one-half hour, strain, allow to cool somewhat, and incorporate the tincture and oil.

III.

Spermacetiav.oz. 2
Yellow waxav.oz. 1
Sweet almond oilfl.oz. 4
Oil of rosedrops 5
Alkanet rootsufficient to color
M 1, 1 C . 11 11 1 1 1 1

Melt the fats, add the almond oil, color with the alkanet, strain and add the rose oil.

IV.

Spermacetiav.oz.	1
Lardav.oz.	2
White waxav.oz.	1/2
White waxav.oz. Sweet almond oilfl.dr.	2
Mir and malts if a salared meanant	

add a small amount of bruised alkanet root. macerate for a short time at a moderate temperature, strain and perfume to suit.

Paraffin waxav.oz	. 2
Cacao butterav.oz.	. 2
Petrolatum, whiteav.oz.	21/2
Eosingr.	
Oil of rosedrops	

Dissolve the last two ingredients in a minimum quantity of alcohol, and add to the fats when melted.

Yellow waxav.oz.	1
Olive oilfl.oz.	
Alkanet root, bruisedgr.	35
Oil of rose drops	2

Melt the wax, add the oil and root, macerate at a moderate heat for an hour, strain, allow to cool somewhat and incorporate the oil.—H.

VII.

White waxav.oz.	1/2
Spermacetiav.oz.	1/2
Sweet almond oilfl.oz.	1
Oil of rosedrop	1

Melt the wax and spermaceti, add the almond oil, and incorporate the oil of rose. -D.

VIII.

Yellow waxav.oz.	1 3/4
Spermacetiav.oz.	1/4
Sweet almond oilfl.oz.	3
Oil of lemondrops	10
Oil of bergamotdrops	
Alkanningr.	
G	

Melt the wax and spermaceti, add the almond oil, and add the remaining ingredients.-D.

IX. To the latter may be added, if desired, 12 grains of salicylic acid.—D.

X.

Cacao	butter									av.oz.	4
White											

Mix by fusion, and perfume to suit.

Cold creamav.oz.	4
Glycerinfl.dr.	
Tincture of benzoinfl.dr.	
Carminesufficient to col	or

Rub the carmine with the glycerin, and incorporate with the cold cream; then add Mix and melt; if a colored preparation, the tincture of benzoin, and rub the ointment until the alcohol of the tincture has evaporated.

### XII.

Carmine, fine	powdergr. 5	)
Glycerin	fl.dr. 8	3
Cold cream	av.oz.	1

Rub the carmine with the glycerin and intimately mix with the cold cream. the shade to suit, more or less carmine may be used.

This and the preceding preparation may be entitled "rose lip cream."

#### Cachous.

These consist of the various aromatics combined with licorice extract, sometimes sugar, the whole being formed into a mass which may be divided into pellets, or it may be rolled out in a thin sheet and cut into little squares, or it may be rolled out into a very thin pill "pipe," which may then be cut into short sections. After dividing the mass, the particles may be dried. If desired, the pellets may be silver-coated.

I.

Oil of peppermintdrops	30
Oil of lemondrops	20
Oil of nerolidrops	20
Oil of cinnamondrops	20
Cloves	40
Cardamomgr.	80
Vanillagr.	120
Orris rootgr.	150
Mace	400
Sugargr.	300
Licorice extract, powderav.oz.	11/4
Mucilage of gum arabic suffici	ent
-	

Reduce the drugs to powder, add the remaining ingredients, make a mass and divide into pills weighing 1 grain each, or roll out flat and cut into small pieces.

#### II.

Licorice extractav.oz.	3
Gum arabicav.oz.	1/2
Catechu av.oz.	1
Mastic	60
Cascarillagr.	60
Charcoal gr.	60
Orris root gr.	60
Oil of peppermintfl.dr.	1/2
Waterfl.oz.	

Dissolve the licorice in the water on a water bath, add the gum arabic and catechu, evapthe remaining solids reduced to fine powder, and finally when of proper consistence re- flat and cut into pieces.

move from the fire, add the oil of peppermint, and divide into small pellets, or roll the mass out flat and cut into small pieces.

#### III.

Licorice extract, powderav.oz.	2
Oil of clovesfl.dr.	1
Oil of cinnamondrops	10

Mix well, add sufficient mucilage of acacia to form a mass, and divide into pellets, or roll out flat and cut into small pieces.

### IV.

Nutmeggr.	192
Cardamomsgr.	
Vanillaav.oz.	1/2
Clovesgr.	64
Orrisgr.	256
Muskgr:	1
Oil of peppermintfl.dr.	1
Oil of lemon	40
Oil of cinnamon drops	10
Oil of nerolidrops	20
Sugarav.oz.	11/4
Licorice	21/4
Extract of licorice soft	~ /4
Waterof each, suffice	iont
wateror each, sume	lent

Reduce the drugs to fine powder, add the remaining ingredients, mix well, make a mass with the extract and water, form into pellets, or roll the mass out and cut into small pieces.

# V.

Extract of licoriceav.oz.	3
Catechu, powderav.oz.	1
Sugar powderav.oz.	
Tragacanth, powderav.oz.	
Oil of clovesfl.dr.	
Oil of cassiafl.dr.	
Oil of nutmegdrops	10

Make a mass with water and divide into 1grain pills.

### VI.

Orris root, powderav.oz. 5
Musk
Coumaringr. 12
Vanillin
Oil of rosedrops 10
Oil of nerolidrops 20
Oil of peppermintdrops 20
Oil of spearmintdrops 20
Oil of ylang ylangdrops 5
Purified extract of licoricesufficient

Mix the orris root with the remaining orate to the consistence of an extract, add ingredients, add enough extract to form a mass. Divide this into pellets, or roll out

VH.
Sugar, powderav.oz. 3
Licorice, powderav.oz. 3
Oil of anisedrops 20
Oil of fenneldrops 5
Purified extract of licoricesufficient

Mix the first four ingredients, then add enough of the latter to form a mass, divide into pellets, or roll out flat and cut into small pieces.—D.

### VIII.

Soft extract of licoriceav.oz.	
Catechu, fine powderav.oz.	3
Sugar, finely pulverizedav.oz.	1
Tragacanth, powder av.oz.	1/2
Oil of cloves fl.dr.	1
Oil of cassiafl.dr.	1/2
Oil of nutmegfl.dr.	1/2
Tincture of ambergrisdrops	$12^{'}$
Orange flower watersufficie	

With the aid of the water heat into a hard pilular mass, which divide into 2-grain cachous.

# SECTION V—PREPARATIONS FOR THE BATH.

# Bath Powder.

I.	
Tartaric acidav.oz.	10
Sodium bicarbonateav.oz.	
Starchav.oz.	6

A few spoonfuls of this when stirred into a bathtubful of water cause a copious liberation of carbon dioxide, which is thought by some to be "refreshing."

Perfume may be added to this powder, volatile oils being a good form. Oil of lavender flowers would be a suitable addition in the proportion of a fluidram or more to the av.pound of powder. A better but more expensive perfume may be obtained by mixing 1 part of oil of rose geranium with 6 parts of oil of lavender flowers.

A perfume still more desirable may be had by adding a mixture of the oils from which cologne water is made. For an ordinary quality the following will suffice:

Oil of lavender flowers fl.dr. 1
Oil of rosemary
Oil of lemonfl.dr. 4
Oil of clovesdrops 8

For the first quality the following may be taken:

Oil of neroli	fl.dr. 2
Oil of rosemary	fl.dr. 1
Oil of bergamot	fl.dr. 1
Oil of cedrat	fl.dr. 21/4
Oil of orange	

A fluidram or more of either of these mixtures may be used to the pound, as in the case of lavender.

I.	
Borax, powderav.oz.	4
Salicylic acidgr.	60
Essence of cassiefl.dr.	1
Essence of jasminefl.dr.	1
Oil of lavender flowersdrops	

Rub the oil and extracts with the borax and acid until the alcohol has evaporated. Use a heaping teaspoonful to the body bath.

#### III.

Borax		 .av.oz. 8
White castile	soap	 .av.oz. 8

Mix both ingredients, first reduced to powder, and perfume, if desired, as directed in No. I.

# Bath Tablets.

These are formed from the preceding powders by moistening with alcohol; No. 3 may be moistened with water to form tablets.

# Artificial Sulphur Baths.

Sulphur baths ordinarily are made by simply dissolving potassium sulphuret (sulphurated potassa) in water, in the proportion of from ½ av.ounce to 2 av.ounces for every 40 gallons of water. But, in order to obtain a bath more closely resembling some of the more noted natural sulphurous springs which have proven so effectual in the treatment of rheumatism and skin diseases of certain types, the following is advised:

Sulphurated potassa or sodaav.oz.	
Sodium bicarbonateav.oz.	1
Sodium chloride gr.	60
Castile soap shavingsgr.	30
Alumgr.	30
Calcium carbonategr.	30
Water gal.	1

These various materials are boiled in a sufficient quantity of the water to dissolve them, and the solution is stirred about with a wooden or glass rod until an odor of sulphuretted hydrogen becomes manifest. The solution is then poured into the patient's ordinary water bath, previously heated to about 35 degs. C.

# PART VI.

# SODA WATER PREPARATIONS.

# Coloring for Syrups.

In coloring either orange syrup or strawberry red, nothing is perhaps equal to a good black raspberry juice. It makes a nice color and is unobjectionable in every way.

Another convenient and excellent preparation is cochineal coloring, N. F., which may be prepared as follows:

Cochineal, powdergr. 48	U
Potassium carbonategr. 24	0
Alumgr. 24	
Cream of tartargr. 48	
Glycerinfl.oz.	
Alcoholfl.oz.	
Water, enough to makefl.oz. 1	

Triturate the cochineal intimately with the potassium carbonate and 8 fluidounces of water. Then add the alum and then the cream of tartar; heat the mixture to boiling in a capacious vessel; set it aside to cool, add the glycerin and alcohol, filter, and pass enough water through the filter to make 16 fluidounces.

# Essence, Birch.

Oil of wintergreen	.fl.oz.	5
Oil of lemon	.fl.dr.	2
Oil of cloves	.fl.dr.	1/2
Extract of vanilla	.fl.oz.	4
Alcohol	.fl.oz.	12

Dissolve the oils in the alcohol, and add the extract of vanilla.

# Essence of Ginger, Soluble.

I.	
Fluid extract of gingerfl.oz.	4
Pumice, fine powderav.oz.	
Water, enough to makefl.oz.	

Introduce the fluid extract into a bottle, add the pumice, and shake the mixture thoroughly and repeatedly during the course of several hours. Then add the water in portions of about 2 fluidounces, shaking well and repeatedly after each addition. When all is added, repeat the agitation occasionally during 24 hours, then filter returning the first portions of the filtrate until it runs through clear, and, if necessary, pass enough

water through the filter to make 12 fluidounces.—N. F.

II.

Jamaica ginger, ground....av.oz. 16 Alcohol.....fl.oz. 8

Mix, let stand for several hours, and with same menstruum percolate to obtain 24 fluidounces. To this tincture, add 2 av.ounces heavy magnesium carbonate, shake well, and add 24 fluidounces of water, shake again, and filter. If the filtrate is turbid, add more magnesium carbonate and filter again. It deposits slightly on standing a few days, but if again filtered, it remains clear.

# Essence, Mead.

Oil of clovesdrops	20
Oil of pimentodrops	
Oil of nutmegdrops	30
Oil of corianderdrops	10
Oil of sassafrasdrops	$\tilde{20}$
Oil of cinnamondrops	
Oil of lemonfl.dr.	
Extract of vanillafl.oz.	4
Alcoholfl.oz.	8
	4
Waterfl.oz.	-
Magnesium carbonatedr.	4

Mix the oils, dissolve in the alcohol, add vanilla and water and rub with magnesia and pass through filter, to make 1 pint; use of this essence 1 fl.ounce to 1 gallon of the finished syrup.

# Essence, Sarsaparilla.

Oil of wintergreenfl.dr. 4
Oil of sassafrasfl.dr. 4
Alcohol
II.
Oil of wintergreenfl.dr. 4
Oil of sassafrasfl.dr. 3
Oil of anisefl.dr. 1
Alcohol
Water, enough to makefl.oz. 16
Dissolve the oils in the alcohol and add the
water.

III.

Oil of wintergreenfl.dr.	2
Oil of anisefl.dr.	
Oil of sassafrasfl.dr.	
Alcohol, enough to makefl oz.	16

### Extract of Vanilla.

It may be said that the process of manufacture has less to do with the quality of a vanilla extract than, first, the quality of the bean employed, and, next, the skill of the operator. Thirdly, it may be added, a vanilla extract greatly improves by aging. "The only requirements are cologne spirits, water, sugar, good beans, and time, especially the latter two." The value of glycerin, advised by some, is doubtful.

I. Vanillaav.oz.	1
Rock candy av.oz.	$^2$
Alcohol, deodorizedfl.oz.	9
Waterfl.oz.	7

Cut the vanilla in small pieces with a sharp knife or scissors, transfer to an iron mortar and beat, with the rock candy, into a fine powder. The sugar should be added in divided portions. Place this in a bottle with the alcohol; allow to macerate, with occasional shaking, for 24 hours, then add the water and continue the maceration for 2 days, or as much ionger as is convenient. Finally express and filter.

II.

Vanilla, cut fineav.oz.	1
Alcohol, deodorizedfl.oz.	10
Waterfl.oz.	6

Mix the liquids. Put one-third of the mixture into a suitable water bath apparatus with the cut beans. Cover closely, and heat to not over 60 degs. C. for 1 hour, and remove the heat. Drain off the liquid, add another third of the liquid, repeat the process, and again with the remaining portion of the menstruum. Put the beans in a percolator and, having mixed 2 fluidounces of menstruum in the proportions given (5:3), percolate to remove the last traces of the extract from the beans. Filter the mixed liquids and pour the percolate on the filter to remove the adherent extract.

It will be an advantage to triturate the beans with rock candy, granulated sugar, or clean sand before adding to the liquid. The ignition of alcoholic vapors must carefully be guarded against.

III.

Vanillaav.oz.	1
Tonkaav.oz.	
Alcohol, deodorizedfl.oz.	32
Syrupfl.oz.	8

Cut and bruise the vanilla, afterward adding and bruising the tonka; macerate for 14 days in 16 fluidounces of the alcohol, with occasional agitation; pour off the clear liquid and set aside; pour the remaining alcohol on the magma, and heat by means of a water bath to about 77 degs. C., in a closely covered vessel. Keep it at that temperature for 2 or 3 hours, then strain through flannel with slight pressure; mix the two portions of liquid and filter through felt. Lastly add the syrup. To render this tincture perfectly clear it may be treated with pulverized magnesium carbonate, using from ½ to 1 av.ounce to each pint.

### Fruit Juices.

Carefully select the fruits, and if necessary, as happens with berries, pick out the unripe or decayed ones. Mash the fruit in a tub or barrel by means of a wooden pounder, and leave the pulp in a cool place, at a temperature of about 21 degs. C, for 12 or 24 hours, or until the liquid, when taken in a silver spoon, appears perfectly bright. This shows that the alcoholic fermentation necessary to separate the pectin and other gummy matters has taken place. Then press out the juice, add to it for each 16 fl.ozs. of liquid one fluidounce of cologne spirit, set aside for one night, and filter through paper. The filtrate is now ready to be made into syrup by the addition of the requisite proportion of sugar. If the juice is to be preserved as such, the following, known as Appert's Process, is recommended.

Collect the juice after expression, and omitting the addition of alcohol and the filtering through paper, introduce it into strong bottles-champagne are very good-taking care to leave ample space for the expansion of the liquid. The bottles being well corked and the corks secured with stout cord, are now placed in a vessel of sufficient depth. To prevent breakage, a cloth or a thin board with holes is laid under the bottles, and straw is packed loosely between them. vessel is then filled with cold water to a height sufficient to cover the bottles up to the shoulder, placed over a gentle fire, and the water slowly brought to ebullition. The boiling is kept up for about 10 minutes,

when the vessel is to be removed from the fire, and the whole allowed to cool down. Lastly, the bottles are sealed by dipping the top in melted sealing wax.

# Fruit Pulp.

To prepare fruit pulp take a quantity of thoroughly ripe fruit; rub and press it to a pulp through a hair sieve into earthen or stoneware pans; add a quarter of a pound of white granulated sugar to each pound of pulp; mix thoroughly; fill the bottle to the neck; cork and tie down with wire; place them in a boiler of cold water as above directed; put over the fire; boil gently for 20 minutes; when cold seal the corks and put the bottles in a cool place, laying them sideways.

# Ginger Ale.

Soluble essence of gingerfl.oz.	6
Citric acid av.oz.	11/
Spirit of lemonfl.dr.	$2^{'}$
Caramel av.oz.	
Syrup	

This is sufficient for a 10-gallon fountain.

### Lemonade Seltzer.

Juice of 1 lemon.
Sugar4 teaspoonfuls
Cracked icesufficient
Waterfl.oz. 1

Mix, shake, strain and fill soda glass with seltzer water. Serve with straws.

# Phosphate, Wild Cherry.

	Cherry juicefl.oz.	4
	Syrup of wild cherryfl.oz.	4
	Syrupy glucosefl.oz.	6
	Diluted phosphoric acidfl.oz.	2
	Oil of bitter almondsdrops	2
H		
	Oil of bitter almonds drops	2
	Alcoholfl.dr.	
	Diluted phosphoric acidfl.oz.	
	Simple syrupfl.oz.	8
	Syrupy glucosefl.oz.	6
	Caramel sufficient to cold	or

### Root Beer.

10
. 10
. 4
. 4
s <b>4</b> 8
s 24
12
. 10

This makes a root beer "extract" which may be mixed with syrup, or it may be diluted with 9 gallons of water containing 1 gallon of refined molasses, and charged in a fountain. If it is preferred to use a fermented article, add the water and molasses, using warm water, also 1 quart yeast, and keep in a warm place until fermentation is complete.

#### II.

Sassafrasav.oz.	4
Yellow dockav.oz.	4
Pimento av.oz.	
Wintergreenav.oz.	4
Wild cherry barkav.oz.	$^2$
Coriander seedav.oz.	
Hopsav.oz.	1

Reduce to powder and percolate with a menstruum composed of 3 volumes of alcohol and 5 volumes of water until 48 fluidounces of liquid have passed. Of this half-strength fluid extract 2 fluidounces are sufficient to make 1 gallon of root beer. Or, exhaust the above drugs with the menstruum indicated, add enough water to make 6 gallons, and start fermentation with 1 pint of yeast.

### III.

Sarsaparillaav.oz.	
Sassafrasav.oz.	21/2
Wild cherry barkav.oz.	$2\frac{1}{2}$
Wintergreen barkav.oz.	21/2

Mix with 5 gallons of lukewarm water, add 4 fluidounces of molasses and 2½ fluidounces of fresh yeast, and allow fermentation to proceed, then draw off and bottle.

IV. In a suitable vessel place 300 grains. each of pipsissewa, dandelion, sassafras, American sarsaparilla, Jamaica ginger, and hops; add 3 gallons of boiling water and keepcovered and hot, but not boiling. for 3 hours; cool partially; strain through a cloth and add 5 pounds of white or coffee sugar (or 5 pints of molasses or syrup) to the colature. When dissolved transfer to a large jar and make up to 5 gallons with water. Add one-half pint fresh brewer's yeast (or sufficient compressed yeast), stir, allow to remain in a moderately warm place, and in from 24 to 72 hours it will be fit for use. The beaten white of 1 egg or a little isinglass is often employed for clarification.

#### Soda Foam. (Gum Foam.)

By the title "soda foam," or the more improper term "gum foam," is meant a liquid to be added to syrups, so that when mixed with carbonated ("soda") water, a certain proportion of gas will be retained in the mixture in the desirable form of foam. Different substances are used in these "foams," and these vary in their gas-retaining or foam -holding qualities. Among the more common substances used in "foams" are gelatin, white of egg, and quillaja tinc-

If gelatin be used, it must be dissolved in the water used in making plain syrup. About one-half av.ounce will be sufficient for 1 gallon of syrup.

In using albumen, the white of 1 egg should be added to 16 fluidounces of water, stirring well, and straining. Or one-half of the water may be replaced by simple syrup. This mixure decomposes very quickly, and should be preserved on ice, or, better yet, it should be prepared only as required.

Quil aja may be used in the form of a tincture which may be prepared as follows:

Quillaja, fine chipsav.oz. 4
Alcoholfl.oz. 8
Watersufficient

Mix the drug with 16 fluidounces of water, boil for 15 minutes, strain, and add enough water through the strainer to make the colature measure 16 fluidounces. Mix the liquid, when cool, with the alcohol, let stand for 12 hours, filter, and to the filtrate add enough water to make it measure 24 fluidounces.

If a cheaper preparation is desired, the alcohol may be replaced by water. The product, which is just as efficient, as a "soda foam "as the preceding, may be preserved by the addition of a small amount of salicylic acid.

One fluidounce of this preparation is required as a "foam" for 1 gallon of syrup.

# Solution of Acid Phosphates.

Bone ash, powder	. av. oz. 8
Sulphuric acid, concentrated	
Water	sufficient

Mix the bone ash with 8 fluidounces of

16 fluidounces of water, mix thoroughly with a porcelain or glass stirrer, add enough water to make the whole weigh 32 av.ounces, and set the mixture aside for 24 hours, agitating occasionally. Then transfer the mixture to a strong muslin strainer, and subject this to pressure, avoiding contact with metals, so as to express as much liquid as possible. Lastly, filter the liquid through paper.

The acid used in this preparation may be the commercial variety, provided it is free from arsenic, and of a specific gravity not less than 1.83.-N. F.

	Calcium carbonate, precipitatedgr.	369
4	Magnesia, calcinedgr.	116
	Potassium carbonategr.	151
	Phosphoric acid, U. S. P., or	
	85 per cent. fl.oz.	34

Water, enough to make.....fl.oz.

Mix the acid with 8 fluidounces of water, add the calcium carbonate gradually with constant stirring. When effervescence has ceased, add the magnesia in the same way, and then the potassium carbonate. Finally add the rest of the water, stir well and filter.

#### Solution of Citric Acid. (Fruit Acid.)

1.		
Citric acid	.av.oz.	8
Water, enough to make	fl. oz.	16
Dissolve and filter.		

Citric acidav.oz.	8
Alcoholfl.oz.	2
Water, enough to makefl.oz.	

Dissolve and filter.

# Spirit of Nutmeg.

Oil of nutmeg.								.fl.dr.	4
Alcohol								.fl.oz.	91/2

# Syrup.

Simple or plain syrup for soda fountain use, or "soda syrup" as it is frequently called, is made of different strengths depending upon the peculiar ideas or notions of the pharmacists. Some use 10 av.pounds to 1 gallon of water, others again use the regular simple syrup of the pharmacopæia, but the most common formula in vogue is the following:

Sugarav.lbs.	12
Watergal.	1

Of course, only the purest granulated sugar water, add the acid previously diluted with should be used. It may be dissolved in the water by means of heat or by the process of percolation which is now so largely employed in making medicinal syrups.

If the heat process be preferred, the water and sugar should positively not be mixed before applying heat, as scorching of the sugar may occur, thus imparting to the product a certain disagreeable taste which is highly objectionable to a discriminating and delicate palate.

The percolation process should be preferred for making this preparation, as it is much more cleanly, it is constant, and requires but little supervision. Any amount may be made by having a large percolator or several percolators, which may be replenished with sugar and water as required. These percolators should be mounted in a substantial rack: a convenient receptacle for the syrup for ordinary drug store use is a clean glycerin can.

In a few instances it may be found that the density of the above syrup is too low; the U. S. P. syrup must then be used.

# Syrup, Ambrosia.

Port winefl.oz.	16
Lemon syrupfl.oz.	
Raspberry syrupfl.oz.	
Soda foamsuffici	ent

# Syrup, Birch.

Birch essence	.fl.oz.	2
Oil of sassafras	.drops	2
Syrup, enough to make	.fl.oz.	64
Soda foam	.suffici	ent

# Syrup, Catawba.

Simple syrup, U. S. P	fl.oz. 16
Catawba wine	.fl.oz. 16
Soda foam	sufficient

# Syrup, Cherry.

Cherry juicepint 1
Syruppints 7
Fruit acidfl.dr. 4
Soda foamsufficient

# Syrup, Wild Cherry.

Wild cherry barkav.oz. 1
Glycerinfl.oz. 1
Sugarav.oz. 6
Watersufficient

Reduce the wild cherry bark to No. 20 powder. Mix the glycerin with 4 fluidounces of water and moisten the powder with sufficient of the liquid, macerate for 24 ground coffee, add 32 fluidounces of water,

hours in a close vessel, then percolate and pour on water until the percolate measures 19 fluidounces, add the sugar and when dissolved strain, add half fluidounce of fruit acid and sufficient water to make 1 pint. This can be dispensed as cherry phosphate, by making an addition of solution of acid phosphate when it is drawn.

# Syrup, Chocolate.

L.		
	Cacao, powderav.oz.	2
	Waterfl.oz.	32
	Sugarav.oz.	52
	Extract of vanillafl.dr.	

Triturate the cacao in a mortar with a portion of the water to a smooth paste, add the remainder of the water, then the sugar, heat the whole in a suitable vessel with constant stirring until it nearly reaches the boiling point, then strain through a fine sieve, and when cold, add the vanilla extract.

II.	
Chocolate powderav.oz.	4
Sugarav.oz.	52
Extract of vanillafl.dr.	6
Water, boilingfl.oz.	24

Mix the chocolate and sugar, triturate the mixed powders with the boiling water added slowly and strain; when cool, add the vanilla extract.

# Syrup, Coffee.

I.	
Mocha coffeeav.oz.	2
Java coffeeav.oz.	2
Sugarav.oz.	60
Soda foam,	
Water of each, suffici	ent

The coffee should be fresh roasted, of the very best quality, and be ground to fine powder. Heat it in a vessel with 16 fluidounces of water to boiling, and boil for 1 minute, set the mixture aside for several minutes, then filter through a double filter, and add gradually hot or nearly boiling water, until the filtrate measures 32 fluidounces. In this filtrate dissolve the sugar by percolation.

I			
	Mocha coffee	.av.oz.	2
	Java coffee		
	Sugar	.av.oz.	56
	Water, enough to make		
	Soda foam		

Mix the previously roasted and finely

macerate in a suitable vessel, a wide-mouth bottle, for example, over night; then, covering the vessel loosely, place in another vessel of water, heat for 2 hours, strain, let stand about 2 hours, pour off clear liquid through muslin strainer, avoiding any of the precipitate, or the liquid may be filtered. Through the filtrate add enough water to make the filtrate measure 32 fluidounces. In the filtrate dissolve the sugar by agitation or percolation, and add the foam.

III.

Mocha coffee	.av.oz. 4
Glycerin	
Soda foam,	
Water, boiling of each,	sufficient
Sugar	.av.oz. 52

Mix the glycerin with the ground coffee, allow to stand for 1 or 2 hours, pack in a percolator, and pour on the water until 32 fluidounces of liquid are obtained. In this dissolve the sugar by percolation.

IV.

Coffee, roasted and reduced to	
fine powderav.oz.	7
Distilled water, hotfl.oz.	8
Brandyfl.oz.	2
Simple syrup, U. S. P., boiling	
hot	20
Soda foamsufficie	nt

Mix the ingredients, cover well and set aside in moderately warm, not hot, place for about 15 minutes. Then allow to stand for 24 hours at the ordinary temperature, and filter.—D.

# Syrup, Cream.

1.							
	Cream,	fresh		 	 .fl.	oz.	16
	Sodium	carbo	nate	 	 	gr.	60
	Sugar.						

Mix and dissolve by frequent stirring with a glass rod.

II.

Cream,	fres	h.								fl.oz.	16
Milk, f	resh									.fl.oz.	16
Sugar.		٠.								av.oz.	32

Dissolve by shaking. Keep in a cool place. The addition of 60 grains of sodium bicarbonate will retard souring.

# Syrup, Egg Cream.

Creamfl.oz.	16
Syrupfl.oz.	48
Extract of vanillafl.dr.	4
Yolks of 16 eggs.	

Rub cream with egg-yolk until perfec.ly smooth, then add the syrup and flavoring. This is to be served like any other soda syrup, but before handing over, sprinkle a little mixed spice on the foam.

# Syrup, Ginger.

Ι.				
	Tincture of	ginger	fl. oz.	2

When greater pungency is desired, 1 fluiddram of tincture of capsicum may be added. For the ordinary tincture of ginger, the soluble essence of ginger may be substituted.

II.

Soluble essence of ginger	fl. oz.	1
Tincture of capsicum		
Syrup	fl. oz.	64
Soda foam	.suffici	ent

For many people ginger is scarcely warm enough without the addition of capsicum.

# Syrup, Kola Coca.

Wine of kola	fl.oz. 4
	fl.oz. 4
	fl.oz. 48
Soda foam	sufficient

Color with caramel and cochineal solution.

# Syrup, Lemon.

1.		
Solution of citric acid	fl.oz.	1
Spirit of lemon	fl.dr.	4
Syrup		
Soda foam	suffici	ent
TT .		

II.	
Citric acidgr.	180
Spirit of lemonfl.dr.	11/2
Ŵaterfl.oz.	6
Syrup, enough to makefl.oz.	64
Soda foamsuffic	ient

Dissolve the acid in the water and add the spirit, syrup and foam.

III.

Oil of	lemondrops	12
Citric	acidgr.	300
Syrup	fl.oz.	64
Soda	foamsuffic	ient

Rub oil with acid and a little syrup, add remainder of syrup, and dissolve, and add the foam.

IV. Grate rind from 3 lemons, rub with 6 av. ounces granulated sugar, add 8 fluidounces of water, macerate a short time, stir fre-

SODA WATER
quently, strain, express lemons, mix juice with other liquid, add one-half gallon of simple syrup, U. S. P., and finally sufficient soda foam.
Syrup, Maple.
Maple sugar
Syrup, Malto.
Extract of malt, thickfl.oz. 4 Solution of acid phosphatefl.oz. 4 Syrup, enough to makefl.oz. 64
Syrup, Mead.
I.
Pineapple syrup
II.
Sarsaparilla rootav.oz.1½Licorice rootav.oz.2Marshmallow rootav.oz.1Gum arabicav.oz.2
Make a decoction with water, strain to 6
pints, add:
Sugarav.lbs. 10
When cold, add:  Oil of lemon
III.
Essence of mead
Syrup, Moxie.
I.
Oil of sassafras drops 12

Oil of wintergreen......drops 12

Alcohol ......fl.dr. 6

Fluid extract of gentian .....fl.dr. 6

Compound syrup of sarsaparilla.fl.oz. 6

Caramel fl.oz. 1 Syrup, enough to make fl.oz. 64

REPARATIONS.	308
II.  Compound tincture of gentian. Sarsaparilla essence Syrup Syrupy glucose Caramelsufficient	.fl.dr. 4 .fl.oz. 32 .fl.oz. 32
Syrup, Nectar.	
I.	
Raspberry syrup Pineapple syrup Vanilla syrup Syrup Soda foam	fl.oz. 2 fl.oz. 1 fl.oz. 2
II.	
Spirit of nutmeg. Spirit of lemon. Extract of vanilla Water. Simple syrup, U. S. P. Soda foam.	.fl.dr. 4 .fl.dr. 4 .fl.oz. 8 .fl.oz. 56
Add coloring if thought desirab	le.
III. Strawberry syrup. Simple syrup, U. S. P. Madeira wine Spirit of bitter almonds. Citric acid. Soda foam.	.fl.oz. 4 .fl.dr. 1 av.oz. ½
Syrup, Orange.	
I.	
Oil of orange (fresh)	.fl.dr. 4 .fl.oz. 64
II.	
Oil of orange	gr. 120

Soda foam.....sufficient

Rub oil with acid and small portion of syrup, add remainder of syrup; dissolve, add the foam and strain.

Take 6 good oranges and rub the oil from the rind by means of cut, loaf sugar. Then express the juice of the oranges and add to about 4 or 5 pints of syrup. The sugar used to extract the oil should be added to the syrup and the whole heated gently to dissolve the sugar; then strain. If desired for blood orange, color with raspberry juice or tincture of cudbear. Now add syrup to make 1 gallon. In case the oranges are unusually sweet acidify with citric acid. Finally add sufficient soda foam.

370	THE STANDAR
Syrup, Orgeat.	
I. Sweet almonds:	av.oz. 8
Bitter almonds	av.oz. 2½
Sugar	av.oz. 48
Water Orange flower water.	fl.oz. 26
Orange flower water .	fl.oz. 4
Blanch the almonds,	rub them in a mortar
to fine paste with 12 a	v.ounces of the sugar
and 2 fluidounces of t	
paste with the remainde	r of the water, strain
with strong expression,	add the remainder of
the sugar, and dissolve	it with the aid of a
gentle heat. Lastly, ac	
water and strain the syr	
II:	1 0
Cream svrup	fl.oz. 8
Vanilla syrup	fl. oz. 16
Simple syrup	fl.oz. 8
Oil of bitter almonds.	drops o
Syrup, Pineapple.	
I. Concentrate	
Take 1 pineapple, cu	
spread these in layers in	a wide shallow vessel
and sprinkle sugar ov	er them, a layer of
sugar for each layer of	
hours, pour off the li	
Wash the pieces with 2	
express. To the expre	essed liquid add 4 av.
pounds of granulated	sugar, and apply a
gentle heat until disso	olved. When nearly
dissolved, add the juic	
simmer, strain, and kee	
tles.	*
II.	
Concentrated pineapp Syrup	ole syrupfl.oz. 4
Syrup	fl.oz. 32
Soda foam	sufficient
This is the diluted s	yrup for fountain use.
Syrup, Raspberry.	
	raspberries as directed
for strawberry syrup, or	
trated fruit juices of the	
Raspberry juice	
Syrup	pints 7
Mix and add	, , , , , , , , , , , , , , , , , , ,
	0.1.4
Fruit acid	n.dr. 4

Soda foam.....sufficient

Essence of sarsaparilla.....fl.dr. 4

Soda foam.....of each, sufficient

Syrup, Sarsaparilla.

Caramel,

Ī	B I CHIM CEMIN I.
	II.  Fluid extract of sarsaparillafl.oz. 1 Fluid extract of licoricefl.dr. 4 Oil of wintergreendrops 10 Oil of sassafrasdrops 6 Waterfl.oz. 8 Simple syrup, U. S. P., enough to makefl.oz. 64 III. Sarsaparilla essencefl.dr. 4
	Compound fluid extract of sar- saparilla (for syrup)fl.dr. 4 Syrupfl.oz. 64 Caramelsufficient to color
	Syrup, Sherbet.
	I.       White wine       fl.oz. 16         Lemon syrup       fl.oz. 16         Pineapple syrup       fl.oz. 32         Soda foam       sufficient         II.         Vanilla syrup       fl.oz. 48
	Pincapple syrup
	Syrup, Strawberry.
	Fresh, ripe strawberriesquarts 5 Sugarav.lbs, 12 Waterpints 1
	Spread a portion of the sugar over the ber
	ries, arranging sugar and berries in layers
	let stand for several hours, express the juice and strain, washing out the mark with water
	Add the remainder of the sugar and water

Spread a portion of the sugar over the berries, arranging sugar and berries in layers, let stand for several hours, express the juice, and strain, washing out the mark with water. Add the remainder of the sugar and water, raise to the boiling point and strain; bottle while hot and cork well. When wanted for use, mix with an equal volume of simple syrup. Add fruit acid, and soda foam sufficient.

# Syrup, Tea.

Orange Pekoe teaav.oz.	11/2
Sugarav.oz.	28
Water,	
Soda foamof each, sufficie	nt

Heat 22 fluidounces of water to boiling, remove vessel from source of heat, add the tea leaves to the water, cover the vessel, and allow leaves to infuse not to exceed one or two minutes; pour the liquid off into a filter, and if the filtrate does not measure 16 fluidounces, pour sufficient cold water on the leaves, stir about for a moment, and decant into filter until filtrate measures 1 pint; in this filtrate dissolve the sugar by agitation

or percolation, and to the solution add the foam.

# Syrup, Vanilla.

This is prepared by adding enough extract of vanilla to impart the desired flavor, coloring the mixture with caramel, and adding 1 fluidounce of soda foam to each gallon of syrup. Some use cream syrup instead of plain syrup.

# Syrup of Violets.

A so-called syrup of violets may be made by adding a little strong tincture of orris root to water, rendering clear or nearly so, by filtration through magnesinm carbonate, and dissolving in the flavored water enough sugar to make a syrup. Tincture of grass may be used as a coloring, if a green tint is desired; but green frequently suggests poison to the lay mind.

# Syrup Walnut or Hickory-Nut Cream.

Take 1 pound of walnut or hickory nut kernels and by blanching remove skin which, if left on, would impart an unpleasant bitter taste; then rub to powder in a wedgewood or porcelain mortar, adding a few drops of lemon juice to prevent separation of oil in kernels; then add water gradually so as to make a When the emulsion is thick emulsion. formed, the whole should be transferred to a cloth and be expressed; the residue should be returned to the mortar and treated as before, pulverizing, triturating again with water, and expressing, repeating this process until all of the nut passes through, occasionally adding a little more lemon juice to the residue. The result of this process, which should measure about 32 fluidounces, should be added to 1/2 gallon of cream syrup. Extract of lemon, vanilla, or other flavoring may be added and possibly some kind of coloring. This syrup is to be served like other soda water syrups.

# Tonic, Calisaya.

	,	
I.	Cinchona barkgr.	120
	Gentian rootav.oz.	3/1
	Orange peelav.oz.	3
	Cochinealgr.	60
	Caraway seedgr.	30
	Dilnted alcoholsuffici	ient
	Quinine sulphategr.	8
	Oil of rosedrop	1
	Simple syrup, enough to makegal.	1

Mix the calisaya, gentian, orange peel, cochineal and caraway, reduce to coarse powder, and extract by percolation by means of diluted alcohol so as to obtain 16 fluidounces of percolate; to this add the remaining ingredients.

In dispensing as a carbonated beverage it is best to draw '' flat,'' without foam.

### H.

Red cinchonaav.oz	z. 4
Gentianav.oz	
Orange peel · av.oa	z. $1\frac{1}{2}$
Cinnamonav.oz	z. 1
Water,	
Alcoholof each, suffice	
Simple syrup, U. S.Pfl.oz	z. 64

Mix the drugs, reduce to coarse powder, and extract by percolation so as to obtain 32 fluidounces of percolate, using a menstruum consisting of 1 volume of water and 2 of alcohol. To this percolate should be added the syrup.

# Tonic, Java.

Compound tincture of cinchona.fl.dr.	6
Coffee syrupfl.oz.	8
Vanilla syrupfl.oz.	
Syrupy glucosefl.oz.	
Syrup, enough to makefl.oz.	

# Vinegar, Raspberry.

Acetic acid	.fl.dr. 4
Raspberry syrup	.fl.oz. 8
Syrup	

# Water, Congress.

Potassium bicarbonateav.oz.	3/4
Sodium bicarbonateav.oz.	51/2
Magnesium sulphateav.oz.	
Sodium chloride (pure)av.oz.	234
Calcium chloride (anhydrous)av.oz.	31/2
Water sufficier	ıt

Dissolve the calcium chloride and magnesium sulphate each in 12 fluidounces of water, mix the solutions and after 10 or 15 minutes strain the liquid through muslin with thorough pressure.

Powder the potassium bicarbonate in a mortar, add the sodium chloride and bicarbonate; mix the whole with 16 fluidounces of water, pass the magma through a No. 50 hair sieve, following it with another 16 fluidounces of water, then with the calcium and magnesium solution first obtained, and finally with more water, until the united liquids measure 4 pints. Shake the mixture, place in a 10-gal-

lon fountain, fill the latter with water, and charge the whole in the usual way with carbonic acid gas.

Inasmuch as the mixture of magnesium sulphate and calcium chloride has for its object the formation of some magnesium chloride, the following solution may be substituted instead:

Calcium chloride (anhydrous)av.oz.	2
Magnesium chloride (anhy-	
drous) av.oz.	1
Waterfl.oz.	16

Dissolve and mix the sodium chloride and bicarbonate and potassium bicarbonate as before.

# Water, Friedrichshall.

Sodium bicarbonategr.	384
Sodium sulphate, crystalav.oz.	11/4
Potassium sulphategr.	165
Magnesium sulphateav.oz.	20
Sodium chloride (pure)av.oz.	101/4
Calcium chloride (anhydrous), av.oz.	1
Watersuffic	ient

Triturate the potassium and sodium sulphates in a mortar, add the magnesium sulphate and then 3 pints of water, and stir until dissolved; now add the sodium chloride and bicarbonate, continue the stirring for a few minutes, pour the mixture on a No. 50 hair sieve, add the calcium chloride, previously dissolved in 8 fluidounces of water, and then enough water to make the whole measure 4 pints. Put this into the usual 10-gallon fountain, fill the latter with water, and charge with carbonic acid gas to moderate pressure only.

# Water, Hunyadi Janos.

TD1 - C-11 - \*

The following makes an excellent imitation	n:
Potassium sulphategr. 6	
Calcium sulphategr. 60	
Sodium sulphateav.oz. 3	1/2
Magnesium sulphateav.oz. 4	1/2
Water, enough to makegal. 1	
Mix, dissolve and filter.	

# Water, Kissingen (Rakoczy).

,
Potassium bicarbonategr. 272
Sodium bicarbonateav.oz. 23/
Magnesium sulphateav.oz. 33/
Sodium chloride, pure
Calcium chloride (anhydrous)av.oz 234
Watersufficient
Pulverize the potassium bicarbonate in a

mortar, add the sodium bicarbonate and magnesium sulphate, and triturate the mixture with 1 pint of water, until the potassium and magnesium salts are dissolved. Pass the magma through a No. 50 hair sieve, washing what may remain on the sieve through with another pint of water.

Next rub the sodium chloride with 24 fluidounces of water until nearly dissolved and pass this liquid through the sieve.

Finally dissolve the calcium chloride in a few fluidounces of water, pass it through the sieve, and add a little more water to dissolve all the salt, using enough water to make the combined liquids measure 4 pints. Shake the whole well and place in the usual 10-gallon fountain, fill the latter with water, and charge with carbonic acid in the usual manner.

# Water, Selters (Seltzer).

Sodium bicarbonate av.oz. 3 gr. 384	4
Sodium chloride (pure) av.oz. 2 gr. 384	4
Calcium chloride (anhydrous)gr. 490	0
Magnesium sulphate av.oz. 1 gr. 165	5
Watersufficien	t

Dissolve the calcium chloride and magnesium sulphate each in 4 fluidounces of water, mix the solution, let stand for 10 or 15 minutes, and strain through muslin with pressure.

Mix the sodium cnloride and bicarbonate with a pint of water, pass the mixture through a No. 50 hair sieve, follow with the preceding liquid and then with enough water to make the liquid measure 4 pints. Shake the whole well, pour into the usual 10-gallon fountain, fill the latter with water, and charge in the usual way with carbonic acid.

The first mixture is for the purpose of forming some magnesium chloride, and hence the following solution may be used instead:

Calcium chloride (anhydrous)av.oz.	1/2
Magnesium chloride (anhydrous). av. oz.	1/2
Waterfl.oz	. 8

Add this to the sodium chloride and bicarbonate as before.

### Water, Pyrmont.

Calcium chloride (anhydrous).av.oz.	21/2
Sodium carbonateav.oz.	31/2
Sodium sulphateav.oz. 3 gr.	55
Magnesium sulphateav.oz. 1 gr.	
Ferrous sulphategr.	
Water suffic	

Dissolve the calcium chloride in 8 fluidounces of water, and the sodium sulphate and carbonate together in 1 pint of water by aid of heat; filter the latter solution, and while yet hot, add to it the calcium chloride solution. After 10 or 15 minutes, the precipitate will have contracted to a heavy mass at the The supernatant bottom of the vessel. liquid should then be decanted without losing any of the precipitate. To the latter, add the magnesium sulphate, shake thoroughly and rinse into a 10-gallon fountain nearly filled with water. Charge with carbonic acid gas to a pressure of 20 pounds, re-open the fountain, throw in the ferrous sulphate, coarsely powdered, close again, and charge to the usual pressure.

The object of charging lightly first before introducing the iron salt is to prevent oxidation of the latter subsequent to its introduction into the fountain.

# Water, Vichy (Grand Grille).

Potassium bicarbonate gr.	272
Sodium bicarbonateav.oz.	10
Sodium phosphate, crystalgr.	220
Magnesium sulphategr.	490
Sodium chloride (pure)gr.	110
Calcium chloride (anhydrous)gr.	272
Water suffic	eient

Triturate sodium phosphate with the potassium bicarbonate, add the sodium chloride. magnesium sulphate, and sodium bicarbonate, stir the mixture with 2 pints of water, pass the magma through a No. 50 hair sieve, rubbing through if necessary with the aid of a little more water.

Dissolve the calcium chloride in 4 fluidounces of water, add it to the other solution, and add enough water if necessary, to make the whole measure 4 pints. Shake the whole well together, pour into a 10 gallon fountain, fill the latter with water, and charge with carbonic acid gas in the usual way.

# PART VII.

# MISCELLANEOUS PREPARATIONS.

# Alcohol Dilution Table.

To make the below-mentioned strengths of alcohol, ordinary alcohol should be mixed with water, as follows:

85 p.c. alcohol=17 vol. of alcohol+2 of water.
80 p.c. alcohol=16 vol. of alcohol+3 of water.
75 p.c. alcohol=15 vol. of alcohol-4 of water.
70 p.c. alcohol=14 vol. of alcohol+5 of water.
65 p.c. alcohol=13 vol. of alcohol+6 of water.
60 p. c. alcohol=12 vol. of alcohol+7 of water.
55 p. c. alcohol=11 vol. of alcohol+8 of water.
50 p.c. alcohol=10 vol. of alcohol+9 of water.
45 p.c. alcohol=9 vol. of alcohol+10 of water.
40 p.c. alcohol—8 vol. of alcohol—11 of water.
35 p.c. alcohol=7 vol. of alcohol+12 of water.
30  p.c. alcohol = 6  vol. of alcohol + 13  of water.
25 p. c. alcohol=5 vol. of alcohol+14 of water.
20 p.c. alcohol=4 vol. of alcohol+15 of water.
15 p.c. alcohol=3 vol. of alcohol+16 of water.
10 p.c. alcohol=2 vol. of alcohol+17 of water.
5 p.c. alcohol=1 vol. of alcohol+18 of water.
s produced of a constrained of a constrained

# Alloys of Low Melting Point.

1. 1	New	ton's	met	al:
------	-----	-------	-----	-----

Bismuth	parts 8
Lead	
Tin	parts 3

This mixture melts at 95 degs. C.

II. Rose's metal:

Bismuth			
Lead	 	 	 part 1
Tin		 	 .part 1

This mixture liquefies at 94 degs. C.

III. Wood's metal:

Bismuthp	arts	15
Leadp		
Tinp		
Cadmiump		
Ошиниши	aits	9

This mixture melts at 68 degs. C.

# Ammonia, Domestic or Household.

I.

۰		
	Boraxgr.	120
	Oil of cinnamondrop	
	Oil of clovesdrop	1
	Oil of citronelladrop	1
	Alcoholfl.dr.	1
	Ammonia waterfl.oz.	32

Dissolve the borax in the ammonia and the oils in the alcohol, and mix the two solutions.

II.

Sodium carbonateav.oz.	20
Water of ammoniafl.oz.	48
Waterfl.oz.	

These are mixed and the clear solution is decanted after 2 or 3 days.

### Axle Greases.

T

Plumbago, very fine powder. Lard	
Mix well.	

II.

Plumbago, very fine powderav.oz. Suetav.oz.	
Mix well.	

III.

Plumbago, very fine powderav.oz.	
Petrolatumav.oz.	10
Mix well	

IV.

-		
	Caustic sodaav.oz.	4
	Waterfl.oz.	16
	Palm oilav.oz.	8
	Tallowav.oz.	8

Dissolve the soda in the water, add the fats, and heat until a homogeneous mixture is produced

V. An excellent lubricant is produced by filtering crude petroleum through animal charcoal (bone black).

VI. Heat together 10 pounds of rosin oil, and 8 pounds of lime, slaked and afterward sifted fine. Stir the mixture constantly while heating, and continue the heating until the mixture is uniform and of the consistency of syrup. The resulting mixture is called rosin soap. Take 1 pound of this and 1 pound of palm oil, melt together, then stir in 50 pounds of rosin oil, and sufficient rosin soap to make the mixture of the consistency of butter. Lastly, add 34 pound of caustic soda, heat and stir until thoroughly combined.

# Axle Grease Stains, Removal of.

See "Stains, Removal of."

# Barometer or Hygrometer Paper.

Cobalt chlorideav.oz.	4
Sodium chloride av.oz.	2
Acaciaav.oz	1
Calcium chloride gr. 175 to	350
Waterfl.oz.	12

Mix, dissolve and filter. In very dry regions, a larger amount of calcium chloride must be employed than in moister regions. Glycerin may be substituted for the calcium chloride, but the latter is to be preferred.

To prepare the paper, soak white blotting paper in this liquid and then dry.

The amount of moisture in the air is indicated by the following colors:

Rose redrain
Pale redvery moist
Bluish red moist
Lavender bluenearly dry
Bluevery dry

# Batteries, Filling for Dry.

Charcoalav.oz. 3
Mineral carbon or graphiteav.oz. 1
Manganese peroxideav.oz. 3
Calcium hydrateav.oz. 1
Arsenic (oxide)av.oz. 1
Glucose, mixed with dextrin or
starch av.oz. 1

These are intimately mixed dry, and then worked into a paste of proper consistence with a fluid composed of equal parts of a saturated solution of chloride of ammonium and chloride of sodium in water, to which are added one-tenth volume of a solution of bichloride of mercury and an equal volume of hydrochloric acid. The fluid is added gradually and the mass well worked up.

# Battery, Fluid.

I. For bichromate batteries:

Mercury bisulphategr.	120
Potassium bichromateav.oz.	21/4
Sulphuric acid, crudefl.oz.	
Waterfl.oz.	16

In the water dissolve first the mercury salt and then the bichromate; then add the sulphuric acid very carefully, stirring constantly with a glass rod. When cool the solution is ready for use. The mercury keeps the zincs well amalgamated.

Sometimes the mercury salt is omitted, and frequently sodium bichromate is substituted for the potassium bichromate.

11		
	Potassium bichromateav oz.	3
	Sulphuric acidfl.oz.	2
	Waterfl.oz.	16

### III. For Leclanche batteries:

Mix and dissolve.

Ammonium chloride......av.oz. 5 Water, enough to make.....fl.oz. 16 Mix and dissolve.—N. F.

III. For gravity batteries:

Use a saturated solution of copper sulphate in water.

# Battery, Storage.

A very satisfactory storage battery may be constructed in the following manner: After procuring two half-round porous cups and a glass jar sufficiently large to hold them both, get two pieces of sheet lead one-sixteenth of an inch thick, wide enough to fit the halfround side of the porous cups, and deep enough to come an inch above the top edge of the cups and jar. Solder a screw post to each lead plate, nearly fill the cup with a paste consisting of red lead and a solution of sodium sulphate thin enough to run like a cement, and put the lead plates in place, one of them being marked with an +. Fill the outer jar to with'n half an inch from the top with a 1:8 solution of sulphuric acid, and the battery is ready for charging. This may be done by attaching for 24 hours to a 12-cell copper sulphate battery, or to a dynamo; but always charge in the same direction. If well charged these storage cells will retain a large volume of electricity for a considerable time, and with a battery consisting of two or more cells small motors, lamps and induction coils may be operated. After the first charge a 5-cell battery suffices to recharge.

### Baume's Scale.

To convert Baume's degrees to specific gravity, the following may be employed:

For liquids lighter than water, add the degree Baume to 130 and divide the sum into 140, viz.:  $45^{\circ}$ = $140 \div (130 + 45)$ = $140 \div 175$ =0.80 sp. gr.

II.

For liquids heavier than water, subtract the degree Eaume from 145 and divide into 145, viz.:  $29^{\circ}B=145\div(145-29)=145\div116$  =1.25 sp. gr.

# Bedbug Exterminators.

The number of "cures" for bedbugs is legion. The following list embraces some of the substances employed for their destruction: Oil of turpentine, kerosene, benzin, mercuric chloride, mercury, paris green, zinc chloride, arsenic, insect powder, Scotch snuff, capsicum, naphthalin, camphor, sulphur fumes, ammonia vapor, soft soap, carbolic acid (both pure and crude), colocynth, wormwood, aloes, pepper, sodium borosalicylate, cimicifuga root; also fresh sprays of strongly-scented plants, such as ledum palustre, pennyroyal, tansy, pine, etc., placed beneath the mattress.

Bedbug exterminators may be in the powder, the liquid or the paste form. The powder may be the well known insect powder, or it may be paris green, or it may be a mixture of different insecticides. Sometimes these powders are made into a paste by moistening and are pressed into cracks containing, or suspected of containing, bedbugs or their eggs.

If the powders are used in the dry form, they may be introduced into the crevices by means of an insect powder blower or "gun."

The liquid exterminators may consist of poisonous solutions like those containing corrosive sublimate or carbolic acid, or they may consist of oil of turpentine, kerosene, benzin, oil of cedar, etc., or they may consist of tinctures of bitter substances like coloycuth or quassia, or they may consist of resinous solutions, or they may consist of soapy solutions, or again they may consist of several of these classes of substances in combination.

Substances like kerosene, benzin, volatile oils, etc., act by dissolving the chitinous coating of insects and thus obstructing the breathing pores and cause death. The resinous substances act largely by cementing over the eggs and thus prevent their hatching. The bitter substances mentioned are usually destructive to insect life.

The objections to these different substances or exterminative purposes are that resins, oil of turpentine, etc., leave stains, benzin, kerosene, etc., are inflammable, corrosive sublimate, paris green, etc., are excessively poisonous, carbolic acid has an unpleasant odor, etc. The evils of the different substances are

therefore often mitigated by combining several of them.

Liquid exterminators may be applied by means of a brush or feather, but a better method is to employ a machinist's oil-can or a bottle containing a perforated cork in which is inserted a quill.

I. One of the most commonly-used bedbug exterminators is the following:

Corrosive sublimate.....av.oz. 1 Alcohol.....fl.oz. 32

Or some of the alcohol may be replaced by water. However, inasmuch as it is the alcohol and not its corrosive consort, that is presumed to be the insecticide, this replacement is not to be recommended. Very frequently a portion of the aicohol, from about 20 to 80 per cent, is replaced by oil of turpentine; this reacts with the corrosive sublimate, precipitating the latter and being itself partially precipitated. Oil of turpentine alone is an excellent bedbug destroyer.

II.

v.

Corrosive sublimategr. Ammonium chloridegr.	
Decoction of quassia (about 1 in 20)fl.oz.	32

Mix and dissolve.—H.

11.		
Sodium chloride	.av.oz.	2
Zinc sulphate	.av.oz.	4
Water		

Mix and dissolve.-H.

IV. A safe and satisfactory method of exterminating bugs in matresses, upholstered furniture, etc., is by fumigation with sulphurous acid gas, that is, by burning sulphur in a closed room where these articles are located. The bleaching effect of the gas may be a disadvantage.

Soft or green soap	av.oz. 1
Caustic soda	gr. 60
Water	
VI.	
Soft or green soap	av.oz. 6

Dissolve the soap in the hot water, incorporate the turpentine, then the kerosene, and stir until cold.—D.

VII.	X
Naphthalin	
This mixture may be used indiscriminately	
on bedding, furniture, textiles of all descrip-	
tions, wall-paper, etc.	n
VIII. There are a number of preparations on the market which are put up in flattened	d
bottles, provided with a perforated metallic	N
top and which consist mainly or entirely of	
benzin or gasoline, flavored with some vola-	
tile oil, and colored with alkanet. These	
preparations are known by such titles as "Bug Dynamite," "Bugine," etc. Like all	
benzin or gasoline preparations, they must	О
be used with great caution to avoid explosion	2
or ignition from contact with light or fire.	
IX.	
Resinav.oz. 1	
Benzin	
Dissolve the resin in the benzin and add	3
the oil.	
X.	
Oil of amber	
Oil of cedar	
Resinav.oz. 1	F
Benzinfl.oz. 64	I
Mix and dissolve.	t
XI. Camphor	-
Paraffin wax av. oz. 2½	12
Oil of poppyfl.oz. 5	
Benzin	
The oil of poppy may be replaced by the	
cheaper cotton seed oil.	a
The paraffin acts like resin in gluing over	,
the eggs of the insect.	
XII.	
Picric acidgr. 270	
Stearic acid	
Oil of cloves	
Kerosene	:
Mix and dissolve.	
XIII.  Acetic acidfl.dr. 10	
Oil of clovesfl.dr. 3	
Oleobalsamic mixturefl.oz. 5	
Alcoholfl.oz. 24	Ι.

### XIV.

Naphthalin, crudeav.oz.	
Tobacco, cut (or Scotch snuff).av.oz.	-3
Benzin	
Oil of melissaenough to flav	

Mix the naphthalin, tobacco and benzin, macerate for 5 days, agitating occasionally, decant the clear liquid, and flavor with the oil.

# XV.

Colocynth, broken		
pieces		
Insect powder		
Benzin	 fl.oz.	32

Mix, macerate for several days, agitating occasionally, and decant the clear liquid.

### XVI.

Sodium borosalicylateav.oz.	4
Water or decoction of quassia	
(1 in 20)fl.oz.	20
Spirit of lavenderfl.oz.	10
Mix and dissolve.	

### XVII.

Savinav.oz.	1
Colocynthav.oz.	
Capsicumav.oz.	
Aloes av.oz.	
Water, hotfl.oz. 40 to	

Mix the drugs, previously reduced to coarse powder, with the water, and keep in a warm place for several hours, stirring occasionally, then allow to cool and decant the clear liquid—H.

# XVIII.

Oil of sage (volatile)	.fl.dr.	3
Lampblack	av.oz.	3/4
Alum, powder	av.oz.	31

This may be made into a paste with water and smeared into the crevices of the wood work.

# XIX.

Tobacco, powder (snuff)av.oz.	10
Insect powderav.oz.	10
Carbolic acidfl.oz.	8
Boric acid, powderav.oz.	21/2
Oil of citronellafl.dr.	
****	

### XX.

·-H.

Insect powder av.oz. 15	
Pellitory, powderav.oz. 15	
Carbolic acidfl.dr. 6	
Oil of citronellafl.dr. 6	
Diluted alcoholsufficient	

Make a thin paste, which is to be brushed into the cracks.—H.

# Benzin Jelly. (Gelatinized Benzin.)

Cocoanut oil soapav.oz.	2
Ammonia water	3
Solution of potassafl.oz.	11/2
Water, enough to makefl.oz.	12

Dissolve the soap with the aid of heat in 4 fluidounces of water, add the ammonia and potassa and the remainder of the water.

If the benzin is added in small portions, and thoroughly agitated,  $2\frac{1}{2}$  fluidounces of the above will be found sufficient to solidify 32 fluidounces of benzin.

# II.

Cocoanut oil soapav.oz.	11/2
Ammonia waterfl.oz.	
Glycerinfl.oz.	1
Ether fl.oz.	
Water, distilledfl.oz.	32

Prepare in a similar manner as the preceding, the finished solution containing only 17 grains of soap to the fluidounce.

# III.

Tincture of quillajafl.oz.	3
Benzin, enough to makefl.oz.	

Mix and shake for half an hour, then allow to stand 12 hours to solidify.

Sixteen fluidounces of benzin may also be jellified with 4 fluidounces of a 20-per cent infusion of quillaja.

### IV.

Castile soap, whiteav.oz.	21/2
Water, boilingfl.oz.	
Water of ammoniafl.dr.	
Benzin, enough to make fl.oz.	16

Dissolve the soap in the water, and when cold, add the other ingredients.

#### V.

Hard soap, whiteav.oz.	3
Water, boilingfl.oz.	5
Stronger water of ammonia fi.oz.	8
Benzin fl.oz.	

Dissolve the soap in the water, and when nearly cold add the ammonia and the benzin, and then perfume to suit.

Soaps with an excess of alkali give the best results.

# Bicycle Oil, Illuminating.

Equal parts of kerosene and lard oil.

### Bicycle Oil, Lubricating.

Equal parts kerosene and castor oil.

# Bicycle Paint (Glossy Black).

Amber	av.oz.	16
Linseed oil, boiling		
Asphaltum, Trinidad	av. oz.	3
Resin		
Oil of turpentine		

Melt the amber in the boiling oil and add the asphaltum and resin. Mix thoroughly, remove to the open air, and gradually add the turpentine oil.

Useful for metallic surfaces, such as on bicycles.

# Blackboard Slating or Paint.

In preparing these paints it is essential that the insoluble substances be reduced to very fine powder and that they be thoroughly incorporated in the mixture, and also that they be kept in a state of suspension, during the process of application, by constant agitation.

Of course, much depends upon the skill of the painter, for unless he prepares the surface of the board or wall well before putting on the paint, the latter cannot be expected to appear to the best advantage. Two coats are usually to be preferred to one, and uneven surfaces, after either coat has been applied should be rendered smooth by rubbing with sandpaper or emery cloth.

# 

Emery flour.....av.oz.

Ultramarine blue......av.oz.

Alcohol.....f.oz. 32
Dissolve the shellac in the alcohol. Place the lampblack, emery and ultramarine blue on a cheese-cloth strainer, pour on part of the shellac solution, stirring constantly, and gradually adding the solution until all of the powders have passed through the strainer.

#### III.

Shellac av.oz.	4
Lampblackav.oz.	3/4
Ultramarine blueav.oz.	
Rottenstone, powderav.oz.	2
Pumice powderav.oz.	
Alcoholfl.oz.	32

Dissolve the shellac in the alcohol, add the other ingredients, and shake well.

MISCELLA NEOUS
IV.
Ivory blackav.oz. 2
Emery flourav.oz. 1
Ultramarine blue av.oz. 1
Shellac
Alcohol
Mix well and agitate until the shellac is
dissolved.
Wood alcohol may be substituted for the
alcohol.
Blacking for Shoes.
I.
Bone blackav.oz. 6
Molasses
Sugar
Sulphuric acid, commercial fl.dr. 5
Mix together and set aside for 10 or 12
hours, giving an occasional shake. Then add,
under constant stirring, the following:
9. ,
Decoction of tan barkfl.oz. 4
Bone black
Which have previously been mixed and
allowed to stand a few hours.
II.
Rape seed oilfl.oz. 5
Simple syrup
Water
Ivory black
Sulphuric acid, commercialfl.oz. 7

Sulphuric acid, commercial
Mix the oil, syrup, and 25 fluidounces of
water, then add slowly, with constant stirring,
the acid, and finally the remainder of the
water.

# III.

Bone black	av. oz.	10
Fish oil	fl.oz.	1
Simple syrup	fl.oz.	4
Water	fl.oz.	25
Sulphuric acid, commercial.		
Muriatic acid, commercial		
Ferrous sulphate	gr.	150

Mix the bone black, oil, syrup, and 20 fluidounces of water, gradually, and with constant stirring, add the sulphuric acid and then add the muriatic acid and the ferrous sulphate, previously dissolved, in the remainder of the water.—H.

V .	
Bone blackav.oz.	10
Molassesav.oz.	6
Waterfl.oz.	5
Muriatic acid, commercial fl.dr.	7
Sulphuric acid, commercialfl.dr.	7
Oleic acidfl.oz.	
	Η.

Bone blackav.c	nz. 10
Rape oilfl.c	
Simple syrupfl.c	
Mucilage of gum arabicfl.c	oz. $1\frac{1}{2}$
Diluted acetic acidfl.c	oz. 2
Waterfl.c	
Alizarinav.c	oz. 4
	—H.
VI.	
5	- 15
Bone black, av.c	)Z. 10
Bone black	
Simple syrupav.c	oz. 9
	oz. 9 oz. 3
Simple syrupav.c	oz. 9 oz. 3 oz. 134

Mix the bone black, syrup and vinegar, stir well and add gradually, with constant agitation, the acid, set aside for 8 days, giving the mixture an occasional stir, and then add the caoutchouc previously dissolved in the oil by the aid of heat.

Rape oil.....fl.oz. 2

# Blacking, Day & Martin's.

Ivory blackav.oz.	16
Sulphuric acid, commercialfl.dr.	
Olive oilfl.oz.	1
Sugarav.oz.	16
Diluted acetic acid, enough to	
makegal.	1

# Bleaching of Linseed and Poppy Seed Oil.

Mix 1 pint of the oil in a bottle with a solution of 150 grains of potassium permanganate in 8 fluidounces of water, shake thoroughly, set aside for 24 hours in a warm place, and then add 225 grains of sodium sulphite in coarse powder. Agitate the whole thoroughly until the latter is dissolved, and incorporate 5 fluidrams of crude hydrochloric acid. Shake frequently until the brown liquid has become quite light in color, and wash the oil with water containing a small amount of chalk until the washings are no longer acid. After separating all the water, the oil may be filtered through exsiccated sodium sulphate.—D.

# Bleaching Sponges.

Soak the sponges in dilute muriatic acid over night; wash well to remove lime; dissolve I pound of hyposulphite of soda in a gallon of water, and immerse in this solution the moist sponges for several hours; then pass the sponges through a bath of dilute muriatic acid; wash in water and dry.

# Bleaching of Sponges.

See "Sponges, Bleaching of."

### Blue Prints.

See "Paper, Blue Print."

# Bluing, Liquid.

Prussian blueav.oz.	5
Oxalic acid	11/4
Waterfl.oz.	10

After solution is effected, dilute as much as desired.

Soluble blue or blue aniline may also be employed for making this preparation.

# Copper, Bluing of.

Dissolve 1 part of Schlippe's salt in 15 of water, heat to boiling in a porcelain or porcelain-lined vessel, then introduce the copper, suspending the latter so it does not touch the sides of the vessel, allow it to remain until sufficiently affected, then remove, wash and dry.—H.

# Boiler Compounds for Preventing Incrustation.

A great many substances are recommended as useful in preventing the lime of the water forming hard scales on the interior of steam boilers, and all act by preventing the agglutination of the particles. Among the best of these may be mentioned potatoes, one-fiftieth of the weight of the water being introduced glycerin, 3 pounds to every ton of coal consumed, is another useful addition. Sodium carbonate, ammonium chloride, molasses, spent tanner's bark, slippery elm bark, glucose, etc., are similarly employed. The following formulas for "boiler compounds" may also be employed:

I.

 $\mathbf{I}$ 

Catechu	av.	.lb.	2
Sal soda, crystal	av.	.lb.	2
Dextrin			
Potash, crude	av.	oz.	8
Alum	av.	oz.	8
Sugar	av.	oz.	8
Gum arabic			
I.			
1,			
m .	-		_

Turmeric	.áv.lb. 2
Sodium bicarbonate	.av.lb. 2
Dextrin	
Potash, crude	.av.oz. 8
Alum	.av.oz. 8
Molasses	av.0z 8

The foregoing amounts are for a 5-horse power boiler, and for water rich in lime. The next is for river water, 100-horse power boiler, and must be renewed whenever the boiler is emptied:

### III.

Sal soda, crystal av.lb. 1	8
Dextrinav.lb. 1	8
Alumav.lb.	6
Sugarav.lb.	6
Potash, crudeav.lb.	3

# Boiling Points of Saturated Aqueous Solutions.

Sodium acetate. 256 degs. F. 125 degs.	C
Sodium nitrate. 246 degs. F. 119 degs.	C
Potassium nitrate. 238 degs. F.115 degs.	C
Ammonium chloride	
	C
Sodium chloride. 224 degs. F. 107 degs.	
Magnesium sulphate	
	C
Alum 220 degs. F. 105 degs.	
Potassium chlorate	
	C
Copper sulphate. 216 degs. F. 102 degs.	
Iron sulphate216 degs. F. 102 degs.	
Lead acetate215 degs. F. 101 degs.	
Sodium sulphate, 213 degs, F. 100 degs.	

### Boric Acid to Powder.

This acid is found very difficult to reduce to a fine powder by ordinary manipulations, but a satisfactory and elegant powder may be made by the following process: First warm a wedgewood mortar by pouring into it a little alcohol and setting fire to it. Then put into the warm mortar the boric acid with a few drops of glycerin, when it will be found to be easily reduced to a fine powder.

# Bottle Capping Mixture.

T

1.		
	Gelatinav.oz.	1
	Gum arabicav.oz.	1
	Boric acidgr.	20
	Starchav.oz.	
	Waterfl.oz.	16

Mix the gelatin, gum and acid with 14 fluidounces of cold water, stir occasionally until the gum is dissolved, heat the mixture to boiling, remove the scum and strain. Also mix the starch intimately with the remainder of the water, and stir this mixture into the hot gelatin mixture until a uniform product results. The latter may be tinted with any suitable aniline dye.

This mixture may be used instead of sealing wax for sealing bottles. In using it must be softened by the application of heat.

—D.

II.

Shellacav.oz.	3
Venice turpentineav.oz.	
Boric acidgr.	72
Talcum, powderav.oz.	
Ether	
Alcoholfl.oz.	121/2

Dissolve the shellac, turpentine, and acid in the mixed alcohol and ether, color with a spirit-soluble aniline dye, and add the talcum.

During use, the mixture must be agitated frequently.—D.

III. Put a weighed amount of dry glue or gelatin in water, and let it stand over night. In the morning drain and press off all the surplus water, and then dissolve the swollen mass by heating in a waterbath. Add while still in the bath about one-half as much glycerin as there is liquefied gelatin, and for every av. pound of gelatin employed add 1 av. ounce of tannic acid, and stir until entirely homogeneous. If it is desired to color the material any of the mineral colors may be used. Test the liquid on a piece of glass, and if when cold, it is too hard or brittle add a little more glycerin, and if too soft more glue and tannin, preserving the proportions indicated.

# Cleaning of Greasy Bottles.

The following preparation is an efficient solvent for grease in obstinately dirty bottles:

Castile soap, in shavingsav.oz. 4
Sodium carbonateav.oz. 2
Boraxav.oz. 1
Aqua ammoniafl.oz. 7
Alcoholfl.oz. 3
Sulphuric etherfl.oz. 2
Soft water, enough to makegal. 1

The soap should be boiled in the water until it is dissolved, and the other ingredients then added.

# Brass Plating.

See "Plating with Gold, Silver," etc.

# Brass, Polish for

ı.		
	Oxalic acidav.oz.	2
	Pumice stone, powderav.oz.	4
	Water, enough to make fl.oz.	16
	95	

Dissolve the acid in the water previous wadding the pumice.

Apply, after shaking, with a rag, and polish with a second dry woolen rag.

II. Dip the article in a mixture of 2 parts of common nitric acid and 1 part of sulphuric acid, contained in a stone jar. The articles, after being dipped in this mixture, are thoroughly washed with water and then rubbed with dry sawdust. A solution of oxalic acid is used in the same way, smooth surfaces being rubbed with prepared chalk, or equal parts of the latter (or whiting) and oxalic acid, made into a paste with water, may be applied. A great many other processes are employed, such as rubbing with rotten stone and sweet oil, and then with whiting; this is particularly effective with copper articles.

III. See also "Putz Pomades" and "Polishing Powders."

# Brick Walls, Removal of Efflorescence

The usual method of treatment for removal of the white efflorescence on brick walls is by painting with dilute muriatic acid.

# Bronze Paints.

The bronze colors as furnished in the pharmacy serve for temporary purposes; that is, they are expected, in addition to drying rapidly, to be fairly permanent, but not so much importance is laid upon their resisting moisture and atmospheric influences.

Where these latter qualities are desired a copal shellac varnish is the best; though the use of any such varnish is objectionable because the fatty or resinous acids, either already present or liable to develop have a chemical action on the copper of the bronze and are apt to cause it to turn green or to deaden the luster of the bronze.

The commercial liquid bronzes consist for the most part of solutions of resins in turpentine oil and should be rejected on the grounds above indicated. Another variety is made of a mixture of gum dammar, rubber and benzin and this does not present the objectionable features above noted in so marked a degree, but has the disadvantage that owing to the very rapid evaporation of the benzin it is difficult to work with.

The following formulas avoid these objections. The liquid bronze is particularly useful for applying to wicker work, plaster figures, frames, leather, etc. With bronze powder no previous coating with varnish is necessary. The bronze paints are used most in the gold, silver and copper colors.

# I. Liquid bronze:

Bronze powderav.oz.	11
Borax shellac solutionfl.oz.	5
Alcoholfl.oz.	2

Rub the powder, adding the liquid very slowly; put in bottles holding about 1 ounce with not too narrow mouths and label with the following directions

Shake before using until the contents are thoroughly mixed. Then apply with a camel's hair pencil, shaking again each time before dipping the pencil in.—D.

# II. Weather-proof bronzing powder:

Bronze powderav.oz.	13
Dextrinav.oz.	8
Potassium bichromategr.	

Powder the bichromate very fine and mix thoroughly with the other powders.—D.

III. Bronze powder, not weather proof:
Bronze powderav.oz. 3
Dextrinav.oz. 1

Dispense in paper parcels of about 150 grains each with the following directions:

Mix the contents of this package with 2 teaspoonfuls of water, set aside until no lumps are left, and then apply with a camel's hair brush.—D.

IV. Bronze paint for cheaper work may be prepared by mixing chrome green, 2 av. pounds; ivory black, 1 av.ounce; chrome yellow, 1 av.ounce; japan, 4 av.ounces. Grind together and thin with linseed oil.

### V. Another formula is as follows:

Prepare a size consisting of benzoin, 60 grains; shellac, 1 av.ounce; alcohol, 4 fluidounces. After dissolving by means of a gentle heat, set aside in a cool place for several days and decant the clear solution. To this size the bronze powder is added in sufficient quantity. This paint is applied with a soft brush to the clean metallic surface, a second coat being given if desirable. Varnish over all. Sometimes the metal is

first grounded by painting an orange or scarlet color.

# Carriage-Top Dressing.

1.		
	Asphaltum varnishfl.oz.	32
	Linseed oil, boiledfl.oz.	1
	Oil of turpentinefl.oz.	. 6
	Benzin	

Mix the varnish with the oil, and add the turpentine and benzin.

#### II.

Burnt umberav.oz. 8	3
Asphaltumav.oz. 4	
Linseed oil, boiledgal.	1
Oil of turpentinesufficien	t

Grind the umber with a little of the oil, add the asphaltum previously dissolved in a small quantity of the linseed oil by the aid of heat; then mix all together and boil; when cool add turpentine oil until a proper consistency is attained.

The carriage top should be thoroughly sponged with hot soap suds, rinsed and dried before the dressing is applied.

#### Cements.

These are frequently entitled "glues" and possibly some of the formulas to be found under "Glues, Liquid" will serve the purpose of a cement.

# Cement, Acid Proof.

A cement which is proof against boiling acids may be made from India-rubber, tallow, lime and red lead. The India-rubber must first be melted by a gentle heat, and then 6 to 8 per cent by weight of tallow is added to the mixture while it is kept well stirred; next day slaked lime is applied, until the fluid mass assumes a consistence similar to that of soft paste; lastly, 20 per cent of red lead is added, in order to make it harden and dry.

### Cement, Aquarium.

I.	
	Lithargeav.oz. 3
	Sand, fine whiteav.oz. 3
]	Plaster of parisav.oz. 3
	Resin, fine powderav.oz. 1
	Linseed oil,
	Drier of each, sufficient

Mix the first 3 ingredients, add sufficient linseed oil to make a homogeneous paste, and then add a small quantity of drier. This should stand a few hours before using.

II.
Lithargeav.oz. 1
Sand, fine whiteav.oz. 1
Plaster of parisav.oz. 1
Manganese borategr. 20
Resin, powderav.oz. $3\frac{1}{2}$
Linseed oil varnish
sufficient to form a paste
D.
Coment for Bicycle Tires.

# Cement for Bicycle Tires.

I.

Gutta p	perchaav.oz.	1/2
Caoutch	nouc av.oz.	1
	bisulphidefl.oz.	

Mix and dissolve.

This cement is dropped into the crevices after they have been properly cleaned. If the rent is very big apply the cement in layers. Bind up the rubber tire lightly with thread, let dry for 24 to 36 hours, cut off the thread and remove the protruding cement with a sharp knife, which must previously have been dipped in water.

II. A very good cement for attaching rubber tires to bicycles can be obtained by placing pulverized shellac in 10 times its weight of strong water of ammonia. The shellac becomes softened, a viscid transparent mass resulting, which, after standing for about 3 or 4 weeks, will liquefy.

This glue is always ready for use, only in cold weather it is necessary to soften it by placing in hot water. It keeps for years without spoiling if well corked. When applied to rubber the surface of the latter becomes soft, but hardens again after the evaporation of the ammonia. This is highly recommended for fastening rubber of any kind to smooth glass or metallic surfaces.

#### III.

Shellacav.oz.	2
Gutta perchaav.oz.	2
Red leadgr.	90
Sulphurgr.	

Melt the shellac and gutta percha, and add, with constant stirring, the red lead and sulphur, melted. Use while hot.

# IV.

•	
Coautahous in fina	shreds av.oz. 1
Caoutenoue, in line	shreus av.oz. 1
Carbon digulahida	0 1
Carbon dishipinde.	fl.oz. 4

Macerate in a well-covered vessel for several days.

V.	
Caoutchouc, in fine shredsav.oz. Chloroformfl.oz. 10	
Dissolve by maceration.	
VI.	
Caoutchouc, in fine shreds av. oz. 2 Resin gr. 140 Shellac gr. 100 Carbon disulphide enough to dissolve the other ingredients	
VII. A.	
Caoutchouc, fine shredsav.oz. 1 Chloroformfl.oz. 20	
Dissolve by maceration.	Ì
В.	
Caoutchouc, fine shreds       .av.oz.       1         Resin       .gr.       180         Venice turpentine       .gr.       90	

For the solution B, the rubber is shaved into small pieces and melted with the resin; the turpentine is then added, and all is dissolved in the turpentine oil. The two solutions, A and B, are then mixed.

Oil of turpentine......fl.oz.

### VIII.

Melt together 2 parts of asphalt or pitch and 1 of gutta percha at a gentle heat. This is to be used. If possible the wheels should be warmed.

IX. See also the rubber and gutta percha

# Cement for Coating, Boiler-Covering, etc.

Litharge	av.oz	15
Boiled linseed	oilfl.oz	. 3

Triturate them in a warmed mortar until a plastic mass results.—D.

### Cement Casein.

Sodium	borate	 		.gr. 25
Water.		 	fl	.oz. 1

Dissolve and add sufficient casein to make a mass of the consistence of honey.

This is used for paper, cloth and leather—D.

See also Nos. II., III. and IV. "Cements for Porcelain, Marble," etc.

# Cement, Celluloid.

To repair broken articles in plaster, a good cement may be prepared by dissolving small pieces of celluloid in ether. Decant the liquid after a short time. The pasty residue is a cement that will dry rapidly and not dissolve in water if the articles should be exposed to it.

# Cement, Dental.

Sorel's cement for filling cavities in teeth is made by adding, rapidly, deliquescent chloride of zinc to enough oxide of zinc to make a thick paste, and applying it immediately.

Zinc phosphate cement is made by mixing zinc oxide with syrupy phosphoric acid made by boiling the 50 per cent phosphoric acid until the temperature rises to 215 degs. C.

# Cement, Diamond or Armenian.

Isinglassgr.	240
Masticgr.	120
Gum ammoniac or galbanum,	
powdergr.	60
Alcoholfl.oz.	
Waterfl.oz.	4

Soak the isinglass in the water for 24 hours, evaporate on a water bath to 2 fluid-ounces, add 2 fluidounces of alcohol, strain, add the mastic dissolved in the remaining alcohol, and add the ammonia by trituration, avoiding loss of alcohol as much as possible.

This cement must be warmed before use.

#### Cement for Gas Burners.

Litharge,

Glycerin, of each sufficient to make a stiff paste

### Cement for Glass.

Ι.

Brown glue, good quality....av.oz. 4
Acetic acid, 96 per cent.....av.oz. 6
Ammonium bichromate, powder..gr. 90

Dissolve the glue in the water by the aid of moderate heat and the ammonium bichromate. The mixture should be preserved from light.

In using, apply to the surfaces to be cemented, tie together, set aside for several days to dry thoroughly, and then expose to strong sunlight.—D.

II. A solution of potassium bichromate and glue yields a superior cement for broken glassware. The moderately strong glue or gelatin solution is mixed in a dark place or in a photographic dark room, with a small amount of concentrated solution of potassium bichromate. The edges of the fracture.

which have been thoroughly cleaned, are then coated with a thin layer of the mixture, strongly pressed together and kept close by tying with twine, or in some other manner. The glass is then exposed to the sun for some hours. This causes the cement to become insoluble even in hot water.

III. The liquid glues are suitable for mending glassware which does not come in contact with liquids. See also "Cements for Porcelain, Marble, Earthenware," etc.

IV.

Caoutchouc, finely cut	av.oz.	1/2
Caoutchouc, finely cut	.fl.oz.	10
Mastic powder	av.oz.	4

Mix and macerate until dissolved. Apply with a brush. A larger proportion of caout-chouc renders the cement elastic.

# Cement, Gutta Percha.

Gutta percha, in piecesav.oz.	2
Carbon disulphide fl.oz.	
Oil of turpentinefl.oz.	
Asphalt, powderav.oz.	

Dissolve the gutta percha in the carbon disulphide and oil, add the asphalt and let stand for several days, when it will be ready for use.

This is used for mending leather, cementing leather upon wood, etc. Before using upon leather, the latter must be freed from fat by treatment with benzin.—D.

### Cement for Iron.

1.												
	Sulphu	ır									.av.oz.	6
	White	lead									.av.oz.	6
	Borax										.av.oz.	1

Mix well and make into a paste with sulphuric acid.

Apply cement to the parts, using pressure and allowing to stand for 5 to 7 hours.

l II

White bole.		 	 	av.oz.	4
Borax		 	 	av.oz.	1
Asbestos, p	owder.	 	 	av.oz.	1

Mix and make into a paste with water.

III.

Powdered ironav.oz.	17
Sublimed sulphurav.oz.	2
Ammonium chlorideav.oz.	1

centrated solution of potassium The ingredients are rubbed with sufficient water to form a thick mass, and applied to

the parts, previously well cleansed. After 8 days the luting becomes as hard as iron, withstanding any temperature. The cement is therefore serviceable for mending distillatory apparatus.—D.

IV.

Manganese dioxide, fine powder.av.oz.	1
Clay, dryav.oz.	4
Borax, powderav.oz.	

Mix well.

In using mix enough water to form a very thick paste, fill this into the cracks, and set aside for 24 hours. This is useful for mending cracks in stoves. Upon the application of heat, the cement is melted and securely seals the crack.—D.

# Cement for Attaching Glass Labels to Bottles.

Resin.												av.oz.	4
Yellow	W	ax										av.oz.	8

# Cement or Glue, Marine.

Macerate 1 to 2 av.ounces of caoutchouc cut into small pieces in 16 fluidounces of benzol (not benzin), promoting solution by the application of heat and by agitation. To the so'ution when formed, and which will have the consistence of thick cream, add 30 av.ounces of powdered shellac, heat the mixture with constant stirring until complete fusion and combination have been effected. Pour this mixture while hot on plates of metal, so that it may cool in sheets like leather.

In using this cement, put some of it into an iron vessel, heat to 120 degs. C., and apply with a brush to the surfaces to be joined.

#### Cement for Meerschaum.

Use No. II. under "Cements for Porcelain, Marble," etc., and add 90 grains of calcined magnesia for every 4 av. ounces of casein.

"Diamond Cement" is also excellent for cementing meerschaum.

# Cement for Fastening Metal to Glass.

Resin			 	 	 	.av.oz.	5
Yellow wax .			 	 	 ٠.	.av.oz.	1
Venetian rec	1	 	 			av.oz.	1

Melt the wax and resin by aid of water bath be dampened with wand add under constant stirring the Venetian will at once adhere.

previously well dried. Stir until nearly cool so as to prevent the Venetian red from settling to the bottom.

# Cement for Mother-of-Pearl.

Isinglass, thin sheetsgr.	240
Mastic gr.	120
Ammonium chloride, powdergr.	60
Alcoholfl.oz.	31/2
Waterfl.oz.	

Steep the isinglass in the water for 1 day, and then dissolve by aid of a gentle heat, add 16 fluidounces of alcohol, pass through a cloth strainer, and to the hot solution add, with constant stirring, the mastic, previously dissolved in 12 fluidrams of alcohol, and the ammonium chloride.

The articles to be repaired should be warmed, the broken edges smeared with the cement and brought together, and so bound for 6 or 8 hours.

# Cement for Mending Pestles and Mortars.

To unite pestles with the handle, both should first be thoroughly cleansed, then melt together equal quantities of gutta percha and shellac (shellac alone is also used), fill the cavity of the pestle with the melted substance, then insert the handle, and retain the latter in proper position until cool.

Pieces of a pestle or mortar may be united in the same manner. It is advisable to warm the pieces before applying the cement.

A mixture of liharge and glycerin may also be employed for inserting loosened handles of pestles.

# Cement for Porcelain, Marble, Alabaster, etc.

Lime ... av.oz. 1

White of egg, fresh av.oz.  $2\frac{1}{2}$ Plaster of paris av.oz.  $5\frac{1}{2}$ Water ... fl.oz. 1

Reduce the lime to powder, and triturate it with the white of egg to a uniform paste. Dilute this with the water, quickly incorporate the plaster of paris, and use the cement at once.—D.

The materials to be cemented must be ready at hand. The broken surfaces should be dampened with water so that the cement will at once adhere. The pieces must be

firmly pressed together and kept in this position for about 12 hours.

 $\Pi$ 

Mix the casein in a mortar with enough of the solution to produce a uniform honey-like mass.

This cement is transparent and keeps for some time. It is not waterproof.—D.

III.

Casein, fresh.	av.oz.	5
Slaked lime	àv.oz.	1
	1	r)

IV.

Portland cementav.oz. 2
Prepared chalkav.oz. 1
Fine sandav.oz. 1
Solution of sodium silicate

..sufficient to form a semi-liquid paste Apply the cement and allow to stand for 24 hours.

# Cement for Porcelain Letters.

Mix the solution of sodium silicate and lime, add the litharge, and then enough glycerin to form a paste.

This should be used immediately.

ΙÍ.

Copal resinav.oz. 1	
White leadav.oz. 1	
Lithargeav.oz. 2	
Linseed oil, boiledfl.oz. 3	

Mix well and use at once.

### Cement or Lute for Retorts, etc.

Clay, powdered and sifted av.oz.	3
Rye flourav.oz.	
Bran av.oz. 1	L

Mix them well. When wanted, take a sufficient quantity and mix it with water to a dough to be applied to the retort or flask.—D.

# Cement, Rubber.

1.	
Carbon bisulphidefl.oz.	8
Gutta perchaav.oz.	1/2
Resingr.	
II.	
India rubbergr.	15
Chloroform fl.oz.	2

First mix the india rubber and chloroform together, and when dissolved the mastic is added in powder. It is then allowed to stand by for a week or two before using.

III. See also "Cements for Bicycle Tires."

### Cements for Rubber Shoes.

The various "rubber cements," many of the "cements for bicycle tires," may be used in mending rubber shoes.

# Cement, Sodium Silicate.

When sodium silicate solution is used as a cement it must be hardened by the gradual application of heat. If the object is heated quickly, air bubbles will form in the cement and weaken the joint. To resist the action of acids, make a paste of the solution with finely powdered glass, apply, and, after drying in a warm place for a day or more, heat if possible, to redness. The gray enameled evaporating dishes may be repaired in this manner when fine holes nave appeared in the enamel, and be made very serviceable.

A cement of great hardness and durability may be made by mixing chalk with a thick solution of sodium silicate. The chalk must be well incorporated with the liquid, and 8 or 10 hours are required for the cement to set. When this occurs, the mass is so hard as to be capable of taking a high polish. Mineral coloring materials may be added to this to match the color of the article to be mended.

# Cement, Starch.

Starch	:	 av.oz.	1
White su	gar	 av.oz.	4
Gum ara	bic	 av.oz.	1
Water		 sufficien	t

Dissolve the gum in a little hot water and the sugar and starch, and boil until the starch is cooked.

Good for repairing china, pottery, etc.

### Cement, Winchell's.

Gum arabic,			
Starch	 	 .av.oz.	1 1/4
Sugar	 	 .av.oz.	1/2

Dissolve the gum in as much water as a laundress would use for the quantity of starch indicated; with this solution thoroughly incorporate the starch and sugar, then cook this mixture in a vessel suspended in boiling water until the starch becomes clear. The

cement should be as thick as tar and should be kept so. It may be preserved by the addition of a little camphor, or oil of cloves, sassafras, or wintergreen. The addition of a small amount of sassafras will increase its effectiveness.

# Cement, Transparent.

Calcium nitrategr.	60
Gum arabic, powderav.oz.	3
Waterfl.oz.	

Dissolve the calcium salt in the water, and in this dissolve the gum arabic.

# Chemical, Garden.

This may be prepared as follows:

Place a quantity of sand in a wide-mouthed bottle (or better, a half gallon fish-globe) to the depth of 2 or 3 inches; in this layer of sand, slightly imbed a few pieces of copper sulphate, aluminium sulphate and iron sulphate; pour over the whole a solution of sodium silicate (commercial water-glass) one part and water three parts, care being taken not to disarrange the chemicals in pouring in the solution. Upon standing a week or so, a dense growth of the silicates of the various bases used will be seen in various colors. Now displace the solution of the sodium silicate with clear water, by conveying a small stream of water through a small rubber tube (such as nursing bottle tubing) into the vessel, which will gradually displace the silicate of soda solution. Care must be taken not to disarrange or break down the growth with the stream of water. When successful this produces a very beautiful scene.

# Cleansing Creams. (Electric Cleansing Compound, Lightning Renovator, or Japanese Cleansing Cream.)

Ι.		
	Chloroform fl.dr.	1
	Alcoholfl.dr.	2
	Ammonia water fl.dr.	10
	Sodium carbonategr.	120
	Castile soapav.oz.	1
	Water, distilledfl.oz.	64

Cut the soap up fine and dissolve in some of the water. Dissolve the soda in the remainder of the water; add this in another bottle to the chloroform, alcohol and ammonia. When the castile soap is entirely dissolved add all these ingredients to it. Shake well and bottle securely.

II.	
Stronger water of ammonia fl.dr.	2
Glycerinfl.oz.	1
Etherfl.oz.	
Castile soap, whiteav.oz.	1
Water, enough to make fl.oz.	

To 16 fluidounces of water add in the following order the glycerin, ammonia and ether. Shake and add enough water to measure 32 fluidounces. Then add the soap in shavings and shake until dissolved.

# III.

11.	
Castile soapav oz.	11/2
Waterfl.oz.	$64^{'}$
Ammonia waterfl.oz.	
Alcoholfl.oz.	11/2
Etherfl.oz.	11/2
Glycerinfl.oz.	1
Oil of citronella or myrbanedrops	5

Prepare this like either of the preceding.

# IV.

١	/.	
	Glycerinfl.oz.	1
	Etherfl.oz.	1
	Alcoholfl.oz.	1
	Ammonia waterfl.oz.	4
	Castile soapav.oz.	1
	Water, enough to makefl.oz.	32

Prepare like the preceding.

V.	
Castile soapav.oz.	2
Sodium carbonateav.oz.	1
Boraxav.oz.	
Ammonia waterfl.oz.	4
Alcoholfl.oz.	2
Etherfl.oz.	
Water, enough to make fl.oz.	64

Prepare like the preceding.

### VI.

Castile soapav.oz.	1
Boraxav.oz.	1
Soap linimentfl.oz.	
Alcoholfl.oz.	
Ammonia waterfl.oz.	-
Water, boilingfl.oz.	48

Dissolve the borax and soap in the hot water, and when cool add the other ingredients.

### Cleansing Liquids.

These liquids are known by such titles as "Glove Detergent," "Peerless Glove Cleaner," "French Cleaning Liquid," "Lightning Renovator," etc. They are employed for removing various stains, but particularly grease and paint stains. Hence

they usually contain such grease solvents as	II. Cohn's:
benzin, chloroform, ether, etc.	Ammonium tartrategr. 150
See also "Stains, Removal of."	Ammonium acetategr. 150
I.	Potassium phosphategr. 1½
Benzingal. 1	Magnesium sulphategr. 1 Calcium chloridegr. 1
Chloroformfl.oz, 4	Distilled water
Ether	Dissolve and filter.—D.
II.	
Oil of turpentinefl.oz. 5	III. Miquel's:
Water of ammoniafl.oz. 4	Peptone
Wood alcoholfl.oz, 5	Gelatingr. 15 Sodium chloridegr. 40
Ether	Potassium carbonategr. 4
Water	Distilled waterfl.oz. 16
III.	Dissolve by aid of heat and filter.
Benzin	Dentist's Amalgam Fillings.
Ether	The exact composition of these fillings
Alcoholfl.dr. 4	is kept secret, but they are supposed to
Oil of wintergreenfl.dr. 4	contain among other things mercury, gold,
IV. Tincture of quillajafl.oz. 3	platinum, etc. The following formula may
Ether	be used:
Spirit of ammoniafl.oz. 1	Tin, fine raspingsparts 31
Oil of lavender flowersfl.dr. 1½	Silver, fine raspingsparts 19
Benzinfl.oz. 26	Mercuryparts 25
Cockroach Exterminators.	Mix, heat gently until dissolved, allow to
See "Roach Exterminators."	cool, and press through chamois leather.
Copper Plating.	Another formula is the following:
See "Plating with Gold, Silver," etc.	Fletcher's platin-gold amalgam.
Copper, Polish for.	Platinumparts 1.30
See "Polish for Brass."	Gold
Counterfeit Coin Detector.	Copperparts 1.65
Silver nitrategr. 24	Tin
Nitric acid, puredrops 15	Another formula is this:
Distilled waterfl.oz. 1	Telschow's gold amalgam.
Mix and dissolve.	Goldparts 4.18
Apply a drop to the suspected coin by	Silverparts 55.00
means of a glass rod. If any other metal	Tinparts 40.00
than silver is present in larger quantities	Dentist's Arsenic Paste or Nerve
than the standard United States alloy, a	Destroyer.
black spot or stain will be produced on the	
coin.	Arsenous acidparts 2 Morphine sulphatepart 1
Croton Bug Exterminators.	Creosotesufficient to form a stiff paste
The same remedies may be employed for	Used by dentists to destroy dental nerves
the extermination of croton bugs as are em-	
ployed against roaches.	It should be used only under the direct super-
Culture Fluids. (Nutrient Fuids.)	vision of a dentist.
I. Pasteur's:	II.
Ammonium tartrategr. 30	Arsenious acidgr. 120
Potassium phosphategr. 6	Morphine sulphate (or cocaine)gr. 4
Sugargr. 600	Gycerin and water
Distilled waterfl.oz. 13½	
Dissolve and filter.	Use like the preceding.

111.		
Arsenious aci	d	gr. 60
Iodoform		gr. 60
Lysol	sufficient to	form a paste
IV.		-

Camphor, phenol and talcum, of each sufficient to make a paste.

# Dentist's Modeling Wax.

I.

Resin	 			av.oz. 1
Olive oil	 			fl.oz. 2
Hard paraffin	 			av.oz. 3
Rose pink	 	<b>.</b>	SI	ifficient

Melt the resin and paraffin and mix with the oil and coloring, stir constantly until cool. The amount of oil may be increased or decreased according to the consistence desired. Liquid petrolatum, cottonseed oil, or other oil may be substituted for the olive oil. V.

Stearinav.oz.	3
Copal resinav.oz.	3
Talcum, powderav.oz.	6
Carmine, powdergr.	15
Oil of rose geraniumdrops	90

Melt the copal by the heat of a sand bath, add the stearin, mix, remove from the fire, add the other ingredients and stir to produce a homogeneous mixture.

#### Diamond Dust. (Powdered Glass.)

Heat glass red hot, throw it into cold water, dry and powder it. This may be used for powdering the hair, and also as a polishing powder or for filtering acids, etc.

#### DISINFECTANTS.

The object of disinfection is to prevent the extension of infectious or contagious diseases by destroying the specific infectious material, known as bacteria, microbes or germs, which give rise to them. This is accomplished by the use of disinfectants.

Popularly the term disinfection is used in a much broader sense. Any chemical agent which destroys or masks bad odors, or which arrests putrefactive decomposition is spoken of as a "disinfectant." And in the absence of any infectious disease it is common to speak of "disinfecting" a foul cess-pool, or badsmelling stable, or privy vault.

This popular use of the term has led to much misapprehension, and the agents which

have been found to destroy bad odors—deodorizers—or to arrest putrefactive decomposition—antiseptics—have been confidently recommended and extensively used for the destruction of disease germs in the excreta of patients with cholera, typhoid fever, etc.

The various consequences which are likely to result from such misapprehension and misuse of the word "disinfectant" will be appreciated when it is known that many of the agents which have been found useful as deodorizers, or as antiseptics, are entirely without value for the destruction of disease germs. This is true, for example, as regards the iron sulphate or copperas, a salt which has been extensively used with the idea that it is a valuable disinfectant. As a matter of fact, iron sulphate in saturated solution does not destroy the vitality of disease germs or the infecting power of material containing This salt is, nevertheless, a very valuable antiseptic, and its low price makes it one of the most available agents for the arrest of putrefactive decomposition in privy vaults, etc.

Antiseptic agents also exercise a restraining influence upon the development of disease germs, and their use during epidemics is to be recommended, when masses of organic material in the vicinity of human habitations cannot be completely destroyed, or removed or disinfected.

While an antiseptic agent is not necessarily a disinfectant, all disinfectants are antiseptics, for putrefactive decomposition is due to the development of "germs" of the same class as that to which disease germs belong, and the agents which destroy the latter also destroy the bacteria of putrefaction, when brought in contact with them in sufficient quantity, or restrain their development when present in smaller amounts.

A large number of the proprietary "disinfectants" so-called, which are in the market, are simply deodorizers or antiseptics, of greater or less value, and are entirely untrustworthy for disinfecting purposes.

Antiseptics are to be used at all times when it is impracticable to remove filth from the vicinity of human habitations, but they are a poor substitute for cleanliness.

During the prevalence of epidemic diseases, such as yellow fever, cholera, typhoid fever, etc., it is better to use in privy-vaults, cesspools, etc., those antiseptics which are also disinfectants-i.e., germicides; and when the contents of such vessels are known to be infected this becomes imperative. Disease germs exist not only in dejecta, but also in the atmosphere; they may be attached to clothing, the germs of tuberculosis may exist in sputa, etc.

In the sick room we have disease germs at a disadvantage, for we know fairly well how to find them as well as how to destroy them. Having this knowledge, our efforts should be directed to restrict the dissemination and propagation of these germs.

The disinfectants that are of the most value depend on the immediate object to be accomplished. Experiments have shown that among the most efficient of all true disinfectants must be ranked corrosive sublimate. this is poisonous also to the higher animals, and cannot, therefore, have universal applica-For disinfecting excrementitious products, it must be considered the best agent there is, and it can be employed also in treating articles of clothing, etc., which should also be boiled before they are again used. Potassium permanganate, which is far less poisonous, is useful, especially from its deodorizing power. While these substances can be employed in the form of a spray, and thus diffused through an apartment, they should be replaced in many cases by gaseous agents, which can more readily pursue the disease germs floating in the air. Of gaseous disinfectants, choice is had between sulphurous acid, chlorine and bromide, and to this list may be added also iodine. The results of recent researches prove that, of the agents available from their cheapness as disinfectants, corrosive sublimate, permanganate of potassium, chlorine, bromire, and perhaps the chloride of zinc, are the only ones having sufficient germicidal power to be worthy of consideration.

# Disinfectants and Antiseptics.-Principles to be Regarded in use of.

1. Seek to prevent the disease germs from

long retain life. To this end, houses, and especially hospitals and pest-houses, must be thoroughly ventilated. Scatter these germs where there is plenty of light and air, and they become harmless; in damp, dark spots they retain their vitality a long time, but sunlight, thorough desiccation, and the oxidizing action of the air, will speedily destroy them. There must be no neglected places about cellars or basements where they can hide themselves and thrive and multiply. Every part of the house, and, most of all, the drains, privy vaults, etc., whose function is to aid in disposing of refuse material, must be kept scrupulously clean. In the instruction issued by the National Board of Health, these points are emphasized and reiterated: "Disinfection cannot compensate for want of cleanliness nor of ventilation." "The most available agents in combating infectious diseases are fresh air and cleanliness."

2. Endeavor to prevent the propagation of these germs by sterilizing the soil on which they fall. Accumulations of refuse matter cannot be altogether avoided, but by the free use of antiseptics, they can be kept in such a condition that spores will not readily germinate in them. It is well known that decomposing organic matter affords the most favorable possible soil for the growth of the lower forms of vegetable and animal organisms. While foul odors are not, in themselves, an evidence of the presence of contagion, they give warning that there is danger, and it is well to heed the warning.

3. Attack the germs themselves, and endeavor to lower or destroy their vitality. This is what is to be accomplished by the use of disinfectants; but the germs are organisms of a very low grade of life, and are therefore The study of the various not easy to kill. disinfectants, with especial reference to their relative value in different diseases or under different circumstances, therefore becomes an It is also essential to know important one. the best means and modes of using them.

# Disinfectants, When and Where to Use.

Disinfection of Excreta.—The infectious character of the dejecta of patients suffering finding lodgment where they can multiply or from cholera and typhoid fever is well estab-

lished, and this is true of mild cases and of the earliest stages of these diseases as well as of severe and fatal cases. It is probable that epidemic dysentery, summer complaint, diphtheria and other diseases are disseminated by means of the alvine discharges of the sick. These should therefore be thoroughly disin-In cholera, diphtheria, yellow fever and scarlet fever, all vomited material should be regarded as infectious and should be dis-As in tuberculosis, diphtheria, scarlet fever and infectious pneumonia, the sputa should be disinfected or destroyed by fire.

Disinfection of the Person.-The surface of the body of a sick person, or of his attendants, when soiled with infectious discharges, should be at once cleansed with a suitable disinfecting agent. For this purpose Labarraque's solution, diluted with five times its volume of water, may be used. After carefully washing soiled surfaces with this solution, the disinfectant itself is to be washed away with a towel wet with water or with diluted alcohol, one part to ten. The surface of the body of the dead may be disinfected by the use of the same solution, and cloths wet with this solution should be placed over orifices from which infectious material is likely to escape.

In diseases like small pox and scarlet fever, in which the infectious agent is given off from the entire surface of the body, occasional ablutions with Labarraque's solution, · diluted with 20 parts of water, will be more suitable than the stronger solution above recommended.

Disinfection of Clothing.—Boiling for half an hour will destroy the vitality of all known disease germs, and there is no better way of disinfecting clothing and bedding which can be washed, than to put it through the ordinary operations of the laundry. delay should occur, however, between the time of removing soiled clothing from the person or bed of the sick and its immersion into boiling water. If circumstances make it impracticable to do this at once, clothing should be immersed in a suitable disinfecting fluid.

Disinfection of Apartments.-In the sick-

free ventilation and cleanliness. axiom in sanitary science that it is impracticable to disinfect an occupied apartment; for the reason that disease germs are notdestroyed by the presence in the atmosphere of any known disinfectant in respirable Bad odors may be neutralized, quantity. but this does not constitute disinfection in the proper sense of the term. odors are, for the most part, an indication of want of cleanliness or of proper ventilation; and it is better to turn contaminated air out of doors than to attempt to purify it by the use of volatile chemical agents, such as carbolic acid, chlorine, etc., which are all more or less offensive to the sick and are useless as far as real disinfection is concerned.

For the complete disinfection of an apartment in which there has been a case of infectious disease, it is necessary to fill the room completely with chlorine, bromine, sulphurous acid gas, or hyponitric acid, and to keep it shut up for several hours, until every crevice is thoroughly penetrated and permeated by the disinfectant. But this thorough use of disinfectants is only admissible for the prevention of contagion after the recovery or death of the patient.

Disinfection of Privy Vaults, Cesspools, e.c.-When the excreta-not previously disinfected—of patients with cholera or typhoid fever, have been thrown into a privy vault this should be infected, and disinfection should be resorted to as soon as the fact is discovered, or whenever there is reasonable suspicion that such is the case. This may be accomplished with corrosive sublimate or with chlorinated lime. The amount used must be in proportion to the amount of material to be disinfected.

Use 1 pound of corrosive sublimate for every 500 pounds-estimated-of fecal matter contained in the vault, or 1 pound of chlorinated lime to every 30 pounds.

Disinfection of Ingesta.-It is well established that cholera and typhoid fever, are very frequently, and perhaps usually transmitted through the medium of infected water or articles of food, and especially milk. Fortunately there is a simple means at hand for disinfecting such infected fluids. This conroom no disinfectant can take the place of sists in the application of heat. The boiling

temperature maintained for half an hour kills all known disease germs. So far as the germs of cholera, yellow fever, and diphtheria are concerned, there is good reason to believe that a temperature considerably below the boiling point of water will destroy them. But in order to keep on the safe side it is best not to trust anything short of the boiling point (212 degs. F.) when the object in view is to disinfect food or drink which is open to the suspicion of containing the germs of any infectious disease.

During the prevalence of an epidemic of cholera it is well to boil all water for drinking purposes, unless it comes from a source which is beyond suspicion, and especially if it is obtained from wells, or from rivers receiving the sewage of towns, etc. After boiling, the water may be filtered, if necessary, and then cooled with ice placed around the water vessel, not put directly into the water.

Forms of Disinfectants. - Disinfectants may be used in the form of gases as in chlorine, bromine or sulphur fumigation, or they may be used as liquids to be sprinkled about the sick room or poured upon cloths suspended about the room or poured upon excreta and other infected or infectious matter, or as liquids to be sprayed about the room by means of an atomizer (so-called aromatic disinfectants are used in this manner); they may be used as powders which consist of inert powder, mix with disinfectants; or the latter mixture may be formed into cakes; the disinfectants may also be used as pastilles, to be ignited in the room. The fumigating pastilles, powders, etc., of Part V. are also used with the idea of providing disinfection, but their disinfecting powers are very weak indeed. The incenses (see "Incense") are also used for a similar purpose and are similarly inefficacious.

Ι.

Aluminium chlorideav.oz.	6
Zinc chlorideav.oz.	
Sodium chlorideav.oz.	
Calcium chlorideav.oz.	3
Water, enough to makefl.oz.	32

Dissolve the aluminium and calcium salts separately, then mix and allow to settle. In the clear supernatant liquid dissolve the other ingredients.

This mixture may be aromatized by the addition of oils of eucalyptus and wintergreen, or oil of rosemary and thymol may be added.

TI.

A.1		10
Alum		
Sodium carbonate	av.oz.	10
Ammonium chloride	.av.oz.	2
Sodium chloride	av.oz.	2
Zinc chloride	av.oz.	1
Muriatic acid, commercial,		
Water of each.	sufficie	en٠

Dissolve the alum in ½ gallon of boiling water, then add the soda which precipitates the aluminium hydrate. Muriatic acid should then be added in sufficient quantity to dissolve the precipitate. The other salts should then be dissolved in 3 pints of water; this should be added to the first solution, and then enough water added to make 1 gallon.

III.

Zinc sulphateav.oz.	7
Ferrous sulphateav.oz.	
Naphtholgr.	
Oil of thymedrops	12
Hypophosphorous acid, diluted.fl.dr.	
Waterfl.oz.	32

Dissolve the zinc and iron sulphates in 32 fluidounces of boiling water, add the naphthol and oil and shake the mixture occasionally in a stoppered bottle until cold. Then add the acid and filter.—N. F.

- IV. Where iron is objectionable, as in the treatment of woven fabrics, the iron sulphate of the preceding formula may be replaced by aluminium sulphate (commercial will do), in which case the acid may be omitted.
- V. This preparation is known as Burnett's disinfecting liquid, and also as Crew's disinfectant.

Zinc, granulated, or scrapsav.oz. 7
Muriatic acidfl.oz. 6 or sufficient
Waterfl.oz. 32
Dissolve avoiding excess of acid.

VI. The following is known as Ledoyen's disinfectant:

A.

Litharge						 .av.oz	. 41/
Nitric acid						fl.dr	. 23
Water							

Dissolve the litharge in the acid and water previously mixed.

В.	XIII. Tar fumigation.
Lead nitrate	Pine tar
	Mix and let simmer in an open vessel
VII.       Ferrous sulphate	standing on a flame or a hot stove.  This is mentioned here, not because of its value, but for the reason that there may still be some demand for it.
Dissolve the iron sulphate in 24 fluidounces	XIV.
of water, and the corrosive sublimate in the alcohol, mix the two solutions, add the ammonium chloride, and then enough water to make 32 fluidounces.	Carbolic acid
In using as a disinfectant, dilute with an	or else dilute with water, moisten cloths with
equal volume of water.	the mixture and suspend in the room.
VIII. The following was recommended by	XV.
the Illinois Board of Health some years ago:	Mentholgr. 30
Sulphate of iron	Oil of lavender.         fl.dr. 1           Oil of lemon         fl.dr. 1           Oil of eucalyptus         fl.dr. ½           Tincture of benzoin         fl.dr. 6           Alcohol         fl.oz. 16
Part I.	Use like the preceding.
X. Chlorine fumigation: Table salt	XVI.  Oil of eucalyptus
lime and dilute muriatic acid may be used.  XI. Nitric or nitrous fumigation:	Crude carbolic acid
Place 4 av. ounces of powdered potassium	Reduce the mixture to fine powder.
nitrate in a saucer and add, little by little, 2 ¼	XVIII.
fluidounces of crude sulphuric acid previously diluted with 2 fluidounces of water.—D.	Lime

# XII. Sulphurous fumigation: Sulphur, sublimed......av.oz. 7

Potassium nitrate.....av.oz. 4 Benzoin.....av.oz. 2 Olibanum.....av.oz. 2 Camphor.....av.oz. 1

Usually, however, simply sublimed or roll sulphur is used for sulphurous fumigation. Latterly "sulphur candles" are used, which consist of sulphur poured in a melted condition upon a coil of wicking.

Mentholgr. 36	0
Oil of lavenderfl.dr.	
Oil of lemon fl.dr.	1
Oil of eucalyptusfl.dr.	1/2
Tincture of benzoinfl.dr.	3
Alcoholfl.oz. 16	3
TT 111 .1	

Oil of eucalyptus	.fl.dr.	1
Oil of bergamot	.fl.dr.	11/2
Acetic ether	.fl.dr.	11/2
Glycerin	.fl.dr.	2
Alcohol	.fl.oz.	16

Limeav.oz.	16
Coal tarfl.oz.	4
Infusorial earth	

.... sufficient to make a powder

#### XIX.

Sulphate of ironav.oz.	17
Plaster of parisav.oz.	3
Infusorial earth or fine clayav.oz.	11/2

XX. A disinfectant tablet may be prepared as follows:

Intimately mix 4 av.ounces of powdered talc with 10 av.ounces of plaster of paris and 2 fluidounces of carbolic acid; sufficient water is then added to form a mass, which is poured into small paper capsules prepared for the purpose. The mass soon becomes hard; each tablet is wrapped in paper and tinfoil, and the whole preserved in a tin box.

For use, the wrapper is removed and the tablet is placed in a suitable place in the room, in which a pretty strong odor of phenol will be perceptible for 10 or 15 days, according to the temperature.

XXI. Tin waste or scraps, such as old or useless tin cans, tin boxes, etc., may be utilized to make a disinfectant fluid by throwing them into a wooden barrel or cask containing dilute muriatic acid; the acid gradually dissolves the tin and iron present.

XXII. The following directions are well adapted for placing on bottles of liquefied crude carbolic acid, to which 5 per cent of soft soap has been added.

MIX ONE PART OF THE SOLUTION WITH TWENTY PARTS OF WATER.

To Purify Sick Rooms.—Moisten with the diluted solution a piece of flannel cloth attached to a long rod, and wave it through the air of the apartment a few minutes. Also, sprinkle it over the floor, and put a small quantity of the solution into the close-stools and bed-pans. .

To Purify the Odor of Night Chairs.—Put a half pint of the dilute solution into the pan previous to its use, and when emptied rinse it out with a small quantity.

To Disinfect Cesspools, Drains, Water Closets.—Pour in a quantity of the solution in proportion to the capacity of the receptacle. For ordinary water closets, 1 gallon of the diluted solution will generally be effectual. For large cesspools the quantity must be increased in proportion to their contents.

To Purify Larders and Stables.—Sprinkle the floor and wash all the woodwork with the dilute solution, avoiding the use of soap or alkali.

To Sweeten Musty Casks, Tubs, etc.— Wash and rinse them well with the dilute solution.

To Extirpate Bugs and Other Vermin.— Wash the floors and all the crevices with the dilute solution. The joints, etc., of the bedsteads should be moistened by a brush, with

a solution consisting of 1 part of the fluid to 2 parts of water.

OBSERVE IN ALL CASES THE DIRECTION TO DILUTE THIS SOLUTION BEFORE USING.

# Doses, Rules for.

Wiggins' rules for doses are as follows:

- The dose of all infusions is 1 to 2 fluidounces, except digitalis, which is 2 to 4 fluiddrams.
- 2. All poisonous tinctures, 5 to 20 minims, except tincture of aconite, which is 1 to 5 minims.
- 3. All wines, from ½ to 1 fluidram, except wine of opium, which is 5 to 15 minims.
- 4. Most solid extracts can be given in doses of ½ grain; the exceptions are the extracts of poisonous drugs.
- 5. All diluted acids, from 5 to 20 minims, except hydrocyanic acid, which is from 2 to 5 minims.
- 6. All waters, from 1 to 2 fluidounces, except chloroform, creosote, cherry-laurel, bitter almond, and ammonia waters.
- 7. Medicated syrups, usually from ½ to 2 fluidrams.
  - 8. Mixtures, from 1 to 4 fluidrams.
- 9. Spirits, from ½ to 1 fluidram, except spirit of glonoin.
  - 10. Essential oils, 1 to 3 minims.

Young's rule to determine the dose of a child is to divide the age of the child by the age plus 12; the quotient represents the portion of the adult dose to be given to the child. For example: The age of the child is 6; divide 6 by 6 plus  $12 \frac{6}{18} \frac{1}{18} \frac{1}{3}$ ; a 6-year-old should receive  $\frac{1}{3}$  of the dose given to an adult.

#### Driers. (Siccative.)

Manganese borate alone may be used as a solid drier; however, its action is usually too powerful, and it is usually mixed with other substances. The following mixtures are used: Zinc oxide, 4 parts, manganese borate, part 1; and equal parts zinc oxide and manganese borate.

The manganese borate may be prepared from the residue remaining after the extraction of chlorine from a mixture of black oxide of manganese and hydrochloric acid.—D.

Cobalt borate may be employed instead of the manganese borate.

Liquid drier may be produced by heating 100 parts of linseed oil over the naked flame, stirring constantly until it weighs 85 parts.

See also "Varnish, Linseed Oil."

# Dyes for Easter Eggs.

See " Easter Egg Dyes."

#### Dyes for Fabrics.

Fabrics cannot be dyed promiscuously—one color upon another—and certain rules must necessarily be observed. Inasmuch as these should be known to the pharmacist, they are stated here. All the directions contained herein refer to the coal tar dyes, or aniline colors, from which the package dyes of the market are prepared exclusively.

- 1. White, that is, uncolored, but unbleached goods, may be dyed any color.
- 2. Yellow goods may be dyed with orange, red, green, brown, gray, or black. Dark blue, violet, or a moderately strong black dye will produce a dark olive-brown color.
- 3. Red goods may be dyed with red, violet, coffee-brown or dark brown. Black, dark blue, or dark green will give a dark brown color.
- 4. Violet goods may be dyed with violet, black, coffee-brown, dark brown, or dark green. Orange will produce a brown, and dark green a dark-bronze brown color.
- 5. Blue goods may be dyed with blue, violet, black, coffee-brown, dark brown, or dark green. Orange dye will produce a brown color.
- 6. Green goods may be dyed with green, coffee-brown, dark brown, or dark gray. Upon dyeing over with black, a dark green to black color will be produced.
- Brown goods may be dyed with brown or black. Red will produce a red-brown color, and black or dark blue a dark brown color.
- 8. Gray goods may be dyed with gray, brown, dark red and dark green. If the goods are light gray, they may be dyed with marine blue. Violet will produce a gray violet, and dark blue a more or less dark blue gray to black color.
- 9. Black goods can only be re-dyed with black.

The following directions will indicate how the various dyes are to be employed:

Free the goods from stains, wash well in warm soap water, rinse thoroughly with clear water, and place into rain or river water contained in an earthen or copper vessel large enough so that the liquid will be several inches above the goods. Now dissolve the dye in another vessel by boiling for several minutes in rain or river water; remove the goods from the first vessel; remove as much of the water as possible by expression, returning the expressed water to the vessel; add the dye solution to the water, and then return the goods to the vessel now containing the dye bath. Heat the whole to boiling, turning about continuously with a stick; allow to boil for several minutes; remove the goods from the bath, wash well in clear water; express lightly, and dry in the air.

If it is desired that the goods shall have a glossy appearance upon drying, the dry goods should be sponged on the inner surface with tragacanth water, after which they should be ironed till dry.

The following mixtures should be wrapped in good paper, the latter to be enclosed in envelopes, which should be sealed securely.

—D.

#### I. Black:

This is sufficient for 1 to 2 pounds of silk or wool. It is not suitable for cotton.

#### II. Blue, dark:

 True blue, R
 gr. 240

 Oxalic acid
 gr. 60

 Dextrin
 gr. 300

This is sufficient for 1 pound of silk or wool; it is not adapted to cotton.

#### III. Blue, imperial:

 Water blue, T B.
 gr. 110

 Oxalic acid
 gr. 28

 Dextrin
 av.oz. 134

This is sufficient for 1 pound of silk, wool cotton or linen.

#### IV. Blue, marine:

New Victoria green, II. gr. 60
Methyl violet, B. gr. 60
Dextrin gr. 180

This is sufficient for 1 pound of silk, wool or cotton.

V. Blue, sky:

· · zinc, chij ·
Water blue, IB.       gr. 70         Oxalic acid       gr. 18         Dextrin.       gr. 500
This is sufficient for 1 pound of silk, wool
or cotton.
VI. Brown, Bismarck:
Vesuvin S         gr. 115           Dextrin         av.oz.         3/4
This is sufficient for 1 pound of silk, wool
or cotton.
VII. Brown, coffee:
Vesuvin B.         gr. 240           Dextrin.         gr. 360
This is sufficient for 1 pound of silk, wool
or cotton.
VIII. Gray:
Nigrosin, W.         gr. 90           Oxalic acid .         gr. 30           Dextrin .         av.oz. 1
This is sufficient for 1 pound of silk or
wool; it is not adapted to cotton.
IX. Green:
New Victoria greengr. 75
Dextringr. 225
This is sufficient for 1 pound of silk, wool or cotton.
X. Orange:
Orange II
This is sufficient for 1 pound of silk, wool or cotton.
XI. Red, amaranth:
Diamond fuchsin I, small crystals.gr. 45
Dextrinav.oz. 1
This is sufficient for 1 pound of silk, wood or cotton.
XII. Red, cherry:
Cerise D, IVgr. 50
Dextringr. 200 This is sufficient for 1 pound of silk, woo
or cotton
XIII. Red, imperial:
Erythrosin, INgr. 90 Dextringr. 360
This is sufficient for 1 pound of silk, woo
or cotton.
XIV. Scarlet:

True ponceau, GGN.....gr. 180

Oxalic acid .....gr. 36

This is sufficient for 1 pound of silk, wool

or cotton.

XV. Violet, bluish:	
Methyl violet, 3 B Dextrin	gr. 75
4	,11

This is sufficient for 1 pound of silk, wool or cotton.

XVI. Violet, reddish:

Methyl, violet R.....gr. 90 

This is sufficient for I pound of silk, woo or cotton.

XVII. Yellow:

Naphthol, yellow S. . . . . . . . . gr. 120 Oxalic acid.....gr. 24

This is sufficient for 1 pound of silk or wool; it is not adapted to cotton.

# Dyeing Leather.

I. Black:

Treat with a solution of ferrous sulphate or iron acetate. The leather may first be mordanted with a solution of extract of logwood.

#### II. Blue:

Extract 150 grains of nutgall with 32 fluidounces of water and brush the leather with this liquid. Then brush over with a solution of 150 grains soluble blue aniline and 75 grains of glue in 32 fluidounces of water. Use each mixture three times, then dry and finish with yelk of egg.

#### III. Brown:

Apply an intimate mixture of 4 av.ounces of umber (raw or burnt), 1/2 av.ounce of lampblack, and 17 fluidounces of oxgall.

#### IV. Green:

Dissolve 1 av.ounce of alum in 1 gallon of water which furnishes the mordant.

The dye consists of 4 av.ounces of indigocarmine dissolved in 7 pints of boiling water, 2 pints of strong decoction of fustic, and 8 fluidounces of logwood.

#### V. Red:

Apply a tincture made from 1 av.ounce of cochineal and 16 fluidounces of 80-per cent alcohol.

#### VI. Yellow:

Extract 1 av.ounce of turmeric and 1/2 av. ounce of gamboge with 24 fluidounces. The leather requires mordanting with a solution of alum or potassium carbonate before applying the dye.

# Easter Egg Colors or Dyes.

The following mixtures containing coal tar colors are intended for the dyeing of easter eggs. About 75 grains may be wrapped in parchment or waxed paper and put up in a package, which will be sufficient for at least 5 eggs. If more than one color is intended for one package, smaller amounts of each dye may be put up in each individual package. The directions on a 75-grain package should be as follows:

Dissolve the dye in a porcelain or earthen vessel in 1 pint of boiling water, stirring until solution is completed. In the meantime boil 5 well-washed eggs in water for 5 minutes, then transfer them to the dye bath, and allow to remain until sufficiently colored, turning the eggs about occasionally. Then dry them with a soft cloth, without pressure, and rub with oil or fat meat until they appear glossy. The remaining dye can be used for 5 or more eggs if desired.—D.

e used for o of more eggs if desired D.
I. Blue:
Marine blue, BNgr. 60
Citric acid
Mix well and divide into 20 parts.
II. Brown:
Vesuvin, S av.oz. 1
Citric acid
Mix well and divide into 20 parts.
III. Green:
Brilliant green, Ogr. 225
Citric acid
Mix well and divide into 20 parts.
·
IV. Orange: Orange, I Igr. 150
Citric acidgr. 300
Dextrinav.oz. 2½
Mix well and divide into 20 parts.
V. Red:
Diamond, fuchsin, I, small crystalsgr. 60
Citric acidgr. 300
Dextrinav.oz. 2½
Mix well and divide into 20 parts.
VI. Rose:
Eosin, Agr. 75 Dextrinav.oz. 3
Mix well and divide into 20 parts.

VII.	Vi	ole	t:	
Meth	vl v	zio'	let.	6

 Methyl violet, 6 B.
 gr. 60

 Citric acid.
 gr. 300

 Dextrin
 av.oz. 2½

Mix well and divide into 20 parts.

# VIII. Yellow:

VIII. I CHOW.	
Naphthol, yellow, Sgr.	225
Citric acidgr.	600
Dextrinav.oz.	21/4

Mix well and divide into 20 parts.

# Eggs, Preservation of.

One pound of lime should be stirred with a gallon of water, and the eggs, perfectly fresh, immersed therein in barrels or jars. This excludes air and any germs that might cause mildew or mold, and prevents evaporation, so that the contents of the eggs are not reduced in bulk. It is important to have a considerable excess of lime to replace any that may become carbonated. The vessels containing the eggs should be kept in a cool, well-ventilated place. A very successful variation in the process consists in imbedding new-laid eggs, warm from the nest, in a thick paste of lime and water. Eggs thus preserved for six months could hardly be distinguished from those newly laid. The contents of eggs evaporate rather rapidly through the shell, and the object of the preserver must be to prevent this evaporation, and at the same time to allow for the expansion and contraction of the natural air-space in the egg due to changes of temperature. The plan of coating the shells with wax or melted paraffin fails in the latter particular. Strong brine fails because the contents of eggs preserved in it become much reduced in bulk.

Eggs for keeping should never be laid on their sides; pack with the small end down.

# Embalming Fluids.

I.			0.00
	Arsenous acid	gr.	360
	Mercuric chloride	av.oz.	1 1/4
	Alcohol	.fl.oz.	9
	Solution of carbolic acid, 5 pe	er	
	cent	floz	120
	cent	. 11.02.	1.00

From 10 to 12 pints of this are injected into the carotid artery—at first slowly and afterward at intervals of from 15 to 30 minutes. The addition of a little red aniline imparts to the corpse a life-like hue.

II. Brunelli's embalming process is as follows:

The circulatory system is cleansed by washing with cold water until it issues quite clear from the body. This may occupy from 2 to · 5 hours. Alcohol is then injected so as to extract as much water as possible. This requires about 15 minutes. Ether is then injected to remove fatty matters, this requiring from 2 to 10 hours. A strong solution of tannin is injected and allowed to be absorbed, which will require from 2 to 10 hours. The body is then dried in a current of warm air, which has passed over calcium chloride. This may occupy from 2 to 5 hours, when the operation is complete.

#### III.

Thymol gr.	15
Alcoholfl.dr.	
Glycerinfl.oz.	
Waterfl.oz.	
701 1 11 11 11 1 1 1 1 1	

Dissolve thymol in the alcohol, then add the glycerin and water.

# Etching Copper, Liquid for.

Solution of ferric chloride, U.	
S. P	13
Sodium chlorideav.oz.	43/4
Mix and dissolve.—H.	

#### Etching of Glass, Tin and Zinc.

See "Ink, Diamond," and "Ink for Writing on Metals."

# Etching Iron and Steel, Liquid for.

١.	Corrosive sublimateav.oz.	1
	Tartaric acidgr.	
	Nitric acidfl.dr.	
	Waterfl.oz.	24
	Mix and dissolve.—H.	

II.

Tincture of iodinefl.oz.	31/2
Potassium iodideav.oz.	(~
Distilled waterfl.oz. 5	5
Mix and dissolve.—H.	

#### Feathers, Bleaching of.

Feathers turned yellow are bleached according to one process by soaking them for a few hours in a warm soap bath (78 to 85 degs. C.) which should not be too strong, rinsing and exposing them, strung upon a thread, for some time to the sun, frequently moistening them meanwhile.

According to another process, the feathers, after having been treated in the warm soap year. This yields excellent results,

bath, should be rinsed off and transferred to water acidulated with sulphurous acid. Here they should be allowed to remain for 20 or 30 hours, then washed, drawn through a weak, lukewarm soap bath, and dried in the sun, or left in the sun for 1 or 2 days, being frequently moistened.

According to Dobereiner, a solution of ammonium carbonate is the best means of bleaching feathers as it effects the purpose much quicker than sulphurous acid.

#### Fertilizers.

These are intended for the fertilization of garden and flowering plants. They are often put up under such names as "Plant Food," "Lawn Fertilizers," "Chemical Food for Plants," etc.

٠.	Sugarav.oz.	2
	Ammonium sulphate av.oz.	8
	Potassium nitrateav.oz.	4

Each ingredient in moderately fine powder is mixed by passing through a sieve. It could be put up in packages to sell for 25 cents per pound. The directions for use are: One teaspoonful in a gallon of water, to be used in sprinkling upon the plant after sundown about twice a week.

II.

Calcium phosphateav.oz.	4
Potassium nitrateav.oz.	1
Potassium phosphateav.oz.	1
Magnesium sulphateav.oz.	1
Iron (ferric) phosphategr. 90 to	175
Mir well	

In using, mix 30 grains with 1 quart of water and with this sprinkle the plants.

A superior mixture is produced by using instead of the insoluble calcium phosphate, the soluble acid "superphosphate."-D.

III.

Ammonium nitrateav.oz.	4
Ammonium phosphate av.oz.	2
Potassium nitrateav.oz.	21/2
Ammonium chloride gr.	220
Calcium sulphategr.	260
Ferrous sulphategr.	175
Mix well and use like the preceding	g.—D.

#### IV. Fertilizing Mixture for Lawns:

Sodium nitrate	.av.oz.	8
Superphosphate of calcium	.av.oz.	10
Guano	.av.oz.	20
Gypsum		

This should be applied once or twice a

#### Filter Paper Toughened.

Filter paper may be toughened by immersing in concentrated nitric acid and washing with water. It is then remarkably toughened, the product being pervious to liquids, and quite different to the parchment paper made by means of sulphuric acid. It can be washed like a piece of linen. So treated it contracts in size, and the ash is diminished, the weight is slightly reduced, and it contains no nitrogen. The toughened paper can be used with the vacuum-pump in ordinary funnels, without extra support, and fits sufficiently close to prevent undue access of air, which is not the case with parchment paper. Another admirable way of preparing filters is this: Dip only the apex of the folded paper into nitric acid, and then wash with water; the weak part is thus effectually toughened:

#### Fires, Colored.

Colored fires are composed essentially of three kinds of ingredients, viz.: a combustible, an oxygen producer, and a coloring agent. The cheaper "fires" usually contain sulphur as a combustible, while the better ones contain shellac, sometimes lycopodium. Those containing sulphur are not adapted to indoor use because of the irrespirable sulphurous acid gas which is produced. It is to be noted, also, that "fires" containing a poison like an arsenic compound or calomel are also not to be employed for in-door use.

The second kind of ingredient, the oxygen producer, is usually potassium chlorate or potassium nitrate.

The coloring agent depends, of course, upon the color desired—a barium salt being used for red, a strontium salt for green, a copper salt for blue, etc.

The ingredients of "fires" should always be quite dry; in fact, it may be necessary to dry them before mixing. They should also always be in quite a fine powder. Shellac, which, as is stated, is a common ingredient of these "fires," may be reduced to powder by grinding in a mill. The ingredients now dried and powdered may be mixed by means of the hand or a wooden (not steel) spatula. Triturating or rubbing in any way likely to

cause much friction, must be avoided because of the danger of spontaneous combustion.

Fire, Blue.
I.
Sulphur
II.
Realgar       av.oz.       ½         Charcoal, wood.       av.oz.       ¾         Potassium chlorate.       av.oz.       1½         Sulphur       av.oz.       3½         Barium uitrate.       av.oz.       19
III.
Potassium nitrate
IV.
Potassium chlorate         av.oz. 6           Alum, dried         av.oz. 3           Shellac         av.oz. 2           Sulphur         av.oz. 1
v.
Potassium chlorate
VI.
Orpiment         gr. 60           Charcoal, wood         gr 60           Black antimony         av.oz. 2           Potassium nitrate         av.oz. 6           Sulphur         av.oz. 8
VII.
Shellac
VIII.
Copper sulphate
IX.
Shellac
X. —D.
Potassium chlorate

Stearin.....av.oz.

XI.	III.
Copper oxideav.oz. 1	Barium nitrateav.oz. 4
Sulphurav.oz. 2	Milk sugarav.oz. 4
Potassium chlorate	Potassium chlorateav.oz. 8
-D.	_D.
Fire, Brilliant Stars.	
	Barium nitrateav.oz. 9
Potassium nitrateav.oz. 8	Potassium chlorateav.oz. 4
Sulphurav.oz. 2	Milk sugarav.oz. 2
Black antimonyav.oz. 2	V.
Make this into a stiff paste with this solu-	Black antimonyav.oz. 1/2
tion:	Sulphurav.oz. 3
Isinglassfl.oz. 1	Potassium chlorateav.oz. 3
Diluted acetic acidfl.oz. 4	Barium nitrateav.oz. 13
Alcohol	VI.
	Potassium chlorateav.oz. 2
Form into small pieces, and while still	Barium nitrateav.oz. 10
moist roll in meal gunpowder.	Sulphurav.oz. 5
Fire Crimson.	VII.
I.	Arsenicav.oz. ½
Strontium nitrateav.oz. 17	Charcoal, wood
Sulphurav.oz. 5½	Potassium chlorateav.oz. 11/4
Charcoal, willowav.oz. 1½	Sulphurav.oz. 3
Potassium chlorateav.oz. 1	Barium nitrateav.oz. 19
II.	VIII.
Potassium chlorateav.oz. 2	Barium nitrateav.oz. 8
Strontium nitrateav.oz. 13	Potassium chlorateav.oz. 4
Charcoalav.oz, 1	Sulphurav.oz. 1
Shellacav.oz. 4	Antimony sulphideav.oz. 1/4
Fire Golden Pain	Charcoalav.oz. 1/4
Fire, Golden Rain.	Time Tiles
I. Potassium nitrateav.oz. 8	Fire, Lilac.
Gunpowderav.oz. 8	Potassium chlorateav.oz. 6
Sulphurav.oz. 5	Shellacav.oz. 3
Charcoalav.oz. 2	Chalk
Lampblackav.oz. 1	Black oxide of copperav.oz. 1
Mix and fill into paper tubes.	Fire, Orange Red.
II.	Chalkav.oz. 8½
Potassium nitrateav.oz. 8	
	Sulphurav.oz. 3½
Sulphurav.oz. 4	Sulphur
Sulphur	Sulphurav.oz. 3½
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1	Sulphur
Sulphur	Sulphur
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.         I.       Copper sulphide       .av.oz.       1         Strontium nitrate       .av.oz.       14
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.         I.       Copper sulphide       .av.oz.       1         Strontium nitrate       .av.oz.       14         Calomel       .av.oz.       14
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1         Fire, Green	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.         I.       Copper sulphide       .av.oz.       1         Strontium nitrate       .av.oz.       14         Calomel       .av.oz.       14         Potassium chlorate       .av.oz.       15
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1         Fire, Green.       I.         Potassium chlorate       av.oz. 4         Barium nitrate       av.oz. 14	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.         I.       Copper sulphide       .av.oz.       1         Strontium nitrate       .av.oz.       14         Calomel       .av.oz.       14
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1         Fire, Green.       I.         Potassium chlorate       av.oz. 4	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.         I.       Copper sulphide       .av.oz.       1         Strontium nitrate       .av.oz.       14         Calomel       .av.oz.       15         Shellac       .av.oz.       5         II.
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1         Fire, Green.         I.       Potassium chlorate       av.oz. 4         Barium nitrate       av.oz. 14         Sulphur       av.zo. 5	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.         I.       Copper sulphide       .av.oz.       1         Strontium nitrate       .av.oz.       14         Calomel       .av.oz.       14         Potassium chlorate       .av.oz.       15         Shellac       .av.oz.       5         II.       Black antimony       .av.oz.       34
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1         Fire, Green.       I.         Potassium chlorate       av.oz. 4         Barium nitrate       av.oz. 14         Sulphur       av.zo. 5         Boric acid may be substituted for the ba-	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.         I.       Copper sulphide       .av.oz.       1         Strontium nitrate       .av.oz.       14         Calomel       .av.oz.       14         Potassium chlorate       .av.oz.       15         Shellac       .av.oz.       5         II.       Black antimony       .av.oz.       34         Copper oxide       .av.oz.       234
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1         Fire, Green.       I.         Potassium chlorate       av.oz. 4         Barium nitrate       av.zo. 5         Boric acid may be substituted for the barium nitrate.	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.       I.         I.       Copper sulphide       .av.oz.       14         Calomel       .av.oz.       14         Potassium chlorate       .av.oz.       15         Shellac       .av.oz.       5         II.       Black antimony       .av.oz.       3¼         Copper oxide       .av.oz.       2¼         Sulphur       .av.oz.       6
Sulphur	Sulphur
Sulphur	Sulphur       .av.oz.       3½         Potassium chlorate       .av.oz.       13         Fire, Purple.         I.       Copper sulphide       .av.oz.       1         Strontium nitrate       .av.oz.       14         Calomel       .av.oz.       14         Potassium chlorate       .av.oz.       15         Shellac       .av.oz.       5         II.       Black antimony       .av.oz.       2½         Copper oxide       .av.oz.       2½         Sulphur       .av.oz.       6         Potassium nitrate       .av.oz.       6         Potassium chlorate       .av.oz.       12
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1         Fire, Green.         I.       Potassium chlorate       av.oz. 4         Barium nitrate       av.oz. 14         Sulphur       av.zo. 5         Boric acid may be substituted for the barium nitrate.       II.         Barium nitrate       av.oz. 12         Potassium chlorate       av.oz. 4	Sulphur
Sulphur         av.oz. 4           Gunpowder         av.oz. 4           Charcoal         av.oz. 1           Lampblack         av.oz. 1           Fire, Green.           I.         Potassium chlorate         av.oz. 4           Barium nitrate         av.oz. 14           Sulphur         av.zo. 5           Boric acid may be substituted for the barium nitrate.         su.zo. 12           Potassium chlorate         av.oz. 4           Shellac         av.oz. 4	Sulphur
Sulphur       av.oz. 4         Gunpowder       av.oz. 4         Charcoal       av.oz. 1         Lampblack       av.oz. 1         Fire, Green.         I.       Potassium chlorate       av.oz. 4         Barium nitrate       av.oz. 14         Sulphur       av.zo. 5         Boric acid may be substituted for the barium nitrate.         II.         Barium nitrate       av.oz. 12         Potassium chlorate       av.oz. 4	Sulphur       av.oz.       3½         Potassium chlorate       av.oz.       13         Fire, Purple.       I.         I.       Copper sulphide       av.oz.       14         Calomel       av.oz.       14         Potassium chlorate       av.oz.       15         Shellac       av.oz.       5         II.       Black antimony       av.oz.       3¼         Copper oxide       av.oz.       2¾         Sulphur       av.oz.       6         Potassium nitrate       av.oz.       6         Potassium chlorate       av.oz.       12         Fire, Red.

INTEGERITATION S	7 1 N 131 11 N 11 1 1 1 1 1 1 1 1 1 1 1 1
II.	II.
Strontium oxalateav.oz. 1	Alumav.oz. 3
Lycopodiumav.oz. 1	Potassium carbonate
Milk sugar av.oz. 4 Potassium nitrate av.oz. 4	Potassium chlorateav.oz. 15
Potassium chlorateav.oz. 13	III.
D.	Charcoalgr. 90
III.	Prepared chalkav.oz. 4
Strontium nitrateav.oz. 8	Sulphur
Sugar	Potassium nitrate av.oz. 6
IV.	_D.
Potassium chlorateav.oz. 4	IV.
Shellac	Strontium nitrateav.oz. 4
V.	Potassium chlorateav.oz. 9 Sulphurav.oz. 5
Lycopodiumav.oz. 1	Copper carbonateav.oz. 1
Strontium nitrateav.oz. 1	Calomelav.oz. 1
Milk sugar	V. Potassium nitrateav.oz. 12
VI.	Potassium chlorate,
Charcoal, woodav.oz. 1/2	Shellac
Black antimonyav.oz. 1	Chalk
Potassium chlorate av.oz. 1½ Sulphur av.oz. 2½	
Strontium nitrateav.oz. 11	Fire, White.
—D.	Stearic acidav.oz. 1
VII.	Barium carbonate av.oz. 1
Charcoal, wood	Milk sugar
Potassium chlorate	Potassium chlorate
Strontium nitrateav.oz. 9½	—D.
—D.	II.
VIII.	Stearic acidav.oz. 1 Sulphurav.oz. 1
Strontium nitrateav.oz. 16 Potassium chlorateav.oz. 4	Black antimonyav.oz. 3
Flaxseed, groundav.oz. 4	Potassium nitrateav.oz. 6
This is a cheap red fire for parades.	—Н.
IX.	Potassium nitrateav.oz. 10
Shellacav.oz. 31/4	Sulphurav.oz. 3
Strontium nitrateav.oz. 17½	Orpimentgr. 375
Magnesium metal, powderav.oz. 1/2	IV.
Melt the shellac, add the strontium nitrate;	Black antimonyav.oz. 1
mix well; allow to cool; powder, and add the	Sulphur av.oz. 4 Potassium nitrate
magnesium.  This makes a slow burning and very bril-	V.
liant fire.—D.	Potassium nitrateav.oz. 16
	Charcoal
Fire, Rose.	VI.
Potassium nitrate	Shellacav.oz. 3
Charcoalav.oz. ½	Barium nitrateav.oz. 18 Magnesium metal, powderav.oz. 1
Fire, Violet.	3
I.	To the shellac, melted, the barium is
Calcium carbonateav.oz. 2	added, and the mixture, when cold, powdered and the metal added.
Malachite	This makes a very brilliant and slow-burn-
Potassium chlorateav.oz. 6	ing light.

VII.	11
Potassium nitrate	II
Fire, Yellow.	
I.	
Sodium nitrate	IV
II.	
Sodium oxalate	v.
Flavor for Tobacco.	F
Cascarilla bark       .gr. 120         Valerian root       .av.oz.       ½         Tonka bean       .gr. 120	F

Alcohol, sufficient.....fl.oz. Reduce to a coarse powder, and exhaust with the alcohol, either by maceration or percolation, and lastly add the lavender and ether, and filter through paper.

Benzoin .....gr.

Stems Havana tobacco.....av.oz.

Compound spirit of lavender..fl.dr. Acetic ether.....fl.dr.

# Flash-Light Powders.

These powders are employed by photographers for taking negatives in imperfectly lighted places, or in the absence of good daylight. When blown into an alcohol flame these powders suddenly flare up, producing an intense actinic light. Finely pulverized magnesium (metal) alone may be used for this purpose; some claiming for this the best results, but it is also combined with gun cotton in the proportion of 2 parts to 1 of the latter. Other combinations of magnesium are also used. Of late, aluminium metal is displacing magnesium, it being more economical and comparatively smokeless, while magnesium produces a dense and persistent smoke.

It must be remembered that these powders are very explosive and must be prepared only in small quantities, as wanted.

Ι.		
	Magnesiumav.oz.	
	Potassium permanganateav.oz.	$^2$

-D.

	II.	
	Magnesium powderav.oz.	4
	Potassium permanganateav.oz.	
	Barium peroxideav.oz.	2
	III.	
	Magnesium powderav.oz.	3
	Antimony sulphideav.oz.	1
	Potassium chlorateav.oz.	6
	IV.	
	Aluminiumav.oz.	6
	Lycopodiumav.oz.	2
	Ammonium nitrateav.oz.	1/2
i	V.	
ı	Aluminium powderav.oz.	2
ľ	Sugarav.oz.	
	Potassium chlorateav.oz.	5
	C 0'	
	Flavor for Cigars.	
	Tincture benzoin comp fl.oz	1/

Balsam of Peru.....gr.

Castoreum .....gr.

Tonka beans.....av oz.

Alcohol.....fl.oz.

Reduce the valerian, castor and tonka to coarse powder; macerate with the menstruum of alcohol and water for two or three weeks; strain; add the tincture, balsam and magnesia; mix and filter through paper.

Carbonate of magnesia ......gr. 120

# Fire Extinguisher.

T.	
Potassium nitrate, powderav.oz.	15
Sulphur, powderav.oz.	9
Charcoal, powderav.oz.	1
Colcothar of rougeav.oz.	1/4

Dry them thoroughly, then mix them, and fill into round pasteboard boxes holding 5 pounds. Through an orifice in the side a fuse or quick-match is introduced, which extends some 4 inches inward and about 6 inches outward. The latter end is fastened on the outside of the box, and a strip of red paper pasted upon it, bearing the inscription "light here."

These extinguishers are intended for use in closed rooms, and act automatically. This acts by absorbing oxygen.-D.

# Fire Extinguishers, Liquid.

П					
	Calcium	chloride,	crude	av.oz.	4
	Sodium	chloride.		av.oz.	1
	Water			fl.oz.	15

The resulting solution is thrown into the

fire by a hand-pump. The burning portions become incrusted and cease to be combustible.—D.

#### III.

Calc um	cl	hl	O	rio	de	Э.						.av.oz.	20
Salt												.av.oz.	-5
Water												, .fl.oz.	75

IV. Fill thin spherical bottles with solution of calcium chloride, ammonium chloride, or borax.

# Fireproofing Fabrics, Wood, etc.

While fireproofing materials are used and formulas are here given, it should not be understood that the articles "fireproofed" really cannot burn; the preparation will simply hinder the rapid progress of the fire so as to permit of its easy and rapid extinction.

# I. For light fabrics:

Ammonium sulphateav.oz.	4
Ammonium carbonateav.oz.	11/4
Boraxav.oz.	
Boric acidav.oz.	11/2
Starchav.oz.	1
Waterfl.oz.	48

Dissolve the salts, which should be pure and particularly free from iron, in a sufficient quantity of the water. Add the starch, previously made into a jelly, with boiling water. Impregnate the fabric with the solution, dry it and iron it. In place of 1 avounce of starch about one-quarter the quantity of gelatin or dextrin may be used.

A quart of the solution will be sufficient for about 16 yards of material.

II. For wood and heavy fabrics, ropes, straw hats, mats, etc., the following is recommended:

Ammonium chlorideav.oz.	
Boric acidav.oz.	3
Boraxav.oz.	1
Waterfl.oz.	48

Immerse the articles for 15 or 20 minutes in the solution, heated to 100 degs. C.

Heat to 50 degs. C., and immerse the paper.

If this mixture be added to starch paste, the latter may be applied to fabrics.—D.

#### IV.

Sodium tungstateav.oz.	15
Common soapav.oz.	2
Waterfl.oz.	80

Dissolve and immerse fabrics in the warm solution.—D.

V.

Ammonium phosphateav.oz.	5
Common soapav.oz.	2
Waterfl.oz.	90

Use like the preceding.—D.

VI. For starching curtains:

Sodium tungstate	.av.oz. 2
Borax	.av.oz. 2
Starch	.av.oz. 6
Starch	

Mix and use like ordinary starch for starching.—D.

VI. For application to stage (theatrical) accessories:

Ammonium chlorideav.oz. 10	
Calcium chlorideav.oz. 33	2
Prepared chalkav.oz. 20	
Waterfl.oz. 60	

Dissolve the first two ingredients in water and add the chalk.—D.

VII. For application or coating for wooden implements, partitions, etc.:

Zinc oxideav.oz. 20	)
Waterfl.oz. 10	)
Solution of sodium silicate sufficient	

Mix the zinc oxide, water and 10 av.ounces of the solution to a smooth, mixture then add enough more of the solution to make a thin paint.

This makes a white application, a yellow one may be produced by substituting yellow other for the zinc oxide.

When these mixtures are applied they become of strong hardness and resist the elements. They will serve excellently, therefore, for all external as well as internal purposes.

—D.

#### Floor Polish or Wax.

I.

•	Yellow wax	av.oz.	8
	Potassium carbonate	av.oz.	1
(	Oil of turpentine	.fl.oz.	1
	Water		

Heat the wax and water to boiling; add the potassium carbonate; boil another minute; remove the vessel from the fire; add the oil, and stir until cold. If the floor is well pre-

served, 16 fluidounces more of water may be
added. A brown color may be produced by
adding sienna or umber, and a very dark
brown, by the further addition of lampblack.
_D

II.

Yellow waxav.oz.	8
Potassium carbonateav.oz.	4
Waterfl.oz.	52

Heat the wax in an iron vessel with 40 fluidounces of water until the wax is melted, then add the potassium carbonate dissolved in the remainder of the water and boil together until solution is effected. If it be desired to color the polish, add ½ to 1 av. ounce of annatto previously dissolved in a little alcohol.—H.

#### III.

Yellow waxav.oz.	5
Paraffin waxav.oz.	2
Stearic acidav.oz.	10
Oil of turpentine fl.oz.	6
Benzin fl.oz.	7

Melt together by a gentle heat (waxes and acid), add a sufficient quantity of burnt sienna, thoroughly triturated, with linseed oil varnish. Then add, with a constant stirring (remote from fire), the oil and benzin.—H. IV.

Yellow waxav.oz.	5
Soapav.oz.	1
Potassium carbonateav.oz.	3/4
Waterfl.oz. 68	, ,

First dissolve the soap in 10 fluidounces of water, heating gently, then add the wax, and boil. When the wax begins to come to the surface, add the potassium carbonate dissolved in 10 fluidounces of water; heat for 20 minutes more, and then add the remainder of the water.

V.

Spermacetiav.oz. 4	
Paraffin waxav.oz. 4	
Talcum, powderav.oz. 8	

Shave the spermaceti and paraffin quite fine, mix with the talcum, and pass through a No. 10 sieve.

#### VI.

Stearin, powderav.oz.	20
Yellow wax, powderav.oz.	5
Soap, powderav.oz.	2
VII.	

Stearin, powder		 		.av.oz.	16
Yellow wax, pov	vder.	 		.av.oz.	4

#### VIII.

White wax, powder	.av.oz.	10
Shellac, powder	.av.oz.	10
Resin powder	.av.oz.	1

IX. To about 4 av.pounds of paraffin, melting at about 40 degs. C., add 1 fluid-ounce of oil of mirbane, and allow to cool.

In using melt the wax, and with an old brush sprinkle the melted mixture over the floor.—D.

# Flea Exterminators. (Flea Powders and

Liquids, or Solutions.)

Castile soap, powderav.oz.	8
Starch, powderav.oz.	8
Oil of pennyroyalfl.dr.	1

TT.

.,	••		
	Naphthalin, fine powder	av.oz.	4
	Starch, powder	av.oz.	12

If desired, this mixture may be colored gray by the addition of 10 grains of lamp-black.

#### III.

Talcum, powderav.oz.	2
Zinc oxideav.oz.	
Starch, powderav.oz.	10
Oil of eucalyptusfl.dr.	21/2
Oil of rose geraniumm.	75

Apply with a powder-puff to the body and bedclothes. The powder should not be kept longer than a week.

#### IV.

•	O'1 - C -1	4
	Oil of clovesfl.dr.	
	Cologne waterfl.oz.	5
	Alcoholfl.oz.	
	Mix and filter —H	

v.

Menthol	gr.	45
Camphor		150
Oil of spearmint		
Oil of wintergreen	fl.dr.	4
Oil of bergamot	fl.dr.	4
Oil of cinnamon	fl.dr.	5
Oil of origanum		1
Oil of pennyroyal	fl.oz.	1
Oil of amber	.fl.oz.	1
Petroleum, crude		27
Mix and filter.		

VI. An ingenious plan which has been recommended to rid a house of fleas, is the following:

Place on the floor of each room several sheets of sticky fly paper with pieces of fresh meat in the center of each sheet. Jumping for the meat the fleas are caught on the paper.

#### Fly Exterminators.

Fly exterminators are of different kinds, such as papers, powders, pastilles, etc. The papers which are now so largely employed are of several varieties, viz.: sticky, poisonous or arsenical, and non-poisonous. Full directions are given below for making these preparations.

# Fly Lime.

What is known as "fly lime," which is sold largely in Europe, consists of the mixtures used in the formulas for sticky fly paper, spreading upon paper. These mixtures are spread upon plates or saucers, which are then set about the room, or are painted upon sticks, which are then hung around in convenient places.

# Fly Paper, Arsenical or Poison.

Potassium arseniate, crystal...av.oz. 1
Sugar .......av.oz. 4
Water ......fl.oz. 48

Dissolve and saturate heavy unsized paper in the solution; afterward hang it up to dry on strings. The paper should be stamped or printed as poisonous previous to immersion in the liquid. Potassium arseniate should be employed, or this prepared from arsenic acid, instead of the arsenite—the form officinal in Fowler's Solution. While arsenic acid is more readily partaken of by flies, it has the further advantage of being non-poisonous to the hands.—D.

II.

White arsenicgr.	120
Potassium carbonateav.oz.	1
Sugarav.oz.	4
Waterfl.oz.	32

Mix; dissolve, and saturate paper as in the preceding.

# Fly Paper, Bichromate.

Potassium bichromate	av.oz.	1
Sugar	fl.dr.	3
Oil of black pepper	fl.dr.	11/2
Alcohol	fl.oz.	2
Water	fl. oz.	12

Mix thoroughly; macerate for several days, and filter off the liquid. In this solution soak unsized absorbent paper, allow to dry, and again soak and allow to dry.

# Fly Paper, Cobalt.

Tartar emeticgr.	45
Cobalt chlorideav.oz. Quassia chipsav.oz.	7
Tincture of long pepper (1 in 4).fl.oz.	
Waterfl.oz.	

Mix, macerate for 7 days, agitating occasionally, and filter. Prepare the paper like the preceding.

# Fly Paper, Non-Poisonous.

As a so-called non-poisonous fly paper may be used either the bichromate, cobalt or quassia fly papers.

# Fly Paper, Quassia.

Quassia	chips	 	 	 	.av.oz. 8
Sugar		 	 	 	.av.oz. 4
Water		 	 	 	sufficient

Pour 32 fluidounces of water over the quassia chips, allow to stand over night, strain and boil the liquid down to 16 fluidounces; then boil the chips with 16 fluidounces of water until 8 fluidounces remain. Mix well together, allow to stand for several days; filter, soak strips of absorbent paper in the filtered liquid and dry the slips.

# Fly Paper, Sticky.

I.		
	Resinav.oz.	6
	Lard oilfl.oz.	2
	Turpentine, Canadaav.oz.	1

Melt the resin upon a water bath, add the other ingredients and spread upon paper. The paper should be ordinary printing paper, which has previously been "sized," by applying a coating of a thin solution of white glue by means of a sponge, and hung up to dry. The sticky compound is put on whilst warm by means of a brush, and the paper is then folded together. The proportion of resin must be varied to suit the changes in the temperature.

II.
Resin, in clean piecesav.oz. 8
Castor oilav.oz. 4
Prepare like the preceding.

#### III.

Resinav.oz.	8
Venice turpentineav.oz.	2
Castor oilav.oz.	2

Prepare like the preceding.

IV.

Resin av.oz. 4	
Castor oilav.oz. 2	,
Syrupy glucoseav.oz. 1	

Melt the resin add the castor oil, incorporate the glucose, and spread the mixture upon heavy paraffined paper or upon sized paper.

V.

Resinav.oz.	6
Yellow waxgr.	90
Linseed oilfl.oz.	4

Melt together and strain. It may be made of a handsome color by adding 45 grains of red saunders.—D.

The wax improves the consistence while its odor, being suggestive of honey, is very attractive to flies. It may be increased, if desired, to 180 grains.

# Fly Pastilles.

Potassium nitrate, powderav.oz.	1 1/2
Mucilage of tragacanthfl.oz.	2
Insect powderav.oz.	2
Althæa, powdergr.	125
Tragacanth, powdergr.	125

Intimately mix the potassium nitrate with the mucilage; also mix the other ingredients together, then incorporate the powdery mixture with the paste, divide the whole into pastilles weighing about 30 grains, and dry at a temperature of 20 to 25 degs. C. The pastilles may be bronzed or gilded, if desired.—D.

# Fly Pencils.

Eucalyptol(or oil of eucalyptus).fl.dr.	11/2
Oil of laurel, essentialdrops	20
Petrolatumav.oz.	234
Paraffin waxav.oz.	2

To the paraffin previously melted the oils should be added and the mixture then molded into sticks.—D.

This is used for penciling the body exposed to the attacks of flies.

#### Fly Powders.

I.

Long pepperav.oz. 4	
Quassiaav.oz. 4	
Sugarav.oz. 8	
Diluted alcoholfl.oz. 4	

The solids should be in very fine powder, should be well mixed, and then mixed with

the diluted alcohol, dried, and reduced to fine powder.

This powder is employed by sprinkling upon a saucer.—D.

II

Crude	arsenic (so	called	"co-	
balt	")		, av, oz.	12
Sugar.			av.oz.	4

Each should be in fine powder and should be well mixed.

III.

Eucalyptol (or oil of eucalyptus). fl. oz.	1
Orris rootav.oz.	4
Starchav.oz.	15

Reduce the drugs to fine powder and mix with the oil.—D.

# Fountains, Charging of.

This table indicates the amounts of sodium bicarbonate and sulphuric acid to be used in charging soda fountains.

To produce a pressure of 8 atmospheres, equaling about 120 pounds to the square inch:

Quantity of water.	of sodium bicar- bonate.	of sulphuric acid.
10 gallons 20 " 30 "	86 av.oz. 123 " 161 "	50 av.oz.
40 ··· 50 ···	198 " 236 "	118 " 137½ "

For 9 atmospheres—about 135 lbs. to the square inch.

10 gallons	96 av.oz.	55½ av.oz.
20 "	134 "	73 "
30 "	171 "	100 "
40 ''	209 "	122 "
50 "	246 "	144 "

The amounts of sulphuric acid are somewhat in excess of the quantity required to decompose the soda.

# Freezing Mixtures.

In using freezing mixtures, the salts should be in very fine powder and should be perfectly dry. The vessel should previously be cooled, and cool water should be employed. If the mixture be produced from one salt and water, the former may be recovered by evaporation, and after thorough drying and powdering, it may be used over again. A mixture of salts may be recovered in the same manner and used over again.—D.

	_
I.	1
Ammonium chlorideav.oz. 3 Potassium nitrateav.oz. 1	l
Potassium chlorideav.oz. 6	
Dry each substance and powder, and then	
to the mixtures add 10 fluidounces of cool	]
water.	
The temperature depression amounts to 30	1
degs. C.—D.	1
II. Ammonium chloride av.oz. 3	
Potassium nitrateav.oz. 3	0
Sodium sulphate, in small crystals	1
Dry the ammonium and potassium nitrate,	0
reduce to powder, add the sodium sulphate,	6
and then mix with 11 fluidounces of cold	1
water.	ı
The temperature depression amounts to	l'i
25 degs. C.—D. and H.	
III.	
Ammonium nitrate, powderav.oz. 10 Cold waterfl.oz. 10	
The temperature depression amounts to 30	1
degs. C.—D.	f
IV.	6
Sulphuric acid	:
Sodium sulphate, powderfl.oz. 5	
Mix the acid and water, allow the mixture	
to cool to the temperature of the atmosphere,	
and add the sodium sulphate.	1
V.	1
Sodium sulphate	1
VI.	1
Ammonium nitrateav.oz. 4	1
Sodium carbonateav.oz. 4	1
Waterfl.oz. 4	1
VII. Ammonium chlorideav.oz. 2	
Potassium nitrateav.oz. 2	
Waterfl.oz, 6	
Depresses temperature from 10 degs. C. to	
112 degs. C.—H.	
VIII.	1
Nitric acid, commercialfl.oz. 2 Water fl.oz. 2	
Water	,
Mix the acid and water, allow to cool, and	1

add the sodium sulphate.

to 125 degs. C.—H.

It depresses temperature from 10 degs. C.

IX.		
Zinc	sulphate	 av. oz. 4
Muri	atic acid	fl oz 4

The temperature depression is from 10 degs. C. to 17 degs. C.—H.

# Fuller's Earth, Improved.

Equal parts fuller's earth and talcum.

Mix and perfume.

# Fungicides.

The term fungicide signifies an agent to destroy fungi or lower forms of parasites. The particular parasites referred to in this connection are such as attack plants, for example, blights, rots, smuts, mildew, etc. Some of the formula under the heading "Insecticides for Agriculturalists," may also be employed as fungicides.

٦.		
	Potassium sulphide(sulphurated	
	potassa)av.lb.	1
	Watergal.	

Dissolve the potassium sulphide in the water, and use. This formula is very efficient for gooseberry and currant mildew. Two or 3 ounces of paris green may be added to each 30 gallons of this mixture.

Copper sulphate	av.lb.	2
Water	gal.	45

Dissolve the sulphate in the water, and use. This solution should never be applied to green foliage. Its proper use is as an early spring wash for the trunk and branches of trees and vines to remove lichens and kill disease spores. Four or 6 ounces of paris green may be added to each 45 gallons of the above solution. For stone fruits use only 2 or 3 ounces of paris green.

#### III.

Copper carbonate av. oz.	6
Ammonium carbonateav.lb.	
Or, ammona waterfl.oz.	32
Watergal.	45

Dissolve the ammonium carbonate in half a gallon of boiling water and pour it upon the copper carbonate. Shake or stir vigorously until all is dissolved. If the ammonia water is used, simply pour it upon the copper carbonate in the same way. Dilute to 45 gallons, and use at once. If kept tightly corked, the undiluted solution will keep for a long time. This is a very efficient fungicide,

adheres well to the foliage, and does not spot the fruit. It should not be used upon the peach, plum or cherry. Arsenites should not be added to this solution.

IV.

Copper sulphateav.oz.	6
Fresh lime (or sodium carbon-	
ate)av.lb.	4 1/2
Molasses	32
Waterga	ıl. 45

Dissolve the copper sulphate in sufficient water using a non-metallic vessel. Use good stone lime, well burnt. Slake the lime or dissolve the soda in 2 gallons of water in a separate vessel. Dilute the molasses with a gallon of water, and stir it into the lime wash or soda solution. After these have been thoroughly mixed add the whole, with vigorous stirring, to the copper sulphate solution. The mixture thus produced should be diluted to 45 gallons and used at once, as it deteriorates on standing. For the earlier treatments, or where the disease is bad, less water may be used, but, as a rule, the full amount will be best. For peach and plum foliage the mixture should never be made stronger than indicated. For the earlier treatments it will be better to use lime instead of soda, and to destroy insects, add to each 45 gallons 3 or 4 ounces of paris green or london purple. The latter arsenite cannot be safely used in the soda mixture. But this has the advantage of not staining the fruit. When an arsenite is used, it should be added to the lime and molasses solution before this is added to the copper sulphate. This mixture is very adhesive to the foliage, and fewer treatments will be required than when the plain Bordeaux mixture is used.

#### Eau Celeste.

Copper sulphateav.lb.	11
Waterpints	38
Ammonia waterpints	11

#### Eau Celeste, Modified.

Copper sulphateav.lb.	4
Sodium carbonateav.lb.	5
Ammonia waterpints	3
Watergal.	45

Dissolve the copper sulphate in sufficient water, using a non-metallic vessel. Dissolve the soda in sufficient water in a separate vessel. Mix these two and then add the am-

monia. Dilute to 45 gallons and use within half a day. It does not keep well. This is a very acrid mixture, and in the hands of careless persons may do much damage to the foliage. It should never be used upon the peach, plum or cherry. It gives best results upon the apple. It does not stain the fruit. Arsenites cannot be safely added to this mixture.

#### Bordeaux Mixture.

Copper sulphate	.lb.	6
Fresh lime		
Water	gal.	45

Dissolve the copper sulphate in a wooden or earthen vessel, using 4 or 5 gallons of water, which, if hot, will act quicker. In a separate vessel slake the lime and rub until all lumps are broken. Then strain and stir slowly into the copper solution. Dilute and use as soon as possible. The mixture should not stand over 20 hours, as it tends to spoil. Where a good quality of lime is used, 4 pounds will satisfy 6 pounds of copper sulphate, but it is best to use plenty of lime, as any free sulphate will burn the foliage. Four ounces of paris green may be added to each 45 gallons of this liquid for all except peach and other stone fruits. For these use only 2 ounces. This makes the best and safest combined insecticide and fungicide for general use. Its chief disadvantage is its spotting the fruit and its liability to be washed off the foliage by rains.

#### Furs, Preserving of.

Carbolic acidfl.c	ir. 6
Oil of clovesfl.o	ir. 3
Oil of mirbanefl.c	lr. 3
Oil of lemonfl.c	
Alcoholfl.c	z. 32

Mix and dissolve.

The articles are moderately sprinkled with the fluid. One sprinkling will suffice for the summer, provided they are stored in closed boxes or closets, but cloth in storerooms will require to be sprinkled twice.

#### Furniture Cream.

Potassium carbonateav.oz.	. 1
Soft or green soapav.oz.	. 2
Yellow waxav.oz.	. 8
Waterfl.oz	. 64

Mix and boil the whole until a uniform cream results.

11.	
Yellow waxav.oz.	31/2
Potassium carbonategr.	
Oil of turpentinefl.dr.	21/2
Oil of lavender	
Watersuffic	

Boil the wax with 16 fluidounces of water over a direct fire, and add to the hot liquid the potassium carbonate. Now remove from the fire, add the two oils, stir until cool, and add enough water to make 32 fluidounces.

In using this cream, apply lightly on a woolen cloth, and then rub with a piece of linen until the furniture has acquired a polish.—D.

#### III.

White castlie soapgr.	270
White waxav.oz.	
Oil of turpentinefl.oz.	
Waterfl.oz.	
Potassium carbonategr.	

Melt the soap in water with the aid of a gentle heat, then add the potassium carbonate and white wax. When thoroughly melted, add gradually the turpentine, and shake thoroughly.

#### IV.

Yellow waxav.oz.	4
Yellow soapav oz.	2
Waterfl.oz.	40
Linseed oilfl.oz.	4
Oil of turpentinefl.oz.	4

Mix the wax, soap and water; boil until of the proper consistence, and add the oils.

# V.

Yellow waxav.oz.	3
Linseed oil, boiledfl.oz.	16

Dissolve the wax in the oil by the aid of heat.

#### VI.

Yellow wax	av.oz.	31/2
Castile soap	gr.	60
Oil of turpenting	nefl.oz.	10
Water, boiling	fl.oz.	10
	onategr.	
	9	

Melt the wax, add the oil, dissolve the soap and potassium carbonate in the water, and mix the two liquids, stirring until cold.

#### Furniture Paste.

Yellow wax	.av.oz.	4
Alkanet, coarse powder	.av.oz.	1/2
Oil of turpentine	fl. oz.	16

Macerate the alkanet in the oil for 24 hours, strain and add the colature to the wax

previously melted. The alkanet may be omitted, if desired.

#### II.

Venice	turp	en	tir	ie.				.av.oz.	- 6
Linseed	loil				 	 		fl.oz.	16

Mix by the aid of heat. The mixture may be colored like the preceding by means of alkanet root.

#### 111.

Yellow waxav.oz.	16
Linseed oil, rawfl.oz.	4
Oil of turpentinefl.oz.	4
Alkanet rootav.oz.	1/2

Mix the wax, linseed oil and alkanet, heat moderately until sufficiently colored, then remove from the fire, and add the oil of turpentine.

#### IV.

Yellow wax	.av.oz. 8
Oil of turpentine	fl.oz. 9
Melt the wax, and add the	
When the mixture has solidified	it may b
cut into rectangular pieces.	

In using, the paste should be rubbed over the furniture, after which a cloth moistened with oil of turpentine should be passed over the latter, and the polishing finished by friction with a soft brush.—D.

#### Furniture Polishes, Liquid.

#### I.

Linseed oil, rawfl.oz.	32
Alcoholfl.oz.	8
Diluted acetic acidfl.oz.	8
Oil of turpentinefl.oz.	8
Solution of antimony chloridefl.oz.	2

#### II.

Linseed oil, rawfl.oz.	40
Alcoholfl.oz.	
Diluted acetic acidfl.oz.	16
Solution of antimony chloride. fl.oz.	2
Ammonium chlorideav.oz.	1
Spirit of camphorfl.oz.	1

Add first the antimony solution, then the spirit of camphor and acid, and lastly the ammonium chloride to the oil, and shake well after each addition.

#### III.

Alcohol fl.oz.	10
Linseed oil, rawfl. oz.	10
Sandaracgr.	120
Diluted acetic acidfl.oz.	5
Nitric acidfl.dr.	4

Mix and dissolve.

THE STANDAN	D FORM OLAR I.
IV.  Linseed oil, raw	XI.  Oil of turpentine
Venice turpentineav.oz. 34 Alcoholfl.oz. 12 Mix and shake occasionally until dissolved, then set aside in a warm place for a few weeks and filter.  X.	XVI.       Shellac       av.oz. 2         Alcohol       fl.oz. 16         Linseed oil, raw       fl.oz. 16         Oil of turpentine       fl.oz. 2         Ether       fl.oz. 2
Resin of guaiac av.oz. 1 Benzoin av.oz. 1 Shellac av.oz. 1 Linseed oil, raw fl.dr. 10 Benzin fl.dr. 4 Alcohol, or wood alcohol fl.oz. 24 Mix and dissolve. The polish is applied with a sponge or brush, and the object is, let stand for a half hour. A linen cloth moistened with oil is then used as a rubber, and a brilliant polish	Ether fl.oz. 2 Ammonia water fl.oz. 4 Dissolve the shellac in the alcohol and add the other ingredients.  XVII. Alcohol fl.oz. 4 Oil of turpentine fl.oz. 2 Damar varnish fl.oz. 1 Linseed oil, raw fl.oz. 8 Acetic acid fl.oz. 1 XVIII.
is obtained which is said to be very lasting and is unaffected by water or any substances which usually injure varnish.	Oil of turpentine       fl.oz. 8         Oil of amber       fl.oz. 8         Cottonseed oil       fl.oz. 8

# Gilding Powder.

Gold, chlorideav.oz.	1
Potassium cyanideav.oz.	3
Potassium bitartrategr.	110
Prepared chalkav.oz.	

Mix the ingredients intimately; add water to make a paste; rub with a bit of flannel. The surface must, of course, be thoroughly cleaned.

By substituting silver nitrate for the chloride of gold, a silvering powder will be obtained. To be employed in the same manner.

#### Glass, Cutting of.

The following may be used for cutting glass, bottles, flasks, etc.:

Charcoal, woodav.oz.	1
Potassium nitrategr.	10
Tragacanthgr.	10
Benzoingr.	5

Reduce each to fine powder, add enough mucilage of tragacanth to form a mass, and roll this out into cylinders of about the thickness and length of a lead pencil.

One end of this cylinder may be ignited and passed over the flask, bottle, etc., wherever the crack or cut is to be made.

Instead of the above, the following may be employed:

Dissolve 1 av. ounce of lead acetate in 7 fluidounces of water; saturate blotting paper with this solution, then dry, cover one surface of the paper with paste containing 10 per cent of potassium nitrate, roll this lightly over a knitting needle and dry. This may be used like the preceding.

#### Glass, Mirrors, etc., Polish for.

Moisten calcined magnesia with pure benzin so as to form a paste sufficiently wet that when pressed some of the benzin will exude. The articles are cleansed by taking this mixture on cotton (not cotton cloth) and rubbing over the glass until dry and all powder is rubbed off.

See also "Window Polishing Paste."

#### Glove Cleaner.

See also "Benzin Jelly," "Cleansing Cream," "Cleansing Liquid," "Stains, Removal of," "Gloves, Dry Process for Cleaning," etc.

I.

Solution of chlorinated soda,fl.oz.	12
Ammonia waterfl.oz.	1
Soap, powderav.oz.	15
Waterfl.oz.	20

Make into a soft paste, and rub on the gloves with a flannel.

TΤ

Oil of	turpentine	 	 fl.oz.	5
	l <del></del> .			

III.

Castile soap, shavingsav.oz. 12	
Water, hotfl.oz. 8	
Solution of chlorinated sodafl.oz. 8	
Water of ammoniafl.dr. 4	

Dissolve the soap in the water, allow to cool, and incorporate with the solution and ammonia so as to form a smooth paste.

In using, rub a small portion over the glove by means of a piece of flannel, always rubbing in one direction until clean.

# Gloves, Dry Process for Cleaning.

Put the gloves on a clean board, make a mixture of dry powdered fuller's earth and powdered alum, and apply the powder to both sides of the glove with an ordinary stiff brush. Then wipe the powder off, cover the glove with dry bran and brush this off. The gloves, if not very badly soiled, will, by this process, become entirely clean.

Should there be grease stains, remove them with crumbs of toasted bread and powdered animal charcoal, and then rub the glove with a clean woolen rag dipped into the powder of fuller's earth and alum.

# Glue, Bookbinder's.

Glue,	best.	 	 	 	 ,	.av.oz.	7
						fl. oz.	
						sufficie	

Pour on the glue more than enough water to cover, allow to macerate for several hours, then decant the greater portion of water; apply heat until the glue is dissolved, and add the glycerin. If the mixture is too thick, more water may be added. It may be colored by means of an aniline dye dissolved in alcohol.

#### Glue, Liquid.

The making of so-called "liquid glue" depends on the fact that when gelatin or glue is mixed with certain substances in

the presence of water, the mixture remains permanently semi-liquid. The most common agents used in this liquefying process are acetic and nitric acids, lime and other substances also being used.

The cheaper kinds are made from glue, the better and handsomer preparations are made from gelatin.

Tom gelam

White glue, broken into	small
pieces	av.oz. 2
Acetic acid	av.oz. 8
Nitric acid	drops 10

Mix and keep in a wide-mouthed vial, well corked.

Mix the glue and acetic acid in a widemouth stoppered bottle; set in a warm place, agitate frequently until dissolved, and then add the nitric acid.

II.

Glue av.oz.	4
Waterfl.oz.	10
Nitric acidfl.dr.	4
Tittle dela	_

Boil together for several hours.

III.

Glue av.oz.	12
Alumgr.	50
Acetic acidfl.oz.	
Waterfl.oz.	13
Alcohol fl.oz.	

Mix all but the alcohol, digest on a water bath till the glue is dissolved, allow to cool, and add the alcohol.

IV.

Glue, white or brownav.oz.	51/2
Acetic acidfl.oz.	
Carbolic acid drops	
Watersufficien	

Macerate the glue in 6 fluidounces of water for 12 hours, heat the mixture on a water bath until the glue is dissolved, add to the solution the two acids, and then enough water to make 16 fluidounces.

V		
	1	

Chloral hydrateav.oz.	5
Gelatinav.oz.	8
Waterfl.oz.	20

Mix; let stand for 48 hours, and decant the clear liquid.

VI.

Glue, best whiteav.oz.	4
Lead carbonateav.oz.	1
Alcohol fl.oz.	8
Waterfl.oz.	24

Dissolve the glue by means of a water bath in the water, then incorporate the lead compound and the alcohol, and bottle while warm.

VII. This preparation has been called "Syndeticon":

Slaked limea	v.oz. 4
Sugara	v.oz. 6
Water	
Gluea	v.oz. 6

Dissolve the lime and sugar in the water heated to 75 degs. C., decant the clear liquid, add the glue, and, after allowing to swell, again apply heat until dissolved.

VIII. This is similar to the preceding, the proportions only differing:

Sugarav.oz.	5
Glue, best brownav.oz.	
Lime, slakedav.oz.	$1\frac{1}{2}$
Waterfl.oz.	
Oxalic acidsufficie	
Carbolic aciddrops	10

In a flask dissolve the sugar in the water, add the lime, and warm the mixture to a temperature of about 75 degs. C. for 3 days, agitating frequently; then allow to cool, decant the clear liquid, and to 8 fluidounces of this liquid add the glue, previously reduced to small pieces; allow to macerate for 3 hours, and then heat in a covered vessel on a water bath for 10 hours, replace the water lost by evaporation, neutralize the lime with oxalic acid, and then add the carbolic acid.

IX. This preparation is also known as "Syndeticon":

Calcium chloride	av.oz. 1
Water	.fl.oz. 1
Glue, best brown	av.oz. 5

Dissolve the calcium chloride in the water, add the glue, macerate until the latter is thoroughly softened, and then heat until completely dissolved.—D.

X

Solution of sodium silicateav.oz.	10
Sugar, powderav.oz.	3
Acacia, powderav.oz.	1

Mix well, adding some water if necessary.

Glue, Stick. (Pocket Glue.—Elastic Glue.
—Mouth Glue.)

See "Stick Mucilage."

Glucose.—(Grape Sugar or "Starch Syrup") Is in the market as a solid and liquid. The latter is a variable mixture of grape sugar and dextrine in water.

#### Grafting Wax.

Resin av.oz.	16
Beef tallowav.oz.	
Oil of turpentinefl.dr.	
Alcohol fl.oz.	- 5

Melt the resin, add the tallow, stir until homogeneous, remove from the fire, allow to cool somewhat, and add the oil and alcohol, little by little, stirring well with each addition. If in adding the alcohol there is a tendency to lump, carefully warm the mixture until it melts.

It should be kept in closely stoppered bottles, and when used warmed up slightly (if not in a liquid condition or, say, the consistency of molasses). Apply with a brush. A very thin coat only is needed.

Ditoh

1 1tcn	
Resinav.oz. 2	
Yellow waxav.oz. 1	
Lardav.oz. 1	
II.	
Yellow waxav.oz. 2	
Resinav.oz. 2	
Gum turpentine	

# Guano, Artificial.

Sodium sulphate, driedav.lb.	1 1
Common saltav.lb.	12
Wood ashesav.lb.	4
Ammonium sulphate, commonav.lb.	16
Bone dustbushel	1

#### Gun Barrels, Staining of.

#### I. A..

В.

Solution of iron chloridefl.oz.	1
Corrosive sublimategr.	180
Copper sulphategr.	180
Fuming nitric acidfl.dr.	2
Distilled waterfl.oz.	10
Mix and dissolve	

Potassium sulphide.....gr. 50

Distilled water.....fl.oz. 10 Mix and dissolve.

Clean off the gun barrel with emery paper, then by means of a sponge or soft hair

in a cool place, so that it may occur rather slowly. Repeat this application and drying twice, or oftener if necessary to secure the shade of stain desired, rubbing over thoroughly before each application with a scratch brush. When the metal is stained deeply enough, lay the barrel in solution B for 20 or 30 minutes, then wash with warm water, and finally with soap water. Then dry and rub over with linseed oil varnish.

Better results will be attained by stoppering closely the gun barrel at both ends by means of corks, and laying for at least 30 minutes in each of the baths, which have previously been warmed.-D.

#### II. A.

Fuming	nitric	acid	 	 fl.dr.	2
Distilled	water		 	 fl.oz.	16

Silver nitrate.....gr. 80 Distilled water.....fl.oz. 16

Rub off the gun barrel with emery paper, then by means of a sponge or soft hair brush apply solution A, dry in a cool place, and rub off with a cloth. Repeat this application, drying and rubbing off until a handsome oxidized surface is produced. Then apply solution B repeatedly, with subsequent exposure to light, until the gun barrel is dark enough, and then anoint with linseed oil varnish.—D.

#### Gutta Percha, Purified.

Gutta perchaav.oz.	8
Carbon disulphidefl.oz.	
Alcohol fl.oz.	108
Distilled water flor	25

Soften the gutta percha in lukewarm water; then pull to pieces, dissolve it in the carbon disulphide, set the mixture aside for 24 hours, filter through glass wool into a suitable vessel containing 60 fluidounces of alcohol. Agitate the whole together, and set aside until the mixture separates into two layers. Decant the upper alcoholic layer, wash the residue with the remainder of the alcohol in the same manner, decant as before; add the water, transfer the mixture to a retort, and distill off the carbon disulphide. Owing to the inflammability of the latter, the utmost precaution must be taken to avoid brush, apply solution A, subsequently drying ignition of its vapors. Finally, wash the residual mass by kneading in water, then express the latter, and dry in thin sticks.

The product weighs about 5 to  $6\frac{1}{2}$  av. ounces.—D.

#### Hardwood Filler.

Use boiled oil and enough corn starch to make a very thick paste. Add a little japan, and reduce with turpentine. Add no color for white oak; for dark ash and chestnut use a little raw sienna; for walnut, burnt umber and a very little venetian red; for baywood, burnt sienna. Use enough color to cover the white of the starch. Apply with brush and rags, let dry 48 hours, then sandpaper. For second coat use less oil but more japan and turpentine.

# Harness Blacking, Polish or Oil.

These preparations are similar to shoe dressings and blackings, and some of the preparations to be found in this part under the latter heading may be made to serve the purpose of harness polishes.

1.	
Glue av.oz.	. 4
Acaciaav.oz.	. 2
Diluted acetic acidfl.oz.	. 24
Black inkfl.oz	. 8
Isinglassgr	. 120

Soften the glue by standing in 16 fluidounces of acid, dissolve the gum in the ink, and the isinglass in a little warm water. Add the rest of the acid to the glue solution, then warm it until solution is obtained; add the gum and ink, and next the isinglass. When all is warm and thoroughly mixed, remove from fire.

In using, warm enough to liquefy, and then apply by means of a sponge.

II.

1	
	Mutton suetav.oz. 2
	Yellow waxav.oz. 6
	Powdered sugarav.oz. 4
	Yellow soapav.oz. 2
	Lampblackav.oz. 1
	Indigo av.oz. 4
	Waterfl.oz. 4
	Oil of turpentinefl.oz. 4

Dissolve the soap in the water, add the other ingredients (except the turpentine) melt and mix well together; finally, àdd the turpentine. The mixture is applied on the harness with a sponge, and polished with a brush.

2.2	
Black anilinegr.	35
Muriatic acid	
Bone blackgr.	175
T 11 1	40

 Lampblack
 gr. 18

 Yellow wax
 av.oz. 2½

 Oil of turpentine
 fl.oz. 22

 IV.
 Oil of turpentine
 ...fl.oz. 8

 Yellow wax.
 .av.oz. 2

 Prussian blue
 .av.oz. ½

Melt the wax, add the turpentine, a portion first to the finely powdered prussian blue and lampblack, and thin with neatsfoot oil.

Lampblack.....av.oz. 1/4

# Hectograph Masses.

Hectographs, also known as copygrams, copygraphs, chromographs, collographs, etc., are employed for the purpose of duplicating writing by taking an impression of writing made with a suitable aniline ink on a receiving pad made essentially from gelatin or glue and glycerin and then obtaining copies by laying fresh sheets of paper upon the pad.

The following formula will make a good pad. The directions with regard to air bubbles, pouring of the mass, etc., must be followed strictly in the succeeding formulas:

Take a pound, or any convenient quantity of pure white glue, free from whiting or other insoluble matter, and macerate in water, until it becomes soft and pliable. With a little manipulation and turning, this may be accomplished by using a pint of water for each pound of glue. Drain off the excess of water, if there be any, and add glycerin in the quantity of from two to three pints for each pound of glue used. The lesser amount is for summer, and the larger amount for winter use. Heat the mixture gently until the glue is dissolved, and the water absorbed by the glue has evaporated. The easiest way of ascertaining when this is accomplished is, to take the weight of the evaporating dish out before commencing; then when the weight of the dish and its contents is equal to its tare and the amounts of glue and glycerin used, the operation may be considered completed. If the water is not driven off, the pad is likely to crack in a dry atmosphere by its spontaneous evaporation.

It will be found somewhat troublesome to avoid air bubbles in the mass. They may be avoided to an extent by the use of only moderate heat in dissolving the gelatin. When bubbles have formed, they may be skimmed off, but it is easier to destroy them by the use of alcohol. The mass when finished is strained into a wide mouthed bottle and allowed to stand in a warm place, or in warm water for an hour or two, when most of the bubbles will have arisen to the top and formed a scum on the surface. A small quantity of alcohol is now poured carefully down the side of the bottle, which instantly destroys them all. The gelatin, which is precipitated by the alcohol on the surface, re-dissolves as the alcohol evaporates. When this has occurred, the mass may be poured into a suitable shallow tray, holding the mouth of the bottle as near the tray as possible, to avoid the formation of fresh bub-The tray may be made by any tinner and should be as large as the largest paper on which copies are to be taken. It should be about 1/2 inch deep. To secure firmness, it should be fastened to a board. melted mass may now be poured, as described, to nearly fill the tray; any air bubble which forms must be removed with a hot wire or other suitable means, and when cold the pad or hectograph is ready for use.

It is difficult to obtain a pad absolutely free from bubbles and one which is not sticky. To avoid the latter, various insoluble powders, sometimes soluble salts, are added to the hectograph mass. The former include clay, chalk, sulphur, barium sulphate, etc.; the latter potassium and other chromates. The former must be rubbed to a smooth paste with a portion of the melted mass before adding to the remainder of the smallest amount of water before adding to the remainder of the mixture.

The modus operandi of taking copies is as follows:

Write upon the paper with a suitable ink, allow it to remain for a minute or so to dry partially, then invert carefully upon the pad and press evenly and uniformly. In a minute or two, sufficient ink will have been absorbed by the pad from the paper so that new sheets

of paper, pressed upon the pad, will receive duplicates of the original writing. The number of copies that may be taken in this way, as well as their distinctness, varies according to the pad and the ink. The original writing, if laid upon a fresh pad or a fresh portion of the same pad after taking the first imprint, will furnish still other copies. After all the copies are taken, the pad should be cleansed with a moist sponge. A trifling amount of ink will remain in the pad, but this will not interfere with subsequent operations. The copies are sometimes improved by moistening the copying paper with water or strong alcohol, and then absorbing the excess of liquid between folds of bibulous paper.

Hectograph inks are mentioned under heading "Inks."

The hectograph mass, instead of being poured into a tray, may be formed into roller by casting in a mold. Copies may be taken by passing the roller over the writing and then over the paper to receive the copies.

The following mixtures may be formed into hectograph pads, as described above:

nectograph pads, as described above.
I.
Gelatinav.oz. 4
Glycerin
Potassium bichromategr. 60
II.
Gelatinav.oz. 4
Water
Glycerin
White clay (Kaolin)av.oz. 2
· · · · · · · · · · · · · · · · · · ·
III.
Gelatinav.oz. 1
Molasses av.oz. 1
Glycerinfl.oz. 9
Watersufficient
IV.
Glue
Glycerin
Corbolic acid A on 1/
Carbolic acidfl.oz. 1/2

The glue should be soaked in water several hours before it is melted with the glycerin. Barium sulphate is also used as an addition. V.

Sulphur.....gr. 60

White glueav.oz.	6
Glycerinfl.oz.	
Dextrinav.oz.	2
Precipitated sulphur, pureav.oz.	1/2
Water suffici	ent.

VI.	
Isinglassav.oz.	4
Glycerinfl.oz.	24
Waterfl.oz.	16

Mix; macerate for a few hours, and then warm sufficiently to dissolve.

#### VII.

Gelatin
Glycerinfl.oz. 32
Water,
Barium sulphateof each, sufficient

Use the latter sufficient to make the mass of proper body.

#### VIII.

Good ordinary glueav.oz.	10
Glycerinfl.oz.	4
Kaolin or barium sulphateav.oz.	21/2
Watersufficie	ent

#### IX.

Best furniture glueav.oz.	4
Waterfl.oz.	8
Glycerinav.oz.	14

Soak the glue in the water until soft, stirring frequently meanwhile; then add the glycerin, and heat on a water bath, with gentle stirring, until the mixture weighs 10 av. ounces.

#### Х.

_		
	Gelatinav.oz.	4
	Dextrinav.oz.	
	Sugarav.oz.	
	Waterfl.oz.	10
	Glycerinfl.oz.	12
	Zinc oxideav.oz.	11/2

#### Horn and Ivory Black, to Color.

First place the horn in an aqueous solution of a lead salt, with a slight excess of sodium hydrate. The duration of this treatment depends on the character of the horn and the strength of the solution—generally half an hour is sufficient. Then wash well and introduce the horn into a solution of 350 grains of wool black and 18 grains of naphthol yellow S in 1 pint of water at 40 degs. C. A longer subjection to this latter treatment appears to be necessary.

For bone and ivory, water-soluble nigrosin will answer. It is only necessary to lay the pieces, previously deprived of fat and mordanted, in a hot aqueous solution of nigrosin until the desired tone is obtained. On account of its cartilaginous components,

ivory cannot be boiled in the nigrosin solution, but the same result can be obtained by allowing it to stand for some hours in a concentrated solution at a temperature of about 30 degs. C.

To deprive the bones of fat before treatment with aniline colors it is sufficient to boil them with frequent changes of water; subsequent treatment with ether being unnecessary.

Kellermann's method of mordanting is especially recommended. This consists of placing the defatted bones for fifteen minutes in the following:

Nitric acid, concentrated fl.oz.	1
Waterfl.oz.	21
Tartaric acidgr.	120

They are then washed and placed in a solution of 7 grains of zinc chloride in 1 pint of water with a few drops of hydrochloric acid.

With ivory it is sufficient to mordant for 15 minutes in 1 per cent hydrochloric acid.

# Incense. (Balsamic Fumigation.)

Benzoin	av.oz.	2
Olibanum	av.oz.	3
Myrrh	av.oz.	3
Cascarilla	av.oz.	11/2
Oil of lavender flowers		
Oil of bergamot	.drops	10
Oil of cloves	.drops	5
Oil of cinnamon		
ŗ.		

# Cascarilla.....av.oz. 1

-	
	Olibanumav oz. 4
	Benzoinav.oz. 1
	Liquid styraxav.oz. 1
	Rose petalsav.oz. 1
	Lavender flowers av.oz. 1

#### IV.

Amberav.oz.	
Masticav.oz.	3
Olibanumav.oz.	
Benzoinav.oz.	
Storaxav.oz.	
Camphorgr.	60

#### Inks.

The characteristics of a good writing ink are as follows:

1. It must flow easily from the pen, but not drop from it nor spread on the paper;

- 2. It should not contain finely suspended matter which will subside in the course of time, but should be perfect solution;
- 3. The color should be dark, and therefore practically saturated, and writing made with it should not fade;
- 4. It should not mold, nor be liable to other decomposition;
- 5. It should copy or it should not, as may be desired, and,
- 6. It should not appreciably attack or corrode steel pens.

Black writing inks are frequently classed according to their use, as follows:

- 1. Office or document inks, which must be prepared from nutgall or tannin, should be permanent, and are intended for documents which are to be preserved;
- 2. Copying inks, which are prepared from nutgall, tannin, logwood, or even coal tar dyes, and which must furnish good copies, and
- 3. Ordinary writing inks, such as are employed as house and school inks, and which should be cheap and from which no special permanence is expected.

According to composition, inks may be classified into

- 1. Aniline inks:
- 2. Logwood inks;
- 3. Nutgall inks;
- 4. Tannin inks, and,
- 5. Miscellaneous inks.

The first class yields copying and writing inks, as well as the various colored inks which are in use for various purposes. The second class yields copying and writing inks, and the third and fourth classes yields document and copying inks. The fifth class embraces hectograph inks, indelible inks, sympathetic inks, etc.

In order that a good ink may retain its excellent qualities, certain cautions should be observed in its use: Before putting a new ink in an ink-well, the old ink should be entirely removed and the ink-well washed. Also no ink container should be used which cannot be closed, and such ink vessel should always be closed when not in use. If an ink is liable to thickening or other change, the ink-well should be cleansed before refilling, even if with the same ink.

Several preparations, which are used in the manufacture of many of the inks which follow, are mentioned here.

Nutgall infusion:

Chinese nut	gall						.av.oz.	7
Talcum, pur	rified.	 					.av.oz.	3/4
Water, disti	lled						sufficien	ť

Reduce the nutgall to coarse powder, moisten (not wet) the powder, and set the latter aside at a temperature of 20 to 25 degs. C. until it is thickly covered with mold. order to hasten this molding, the drug should be moistened daily with water, so that it will always have about the same proportion of moisture. At the end of from 8 to 10 days, fermentation will have advanced sufficiently to admit of extraction of the drug. To the latter should be added 14 fluidounces of water, and the mixture heated for an hour on the water bath. Then express, treat the residue in the same manner with the same amount of water, and then again with 7 fluidounces of water. Mix the three liquids obtained, add the talcum, shake well, set aside for 24 hours, filter, and add enough water, if necessary, through the filter to make the filtrate measure 32 fluidounces.

The infusion will keep for several days. Tannin solution:

Tannic acid	.av.oz. 3
Muriatic acid, commercial	fl.dr. 4
Distilled water	sufficient

Mix the two acids and 3 fluidounces of water in a flask, and heat on a water bath to a temperature of 80 to 90 degs. C. for 3 hours, adding from time to time hot water until 27 fluidounces are added.

This solution should not be kept longer than 7 days.

Aniline Inks.—Many of the coal tar dyes (misnamed "anilines,") which are now manufactured, produce excellent copying inks. Compared to nutgall, tannin, and logwood inks, they are less permanent, but they will serve excellently where no especial permanence is required. They are especially useful as hectograph inks. Where permanence of writing is demanded, as in the draughting of documents, aniline inks cannot be employed as they soon become bleached from the action of air and light.

It is to be noted here that water containing

lime decomposes many aniline colors, and solutions of these dyes in calcareous water may thicken in the course of time, hence only distilled water should be employed in the manufacture of these inks.

In using, only the best obtainable dye of the kind mentioned should be used, as otherwise good results cannot be obtained.

Logwood Inks.— These might also be designated as chrome inks, for they always contain potassium bichromate or chrome alum; also some acid, with the object of producing acid salts of the chromium compound. The greater the amount of acid in proportion to the chromium, the paler or redder and thinner the product, and, conversely, the greater the proportion of chromium, the darker and thicker the ink will be. Most logwood inks copy with great facility—writing produced sometimes being copyable even after weeks and months.

Logwood inks can be more easily erased from paper than nutgall inks. They have the advantage of furnishing several copies if desired. All inks lose their copying qualities when exposed to air containing even traces of ammonia. In order to facilitate copying with an exposed logwood ink, the copying paper should be moistened with a one-tenth per cent aqueous solution of potassium chromate. Old writing made with logwood ink may be copied in the same manner, even after the lapse of years.

Logwood inks may be prepared from the following solution:

Logwood extract solution:

Logwood	extract,	best	av.oz.	41/2
Distilled	water		fl. oz.	20

Dissolve the extract in the water on a water bath, set the solution aside for 8 days, and decant the clear liquid.

Nutgall Copying Inks.—Copying inks differ from non-copying in that they retain their copying qualities. Inks which are made with ferric salts soon become non-copyable, while those made with ferrous salts, especially ferrous sulphate, copy the best. Hence copying inks are made with ferrous salts, and non-copying inks with ferric salts. The copying qualities of ink are improved by the addition of pure sugar or of glucose

In preparing the different nutgall copying inks, the following mixture, which may be known as "Nutgall Ink Body I," forms the basic ingredient:

# Nutgall Ink Body I.:

Nutgall infusionfl.oz.	30
Sulphuric acid, concentratedm.	
Ferrous sulphate, puregr.	100
Distilled watersuffic	

Mix the infusion and acid; heat for 15 minutes on a water bath, dissolve the iron salt in the mixture, transfer the latter to a bottle, cork well, set aside for 2 weeks, filter and add through the filter enough water to make the filtrate measure 32 fluidounces.

Nutgall Non-Copying Inks.—The following mixture, known in these pages as "Nutgall Ink Body II.," serves as a body for many of the inks mentioned below:

# Nutgall Ink Body II.:

Nutgail infusion	9%
Solution of chloride of iron, U.	
S. P	21/
Distilled water fl.dr.	

Allow this mixture to stand for 2 weeks in a closed vessel, and then filter.

Tannin Copying Inks.—Tannin copying ink', like the nutgall copying inks, are preferably made with ferrous salts and contain an addition of sugar or glucose. The following mixture, which may be known as tannin "ink body I.," is the basic ingredient of the tannin copying inks:

#### Tannin Ink Body I.:

Tannin	solutionfl.oz.	15
Ferrous	sulphate, pureav.oz.	13/
Distilled	l watersuffici	ent

Heat the tannin solution to about 70 or 80 degs. C., also dissolve the iron salt in 9 fluidounces of hot water; mix the hot solutions by pouring iron solution gradually into the tannin solution, set the mixture aside for 3 weeks, filter and add enough water through the filter to make the filtrate measure 25 fluidounces.

Tannin Non-Copying Inks.—The tannin inks herein mentioned are frequently prepared by the use of the following mixture, which may conveniently be termed "tannin ink body II."

Tannin	Ink	Body,	II.:
--------	-----	-------	------

Tannic acidav.oz.	31/2
Solution of iron chloride, U.	
S. P	4
Muriatic acid, commercial fl.dr.	
Water sufficie	nt

Mix the two acids, the solution and 7 fluidounces of water in a flask on a water bath, at a temperature of 80 to 90 degs. C., for a period of 10 hours. Then add 20 fluidounces of hot water; continue the heat for another hour, transfer to a bottle, cork well, set aside in a cool place for 2 weeks, filter and add through the filter enough water to make the filtrate measure 32 fluidounces.—D.

#### Ink, Alizarin.

Alizarin pastegr.	225
Sodium carbonate, puregr.	105
Extract of logwoodgr.	
Carbolic acidfl.dr.	
Waterfl.oz.	

Dissolve the sodium carbonate in a small amount of water, add the alizarin paste, then the extract of logwood, previously dissolved, in the remainder of the water. Filter and transfer the liquor to a rather large bottle, drop in a few nails or iron filings, and expose the whole to the sunlight for a week, with occasional agitation. Lastly, decant and preserve by incorporating the carbolic acid. This ink does not corrode the pen, is not affected by light, and does not gum, but it is not acid proof.

# Ink, Alizarin, Copying

Indigotingr.	95
Aniline green, Dgr.	
Sugarav.oz.	
Distilled waterfl.oz.	
Nutgall ink body Ifl.oz.	
Carbolic aciddrops	

Dissolve the dyes and sugar in the water by the aid of heat, add the remaining ingredients, transfer the mixture to a bottle; tie over the latter a piece of paper, set aside in a cool place for a week, and decant the clear liquid from the triding precipitate.—D.

# II.

Indigotingr. 70
Aniline green, Dgr. 42
Glucoseav.oz. 11/4
Distilled waterfl.oz, 2
Tannin ink body Ifl.oz. 25
Carbolic aciddrops 15
Dissolve the dyes and glucose in the water

by the aid of heat, add the remaining ingredients, transfer to a bottle; tie over the latter a piece of paper, set aside in a cool place for one week, and decant the clear liquid from the trifling precipitate.—D.

# Ink, Alizarin, Non-Copying.

•	
Aniline green, Dgr	. 76
Indigotingr	. 128
Waterfl.oz	
Nutgall ink body IIfl.oz	
Carbolic acidfl.di	

Dissolve the two dyes in the water by the aid of heat, add the other ingredients, transfer the mixture to a bottle, tie over the mouth of the latter a piece of paper, set aside for one week in a cool place, and decant the clear liquid from the trifling sediment.—D.

#### TT

Indigotingr.	80
Aniline green, Dgr.	48
Tannin ink body IIfl.oz.	
Distilled waterfl.oz.	
Carbolic acid drops	
Sugargr.	
	10

Dissolve the dyes in the water by the aid of heat, add the other ingredients, transfer to a bottle, tie over the latter a piece of paper, set aside for one week in a cool place, and decant the clear liquid from the trifling precipitate.—D.

# Ink, Black, Copying.

I.
Phenol black (coal tar dye)gr. 190
Sugarav.oz. 11/4
Distilled waterfl.oz. 3
Nutgall ink body IIfl.oz, 32
Carbolic aciddrops 20
Prepare like alizarin copying ink, No. I.
—D.

# II.

Phenol black, B. (coal tar dye)gr.	140
Glucoseav.oz.	
Distilled waterfl.oz.	2
Tannin ink body Ifl.oz.	25
Carbolic aciddrops	15
Prepare like alizarin copying ink, N	o. II.

# —D.

•		
	Aleppo gallsav.oz.	51/2
	Cloves gr.	60
	Distilled waterfl.oz.	40
	Ferrous sulphate, puregr.	720
	Sulphuric acid, purem.	35
	Neutral sulphate of indigogr.	120
	The galls and cloves, coarsely ground	l, may

be exhausted by percolation until 40 fluidounces are obtained, or they may be macerated with sufficient water. In either case it is intended to produce 40 fluidounces of the fluid, and allowance must be made for the water absorbed by the marc. To this, when filtered, add the iron, and when dissolved filter again, then add the acid and, after mixing thoroughly, the indigo paste, after which it may be again filtered.

This produces a blue-black fluid, not apt to mold. To insure a superior product, careful attention must be paid to manipulation, details, and to the quality of its ingredients. The galls must be free from insect perforations, and the iron, selected crystals free from efflorescence or ferric salt, and the indigo neutral, or nearly so. If the article sold as "indigo paste" is not at hand, it may be prepared by carefully adding to the ordinary sulphate of indigo a solution of potassic or sodic carbonate until effervescence ceases.

IV.

Galls, groundav.oz.	8
Ferrous sulphate, pureav.oz.	
Gum arabicav.oz.	2
Sugarav.oz.	
Distilled waterfl.oz.	

Macerate the galls, with frequent agitation, in 40 fluidounces of the water for one week: filter, and to this infusion add the iron, previously dissolved in the remaining pint of water. Dissolve in this mixture the gum and sugar; filter, and the ink is ready for use. A better product is obtained if the iron is dissolved in water made slightly acid with sulphuric acid.

V.

Extract of logwoodav.oz.	51/3
Sodium carbonate gr.	525
Waterfl.oz.	
Glycerinfl.dr.	4
Potassium chromate gr.	
Mucilage of acaciafl.dr.	

Heat the extract of sodium carbonate with the water by the aid of heat, add the glycerin, mucilage and potassium bichromate, the latter first dissolved in some water, and then add enough water, if necessary, to make up 40 fluidounces.

This ink will give a good copy without a press, by simply laying a sheet of moist copy- reducing the dye even to 1½ av.ounces.—D.

ing paper over the written page, covering with a sheet of letter paper and pressing evenly with the hand or paper knife

# Ink, Black, Non-Copying.

•	Phenol black, B (coal tar dry)gr.	320
	Waterfl.oz.	28
	Nutgall ink body IIfl.oz.	
	Carbolic acidfl.dr.	1/2

Prepare like alizarin non-copying ink, No. II.-D.

II.

Phenol black, B (coal tar dye)gr.	
Tannin ink body IIfl.oz.	16
Distilled waterfl.oz.	25
Carbolic aciddrops	20
Sugargr.	40

Prepare like alizarin non-copying ink, No. II.—D.

# Ink, Black, Writing.

1.			
	Logwood extract solution	av.oz.	20
	Potassium bichromate	gr.	90
	Chrome alum	av.oz.	.,
	Oxalic acid	av.oz.	1
	Carbolic acid	.fl.dr.	1
	Distilled water	sufficie	enl

Mix the extract solution with 50 fluidounces of water, heat on a water bath to 90 degs. C., add the potassium bichromate, chrome alum, and oxalic acid previously dissolved in 15 fl.ozs. of water; continue the temperature of 90 degs. C. for one-half hour, then add enough water to make the mixture weigh 100 av.ounces, and the carbolic acid; set aside for 2 or 3 days, and decant the clear liquid.

This ink is black in color, and the writing is of the same tint. It is very cheap, and hence is adapted to school purposes.—D.

11.	
Phenol black, B (coal tar dye)av. Sugarav.	oz. 21/4
Sugar av.	oz. 21/4
Carbolic acidfl.	dr. 1
Sulphuric acid, pure	m. 25
Distilled water	oz. 96

Mix the dye with 6 fluidounces of cold water, allow to stand for 2 hours, then add the remainder of the water, in the boiling condition, and the other ingredients, and stir about until dissolved.

This ink writes a handsome blue-black. For school purposes it may be cheapened by

1415055511115003	TREFFICATIONS. 451
Logwood chips	Dissolve the dye in the water by the aid of heat; add the other ingredients, and again dissolve —D.  Ink, Blue, Non-Copying.
fluidounces of decoction, and to it add the potassium salt previously dissolved in water. This makes a very cheap ink.  IV.  Extract of logwood	I
Lime water	I.—D.  II.  Phenol blue, 3 F (coal tar dye)gr. 60 Tannin ink body IIfl.oz. 16 Distilled waterfl.oz. 25 Carbolic aciddrops 20 Sugar
water bath, stirring constantly, and then add the two acids, which change the color of the	Prepare like alizarin non-copying ink, No. II.—D.
solution from red to brownish yellow.  Set the mixture aside until cool, then filter;	Ink, Blue-Green, Copying.
add the potassium salt, first dissolved in some water, then the potassium bichromate, and finally, the remainder of the water.	I. Phenol blue, 3 F (coal tar dye)gr. 38 Aniline green, Dgr. 95 Sugarav.oz. 114
Ink, Blue-Black, Copying.  Prepare like red copying ink, No. III, but	Distilled water
decreasing the sulphuric acid to 13 drops, and increasing the potassium bichromate to	Prepare like alizarin copying ink, No. I.— D.
70 grains.	II.
This ink is of a dark-blue color; the writing and copies are blue-black.—D.	Phenol blue, 3 F (coal tar dye)gr. 28 Aniline green Dgr. 70
Ink, Blue, Copying.	Glucose.       av. oz.       1½         Distilled water       fl. oz.       2         Tannin ink body I       fl. oz.       25
Phenol blue, 3 F (coal tar dye)gr. 48         Sugar	Carbolic acid
Carbolic aciddrops 20	Ink, Blue, Writing.
Prepare like alizarin copying ink, No. I.— D.	Resorcin blue, M (coal tar dye)gr. 48
II. Phenol blue, 3 F (coal tar dye)gr. 42	Sugar       gr. 192         Oxalic acid       gr. 10         Distilled water       fl. oz. 19½
Glucose	Mix the dye with 1 fluidounce of cold water, set aside for 2 hours, then add the remainder of the water, in the hot condition,
Prepare like alizarin copying ink, No. II.	and the other ingredients, and stir about until dissolved.
—D.	This ink writes a handsome blue and flows
III.	111 1 1 1 1 1 1 1 1 1

Resorcin blue, M (coal tar dye)..gr. 100

Sugar.....gr. 100

Oxalic acid.....gr. 20 Distilled water.....fl.oz.

what corroding the pen, and hence the latter should be cleaned frequently.—D. II. A cheap blue ink may also be prepared

readily, but has the disadvantage of some-

from soluble prussian blue by solution in water. A little mucilage of acacia may be added.

#### III.

Boil for an hour, let stand 2 or 3 days, and strain.

# Ink, Blue-Green, Non-Copying.

I.				
Phenol blue,	3 F (coal	tar dye)	gr. 4	8
Aniline gree	n D		gr. 6	0
Water			fl. oz. 2	8
Nutgall ink	body II		fl.oz. 3	8
Carbolic acid	1		fl.dr.	1/2
Prepare like				
I.—D.				

# II.

Phenol blue, 3 F (coal tar dye)gr. 30
Aniline green, Dgr. 50
Tannin ink body IIfl.oz. 16
Distilled waterfl.oz. 25
Carbolic aciddrops 20
Sugargr. 40
Prepare like alizarin copying ink, No. II
6

# Ink, Eosin. (Scarlet Ink.—Coral Ink.)

Ec	sin A,	yellowish	 		gr.	144
Su	gar		 		gr.	288
Di	stilled	water	 	fl	.oz.	20

Mix the dye with 1 fluidounce of cold water, set aside for 2 hours, add the remainder of the water, in the hot condition, and the sugar, and stir until dissolved.—D.

#### II.

Eosin (water soluble)gr.	120
Alcoholfl.oz.	2
Mucilage of acaciafl.oz.	1
Water, enough to makefl.oz.	16

Dissolve the eosin in about 12 fluidounces of water, a small portion of this being poured hot upon the eosin contained in a bottle; next add the alcohol, and shake; finally, add the mucilage and enough water to make 16 fluidounces.

III. See also No. IV., "Red Copying Ink."

#### Inks, Glossy.

Ink may be made glossy by the addition of mucilage of gum arabic or of a solution pre- II.—D.

pared by heating a mixture of borax, 180 grains; shellac, 60 grains; sugar, 120 grains, and water, 16 fluidounces.

#### Ink, Gold.

I. This may be prepared by mixing equal parts of potassium iodide and lead acetate, placing them upon a filter, and then pouring on twenty times the quantity of boiling distilled water. As the filtrate cools the lead iodide separates in golden scales. After the filtrate has cooled the precipitate should be collected on a filter, washed with a little cold water, and rubbed up to an ink with mucilage of acacia. The ink must be shaken before using.

II. Reduce gold foil to powder by triturating in a mortar with honey or syrup, dilute with water, decant the liquid, wash the gold several times with water; dry and mix with mucilage of acacia.

# Ink, Green, Copying.

l.		
	Aniline green, Dgr.	114
	Sugarav.oz.	11/4
	Distilled waterfl.oz.	3
	Nutgall ink body Ifl.oz.	32
	Carbolic aciddrops	20

Prepare like alizarin copying ink, No. I.— D.

# II.

Į,		
	Aniline green Dgr.	70
	Glucoseav.oz.	11/4
	Distilled water fl.oz.	2
	Tannin ink body Ifl.oz.	25
	Carbolic aciddrops	15
	•	

Prepare like alizarin copying ink, No. II.

--D.

# Ink, Green, Non-Copying.

1.	
Aniline green, Dgr.	192
Waterfl.oz.	
Nutgall ink body IIfl.oz.	
Carbolic acidfl.dr.	
	/

Prepare like alizarin non-copying ink, No. I.—D.

# II. Aniline green, D. gr. 100 Tannin ink body II. fl.oz. 15 Distilled water fl.oz. 25 Carbolic acid drops 20

Prepare like alizarin non-copying ink, No. II.—D.

# Ink, Green, Writing.

1.	Methyl green, blu	ish (water		0.0
	soluble)		gr.	96
	Sugar		gr.	192
	Distilled water		11. OZ.	10/2

Mix the dye with 1 fluidounce of cold water, set aside for 2 hours, then add the remainder of the water, in the hot condition, and the sugar, and stir about until dissolved.

—D.

# Ink, Orange.

Aniline orangegr.	144
Sugargr.	288
Distilled waterfl.oz.	20

Lix the dye with 1 fluidounce of water, set aside for 2 hours; then add the sugar and the remainder of the water, in the hot condition, and stir until dissolved.—D.

#### Ink, Purple.

Aniline purplegr. 8	30
Alcoholfl.dr. 1	$^{2}$
Mucilage of acaciafl.dr.	
Water	7

This color is brilliant at first, but is liable to fade.

#### Ink, Red Aniline.

See " Eosin Ink."

A red, inclining to purple, is made by dissolving fuchsin (ordinary aniline red) in water in the proportion of about 25 grains to the pint. Solution may be more readily effected by first dissolving the color in a little alcohol (about 5 fluidrams), and then adding the water. A small proportion of gum arabic is sometimes added to give the ink more "body." Two fluidrams to the pint is sufficient.

Another good formula is the following:

Erythrosingr.	75
Waterfl.oz. 1	6

Thicken with gum arabic, and add a little boric acid or other preservative.

#### Ink, Red, Carmine.

,	
Carminegr.	192
Ammonium carbonategr.	192
Water of ammoniafl.oz.	
Mucilage of acaciafl.oz.	3
Distilled waterfl.oz.	13
	Ammonium carbonategr. Water of ammoniafl.oz. Mucilage of acaciafl.oz.

Mix the carmine and ammonium carbonate, dissolve in the ammonia water, and add the remaining ingredients.—D.

Carmineav.oz. Ammonia waterfl.oz.	1/2
Ammonia waterfl.oz.	1
Dextringr.	120
Waterfl.o2.	

Triturate the carmine, add to the ammonia and water, dissolve by agitation; add the dextrin, and again dissolve.

#### III.

Carmine	.gr.	128
Ammonia waterf		
Distilled waterf	.oz.	8
Gum arabicav	·oz.	3/4
	_	–H.

IV

Carmine	gr.	96
Animonia waterfl.	oz.	41/2
Mucilage of acaciafl.		
Waterfl.	OZ.	13

Dissolve the carmine in the ammonia and add the other ingredients.

#### V.

Carmine	gr. 240
Water of ammonia	fl.oz. 3
Glycerin	fl.oz. 3
Water, q s	fl.oz. 8

Rub the carmine into a fine powder in a wedgewood mortar; make a paste with and dissolve in the water of ammonia, and then add, with constant trituration, the glycerin. Transfer to a porcelain capsule, and heat upon a water bath until the liquid is entirely destitute of ammoniacai odor; cool and add the water. The entire removal of the ammonia gas requires the constant stirring of the liquid with a glass rod, and rather lengthy heating.

This should be diluted with water.

Many of the carmine inks of the market are really eosin inks. See "Eosin Ink."

#### Ink, Red, Cochineal.

Cochineal, powdergr.	490
Potassium carbonateav.oz.	
Cream of tartarav.oz.	63/
Potassa alumgr.	196
Mucilage of acaciafl.oz	21/2
Alcohol	
Oil of clovesdrops	15
Distilled waterfl.oz	. 22

Macerate the cochineal and potassium carbonate with 19 fluidounces of water in a flask for 2 days; then add the cream of tartar and alum, heat on a water bath until all the carbonic acid gas is expelled, add the alcohol and filter. Wash the filter with 1 fluidounce

of water, and to the filtrate add the mucilage Ink, Red, Non-Copying. and the oil.

Writing with cochineal ink is very permanent.-D.

# Ink, Red, Copying.

Ponceau, R. R. (coal tar dye) gr.	152
Sugar av.oz.	1 1/4
Distilled waterfl.oz.	3
Nutgall ink body Ifl.oz.	32
Carbolic aciddrops	20
Prepare like alizarin copying ink, No	o. I.—
).	
r. \	

D II

Ponceau R. R. (coal tar dye)gr.	112
Glucoseav.oz.	11/4
Distilled waterfl.oz.	2
Tannin ink body Ifl.oz.	25
Carbolic aciddrops	
•	

Prepare like alizarin copying ink, No. II. —D.

TIT.

Logwood extract solution, av.oz.	24
Sulphuric acid, concentrateddrops	40
Aluminium sulphategr.	700
Oxalic acidgr.	700
Potassium carbonategr.	700
Potassium bichromategr.	52
Carbolic acid drops	20
Distilled watersuffic	ient

Heat the extract solution with the acid on a water bath for 15 minutes. In the meantime dissolve the aluminium salt in the water at a moderate heat; add the potassium carbonate, stir until there is no further evolution of carbonic acid gas, then add the oxalic acid, stir until all the precipitate is dissolved and there is no further evolution of gas, and now add the potassium bichromate, and dissolve. Incorporate the latter solution with the extract mixture by pouring the former slowly into the latter, continue the heat for 15 minutes more, add enough water to make the liquid weight 40 av.ounces, and finally, add the acid.

This ink is of a handsome red color; it writes red, and the writing speedily darkens. It is the best copying ink here mentioned. -D. IV.

Eosin A, yellowishgr. 240
Sugargr. 288
Distilled waterfl.oz. 20
Mix and dissolve without heat.—D.

١.	Ponceau R. R. (coal tar dye)gr.	192
	Waterfl.oz.	28
	Nutgall ink body IIfl.oz.	38
	Carbolic acidfl.dr.	
	Prepare like alizarin non-copying in	k, No.

I.--D.

1 1	.•	
	Ponceau R. R. (coal tar dye) gr.	100
	Tannin ink body II fl.oz.	
	Distilled waterfl.oz.	
	Carbolic acidgr.	20
	Sugargr.	40

Prepare like alizarin non-copying ink, No. II.-D.

# Ink, Red.

In addition to the red inks previously mentioned, the following may also be of service:

		11/
Pernambuco wood		
Alum	gr.	480
Acacia	gr.	480
Tin muriate (crystals)		
Diluted acetic acid	fl.oz.	16
Distilled water	fl.oz.	16

Mix the wood, water and acid, boil together until 24 fluidounces remain, add the alum; evaporate to 16 fluidounces, strain, add the acacia, dissolve, and then add the tin crystals.

II.

Brazil woodav.oz.	
Water	; 1/
Tin chlorideav.oz.  Mucilage of acaciafl.dr. 1	$\frac{72}{1/2}$

Boil the wood and water, strain, add the tin chloride, evaporate to 16 fluidounces, and add the mucilage.

#### Ink, Scarlet.

The so-called scarlet inks are usually the same as the eosin inks.

#### Ink, Silver.

This may be prepared from silver leaf by a process similar to making gold ink from gold

#### Ink, Vanadium.

Vanadium tannate was first proposed for use as a writing ink by Berzelius, because the writing is not affected by acids, but the high price of vanadium salts was a great obstacle to its introduction. Since these salts have been more largely prepared for use in the manufacture of aniline black and other dyes, vanadium ink has again been proposed. The following formula may be used:

Tannic acidgr.	480
Ammonium vanadate gr.	19
Waterfl.oz.	11

Dissolve the acid in 10 fluidounces of water and the vanadate in 1 fluidounce of water, and mix the solutions.

This ink flows with a deep-black color from the pen, without spreading or striking through the paper, although it contains no gum. has a pleasant gloss, cannot be copied, dries quickly, and, even if the writing is laid in water for 24 hours, does not change its black color. It is very useful for writing addresses of letters, postal cards, etc., when used fresh. Dilute acids do not alter it, but solutions of chlorinated potassa (or soda) bleach it com-After a few weeks the tint of the ink begins to change, writing executed with it becomes lighter and somewhat yellowish, and in about 3 months the change is completed, when it has a fox-yellow tint. writing is still plainly legible, however, and cannot be removed either by water or by acids.

## Ink, Violet, Copying.

Phenol blue, 3 F (coal tar dye)gr.	38
Ponceau R. R. (coal tar dye)gr.	57
Sugarav.oz.	
Distilled waterfl.oz.	~ s′ <sup>⊤</sup>
Nutgall ink body Ifl.oz.	32
Carbolic aciddrops	20
Prepare like alizarin copying ink, No.	

D II

II.
Phenol blue, 3 F (coal tar dye)gr. 28
Ponceau R. R. (coal tar dye)gr. 42
Glucose
Distilled waterfl.oz. 2
Tannin ink body I
Carbolic aciddrops 15
Prepare like alizarin copying ink, No. II
D.

III. Prepare like red copying ink, No. III., but decrease the amount of sulphuric acid to 27 drops and increase the potassium bichromate to 60 grains.

This ink is of a dark violet color, and the

Methyl violet, 3 B (coal tar	
dye)gr.	200
Sugargr.	
Oxalic acidgr.	
Distilled waterfl.oz.	

Dissolve the dye by the aid of heat in the water, add the other ingredients, and again dissolve.—D.

## Ink, Violet, Non-Copying.

Phenol blue, 3 F (coal tar dye)gr.	$4\overset{\circ}{8}$
Ponceau R. R. (coal tar dye)gr.	
Waterfl.oz.	28
Nutgall ink body IIfl.oz.	38
Carbolic acidfl.dr.	1/2

Prepare like alizarin non-copying ink, No. I.-D.

## II.

Phenol blue, 3 F (coal tar dye)gr.	30
Ponceau R. R. (coal tar dye)gr.	40
Tannin ink body IIfl.oz.	16
Distilled water fl.oz.	
Carbolic acid drops	20
Sugargr.	40

Prepare like alizarin non-copying ink, No. II.—D.

#### Ink, Violet, Writing.

Methyl violet, 3 B (coal tar dye)gr.	96
Sugargr.	
Oxalic acidgr.	20
Distilled waterfl.oz.	191/2

Mix the dye with 1 fluidounce of cold water, set aside for 2 hours, then add the remainder of the water, in the hot condition, and the other ingredients, and stir about until dissolved.-D.

#### Ink, White.

White inks, for writing on colored surfaces, consist either of a white mineral suspended in a viscid medium, or of chemicals affecting the coloring material in the paper. This, in the case of ultramarine, is an oxalic acid solution, or hydrochloric acid, according to this formula:

#### Ink for Blue Paper.

Hydrochloric acidfl.d	r. 1
Mucilager	n. 30
Waterfl.d	r. 7

To produce white writing on photographs, writing and copies are of the same hue. -D. | iodine is employed, as in the next formula:

#### Ink for Silver Prints.

lodinegr.	15
Acaciagr.	15
Potassium iodidegr.	
Waterfl.oz.	1

For preparations of the first order take lightest zinc white, or lead white, or magnesium carbonate, or freshly precipitated barium sulphate, or starch (all in an impalpable powder) and suspend in a diluted solution of gum arabic, dextrin, or tragacanth. The mixture requires shaking from time to time to keep the pigments from separating. The "ink" may be preserved by addition of oil of cloves or other antiseptic to prevent decomposition of the mucilage.

## Ink, Branding.

By this is meant an ink used for marking boxes, bales, packages, etc., by means of a small brush. They are frequently termed marking inks.

I.

Shellac av.oz.	2
Boraxav.oz.	2
Gum arabicav.oz.	2
Waterfl.oz.	25
Pigmentsufficie	nt

Boil the borax and shellac in the water until they are dissolved, add the gum arabic and allow to cool. Add water to complete 25 fluidounces and then stir in the pigment, using either venetian red, lampblack, ultramarine, or prussian blue. Black is improved by the addition of blue. Green may be produced from a mixture of blue and chrome yellow.

II.

Extract of logwoodav.oz.	2
Potassium bichromategr.	
Water, hotgal.	1

Dissolve the extract of logwood in part of the water, and the potassium bichromate in another portion; mix the two solutions and allow to stand for 1 or 2 weeks.

III.

Prussian blueav.oz. 2
Lampblackav.oz. 1
Gum arabic
Glycerin sufficient

Triturate together the dry powders and then make into a suitable paste with glycerin.

IV. Mix boiled linseed oil with sufficient lampblack or other suitable pigment.

#### Ink, Diamond.

These are liquids used for etching glass. Commercial strong hydrofluoric acid often gives negative results, because when applied in its pure state, it produces such a smooth corrosion of the glass that it may elude superficial inspection. The most common method consists in mixing ammonium fluoride with precipitated barium sulphate and decomposing with sulphuric acid, and is as follows:

Α	mmor	nium	fluor	ide.				.av.oz.	1
$\mathbf{B}$	arium	sulp	hate.					.av.oz.	3
S	ulphu	ric a	cid		 			sufficier	nt

Rub the two solids together, transfer to a platinum, lead or gutta-percha vessel, and add sufficient sulphuric acid to produce a cream-like paste. Operators must be cautioned against inhaling the exceedingly acrid vapors of hydrofluoric acid. Apply with a quill or camel's-hair pencil.

A second formula is as follows:

Ammonium fluorideav.oz. 2
Barium sulphateav.oz. 2
Hydrofluoric acid, fuming sufficient

Mix the 2 salts in a porcelain mortar, transfer to a platinum or lead vessel, and by means of a platinum wire stir in enough of the àcid to make a thin paste.

Writing may be performed with a steel pen, allow it to remain for one-half hour, and then wash off with water. To make etching more visible, rub in a little printer's ink.—D.

#### Ink, Enamel.

An ink, or rather varnish, for writing labels which are intended to resist the action of acids, etc., may be prepared as follows:

Shellacav.oz.	1
Borax	11/2
Nigrosin, water—solubleav.oz.	
Tannic acidgr.	15
Picric acid gr	. 5
Ammonia waterfl.dr.	12
Watersufficie	ent

Dissolve the shellac and borax in 15 to 20 fluidounces of water by the aid of heat, and filter hot; to the filtrate add the nigrosin, acids and ammonia, and then enough water to reduce the mixture to the required dilution.

The ink should be of such consistence that it will readily flow from the pen.

#### Ink Erasives.

Fresh ink spots are removed with comparative ease; old spots, especially after passing repeatedly through the laundry, are usually extinguished with considerable difficulty. The ink erasives are intended for the removal of ink spots from paper as well as from fabrics.

Τ.

The following composition will remove ink or writing fluids from paper, cloth, etc.:

#### No. 1.

Citric acidav.oz.	2
Waterfl.oz.	16
Saturated aqueous solution of	
boraxfl.oz. 3 or	4

Dissolve the acid in the water and add the borax solution.

No. 2.

Chlorinated limeav oz.	6
Waterfl.oz.	16
Saturated aqueous solution of	
boraxfl.oz. 3 or	4

Mix the lime and water, shake well, set aside for 1 week in a well stoppered bottle, decant the clear liquid, and add the borax solution.

This composition is used by saturating the ink spot with solution No. 1, removing excess of liquid with a blotter, and then applying solution No. 2. When the stain has disappeared, apply the blotter, and wash the spot by the alternate use of clear water and blotting paper. The above composition, we believe, is patented.

Ink destroyed in such a manner cannot well be brought to view again by chemicals. Tannic or gallic acids will sometimes restore obliterated writing.

H.

Take of chlorinated lime 4 av. ounces, thoroughly pulverized, and 32 fluidounces distilled water. Shake well and set the mixture aside for 24 hours, in order to dissolve the lime, then strain through a cotton cloth, after which add 2 fluidounces of acetic acid to every pint of chloride of lime water.

The eraser is used by reversing the penholder in the hand, dipping the end of the reversed penholder into the fluid and applying it without rubbing, to the word, figure or blot required to be erased. When the ink

has disappeared absorb the fluid with a biotter, and the paper is immediately ready to write upon again.

III. Treat the stain with stannous chloride to reduce the ferric salt to the ferrous condition and then treat with oxalic acid solution.

## Inks, Hectograph.

These are inks which are used in connection with hectographs (See "Hectograph Masses"). As will be observed, they are made with coal tar, so-called aniline dyes.

## Ink, Hectograph, Black.

Dissolve nigrosin in water in the proportion of 1 of the former to 5 or 7 of the latter.

## Ink, Hectograph, Blue.

I.

Resorcin blue, Mgr.	50
Glycerinfl.dr.	
Alcoholfl.dr.	1
Acetic acid, glacialdrops	5
Distilled waterfl.dr.	7

Dissolve the dye in a mixture of the other ingredients by the aid of heat.—D.

H

Brilliant green, crystalsgr.	
Hoffmann's violet, 4 Bgr.	
Glycerin	

Mix the two coloring matters in a mortar, and reduce them, if in lumps, to a moderately fine powder. Transfer this to a tared flask, add the glycerin and water, and heat the flask on a water bath, frequently agitating, until the colors are dissolved. Then allow the flask to cool, replace it on the scale and restore the loss of water.

## Ink, Hectograph, Green.

Aniline green Dgr.	100
Acetic acid, glacialdrops	
Glycerin fl dr.	1 1/2
Alcoholfl.dr.	1
Distilled water fl.dr.	7

Dissolve the dye in a mixture of the other ingredients by the aid of heat. - D.

## Ink, Hectograph, Red.

—D.

II.       Aniline red       gr. 120         Alcohol.       fl.dr. 4         Acetic acid, diluted       fl.dr. 1         Water       fl.dr. 4
III.       Rosaniline
IV. A satisfactory ink can also be prepared by dissolving rosaniline in a cold saturated solution of oxalic acid.
Ink, Hectograph, Violet.         I.         Methyl violet, 3 B
Dissolve the methyl violet in the alcohol and water by the aid of gentle heat.—D.
II.  Aniline violet
Violet anilinegr. 40

# Dissolve by trituration. IV.

Hoffmann's violet, BBBgr.	120
Alcoholfl.oz. 2 or	3
Glycerinfl.dr.	
Waterfl.dr	4

Glycerin .....fl.dr.

Water .....fl.dr.

Dissolve the dye in the alcohol, add the other ingredients and evaporate on a water bath to 1½ av.ounces.

V.	
Aniline violetgr.	60
Distilled water, hotfl.dr.	7
Alcohol	1
Glycerinfl.dr.	1/2
Carbolic aciddrops	5

## Inks, Indelible.

All the inks under the heading of "Inks, Marking," are indelible and are intended for marking clothes. There are also a number of the formulas under the titles of "Inks, Stamping" and "Inks for Rubber Stamps," which are indelible and which differ from the marking inks only in the fact that the former are applied with a pen and the latter with a stamp.

## Inks, Invisible or Sympathetic.

Invisible or sympathetic inks are fluids used for writing purposes, the characters made with them being invisible, but becoming visible upon the application of heat or of some suitable reagent. Writing made with those inks which becomes visible upon the application of heat, again becomes invisible upon cooling; on the contrary, writing developed by chemical action remains permanent.

I. Cobalt chloridegr.	150
Glycerinfl.dr.	
Waterfl.oz.	

Mix and dissolve the ingredients.

The characters traced with this ink become blue on gently heating the paper.—D.

#### II.

Linseed oil m.	25
Water of ammoniafl.oz.	1
Waterfl.oz.	5

To make the writing or the drawing appear which has been made upon paper with the ink, it is sufficient to dip it into water. On drying, the traces disappear again, and reappear by each succeeding immersion.

The mixture must be agitated each time before the pen is dipped into it, as a little of the oil may separate and float on top, which would of course, leave an oily stain upon the paper.

III. Write with tincture of iron chloride, diluted with 10 parts of water, and develop with a blotter moistened with a solution of tannin or decoction of nutgalls or strong tea.

This may be reversed by writing with a decoction of nutgalls and developing with the blotter moistened with tincture of iron.

The characters when developed are black. Instead of using nutgalls to develop the iron, potassium or ammonium, sulphocyanide solution may be employed. The writing will then appear red.

IV. Write with a solution of ferrocyanide of potassium in 20 parts of hot water. Develop with a blotter moistened with a solution of iron chloride.

This operation may, like the preceding, be reversed.

The characters when developed are blue.

V. Copper sulphate and ammonia, equal parts, dissolved in water. The writing turns yellow when exposed to heat.

The ammonium chloride may be replaced by potassium bromide.

Lemon juice, or the mineral acids diluted, solution of salt, saltpetre and many other substances, when the writing is exposed to heat, turn yellow or brown.

VI. A weak solution of nickel chloride, mixed with chloride of cobalt, turns a beautiful green when exposed to heat.

VII. Copper, dissolved in muriatic acid and diluted (or a solution of copper chloride), becomes yellow when the writing is heated, and disappears when cold.

VIII. A solution of cobalt acetate, to which a little nitrate of cobalt is added, becomes rose color when the writing is heated, and disappears when cold.

## Inks, Marking.

These are the inks intended for marking clothes prior to passing them through the laundry.

In order to distinguish from branding inks, read article under "Inks, Branding." See also "Inks, Indelible."

# Ink, Black, Marking.

1.		
Silver nitrate	av.oz.	5
Gum arabic	av.oz.	3
Stronger water of ammoni	a fl.oz.	12
Lampblack	gr.	180

Dissolve the silver nitrate and gum arabic in the ammonia water, by frequent agitation in a dark amber bottle, then incorporate the lampblack with this solution by trituration.

In using, write with a quill pen, allow the writing to dry, then pass a hot iron over it.

If the quantity of gum be inreacsed to 5 fluidounces, the ink may be spread on a plate of glass and a rubber stamp may then be used to apply it. When the mark is dry, it hould be ironed as in the preceding case.—D.

II. The indelibility of this ink depends on the fact that when potassium bichromate and gelatin come together, particularly in the form of a thin film, in the presence of daylight, the film becomes insoluble in hot or cold water.

Gelatingr.	2
Potassium bichromate	2 10
Waterfl.oz.	1

Dissolve the gelatin and the nigrosin in most of the water, and the potassium bichromate in the remainder. Mix the two solutions in an amber-colored bottle.

If it is found that the ink "gums" in the pen, the quantity of gelatin and bichromate may be somewhat reduced.

The ink, when properly made, cannot be entirely removed by hot or cold water, acids or alkalies.

III.

Asphalt, in	pieces	.av.oz. 2
Benzol	-	fl. oz. 8
Coal tar		.av.oz. 8

Dissolve the asphalt in the benzol and add the coal tar.

In using the ink, employ an ordinary pen, and if it should be too thick, thin it with a little oil of turpentine. This ink does not spread, and requires no heating. It never fades, and it is not affected by anything.

IV

Silver nitrategr.	240
Infusion of nutgallsfl.dr.	
Acacia	
Distilled water	

Dissolve the silver nitrate in 4 fluidounces of water. In the remainder of the water dissolve the acacia and add the infusion. Then mix the two liquids. The infusion of nutgalls should be made by pouring 4 fluidrams of boiling distilled water upon 15 grains of powdered nutgalls.

This is Woodhouse's indelible ink.

v.

•	
Shellacg	r. 480
Boraxg	
Gum arabicg	
Lampblack sufficient to	
Waterfl.oz	

Dissolve the borax in 9 fluidounces of water, and boil with the shellac until nearly all is dissolved; filter, and add the gum, previously dissolved in 2 ounces of water. Then color with sufficient lampblack.

VI. The following is Boettger's formula for an indelible marking ink:

Aniline blackgr.	60
Acacia	90
Hydrochloric acid, concentrated.fl.dr.	1
Alcoholfl.oz.	
Waterfl.oz.	31/2
	/-

Triturate the aniline with the alcohol and

acid mixed, then add a warm solution of the gum in the water.

This ink does not act upon steel pens, and is indestructible by strong acids or alkalies.

If the deep-blue liquid first produced is diluted (instead of with solution of acacia) with a solution of 75 to 90 grains of shellac in alcohol, a varnish is obtained which may be used to apply a jet-black coat to wood, metal, or rubber.

#### VII.

Silver nitrateav.oz. 1
Ammonia water, 10 per centsufficient
Acaciaav.oz. 1
Sodium hydrate gr. 285
Soluble black anilinegr. 190
Distilled waterfl.oz. 6

Dissolve the silver nitrate in 3 fluidounces of distilled water and add 2 fluidounces of ammonia water, or sufficient to dissolve the brown precipitate formed at first. To this liquid add the acacia and sodium hydrate, first dissolved in the remainder of the water; heat gently for 10 minutes, and finally, dissolve in the mixture the aniline.

#### VIII.

Aniline oilav.oz.	17
Potassium chlorateav.oz.	1
Hydrochloric acid, purefl.oz.	
Copper chloride, puregr.	
Distilled waterfl.oz.	

Mix the aniline oil, potassium chlorate, and 5 fluidounces of distilled water in a capacious flask and heat on a water bath to a temperature of 80 to 90 degs. C., until the chlorate is completely dissolved. Now add one-half the acid, heat again until the liquid begins to darken; to the liquid add the copper chloride dissolved in the remainder of the water, and, finally, add the remainder of the acid, and heat again on a water bath until the mixture has acquired a fine red-violet color. Set the mixture aside for several days in a well-stoppered vessel and decant the clear liquid from the trifling precipitate.

This ink must be applied by means of a quill—a steel pen is quickly corroded by it. It is suited only for marking fabrics made of vegetable fibers (linen, cotton, etc.), and cannot be used on wool or silk. Writing made with it appears at first pale reddish, turning green on exposure to light, and, when washed

with water containing soap or alkali, changes to deep black.

This ink may also be used with a rubber stamp as a stamping ink.—D.

## Ink, Marking, Blue.

1.		
	Silver nitrategr.	240
	Stronger ammonia waterfl.dr.	12
	Sodium bicarbonategr.	240
	Copper sulphategr.	120
	Mucilage of acaciafl.oz.	
	Distilled water, enough to make. fl. oz.	

Dissolve the silver salt in the ammonia, and the soda and copper salt in a portion of the water; mix the two solutions and add the mucilage and the remainder of the water. If the ammonia water mentioned above is insufficient for solution, more should be added.

# Ink, Marking, Red (Crimson or Carmine.)

l.		
	Silver nitrategr.	120
	Sodium carbonate, puregr.	180
	Tartaric acidgr.	40
	Stronger water of ammoniafl.dr.	4
	Carminegr.	$1\frac{1}{2}$
	Sugargr.	
	Gum arabicgr.	
	Distilled watersuffice	

Dissolve the silver nitrate and sodium carbonate separately in 16 fluidounces of distilled water, and mix the solutions. Wash the precipitate by decantation with 32 fluidounces of water three times; collect on a filter, and wash with a fourth pint of water; drain well; transfer the precipitate to a mortar and rub up with the tartaric acid; when effervescence ceases, add the ammonia (in which the carmine has been dissolved), then the sugar and gum (previously made into a cream with water). Finally, make up to 16 fluidounces with distilled water.

#### II.

Silver nitrategr. 120
Tartaric acidgr. 120
Ammonia waterfl.oz. 1
Gum arabicgr. 180
Sugargr. 120
Carminegr. 5
Distilled water, enough to make. fl. oz. 4

not be used on wool or silk. Writing made with it appears at first pale reddish, turning green on exposure to light, and, when washed solutions, and add the ammonia. Dissolve

the gum and sugar in water, triturate this with the carmine, add the first solution, mix well, and add the remainder of the water.

## Ink Stains, Indelible, to Remove.

Inasmuch as these inks usually contain silver, the staining is caused by the latter. One method of removing the stain consists in soaking in solution of common salt, which forms silver chloride, and then washing with ammonia.

Another method recommended is to treat the stains with iodine, and then with potassium iodide solution.

## Ink, Stamping. (Rubber-Stamp Inks.)

Some of these inks are used as marking inks for clothes; in general they are intended for stamping paper. See also "Inks, Metal Stamp."

Mix the dye and dextrin, dissolve by the aid of a water bath in the water, add the glycerin, and replace the water lost by evaporation.

Other colors are produced by substituting for the blue any one of the following:

Methyl violet B, violet colorav.oz.	3/4
Diamond fuchsin I, red colorav.oz.	3/4 1/2
Methyl green, yellowish-green	·
colorav.oz.	1
Vesuvin B, brown colorav.oz.	
Phenol black Bav.oz.	3/4
Eosin, BBN red colorav.oz.	3/4
-I	).

	-1	<i>)</i> .
II. ~		
Sodium carbonate	av.oz.	1
Glycerin	.fl.oz.	3
Acacia		
Silver nitrate	av. oz.	1/2
Ammonia water	.fl.oz.	1
Venice turpentine	av.oz.	1/2
•		/-

Triturate the sodium carbonate, gum arabic and glycerin together. In a separate flask dissolve the silver nitrate in the ammonia water, mix the solution with the triturate and heat to boiling, when the turpentine is to be added with constant stirring. After stamping, expose to the sunlight, or use a hot iron. The quantity of glycerin and gum arabic may be varied to suit circumstances.

т		

Borax	av. oz. 1
Shellac, bleached	av.oz. 1
Mucilage of acacia	
Ultramarine blue	
Water,	/-
Glycerin	of each sufficent

Dissolve the borax in 10 fluidounces of water, heat to boiling, add the shellac, and dissolve. Now triturate the mucilage with the ultramarine, add this to the shellac solution, and then enough of equal parts of water and glycerin to make 12 fluidounces.

IV. Take of the finest quality of lampblack and rub into a smooth paste with a sufficient quantity of glycerin; or take a good black printer's ink and thin the same with kerosene.

## Ink, Stamping (for use with metal stamps).

Rubber-stamp inks usually have glycerin as a vehicle; metal-stamp inks may have oil or glycerin as a vehicle, and hence the preceding rubber stamp inks may be used as metal-stamp inks, or the formulas given below. The formulas mentioned are of two varieties, one containing the coloring matter in suspension, the other in solution.

I. Mixture of pigments with oil:

A	۸.	
	Ultramarine blueav.oz.	1
	Olive oilfl.oz.	3
	Mix intimately.	

D

J.		
Prussian blue	av.oz.	1/9
Ultramarine blue	av.oz.	1/1
Olive oil	fl.oz.	4

Prepare like the preceding. The prussian blue alone does not mix readily with oil, and nence the ultramarine is added.

С.	
Verdigrisav.oz.	21/2
Oleic acidfl.oz.	1/2
Olive oilfl.oz.	7

Prepare like the preceding.

D.	
Cinnabar or vermillionav.oz.	4
Olive oilfl.oz.	6

Prepare like the preceding.

## II. Solution of dye in oil:

Bordeaux red aniline, oil solublegr.	72
Scarlet red aniline, oil solublegr.	72
Oleic acid, crudefl.dr.	
Castor oilfl.oz.	

Mix the dyes intimately with the acid, gradually incorporate the oil, and heat the whole to 40 degs. C., agitating constantly meanwhile.

This makes a nice red color.

#### В.

Aniline blue, oil solublegr. 1	44
Oleic acid, crudefl.dr.	5
Castor oilfl.oz.	10

Prepare like the preceding.

## C.

Aniline violet, oil solublegr.	144
Oleic acid, crudefl.dr.	5
Castor oil	10

Prepare like the preceding.

#### D.

Aniline black, oil solublegr.	240
Oleic acid, crude fl.dr.	6
Castor oilfl.oz.	10

This mixture stamps blue-black.

#### E.

Aniline blue, oil solublegr.	120
Aniline lemon yellow, oil soluble.gr.	72
Oleic acid, crudefl.dr.	5
Castor oilfl.oz.	10

Prepare like the preceding.

This mixture stamps green.—D.

#### Ink, Stencil.

As a stencil ink, use No. I., Branding Ink, and increase the amount of pigment.

## Ink, Typewriter.

Typewriter ink is of two kinds, one being made with petrolatum and lampblack or similar pigment, the other with glycerin and alcohol and aniline color.

In making the first kind, put some best-quality petrolatum into a suitable vessel, and melt it by placing the vessel on a fire; then put in as much lampblack or bone black as the petrolatum will take up without becoming granular. To effect this incorporation the black pigment should be put in a little at a time, and the whole thoroughly stirred while making the additions. Be careful not to let the petrolatum be in excess, as it will cause

the print to have a greasy outline; while on the other hand, if the pigment be in excess, the print will not be clear. Ordinary bone black does not make a good product; purified animal charcoal will do better.

When a proper mixture of petrolatum and pigment has been made, remove the vessel from the fire, and while it is cooling mix equal parts of petroleum, benzin and oil of turpentine, and in this mixture put the black petrolatum compound, mixing in a little of the other, with constant stirring, so as to effect a thorough combination, and the petrolatum compound becomes dissolved. The quantity of the volatile solvent should be sufficient to render the fluid ink of the consistence of oil paint: the result will be a good, permanent black ink, which will not be rubbed off with water like aniline inks. For colored inks of this class use prussian blue, red lead, chrome yellow; and for inks of the aniline class use these dyes dissolved in equal parts of alcohol and glycerin; thus, for a black aniline ink dissolve one-half av-ounce of aniline black in 13 fluidounces alcohol, and then add the glycerin. Ink the ribbon in the usual way.

Having prepared the ink, proceed to ink the ribbon; the secret of success lies in the proper application of the ink to the ribbon. Thus: wind the ribbon on a piece of cardboard, spread on a table several layers of newspapers, then unwind the ribbon in such lengths as may be most convenient, and lay it flat on the paper; apply the ink, after well shaking it, by means of a soft brush, and rub it well into the interstices of the ribbon with a toothbrush. Hardly any ink should remain visible on the surface.

For inking typewriter ribbons the following process is also recommended: Into 2 fluidounces of any aniline writing ink put a teaspoonful of mucilage of acacia, and a teaspoonful of brown sugar, warm the mixture, and immerse the ribbon from the typewriter long enough for it to become well saturated. When dry, spread the ribbon on a board and brush it well with glycerin. Should there be too much color in the ribbon, press it out between absorbent papers with a warm flatiron; or if too dry, brush it again with glycerin.

The secret of the ribbon giving out its

color is the glycerin, and if there is body enough in the color there is no danger that it cannot be made to work well. A ribbon so prepared is not affected by the dryness or humidity of the atmosphere.

It is necessary that the ribbon should retain a certain degree of moisture, for the gum and sugar make it dry and harsh, so the glycerin coating is put on; but there is danger of smearing the paper with too much moisture, or a wrinkled surface, and the ironing obviates this.

Users of the typewriter should so set a fresh ribbon as to start at the edge nearest the operator, allowing it to run back and forth with the same adjustment until exhausted along that strip; then shift the ribbon forward with the width of one letter, running until exhausted, and so on. Finally, when the whole ribbon is exhausted the color will have been equally used up, and on re-inking the work will appear even in color, while it will look patchy if some of the old ink has been left here and there, and fresh ink applied over it.

The following formula may also be used to make a typewriter ink:

Transparent soapav.oz.	1
Glycerinfl.oz.	
Waterfl.oz.	12
Alcohol fl.oz.	24
Aniline color sufficie	ent

Dissolve the soap in the water and glycerin by the aid of heat; dissolve the aniline color in the alcohol and mix the solution. If the ink is too soft, add more soap.

For the aniline color, use any suitable coal tar dye.

## Insect Destroyers.

Insect destroyers are mentioned under the headings "Bedbug Exterminators," Croton Bug Exterminators," "Flea Exterminators," "Fly Exterminators," "Insecticides for Agriculturists," "Insects in Drugs," "Lice Exterminators," "Mosquito Essence," "Mosquito Pastilles," "Mosquito Powder," "Moth Essence," "Moth Paper," "Moth Powder," "Moth Species," "Roach Pastes," and "Roach Powders." Other insecticides are mentioned here:

1.	
Resin av.oz.	1
Oil of amber, crude fl.oz.	
Benzingal.	
II.	~
11.	

Sodium borosalicylate.....gr. 150
Decoction of quassia.....fl.oz. 16

This bitter solution, in Germany known as "liquor insecta fngans," is claimed to be one of the very best applications to ward off biting insects of any kind.

III. The following preparation is known as "tincture of insect flowers."

Make 10 fluidounces of tincture.

It may be applied as a preventive of insect stings; it may also be used, when mixed with an equal volume of alcohol, as a spray for the destruction of flies.—D. modified.

An ethereal tincture may be prepared in a similar manner, using spirit of ether as a menstruum. It is to be used like the preceding.—D.

A perfumed or compound tincture may be prepared as follows:

Eucalyptolav.oz.	80
Oil of anise	80
Camphorgr.	400
Coumarin gr.	
Tincture of insect flowersfl.oz.	

Mix, set aside for several days, and filter. IV. It has been proposed to abandon arsenic and its compounds as an insect destroyer, and to replace them with naphthalin, benzin, petroleum, carbon disulphide and chloroform. The latter liquid (which is not itself inflammable at ordinary temperatures) is capable of materially reducing, or altogether neutralizing the inflammability of the three liquids previously mentioned.

It is necessary to get the naphthalin into solution. For this purpose Hager recommends to mix 2 pounds of soft soap with 1 quart of boiling water, and to gradually add, under energetic agitation, 300 grains of oleic acid. If a sample of this is diluted and shaken with an equal volume of alcohol, and if it then still possesses an alkaline reaction, a little more oleic acid should be added, 150 grains being probably amply sufficient. The resulting product might be called "Mitigated Green Soap."

The parasiticide liquids may be prepared as follows:

## 1. Liquor Naphthalini Benzinatus:

Naphthalin av.oz.	3
Chloroformfl.oz.	
Benzin	11 1/2

Mix at a temperature between 18 and 20 degs. C., and shake until solution has been effected.

## 2. Liquor Naphthalini Sulphocarbonatus:

Naphthalinav.oz.	6
Chloroformfl.oz.	
Carbon disulphide fl.oz.	7

Prepare like No. 1.

For use in a more fluid form, either of these liquids is to be properly diluted; the following being a good formula:

Common family soap, dryav.oz.	1/2
Castile soap, dryav.oz.	1/2
Waterfl.oz.	18
Alcohol fl.oz.	11
Liquor naphthalini benzinatusfl.oz.	3

Dissolve the soaps in the water and alcohol, previously mixed, allow the liquid to become cold, and then add the naphthalin solution. Before using the liquid, shake it thoroughly.

If an ointment is required,  $8\frac{1}{2}$  av.ounces of petrolatum and  $1\frac{1}{2}$  av.ounces of ceresin are melted together, and before the mass sets 18 fluidounces of the liquor naphthalini benzinatus mixed with it.

If either of these is to be used as a parasiticide upon animals, it should be applied with a stiff brush, in quantity only large enough to moisten the skin or to render the hair or fur slightly glossy. Under all circumstances is it preferable to avoid using these compounds at night time, as accidents may occur by approach to or contact with flames.

#### Insects in Drugs.

Insects attack many drugs, and an excellent manner to destroy these insects, cs well as to prevent their ravages is to pour some chloroform into the container, which should be well closed. Upon exposure of the drug to the atmosphere, the chloroform will be volatilized.

#### Insecticides for Agriculturalists.

Under this heading are mentioned various mixtures which are useful to the farmer for

the destruction of various insects that infest and destroy plants.

## Riley Hubbard Kerosene Emulsion:

This insecticide acts by contact, and is applicable to all non-masticating insects (sucking insects, such as the true bugs, and especially plant lice and scale insects), and also to many of the mandibulate insects when the use of arsenites is not advisable. Kerosene emulsion may be made by means of various emulsifying agents, but the most satisfactory sbustances, and those most available to the average farmer and fruit-grower, are milk and soapsuds. In each of these cases the amount of emulsifying agent should be one-half the quantity of kerosene.

One of the most satisfactory formulas is as follows:

Kerosenegals.	2
Common soap or whale-oil soap.av.oz.	8
Watergal.	

Dissolve the soap in the water by the aid of heat, and add the solution boiling hot to the kerosene. Churn the mixture by means of a force pump and spray nozzle for 5 to 10 minutes. The emulsion, if perfect, forms a cream which thickens upon cooling, and should adhere without oiliness to the surface of glass. No free oil should rise from surface of the liquid, as this would injure the foliage.

For use against scale insects, dilute one part of the emulsion with 9 parts of water. For most other insects, dilute one part of the emulsion with 15 parts of water. For soft insects, like plant lice, the dilution may be carried to from 20 to 25 parts of water. For most insects the proper dilution is with 15 parts of water. This liquid should be applied by force, and enough used to thoroughly wet the insects.

The milk emulsion may be produced by the same methods as the above.

#### Cook's Kerosene Emulsion:

Soft soap.	 	 		.quart 2
Kerosene	 	 		.quart 1
Water	 	 	<i>.</i>	gal. 1

Boil the soap with the water until all is dissolved; remove from the fire, add the kerosene, and churn vigorously for 10 minutes to emulsify the kerosene.

This should be diluted with an equal bulk of cold water before using. It should be applied like the preceding.

One-half pound of hard soap may be substituted for the soft soap.

## Glaser's Tobacco Insecticide:

Soft or green soapav.oz.	2
Tobaccoav.oz.	
Fusel oilfl.oz.	21/2
Alcohol fl.oz.	$8^{'-}$
Watersufficien	t

Disselve the soap in 8 fluidounces of water; make about 8 fluidounces of infusion from the tobacco, mix the two liquids, add the remaining ingredients, and then enough water to make 1 quart.

This is to be sprinkled on the leaves of the infected trees.

#### Nessler's Tobacco Insecticide:

Green soapav.oz.	4
Extract of tobaccoav.oz.	
Animal oilfl.oz.	5
Alcoholfl.oz.	20
Waterfl.oz.	65

Dilute with 5 volumes of water before use.

## Tobacco Decoction:

Tobacco	leaves or stemsav.oz.	16
Water	gal.	3

Boil the tobacco with the water for one-half hour. Replace the water lost by evaporation and use without further dilution.

This is very effective against plant lice and soft caterpillars. Unlike the kerosene emulsion, it never burns the foliage. It is also an excellent fertilizer. It is especially recommended for indoor plants and small gardens.

The Resin Washes:

These insecticides act by contact, and also in the case of scale insects, by forming an impervious coating which effectually smothers the insects treated. These resin washes vary in efficacy according to the insect treated. Experience has shown that the best formula for the red scale (Aonidia aurantii Maskell) and its yellow variety (A. citrinus Coquillett) is as follows:

Resinav.lb.	31/2
Caustic sodaav.lb.	
Fish oilfl.oz.	8
Water to make gal.	20

The necessary ingredients are placed in a kettle and a sufficient quantity of cold water

added to cover them; they are then boiled until dissolved, being occasionally stirred in the meantime, and, after the materials are dissolved, the boiling should be continued for about an hour, and a considerable degree of heat should be employed, so as to keep the preparation in a brisk state of ebullitioncold water being added in small quantities whenever there are indications of the preparation boiling over. Too much cold water, however, should not be added at one time or the boiling process will be arrested and thereby delayed; but, by a little practice the operator will learn how much water to add so as to keep the preparation boiling actively. Stirring the preparation is quite unnecessary during this stage of the work. When boiled sufficiently it will assimilate perfectly with water, and should then be diluted with the proper quantity of cold water, adding it slowly at first, and stirring occasionally during the process. The undiluted preparation is pale-yellowish in color, but by the addition of water it becomes a very dark brown. Before being sprayed on the trees it should be strained through a fine wire sieve, or through a piece of Swiss muslin, and this is usually accomplished when pouring the liquid into the spraying tank, by means of a strainer placed over the opening, through which the preparation is introduced into the tank.

The preparing of this compound will be greatly accelerated if the resin and caustic soda are first pulverized before being placed in the boiler, but this is quite a difficult task to perform, and is unnecessary.

This insecticide may be applied at any time during the growing season.

A stronger wash is required for the San Jose scale (Aspidiotus perniciosus Comstock), and the following gives the best results:

Resinav.lb.	6
Caustic sodaav.oz.	29
Fish oilfl.oz.	15
Water, enough to make gal.	20

Place all the ingredients in a kettle and cover with water to the depth of 4 or 5 inches, boil briskly for about 2 hours, or until the compound can be perfectly dissolved with water. When this stage is reached the kettle should be filled up with water, care being taken not to chill the wash by adding large

quantities of cold water at once. It may be thus diluted to about 8 gallons, the additional water being added from time to time as it is used.

This preparation should only be applied during winter or during the dormant period. Applied in the growing season, it will cause the loss of foliage and fruit.

In the application of both these washes a very fine spray is not essential, as the object is not simply to wet the tree, but to thoroughly coat it over with the compound, and this can be best accomplished by the use of a rather coarse spray, which can be thrown upon the tree with considerable force.

#### For Subterranean Insects:

Recent experiments have shown the practical value of the resin compounds against the grape phylloxera, and they will also be applicable to the apple-root louse and other underground insects. The cheapest, and at the same time one of the most satisfactory compounds experimented with is the following:

Caustic sodaav.lb.	1
Resin av.lb.	8
Water to makegal.	10

Dissolve the soda over fire with 1 gallon of water, add the resin, and after it is dissolved, and while boiling add water (slowly) to make 10 gallons of compound. For use dilute to 100 gallons. Excavate basins about the vines 6 inches deep and about 2 feet in diameter, and apply to each vine 5 gallons. The result will be more satisfactory if the treatment is made early in the spring, so that the rain of the season will assist in disseminating the wash about the roots.

The kerosene emulsion made according to the formula given above is also applicable to certain underground insects in cases where it will not prove too expensive; as, for instance, the grape phylloxera, or where white grubs are infesting a valuable lawn. It may then be used in the proportion of 1 part of the emulsion to 15 gallons of water, applied liberally to the soil, and afterward washed down at frequent intervals with large quantities of water for several days. This can be done only where there is plenty of water at

hand, but will be found of great value in special cases.

In other cases carbon bisulphide may be used for specific and local underground forms. Nests of ants, for instance, may be destroyed by pouring an ounce of this substance into several holes, covering them with a wet blanket for 10 minutes, and afterward exploding the vapor at the holes with a torch. Against onion, cabbage, and radish maggots this substance may also be used, by punching a hole with a sharp stick at the base of the plant and pouring in a teaspoonful of the liquid, covering afterward with earth.

The Arsenites:

These poisons (paris green or london purple) are of the greatest service against all mandibulate insects, as larvæ and beetles, and they furnish the most satisfactory means of controlling most leaf-feeders, and the best wholesale remedy against the codling moth. Caution must be used in applying them on account of the liability of burning or scalding the foliage.

The poisons should be thoroughly mixed with water at the rate of from 1 pound to 100-250 gallons of water, and applied with a force-pump or hand spray-nozzle. In preparing the wash it will be best to first mix the poison with a small quantity of water, making a thick batter, and then dilute the latter and add to the reservoir or spray-tank, mixing the whole thoroughly. When freshly mixed, either london purple or paris green may be applied to apple, plum and other fruit trees (except the peach) at the rate of 1 pound to 150-200 gallons, the latter amount being recommended for the plum, which is somewhat more susceptible to scalding than the apple. White arsenic does little if any injury at the rate of 1 pound to 50 gallons of water. It has been shown, however, that when allowed to remain for some time (two weeks or more) in water the white arsenic acts with wonderful energy, scalding when used at the rate of 1 pound to 100 gallons from 10 to 90 per cent of the foliage. The action of the other arsenites remains practically the same, with perhaps a slight increase in the case of london purple.

With the peach, these poisons, when ap-

plied alone, even at the rate of 1 pound to 300 or more gallons of water, are injurious in their action, causing the loss of much of the

By the addition of a little lime to the mixture, london purple and paris green may be safely (applied at the rate of 1 pound to 125 to 150 gallons of water) to the peach of the tenderest foliage, or in much greater strength to strong foliage, such as that of the apple or most shade trees.

Whenever, therefore, the application is made to tender foliage, or when the treating with a strong mixture is desirable, lime water, (milky, but not heavy enough to close the nozzle) should be added at the rate of about 2 gallons to 100 gallons of the poison.

Pure arsenic, however, should never be used with lime, as the latter greatly increases its action.

With the apple, in spraying for the codling moth, at least two applications should be made-the first on the falling of the blossoms, the apples being about the size of peas, and the second a week or 10 days later-but the poison should never be applied after the fruit turns down on the stem, on account of the danger of the poison collecting and remaining permanently in the stem cavity.

For the plum curculio on the plum, cherry, peach, etc., two or three applications should be made during the latter part of May and the first half of June. In the case of most leaf-feeders, spray on the first indication of their presence.

The following formula may also be employed:

Paris green or london purple.	
av.oz. 3 to 4	
Fresh limeav.oz. 8	
Flour	
Watergal. 45	

Shake the lime in a gallon of water and rub till smooth; then strain and stir in the arsenite. Boil the flour to a thin paste. Dilute the arsenite with the necessary water, then add the flour paste and use. This is the standard remedy for all kinds of leaf-gnawing insects. A good sample of london purple is just as effective as paris green, and usually costs less. The lime is added to neutralize with the other powders. any soluble arsenic compounds, and the flour sional bug exterminators

is used to make the arsenite adhere better to the foliage.

Caution necessary in use.

The relative susceptibility of apple, plum and peach has just been indicated under the head of arsenical poisons, and these remarks apply equally well to the use of the kerosene emulsions. In the case of other plants thorough experiments are necessary, and all insecticides should be first used in comparatively high dilution. In general it may be said that tender young foliage is more susceptible, and must be carefully treated. Thin-leaved pilose plants are more readily injured, while thick leaved, glabrous species are least affected. Annual plants, such as cabbages and other garden vegetables, are more susceptible than perennials, but in the case of root crops, such as beets, turnips, radishes and potatoes, there is not the same need of caution as to damage to foliage. Damage to foliage is not shown at once, and, in case of rain following an application, another application should not be made for several days. Fruit trees should not be sprayed with arsenical poison before the blossoms fall, on account of the danger of poisoning honey bees.

## Insect Powders.

I.
Insect powderav.oz. 14
Quassia, fine powderav.oz. 6
II.
Insect powderav.oz. 14
Quassia, fine powderav.oz. 4
Naphthalin
III.
Insect powderav.oz 8
Borax, powderav.oz. 8
Oil of cedar
Oil of pennyrovalfl.dr. 2
Persian insect powderav.oz. 8
Boraxav.oz. 8
Sulphur av.oz. 4
Oil eucalyptus fl dr. 2
Mix. Excellent for cockroaches.
V.
Paris green
Plaster parisav.oz. 2
Borax, powderedav.oz. 12
German chamomileav.oz. 1

Powder the chamomile flowers and mix Used by profes-

#### Javelle Water.

Bicarbonate of sodiumav.oz.	
Chlorinated limeav.oz.	2
Water fl.oz.	16
Boil the soda in the water for a few	min

Boil the soda in the water for a few minutes, add the lime, and when cold, strain.

# Jeweler's Rouge. (Colcothar.—Crocus Martis.)

This is usually prepared by heating ferrous sulphate to a high temperature, but a simpler method is this:

Make a tolerably strong solution of ferrous sulphate, also one of oxalic acid, filter each; add the former to the latter, with constant stirring, let stand a few hours, collect the precipitate, wash it thoroughly with water, dry, and expose to the direct flame until there is no further change of color.

## Lacquer for Brass.

Before applying lacquer to brass it must be well cleaned. This may be done by immersing in a bath of strong caustic potash, followed, after rinsing, by a bath of dilute nitric acid. Rinse in water, rub dry with chamois, and place on hot iron plate, or on top of stove, until warm. Then apply the lacquer with a soft camel's-hair pencil, making all the strokes in one direction. Some little practice is necessary in order to apply the lacquer nicely.

Similar to the lacquers are the varnishes, both being resinous solutions, intended as protectives for metals, wood, etc.

## Lacquer for Brass, Dark.

H

Ι.		
	Turmeric, powdergr.	480
	Annatto, bestgr.	
	Saffron, Spanishgr.	120
	Shellac av.oz.	
	Alcohol	16

Digest the first 3 ingredients with the alcohol for 24 hours; then dissolve the shellac in the liquid, and strain.

ſ.	
Shellacav.oz.	21/2
Masticav.oz.	1/4
Sandaracav.oz.	
Aloesav.oz.	1/2
Turmeric, powder	11/4
Saffrongr.	60
Dragon's bloodav.oz.	1/4
Venice turpentinegr.	
Alcoholfl.oz.	25

Mix; macerate for several days, agitating occasionally, and filter.

#### III.

Seed lac av.oz.	
Turmeric, powderav.oz.	1
Dragon's blood, powderav.oz. Alcoholfl.oz.	1/4
Alcoholfl.oz.	16

Macerate a week, frequently shaking; decant or filter.

## Lacquer for Brass, Gold.

I.		
Orange shellac	.av.oz.	8
Alcohol	fl. oz.	12
Water		

Dissolve with heat. The solution is milky from the insoluble waxy portion of the shellac. Clarify by shaking with an ounce of precipitated chalk, or by shaking with an equal volume of benzin. The benzin will separate from alcohol of this strength, and may be poured off.

#### II.

Gum copalav.oz.	2
Shellacav.oz.	1
Boiled linseed oil,fl.oz.	2
Oil of turpentinefl.oz.	10

Melt the copal and shellac; add the linseed oil, remove the vessel from the fire, and gradually add the turpentine.

#### Lacquer for Brass, Red.

T.	
Alcohol fl. oz.	16
Dragon's blood av.oz.	1/2
Annatto av.oz.	$2\frac{1}{2}$
Sandaracav.oz.	. 4
Oil of turpentine fl.oz.	. 2

Macerate, with frequent agitation, for a week; decant and filter.

#### II.

•	•		
	Sandarac	av.oz.	6
	Mastic	av.oz.	3
	Balsam of copaiba	.av.oz.	1
	Venice turpentine	av.oz.	1 1/2
	Oil of turpentine	, .fl. oz.	2
	Absolute alcohol	.H.oz.	18
	Shellac	av. oz.	21/2
	Dragon's blood	.av.oz.	25
	Alcohol	.fl.oz.	25

Dissolve the sandarac, mastic, copaiba and venice turpentine in the absolute alcohol and oil, and mix with a filtered macerate of the dragon's blood and shellac in the alcohol.

III.	
Shellacav.oz.	5
Sandarac	2
Masticav.oz.	2
Gambogeav.oz.	1/2
Dragon's bloodav.oz.	1/2
Annatto av.oz.	1/2
Red saundersav.oz.	3/4
Venice turpentineav.oz.	11/2
Alcoholfl.oz.	25

Mix, macerate for several days, and filter.

## Lacquer for Leather, Black.

Shellacav.oz.	$1\frac{1}{4}$
Sandaracgr.	180
Mastic	
Venice turpentineav.oz. Alcoholfl.oz.	16

Mix and dissolve, and color deep black with nigrosin.

## Lacquer for Tin.

Alcohol	fl.oz.	16
Turmeric	av.oz.	1
Saffron	gr.	80
Dragon's blood		
Red saunders		
Shellac		
Sandarac	av.oz.	1/2
Mastic	av.oz.	1/2
Balsam of fir	av.oz.	1/2

Reduce the drugs to powder; mix all, macerate for 7 days, agitating occasionally, and filter.

#### Leather, Dyeing of.

See "Dyeing of Leather."

## Leather Polish, Blacking or Dressing.

See "Harness Blackings," "Shoe Grease," "Shoe Polish," "Shoe Dressing," and "Shoe Varnish."

#### Lice Exterminators.

Pharmacists should always avoid selling fish berries (cocculus), on account of their poisonous nature, while mercurial ointment is not always desirable. The following mixture is highly recommended for killing lice and similar vermin.

ί.	
	Boraxgr. 360
	Glycerin
	Decoction of quassia (1 in 5)fl.oz. 15
	Mix and dissolve.
	Apply to the head once daily

II.	
Naphthalinav.oz.	31/2
White wax or ceresinav.oz.	11/2
Cocoanut cilav.oz.	5 3/4
Petrolatumav.oz.	534
Oil of bergamotfl.dr.	11/2
Oil of clovesfl.dr.	11/2
Oil of cinnamonfl.dr.	11/2
Oil of lemonm.	50

Melt the fats, add the naphthalin, stir until the latter is dissolved, allow to cool, and incorporate the oils.—H.

Use like the preceding.

III. The following may be recommended, especially as a powder for exterminating lice on the body.

Sabadilla	powder			,			.av.oz.	16
Orris, po	wder						.av.oz.	4

## Linoleum, for Polishing.

I.	
Yellow waxav.oz	. 1
Carnauba waxav.oz	. 2
Oil of turpentine fl.oz	. 10
Benzin	

Melt the two waxes, carefully add the oil and benzin, and stir until solid.—D.

II.	
Yellow waxav.oz.	5
Oil of turpentinefl.oz.	11
Amber varnishav.oz.	5

Melt the wax, add the oil, and then the varnish.

Apply with a woolen rag.—D.

## Linseed Oil, Boiled.

Linseed oil.....gal. 1
Litharge, powder.....av.oz. 16

Mix, heat and simmer, with frequent stirring, until a pellicle begins to form; remove the scum, and when it has become cold and has settled, decant the clear portion.

#### Lubricants.

See "Axle Greases."

#### Magnesium Lights.

See under "Colored Fires" for such of the fires as contain metallic magnesium.

#### Marble, Cleansing and Polishing.

The marble of soda fountains may be cleansed with the following:

Sodium carbonateav.oz.	2
Chlorinated limeav.oz.	1
Waterfl.oz.	14

Mix well, and apply the mixture (magma

and liquid) to the marble with a cloth, rubbing well in, and finally rubbing dry. It may be necessary to repeat this operation.

The marble may now be polished by rubbing over with kerosene. This should, however, not be applied to white marble.

To remove grease stains from marble, it is recommended to cover the spot with a little pile of powdered talcum or some fine clay, saturating with benzin, and allowing to remain for some time.

Care must be taken in removing stains or discolorations in marble by the use of chemicals as the polish is liable to be injured. The following powder may be used for cleaning marble:

Common salt	.av.oz.	8
Pumice stone, powder	.av.oz.	4
Chalk, powder	.av.oz.	4

Mix, moisten with water, and rub over the marble, allowing to remain for some time; then wash off with soap and water.

## Marble Busts, to Clean.

First free from all dust and then wash with very weak hydrochloric acid; do not use soap.

#### Matches, Japanese.

These can be imitated very closely by making a mixture of 5 parts of lampblack, 11 parts of sulphur and 27 parts of gun powder, rubbing each substance separately to an impalpable powder, and making into a paste with absolute alcohol. Dip the sticks into the paste and let dry slowly, away from a fire. Another plan is to add only enough alcohol to make a doughy mass and to roll this out into a sheet about 1/4-inch thick. Cut into cubes and let dry as before. When desired for use, a cube is stuck into a split straw or splints of wood, and lit. burning a moment the material collects in a ball of molten matter which sends out sparks and scintillations much more brilliant than those of the matches.

#### Mathematics.

To calculate the area of a circle, square the diameter, and multiply by 0.7854 (or 11-14); or multiply the diameter by the circumference, and divide by 4.

To estimate the capacity of a can or any

vessel with straight sides (of the same diameter throughout), multiply the diameter by 0.7854, and this result by the height or depth of the can.

Where the vessel is larger at one end than at the other, and the sides are straight, add the ends together and divide by 2 to get the mean diameter, and proceed as before.

To measure the cubic contents of a barrel or keg with curved staves, add the diameter at the bung and that of the head (both measured from the inner rim of the staves) together; divide by 2 to get the mean diameter; multiply as before by 0.7854, and finally multiply by the length of the stave, taken from the inside of the heads. This latter process is not absolutely mathematically correct, but is so close as to answer for all practical purposes.

## Mildew, To Remove.

Mix equal parts of soft soap and starch, add half as much salt and the juice of a lemon. Apply this to both sides of the fabric and expose to sunlight.

# Molds for Taking Impressions.

т т

Dark	resin								.av.oz	. 12
Beef	tallow								.av.oz	. 8

For coarse work, such as architectural ornaments.

III. Flexible or elastic molds may be made of gutta percha softened in boiling water, and after being freed from moisture, pressed strongly against the objects to be copied.

The same can be produced by the use of gelatin or glue, which has been dissolved in sufficient hot water and passed over the object previously oiled.

# Mosquito Essence. (Mosquito Tincture or Lotion.)

in 5)......fl.oz. 7½

Diluted with from 3 to 6 parts of water.

This may be used as an application to the skin to prevent the attacks of mosquitoes. It may also be sprayed about the room to destroy or expel mosquitoes.

II.	
Carbolic acidfl.oz.	1
Oil of peppermintfl.oz.	2
Oil of camphor, volatilefl.oz.	2
Glycerinfl.oz.	2
Oil of tarfl.oz.	4
Olive oilfl.oz.	4
III.	

.11.		
Ammonia water	fl.dr.	11
Glycerin	il.oz.	2
Oil of pennyroyal		
Olive oil		
V.		
C - 1 - 1' 1 1	Д	-1

Carbolic acid	fl.oz. 1
Oil of pennyroyal	fl.oz. 2
Spirit of camphor	
Glycerin	
Oil of tar	
Lard oil	
Zara ominimi in	

This is an effective application for keeping flies and mosquitoes off horses.

V. The following may be sold as "essence of pennyroyal":

Oil of pennyroyalfl.oz.	1
Alcoholfl.oz.	

VI. The following, known as "fulvis capucinorum," is also useful:

Sabadilla
Cocculus indicusav.oz. 2
Parsley seedav.oz. 2
Anise seedav.oz. 2
Tobacco powder or snuffav.oz. 2

# Mosquito Pastilles.

I.	
Carbolic acidfl.dr.	6
Potassium nitrate, powderav.oz.	11/2
Insect powderav.oz.	5
Charcoal, powderav.oz.	10

Make a paste with powdered tragacanth and water and mold into pastilles.

#### II.

Thyme leaves, coarse powder.av.oz.	2
Lavender flowersav.oz.	2
Insect powderav.oz.	2
Potassium nitrate, powder av.oz.	134
Potassium chlorate, powdergr.	72
Tragacanth, powdergr.	175

Mix well, add sufficient water to form a mass; divide into pastilles, and dry.

## Mosquito Powder.

1.	
Eucalyptol (or oil of eucalyptus).fl.oz.	1
Talcum, powderav.oz.	2
Starch, powderav.oz.	17
Mix well and sift.	

This powder is to be rubbed into the exposed parts of the body to prevent the attacks of the insects.

The mixture may be rendered more effective by replacing 50 per cent or more of the starch by naphthalin.

#### II.

Oil of pennyroyalfl.dr.	4
Naphthalingr.	
Starchav.oz.	

Mix well and sift.

This is to be used like the preceding

## Moth Essence or Tincture.

i.		
	Oil of patchoulydrops	15
	Oil of mirbanefl.dr.	21/4
	Naphthalingr.	
	Carbolic acid, crystal gr.	300
	Camphorav.oz.	13/4
	Oil of turpentinefl.oz.	2
	Alcohoifl.oz.	27

Mix, allow to stand for several days, and filter.

In using, moisten blotting or other absorbent paper with this liquid; then lay the paper between the goods to be protected, and then pack securely in a suitable receptacle.—D.

#### II.

Capsicumav.oz.	31/2
Alcoholfl.oz.	36
Oil of turpentinefl.oz.	
Naphthalingr.	360
Camphor	360
Oil of clovesfl.dr.	21/2

Mix the capsicum with the alcohol and oil of turpentine, macerate for 8 days, filter; to the filtrate add the remaining ingredients, and dissolve.

This is to be used like the preceding.—D.

## Moth Paper.

I.												
	Naphthalin.									av.oz	. 4	4
	Paraffin wax.									av.oz	. :	8

Melt together, and while still warm paint with a rather broad brush upon unsized paper.—H.

II.	
Naphth	alinav.oz. 4
	c acidav.oz. 2
Ceresin	av.oz. 2
Melt to	gether, and spread the mixture b
means of	a wide brush upon unsized pape
loid unon	a hot surface care being taken the

means of a wide brush upon unsized paper laid upon a hot surface, care being taken that this is not done near a light or fire.

If it is desired to avoid the use of a hot surface, and thus preclude any possibility of

surface, and thus preclude any possibility of contact with light or fire, the mixture may be diluted with 1 fluidounce of alcohol. As this does not dissolve the ceresin, the mixture must be well stirred with the brush before each application.—D.

#### III.

Naphthalinav.oz.	4
Camphorav.oz.	
Ceresin	
Oil of mirbanem.	40

Prepare like the preceding.—D.

#### Moth Powder.

I.

Naphthalin	2
Patchouly herb	2

Reduce all to powder and mix well.

II.

Patchoulyav.oz. 5	
Valerianav.oz. 2½	
Camphor	
Napthalinav.oz. 1	
Scotch snuffav.oz. 1	
Orris rootav.oz. 2½	
Sumbul rootav.oz. 2½	
Oil of cassiafl.oz. 1	
Oil of eucalyptus fl.oz. 1	

Reduce the solids to fine powder and incorporate with the remaining ingredients. III.

Insect powderav.oz.	5
Camphorav.oz.	<b>5</b>
Colocynthav.oz.	5
Oil of lavenderfl.dr.	
IV.	

Naphthalinav.oz.	3
Capsicumav.oz.	3
Insect powderav.oz.	
Oil of rosemary fl dr	1

Reduce the capsicum and naphthalin to fine powder and mix well with the other ingredients.

## Moth Species.

Patchouly, cut	av.oz.	2
Rosemary, cut	av.oz.	4
Thyme, cut		
Sage, cut		
Naphthalin		
Oil of mirbane		
Oil of turpentine		
Alcohol		

Dissolve the naphthalin and oils in the alcohol by the aid of heat, and sprinkle the solution while hot upon the mixed herbs or leaves. Introduce the mixture into bags of suitable size and lay them between the clothes to be preserved, which should be firmly wrapped, securely packed, and kept in a cool place.—D.

## Mouse Destroyers.

See "Raticides" and "Phosphorus Pastes."

## Mucilages and Pastes.

l
L
F
l
L
}
3

Mix; allow the tragacanth to soften, and add more water, if desired. The mixture may be preserved by the addition of carbolic acid or of some essential oil, such as oil of cloves or wintergreen.

#### III.

17

Lime,	slakedav.oz	. 1
Sugar,	granulatedav.oz	. 4
Glue	av.oz	. 3
Water	fl.oz.	12

Dissolve the sugar in the water, add the lime, heat nearly to boiling, set aside for several days, decant the clear liquid, and in it dissolve the glue by the aid of a moderate heat.

IV. Take the curd of skim milk (carefully freed from cream), wash it thoroughly, and dissolve it to saturation in a cold concentrated solution of borax.

* *						
Dextrin,	vellow.	 	 		.av.oz.	4
Water	,		 		fl.oz.	8

Heat the water, add the dextrin gradually with constant stirring, and continue the heat

until the dextrin is dissolved. One fluidounce of acetic acid may be added to the water before heating, and to the solution may be added the same amount of alcohol to insure preservation. However, neither one is necessary if carbolic acid or an essential oil, like oil of cloves or wintergreen be added. This mucilage may be thinned when it becomes thickened by exposure, by dilution with water. A small amount of glycerin added to the mucilage will prevent curling of the paper.

Dextrin, yellowav.oz.	4
Waterfl.oz.	6
Syrupy glucoseav.oz.	1/2
Syrupy glucoseav.oz. Aluminium sulphate (not alum)gr.	120

Mix the dextrin with the water, add the glucose and aluminium sulphate, and heat the mixture to about 90 degs. C., when it will become transparent and thin.

#### VII.

Rye flourav.oz.	4
Acacia, powderav.oz.	1/2
Waterfl.oz.	24
Glycerinfl.oz.	1
Oil of cloves drops	
1	

Rub the flour and acacia to a smooth paste with 8 fluidounces of cold water, strain through cheese cloth, add the mixture to the remainder of the water (also cold) and apply heat until thickening ensues; then allow to cool and add the remaining ingredients.

## VIII.

Rye flourav.oz.	4
Waterfl.oz.	16
Nitric acidfl.dr.	
Oil of clovesdrops	
Glycerinfl.oz.	

Mix the flour and water, make a smooth paste, strain through cheese cloth, add the acid, heat until suitably thickened, and the other ingredients when cool.

#### IX.

Wheat flourav.oz.	41/2
Nitric acidfl.dr.	1
Oil of clovesdrops	5
Boric acidgr.	10
Waterfl.oz.	16

Mix the flour thoroughly with the boric acid and water, and strain through a sieve to avoid lumps; add the nitric acid and heat. with constant stirring, until the mixture has thickened. When nearly cold, add the oil of cloves and stir.

X.
Wheat flourav.oz. 16
Corn starchav.oz. 1
Alum
Boraxgr. 120
Water sufficient

Mix the solids with cold water, make a smooth mixture, add boiling water to this, stirring briskly until the whole is of about the required consistency; apply heat until a uniform paste is produced, and when cold add about 1/2 fluidram of fusel oil, which will preserve it indefinitely.

XI.	
Tragacanth, powdergr.	240
Acacia, powder gr.	240
Wheat flourgr.	480
Salicylic acidgr.	30
Waterfl oz.	
Oil of wintergreendrops	

Mix all but the oil; make a smooth paste, bring this to the boiling point, simmer for 20 minutes, stirring frequently, allow to cool, and add the oil.

## XII.

Dextrin, white av.oz.	4
Acaciaav.oz.	2
Sugarav.oz.	
Waterfl.oz.	12

Dissolve the acacia and sugar in 6 fluidounces of cold water, and the dextrin in remainder of the water by the aid of heat; mix the solutions when cool.

#### XIII.

Starchav.oz.	5
Nitric acid	21%
Gum arabicav.oz.	
Sugarav.oz.	
Watersufficie	

Make a smooth mixture of the starch, 2 fluidrams of acid, and 6 fluidounces of water, and set aside in a warm place for 48 hours, stirring frequently; then boil the mixture until it becomes thick and translucent, dilute with water, if necessary, and strain. Now dissolve the gum and sugar in 5 fluidounces of water, add the remainder of the acid, heat to boiling, and add this mixture to the preceding.

This mucilage may be used by bookbinders

XIV. Mucilage may be made to adhere to tin by first roughening the latter by rubbing with emery paper. Or to the mucilage may be added a small amount, of solution of antimony chloride.

## Mucilage, Stick or Bar.

Mucilage, in the form of sticks, is much used in architectural and mechanical drawing for attaching the drawing paper to a board, and is generally spoken of as mouth or lip glue. In making such a glue, only a very pure form of gelatin or glue should be used, as the least odor would prove disgusting when the glue is moistened with the lips. Sugar is generally added, not for the purpose of sweetening the glue, but in order to render it more easily soluble when it is to be used. This probably is brought about by the sugar preventing the glue from becoming too dry and hard. Some even use a good quality of glue without any admixture whatever, but this requires more rubbing when it is applied, although it holds better than that to which sugar has been added. The sugar may be replaced by glycerin.

I.

Glue,	best		 	 	 	 .av.oz. 4
Isingl	lass			 	 	 .av.oz. 1
Brown	n sug	gar.	 	 	 	 .av.oz. 1
Water	r		 	 	 	 sufficient

Soak the glue and isinglass in water until soft. Pour off the superfluous water, and add the sugar. Melt the whole together with a gentle heat and allow to evaporate until quite thick. Pour into a flat-bottomed dish that is quite cold, preferably placed on ice, and when solid cut the glue into the desired shape.

II.

Isinglassav.oz. White glueav.oz.	1
Rock candy gr. Tragacanth gr.	120
Waterfl.oz.	

Boil the whole together until when cold the mixture has the appearance of glue. Then form into rolls for use.

If desired the glue, made according to either of the above formulas, may, while hot, be poured into suitable molds that have been previously well chilled.

III.

Glue,	ŧ	)(	es	t									.a	V. O2	12	,
Sugar													av	·OZ.	. 5	
Water	٠.												.su	ffic	ient	

Soak the glue in water over night, and dissolve it by heat in the smallest possible quantity of water. Add the sugar to the hot solution, and dry the composition, like jujube paste, in oiled molds.

IV.

• •			
Gelatin		 	av.oz. 4
Sugar, wh	ite	 	av.oz. 2
Water			

Mix, dissolve by aid of heat, and continue heating until the mixture weighs about 8 av. ounces, when it may be formed into sticks.

#### Nutrient Gelatin.

Gelatinav.oz.	1
Extract of beefgr.	175
Distilled waterfl.oz.	29

Dissolve the gelatin and extract in the water, filter, heat to boiling, and divide among test tubes which have previously been treated with boiling water. Close the cylinders with plugs of cotton, which has previously been heated for some time to a temperature of 150 degs. C. then set aside for 4 weeks.

Only the gelatin mixture which remains clear is to be used; if it becomes turbid, it is to be boiled again and again until it remains clear.

Another nutrient gelatin is produced by dissolving 1 part of gelatin in 20 parts of infusion of hay.—D.

#### Oil for Watchmakers.

Place a clean strip of lead in a small white glass bottle filled with pure almond (or olive) oil and expose it to the sun's rays at a window for some time, till a curdy matter ceases to be deposited and the oil has become quite limpid and colorless. Used for fine work; does not become thick by age. The finer grades of paraffin oil also are used, at least for clocks.

## Paint, Glossy or White Enamel.

Orange shellac	5
Copal resinav.oz.	5
Venice turpentineav.oz.	1
Linseed oil, rawfl.oz. 4	
Alcoholpints 10	)

Mix, and add 4 or 5 pounds of zinc white or other white pigment.

Another enamel paint may be made by mixing the pigment with a good varnish.

#### Paint Eradicators.

See "Cleansing Creams," "Cleansing Liquids," Benzin Jelly," and "Stains, Removal of."

# Paper, Barcmeter or Hygrometer.

See "Barometer Paper."

## Paper, Blue Print.

The mixture which is to be applied to the paper consists of 2 (sometimes 3) solutions which are to be mixed just prior to use.

I.

#### No. 1.

Red prussiate of potassiumav.oz. Distilled waterfl.oz.	
No. 2.	
Citrate of iron and ammonium.av.oz.	3

Keep these solutions in separate well-stoppered bottles, which exclude actinic light. In using, mix equal parts of Nos. 1 and 2.

In preparing the sensitized paper, take a solid, firm paper, free from impurities, and apply the solution to the surface of the paper with a soft sponge or a broad, soft brush, being careful not to have the sponge or brush charged too heavily with the solution, or else the paper will have a streaked appearance, which will show in the finished print. Go over the surface of the paper in two directions at right angles to each other so as to insure an even coating. The paper must be allowed to dry in the dark, and in a horizontal position.

IT.

#### No. 1.

Iron citrate, solublegr.	96
Distilled water, enough to make.fl.oz.	
No. 2.	

Red prussiate of potash . . . . . . . gr. 72
Distilled water, enough to make fl. oz. 1
No. 3.

Potassium bichromate.....gr. 5
Distilled water.....fl.oz. 1

Mix Nos. 1 and 2, add No. 3; filter quickly, and use immediately. The iron citrate should be in scales, free from powder, and should not have been exposed to light. The prussiate must also be free from any adherent powder.

To coat the paper.—This must be done by ounces of benzin; gas light. Pour some of the solution into a this, and then dry.

saucer, dip a soft pad of absorbent lint into it and pass quickly across the paper; again dip the pad in solution and pass across the paper from where you left off. When all the paper has been thus coated take an artist mop varnish brush and remove the excess of liquid.

#### Paper, Carbolized.

Carbolized paper, suitable for the preservation of furs, etc. from moths, can be readily prepared by applying a strong solution of carbolic acid, with a brush or sponge to any unsized paper. A heavy paper will absorb more, and consequently last longer. The paper should be kept in close boxes until wanted, and the consumer directed to place the sheets freely among the articles to be protected, and wrap them tightly in ordinary paper.

Carbolic paper may also be prepared as follows:

Petrolatum .										av.oz.	4
Paraffin wax.										av.oz.	4
Carbolic acid	١.									av.oz.	2

Melt the paraffin and petrolatum, add the carbolic acid, allow to cool and solidify, and with this prepared carbolized paper, as directed for making waxed and ceresin paper.—D.

#### Paper, Ceresin.

This may be prepared similarly to waxed paper.

## Paper, Copying.

Make a stiff ointment with lard and black lead or lampblack, and smear it thinly and evenly over soft writing paper by means of a piece of flannel; let remain for a day, and wipe off the superfluous grease. Petrolatum may be substituted for lard, and forms the "Manifold Writer" of the stationers.

## Paper, Oiled.

Brush sheets of paper over with boiled oil and suspend them on a line to dry.

#### Paper, Paraffined.

This may be prepared like waxed paper, or the paper may be drawn through melted paraffin; or, a better way is to melt 8 av. ounces of paraffin, remove from the fire, add 16 fluidounces of benzin; draw the paper through this, and then dry.

## Paper, Parchment.

See " Parchment Paper."

#### Paper, Razor.

See "Razor Paper."

## Papers, Test.

See "Test Papers."

## Paper Tracing.

Τ.

Apply with a brush a varnish compound of equal parts of balsam of fir and oil of turpentine to smooth unsized white paper, and hang up the sheets to dry.

TT.

Rub the paper with a mixture of equal parts of cottonseed oil and oil of turpentine; dry immediately, by rubbing it with wheat flour, and then hang up for 24 hours to dry. If washed over with ox-gall, and dried it may be written upon with ink or water colors.

#### III.

Lardav.oz.	6
Yellow waxav.oz.	1
Lampblackav.oz.	1

Melt the wax, add the lard, thoroughly incorporate the lampblack, making a smooth mixture by trituration, and, while still in a fluid condition apply this mixture to suitable paper by means of a brush.

#### Paper, Waxed.

Place strong white paper on a hot iron plate and rub it well with a lump of white wax, the excess to be removed by means of a cloth pad.

## Paper, Wrapping, to Cut.

Wrapping paper of the size usually employed in pharmacies, viz.,  $24 \times 36$ , may be cut to suit the various sizes of bottles, as follows, according to Jacoby's gauge: Eight ounce,  $9 \times 10$  inches; 6 ounce,  $8 \times 9$  inches; 4 ounce,  $7\frac{1}{2} \times 8$  inches; 3 ounce,  $6\frac{1}{2} \times 7\frac{1}{2}$  inches; 2 ounce,  $6 \times 7$  inches; 1 ounce,  $4\frac{1}{2} \times 6$  inches, and  $\frac{1}{2} \times 6$  ounce,  $4 \times 5$  inches.

What remains after cutting a certain size from a sheet may be used for a smaller size, or it may be used for wrapping pill boxes and similar small packages.

## Parchment Paper.

Dip white unsized paper for one-half minute in sulphuric acid diluted with onehalf its bulk of water, then wash well with weak ammonia water.

## Pastes.

See "Mucilages."

## Paste, Library.

Rice starchav.oz.	2
Gelatinav.oz.	3/4
Waterfl.oz.	
Oil of clovesdrops	16

Incorporate the starch powder with the water, add the gelatin and heat gently over a water bath until a jelly-like compound results.

## Phosphorus Pastes.

I.

Phosphorus	.av.oz.	1
Water (38 degs. C.)	fl.oz.	16
Molasses		
Lard	.av.oz.	16
Oat or barley meal or flour	.suffici	ent

Reduce the phosphorus to fine globules by shaking vigorously with the water contained in a suitable bottle, taking care to have the hand protected with a glove, or the bottle wrapped up in a cloth, for fear of accident. When nearly cool, add the molasses, and then the liquefied lard; finally, incorporate sufficient meal or flour to form a stiff paste. II.

Phosphorusav.oz. 1Water, hot..fl.oz.  $4\frac{1}{2}$ Butter, freshav.oz. 6Starch, powderav.oz. 6

Proceed as in the foregoing, finally adding enough boiling water to make a homogeneous paste.

#### III.

Phosphorus	.av.oz.	1
Sulphur	.av.oz.	1/4
Mustard, powder	.av.oz.	1/2
Sugar, powder	.av.oz.	15
Wheat flour	.av.oz.	20
Carbon bisulphide,		
Waterof each,	sufficie	ent

Mix the sulphur and phosphorus in a suitable vessel, not metailic; add enough water to cover the two, and then mix with enough carbon bisulphide to dissolve both the phosphorus and sulphur; then add the mustard,

sugar, flour and enough water to make a suitable paste.

IV.

Phosphorus av.oz.	2
Carbon bisulphidefl.oz.	
Lardav.oz.	
Wheat flourav.oz.	24

Dissolve the phosphorus in the carbon bisulphide; add the lard, and then incorporate the flour.

V.

Phosphorusav.oz.	6
Sulphurav.oz.	1
Mustard, powder av.oz.	2
Sugar av.oz.	8
Rye flourav.oz.	12
Waterfl.oz.	16

Mix the phosphorus and sulphur with 6 fluidounces of water, triturate until liquefied; add the mustard, sugar and flour, and the remainder of the water.

VI.

Phosphorus av.oz.	1
Bisulphide of carbon fl.oz.	1
Lardav.oz.	8
Wheat flourav.oz.	12

Dissolve the phosphorus in the bisulphide of carbon, add the lard, and lastly make a uniform paste with the flour. This paste will not ferment and spoil.

## Plant Insect Exterminators.

See "Insecticides for Agriculturalists."

## Plating with Gold, Silver, Tin, etc.

The deposition of one metal upon another may be made in several ways. What is now the most common and usually most satisfactory method of deposition, generally known as "plating," is by means of the electric current; this method being known as "electroplating." Another method of deposition, which is a very inferior process, is what is technically known as "washing." This consists in the application of a solution of a salt of a metal which is to be deposited by means of a cloth. The coating of metal deposited is so infinitely thin that it very quickly wears away, revealing the inferior material beneath.

I. Gold "washing:"

Gold chloride av.oz. I	
Potassium cyanideav.oz. 3	
Distilled waterfl.oz. 5	
Precipitated chalksufficient	

Dissolve the gold chloride in 1 fluidounce of water, and the potassium in the remainder of the water; mix the two solutions and add enough precipitated chalk to make a thin paste.

After thoroughly cleaning the object to be gilded, and freeing it from grease, etc., apply this paste with a camels-hair pencil evenly over the surface. Let dry slowly at ordinary temperature. When dry, put in an oven and heat to 60 or 70 degs. C. Wash off with clean water, dry and finish by going over the surface with a burnisher.

Silver "washing:"

A-For brass only.

Silver nitrategr.	60
Potassium cyanidegr.	360
Precipitated chalkgr.	
Distilled waterfl.oz.	$6\frac{1}{2}$

B.—For brass, copper, iron, steel, etc.:

1.

Silver oxyhyposulphiteav.oz.	1/2
Ammonium chlorideav.oz.	1/4
Distilled waterfl.oz.	6

2

•		
	Silver nitrateav.oz.	11/4
	Sodium chlorideav.oz.	5
	Cream of tartarav.oz.	
	Water, distilledsufficier	ıt

Dissolve the silver nitrate in the smallest amount of water, add the other ingredients; rub in a mortar to a smooth paste, adding sufficient water. Preserve the mixture from light.

Apply the paste by rubbing on the cleaned copper or brass until the silver layer is thick enough; then wipe with chamois skin.

3

Silver chloride gr.	60
Potassium bitartrategr.	390
Sodium chloridegr.	180

Mix. The powder is made into a cream with water, and the article to be plated is either covered with the paint by means of a brush or immersed in the mixtures for a short time; then, after being dried, it is rubbed off and the article polished with prepared chalk.—H.

Nickeling:

There is no reliable method of depositing nickel from its cold solution, as in the foregoing cases, but a thin and adhesive coating may be given articles of brass, iron, etc., by the following process: Boil in a copper vessel a saturated solution of zinc chloride and an equal quantity of water. While boiling add hydrochloric acid, drop by drop, until the precipitate at first thrown down is again completely redissolved. Now add zinc in powder, until the bottom of the kettle is nearly covered with a precipitate of zinc. The bath is now ready for the addition of a salt of nickel, and you may use either the sulphate or the nitrate. Add it in sufficient quantity to give the bath a strong green color. The articles to be nickeled are now hung in the bath by means of a zinc wire, or a strip of sheet zinc, and a few pieces of the latter are thrown in along with them. Raise the heat to a strong boil and continue it for several minutes, or until the articles are covered with a bright coating of nickel. The articles should be thoroughly cleaned and free from grease before being put in the bath. When finished, rinse and then rub well with precipitated chalk.

## Platinizing:

Platinum chloride, 1 part; sodium chloride, 8 parts; distilled water, 100 parts. Bring to a boil, and put the articles to be platinized, first thoroughly cleaned, in the vessel. Keep at a moderate temperature, and in the course of 3 or 4 hours the platinizing will be completed. Polish with chamois.

There is as yet no known method of depositing aluminium by a process similar to any of the above.

II. Brass scale pans, or any other metallic substance capable of taking a deposit of silver may be plated in various ways. Here is a method recommended by Kayser:

It is absolutely necessary that the article to be plated shall present a perfect metallic surface, free from oxides, dirt, grease, etc.; it must be thoroughly scoured, if necessary, with the intervention of acids, and afterwards carefully washed. It is then to be dipped into a solution prepared by making a saturated aqueous solution of bisulphite of sodium, and adding to the latter so much of a solution of nitrate of silver (30 parts in 100) that there are 6 parts of the silver salt for every 100 of the bisulphite. The follow- servative Fluid, which is adapted to almost all

ing would be a more simple way to state the proportions:

Sodium bisulphiteav.oz.	10
Distilled water enough to disso	lve
Silver nitrategr.	264
Distilled waterfl.oz.	2

Dissolve and mix.

Allow the article to remain in the mixture until it is properly coated, then take it out; wash it with water in which a little carbonate of sodium had previously been dissolved; finally wash with pure water, and dry in sawdust.

## Polishing Paste.

See " Putz Pomades."

## Polishing Powders.

Under this heading are mentioned powdery mixtures used in polishing different metals. These powders must always be impalpably fine, particularly such as are used to polish silver and gold ware.

Chalka	v.oz. 10
White bolea	v.oz. 4
Lead carbonatea	v.oz. 5
Magnesium carbonate a	v.oz. 1
Iron oxidea	v.oz. 1

This mixture is best adapted to brass and copper.

II.				
Calcined	magnesia.	 	av.oz.	9
Leweler's	rouge	 	av. cz.	1

This mixture is recommended for polishing silver; it should be used dry.-D.

	magnesia.							
Teweler's	rouge						.av.oz.	8

This mixture is recommended for polishing gold; it should be used dry.-D. IV.

Magnesium	carbonate.	 av. oz. 4
Chalk		
Jeweler's ro	uge	 av.oz. 7

#### Polishes for Shoes and Leather.

See "Blackings," "Shoe Dressings," "Patent Leather Polish," "Shoe Varnish," and "Harness Blackings."

#### Preservative Fluid.

HI.

The composition of Wickersheimer's Pre-

purposes (excepting the preservation of animal tissues to be used for food) and which is used in medical colleges, is as follows:

Arsenious acidgr. 96	)
Potassium carbonategr. 22	)
Potassium nitrategr. 73	
Potassium sulphategr. 96	-6
Sodium chloridegr. 120	
Sodium borategr. 24	0
Glycerin	3
Wood alcoholfl.oz.	31/
Waterfl.oz. 3-	1

Dissolve the arsenious acid and potassium carbonate in 7 fluidounces of the water, using a gentle heat to accelerate the reaction, and add the remaining portion of the water, in which dissolve the other salts; add the glycerin and alcohol.

If large quantities of the liquid are to be used, it will be economical to use methylic alcohol, as ordered in the formula, otherwise common alcohol may be substituted.

Specimens may be preserved by simply immersing in the fluid, or by injecting it into the veins and intestines of the body.

The following is suggested as a substitute for Wickershiemer's preparation:

Salicylic acidgr.	240
Boric acidgr.	300
Potassium carbonategr.	60
Oil of cinnamonfl.dr.	
Oil of clovesfl.dr.	3
Glycerinfl.oz.	5
Water, hot	121/2
Alcohol	121/2

Dissolve the acids and potassium carbonate in the water; when effervescence ceases add the glycerin, and then the oils dissolved in alcohol.

This fluid is not poisonous, and possesses the desirable property of acting as an-antiseptic and of having a pleasant odor.—H.

#### Putz Pomades.

By this term are signified pasty or fatty mixtures intended especially for polishing copper and brass; these mixtures may be used on steel, but should never be used on silver or gold.

The fatty substance present consists of a fixed oil—oleic acid, lard, petrolatum, palm oil, lard oil, etc., mixed with tripoli, rotten stone, emery, jeweler's rouge, pumice stone, etc.; the whole being flavored, as a rule, with

oil of mirbane (nitro-benzol or artificial oil of bitter almonds). The powders must be in a very finely divided condition; it is, in fact, advisable to pass the mixture of fat and powder through a paint mill to insure fineness and smoothness.

I.
Rotten stoneav.oz. 16
Stearin
Cottonseed oilfl.oz. 4
Oil of mirbane enough to flavor

Melt the fats, incorporate the rotten stone with them, and add the oil of mirbane when cool.

11.	
Oxalic acidav	.oz. 1
Peroxide of iron (jeweler's	•
rouge)av	.oz. 8
Rotten stoneav	.oz. 10
Palm oilav	
Petrolatumav	.oz. 2

Pulverize the acid and add the rouge and rotten stone, mixing thoroughly. Sift to remove all grit; then gradually add the palm oil and petrolatum, and incorporate. Add oil of mirbane or oil of lavender to flavor.

oil of mirbane or oil of lavender to flavor.	
III.	
Rotten stone, levigatedav.oz.	2
Iron subcarbonateav.oz.	
Oil of bitter almonds or mir-	
baneenough to flavo	r
Lard, olive or cottonseed oil.	
enough to form a past	e
IV.	
Charcoal, fine powderav.oz. 1	4
Iron oxide (subcarbonate)av.oz.	3
	6
	3
	6
Oil of mirbanedr.	3

	On or chronena	1
7.		
	Pumice, powderav.oz.	2
	Rotten stone, powderav.oz.	2
	Iron subcarbonate av.oz.	4
	Olive or cottonseed oil, or oleic	
	acidenough to form a pas	
	Oil of mirbane enough to flav	Or

VI.

Oleic acid	. fl.oz. 2
Lard	av.oz. 6
Jeweler's rouge	
Emery, powder	
Rotten stone, powder	
Oil of mirbane, sufficient to give f	

VII.
Rotten stone, fine powderav.oz. 8
Oxalic acidav.oz. 2
Cottonseed oilfl.oz. 3
Oil of turpentine, enough to make a paste
VIII.
Rotten stone, fine powderav.oz. 12
Soft or green soapav.oz. 6
IX.
Rouge (iron oxide)av.oz. 3
Lard or petrolatumav.oz. 15
X.
Palm oilav.oz. 8
Petrolatumav.oz. 8
Rouge (iron oxide) av.oz. 4
Tripoliav.oz. 3
Oxalic acidgr. 80
XI.
Japan, waxav.oz. 2
Oleic acid, crudeav.oz. 11
Tripoliav.oz. 7
Oil of mirbaneenough to flavor
_
If desired, armenian bole, iron oxide,

If desired, armenian bole, iron oxide, or venetian red may be added to this mixture to impart color.—D. modified.

#### Putz Tablets.

Soap, cut fine or powderav.oz.	12
Precipitated chalkav.oz.	1 1/2
Jeweler's rougeav.oz.	3/4
Cream of tartarav.oz.	1
Magnesium carbonateav.oz.	3/4
Watersufficie	ent
This, like Putz pomade, is used for pe	olish-

ing purposes.

Quinine Salts, Extempore Prepara-

# dumine Salts, Extempore Preparation.

Quinine Carbolate.—Quinine alkaloid, 10 grains; carbolic acid, 5 grains.

Quinine Citrate.—Quinine alkaloid, 15 gr.; citric acid, 8 grains. This product is equivalent to 20 grains quinine citrate.

Quinine Hydrobromate.—Quinine sulphate, 100 grains; potassium bromide, 28 grains. This product is equivalent to 100 grains quinine bromide.

Quinine Hydriodate.—Quinine sulphate, 95 grains; potassium iodide, 40 grains. This product corresponds to 100 grains quinine iodide.

Quinine Iodo-hydriodate.—Quinine hydrochlorate, 70 grains; potassium iodide, 50 grains; iodine, 20 grains. These constituents are triturated together with a little alcohol.

This product corresponds to 100 grains quinine iodo-hydriodate.

Quinine Hypophosphite.—Quinine hydrochlorate, 100 grains; calcium hypophosphite, 24 grains. This product corresponds to 100 grains quinine hypophosphite.

Quinine Lactate.—Quinine alkaloid, 70 grains; lactic acid, 35 grains. If necessary, these are triturated together with a little alcohol. This product corresponds to 100 grains quinine lactate.

Quinine Phosphate.—Quinine sulphate, 94 grains; sodium phosphate, 80 grains. This product corresponds to 100 grains quinine phosphate.

## Raticides.

I.		
	Wheat flourav.oz.	5
	Fresh milkfl.oz.	10
	Mutton tallowav.oz.	1
	Sodium chloridegr.	50
	Squill, coarse powderav.oz.	4

Mix the wheat flour and milk, then add the tallow and salt, and heat for 20 minutes over a steam-bath; lastly, incorporate the squill.

—D.

This preparation is known as "Gliricin."

[.	
Strychnine sulphateav.oz.	11/2
Milk sugarav.oz.	11/2
Prussian bluegr.	20
Arsenicav.oz.	3
Wheat flourav.oz.	12

Rub up the strychnine and milk sugar together, add the prussian blue and arsenic, and finally, add the flour, and mix thoroughly. When required for use, moisten and make a dough; divide into small pellets and dry.

#### III.

Barium carbonate (freshly pre-	
cipitated)av.oz.	2
Sugarav.oz.	1/2
Bread av.oz.	8
Form into 100 pills.	

V.		•
Tartar emetic	av.oz.	11/2
Squill, powder		
Barium carbonate, precipitated	.av.oz.	3
Roasted meat		

V. Poisoned wheat (arsenic):	
Potassium arseniateav.oz.	1
Waterfl.oz. 1	0
Fuchsingr.	5

Dissolve the potassium salt in the water; add the fuchsin and dissolve, and then mix with the wheat. The latter may best be incorporated with the poison by introducing into a wide-mouth bottle, and gradually adding the poisonous solution, shaking frequently.—D.

The coloring matter may be omitted if desired.

#### VI. Poisoned wheat (strychnine):

Strychnine nitrategr.	. 18
Water fl.oz.	. 10
Methyl violet gr.	
Wheatav.oz.	

Dissolve the alkaloidal salt in the water, add the methyl violet, introduce into a widemouth bottle, add the wheat; mix well by agitation, set aside for 6 hours, and then take out and dry at a temperature not exceeding 30 degs. C.—D.

Strychnine sulphate may be substituted for the nitrate, and the methyl violet may be omitted.

VII. See also under heading "Phosphorus Pastes."

#### Razor Pastes.

I.

Razor paste can be easily made by taking emery flour and shaking up with water and allowing to stand a moment to allow the coarse particles to subside; then pour off the remainder into a paper filter and allow to drain and dry. When dry mix with enough petrolatum or simple ointment to make a paste.

#### II.

Emery flour, Jeweler's rouge, Spermaceti ointment, of each, equal parts

#### III.

Emery flour					, .	.av.oz	. 2
Spermaceti ointment	•					.av.oz	. 1

#### IV.

Jeweler's rouge, Black lead,

Suet......of each, equal parts

# V.

Levigated oxide of tinav.oz.	4
Oxalic acid, powderav.oz.	1
Gum arabic, powdergr.	80
Waterenough to form a pas	

#### Roach Pastes.

Many of the roach pastes are "phosphorus pastes," or so-called "electric pastes." Others are made with red lead or other ingredients. These pastes may be distributed on papers in the haunting places of the roaches, or they may be fed into cracks or crevices in which the insects hide.

#### I.

Red leada	v.oz. 4
Floura	
Molassessufficient to make a se	oft paste

#### H.

Red lead.			av	7.oz. 4
Corn mea	l, powder.		av	.oz. 8
Molasses.	.sufficient	to form	a soft	paste
TTT 0 -			_	

# III. See also "Phosphorus Pastes."

#### Roach Powders.

These powders should be distributed liberally in the localities frequented by the insects; a blower or "gun" is excellent for the purpose.

## I.

Wheat	flourav.oz.	4
Sugar,	powderav.oz.	8
Borax,	powder av.oz.	2
Unslak	ed limeav.oz.	2

## Keep dry.

This should be strewed about on paper, taking care that no liquids are left uncovered.

#### II.

Богах	 	 	 	 .av.oz.	13
Starch	 	 	 	 .av.oz.	3
Cacao.	 	 	 	 .av.oz.	11/4

All should be in fine powder and be well mixed.

## III.

Plaster of	paris	 	 .av.oz. 4
Oatmeal .			
Sugar		 	 .av.oz. 2

All should be in fine powder and should be well mixed.

## IV.

Angelica root, powderav.oz. Oil of eucalyptusfl.dr.	
Mix well.	

## V.

Tartar	r emeticgr.	140
Insect	powderav.oz.	16

#### VI.

Insect	powder		av.oz.	8
Levant	wormseed.	powder	av.02.	8

·
VII.
Chamomile.       av.oz. 2         Borax.       av.oz. 12         Insect powder.       av.oz. 2         Plaster of paris.       av.oz. 1         Sulphur.       av.oz. 3         Crude arsenic (so-called "co-balt")      gr. 120
All should be in powder and should l
well mixed.
VIII.
Insect powderav.oz. 3 Quillaja, powderav.oz. 1
IX.
Borax       av.oz. 13         Sugar       av.oz. 4         Cacao       .gr. 300
Reduce all to powder and mix well.
X.
Borax       av.oz. 4         Paris green       av.oz. 4         Sugar       av.oz. 2
Reduce all to powder and mix well.
XI.
Wheat flour
XII.
Borax, av.oz. 8 Flour av.oz. 8 —D.

<sup>1</sup> XIII. A very common roach powder is a mixture of insect powder with borax. Thymol in alcoholic solution may also be added to insect powder.

Another mixture often used is a combination of insect powder and paris green.

#### Rust Stains, Removal of.

I.	
Tartaric acidav.oz.	1
Alumav.oz.	
Water, enough to makefl.oz.	16
Mix, dissolve and filter.	

II. A mixture of 2 parts of powdered cream of tartar with 1 part of powdered oxalic acid will remove stains from cotton and linen. This mixture is sometimes sold under the name of salts of lemon. The poisonous character of the acid must not be overlooked, for accidents have occurred from its careless use.

III. Rust Stains on White Goods.—Soak the stains in a solution of tin chloride, and rinse immediately with much water. The tin salt is much more reliable in removing iron rust, and quicker in its action than oxalic acid, unless the stains are soaked in a solution of the latter, contained in a tin spoon, when the stains disappear in a short time.

## Sealing Waxes.

These consist of resinous substances in combination with coloring agents, the mixture frequently being cheapened by the addition of mineral substances. In preparing these, the less fusible resins, such as rosin and shellac should be melted, then the turpentine. Venice turpentine or similar more fusible agent should be added, and with this mixture should be incorporated the remaining ingredients. The powdery substances should be added in the very finely divided form, and should first be added; if there are several powders, they should be well mixed before adding to the liquefied resins. When the ingredients have been mixed, the heating should be continued for a few moments to permit the escape of air bubbles which have been introduced during mixing of the ingredi-

The mixture may be cast into sticks, if desired, by pouring into suitable molds which have previously been moistened.

To cool off to the consistency required for molding, pour off about one-sixth of the mixture on a piece of wetted parchment paper, and as soon as this mass has hardened it should be returned to the vessel and stirred until the cooled portion is dissolved; the mixture may be poured into the molds.

## Sealing Wax, Black.

	Venice turpentineav.oz. 8
	Resin av.oz. 1½
	Lampblacksufficient to color
11	
	Gum turpentineav.oz. 1½
	Resin av.oz. 6
	Stearingr. 260
	Lampblack gr. 20
	Heavy spar(barium sulphate). av.oz. 12
	—D.
TT	T

Venice turpentine.....av.oz.

Chalk .....av.oz.

Lampblack.....av.oz.

Shellac.....av.oz. 12

Melt the resin, add the venice	turpentine
and incorporate with the chalk and	lampblack
which have previously been well m	ixed.

# 

I.	
Gum turpentineav.oz.	4
Resin, whiteav.oz.	15
Ultramarine blue av.oz.	
Heavy spar (barium sulphate)av.oz.	21/2
	D.

111.	
Gum turpentineav.oz.	4
Resin, whiteav.oz.	15
Ultramarine blueav.oz.	2
Zinc oxideav.oz.	2
Heavy spar (barium sulphate)av.oz.	21/2
_	D.

## Sealing Wax, Green.

Ι.		
	Shellacav.oz.	10
	Venice turpentineav.oz.	5
	Resin av.oz.	3
	Magnesiaav.oz.	1/2
	King's yellow (yellow litharge).av.oz.	114
	Mountain (Sander's) bluegr.	260
	Oil of turpentinefl.dr.	3

Melt the shellac and resin, add the venice turpentine, and then incorporate the colors, which have previously been mixed to a paste with the oil.

#### II.

Resin av.oz.	10
Venice turpentineav.oz.	11/2
Chalk	21/2
Chrome greenav.oz.	1

Melt the resin, add the turpentine, and then incorporate the chalk and chrome green which have previously been well mixed.

#### III.

Shellacav.oz.	8
Masticav.oz.	
Green turpentineav.oz.	11/2
Verdigris av.oz.	2

IV. Paris green may also be employed as the coloring agent.

## Sealing Wax, Red.

The coloring agents used may be vermilion or red lead, or even venetian red.

Shellac, bleached
II. Shellac av.oz. 10 Venice turpentine av.oz. 2 Vermilion av.oz. 4
Bleached shellac should be used in making
this wax.  III.  Resin
Melt together and color with red lead
venetian red, or vermilion.
Shellac av.oz. 8 Venice turpentine av.oz. 3 American vermilion av.oz. 2 Balsam of peru av.oz. ½
V. Venice turpentine
VI. Venice turpentine
Melt the shellac and turpentine, add the

Melt the shellac and turpentine, add the cinnabar, and finally, the magnesia in fine powder, triturated with a little oil of turpentine. Cheaper grades may be made by adding resin and reducing the quantity of cinnabar.

## VII.

Venice turpentin	ie	 	 av.oz.	4
Vermilion		 	 av.oz.	2
VIII.	٠			
Gum turpentine				

Orange shellac ......av.oz. 8

Cam tarpentine	
Resinav.oz.	6
Stearinav.oz.	1
Vermilionav.oz.	3/4
Vermilionav.oz. Heavy spar (barium sulphate).av.oz.	12
<u> </u>	D.

## Sealing Wax, Violet.

Gum turpentineav.oz.	2
Resin, whiteav.oz.	71/2
Carmine av.oz.	1/2
Zinc oxideav.oz.	2
Ultramarine bluegr.	110
Heavy spar (barium sulphate), av. oz.	
_	-D.

Sealing Wax, White.	
I.	
Bleached shellacav.oz.	יא
Venice turpentineav.oz.	7 31/4
Plaster of paris	2
Magnesia gr. 12	
Bismuth subnitrateav.oz	3
Lead Carbonateav.oz.	41/2
	1/2
II.	
Gum turpentineav.oz.	2
Resin, whiteav.oz.	71/2
Zinc whiteav.oz.	2
Heavy spar (barium sulphate). av.oz.	
—I	,
Sealing Wax, Yellow.	•
Seaning wax, renow.	
I.	
Shellac, bleachedav.oz.	8
Resin av.oz.	21/2
Venice turpentineav.oz.	4
King's yellow (sulphide of ar-	
senic)av.oz.	11/2
	12
II.	
Shellacav.oz.	71/2
Venice turpentineav.oz.	61/2
Resinav.oz.	31/4
Plaster of parisav.oz.	1
Magnesiagr. 9	, 10
Chrome yellowav.oz.	$1\frac{1}{2}$
III.	
Gum turpentineav.oz.	4
	.5
Chrome yellow av.oz.	2
Heavy spar (barium sulphate)av.oz.	3
	•
—I	).
Sewing Machine Oil.	
I.	
Paraffin oil, bestfl.oz.	4
Olive oil, bestfl.oz. 12 or 1	6
II.	
	4
Petrolatumav.oz.	2

Melt the petrolatum, and add the oil; cool thoroughly, and allow the cloudiness which takes place to clear off by depositing. Decant, and use the clear supernatant oil.

## Shoe Dressing, Kid.

I.	
Ceresinav.o	oz. 4
Oil of turpentineav.o	z. 4
Castor oilav.c	oz. 4
Linseed oil, rawav.d	oz. 20
Pine tarav.c	oz. 1
701 1 1 1 1 1 1 1 1	

Dissolve the ceresin and tar in the oil of turpentine; then add the heavy oils while stirring. Perfume with oil of mirbane.

Shellacav. ɔz.	4
Aqua ammoniafl.oz.	2
Waterfl.oz.	16
Black aniline sufficie	ent

Heat the ingredients slowly together (except the aniline) until the whole is near boiling and the shellac dissolves; (It may be necessary to add a little more ammonia during the boiling.) then add the aniline, and water enough to make the whole measure 16 fluidounces.

## Shoe Dressing, Ladies' or Liquid.

These preparations are usually resinous solutions colored black, and intended for application to shoes by means of sponge. They dry quickly, and give a polish without friction with a brush.

Whenever bone or ivory black is directed in a formula, the purified article should be preferred, as it gives a dead-black color; whereas the unpurified may give but a brownish or grayish black.

Shoe blackings are mentioned under heading "Blacking for Shoes."

I.

٠.			
	Caoutchouc	.av.oz.	1
	Petroleum	av.oz.	1
	Carbon disulphide	fl.oz.	1
	Shellac		
	Lampblack		
	Oil of lavender		
	Alcohol		

Upon the caoutchouc, contained in a bottle, pour the carbon disulphide; cork well and let it stand a few days, or until the caoutchouc has become thoroughly gelatinized or partly dissolved; then add the petroleum, lavender oil and alcohol; next the shellac in fine powder, and heat it to about 50 degs. C., taking care that as little as possible is lost by evaporation. When the substances are all dissolved, and the liquid is tolerably clear, add the lampblack, and fill at once into small bottles.

II.

±.	
Extract of logwoodgr	. 5
Gallic acidgr	. 10
Boraxgr	. 10
Aniline blackgr	. 20
Ammonia waterm.	
Water, hotfl.oz	. 1
Aqueous shellac solutiongal.	. 1
Mix and dissolve.	

The aqueous shellac	solution	should	be pre-
pared as follows:			

Water .		.oz. 5
Borax		gr. 100
Shellac,	powder	gr. 300

The shellac should be gradually added to the hot borax solution.—H.

#### ITT.

Indigogr.	120
Tragacanthgr.	
Glueav.oz.	
Logwoodav.oz.	8
Glycerinfl.oz.	3
Waterfl.oz.	16
Diluted acetic acidfl.oz.	32
Poil together and strain	

Boil together and strain.

## IV.

Aniline blackgr.	400
Camphorav.oz.	1 1/2
Shellacav.oz.	21
Wood alcoholfl.oz.	64

The wood alcohol is used only because it is cheaper than grain alcohol; the latter may be employed if desired.

# v.

Shellac
Ammonia waterfl.oz. 1
Waterfl.oz. 6
Aniline blacksufficient to color

Boil all the ingredients together, except the aniline, until the shellac is dissolved; then add the aniline, and sufficient water to make the liquid up to the measure of 16 fluid-ounces.

#### VI.

Ivory black, very fineav.oz.	32
Molasses av.oz.	24
Sperm oilfl.oz.	4
Gum arabic	1
Diluted acetic acidfl.oz.	8

Mix the first three ingredients, then add the gum dissolved in the acid; mix again, set aside for 24 hours, and add 3 or 4 pints of vinegar or sour beer.

#### VII.

Ivory black av.oz. 8
Molasses
Sweet oilav.oz. 1
Hydrochloric acid, commercialav.oz. 4
Sulphuric acid, commercialav.oz. 2
Water sufficient

Mix the ivory black with the molasses and oil, add the hydrochloric and sulphuri cacids, first mixing the latter with 3 or 4 fluid-ounces of water, and adding while hot. Let

stand until all effervescence ceases, stirring occasionally, and finally, thin to the desired consistence with stale beer.

## VIII.

* * * * * * * * * * * * * * * * * * * *	
Castile soap, whiteav.oz.	1
Sandaracav.oz.	1
Mastic	:/2
Venice turpentine	2
Shellacav.oz.	334
Aniline black, Eav.oz.	11/2
Glycerin fl.oz.	2
Waterfl.oz.	3
Alcoholfl.oz.	

Dissolve the soap in the water, mixed with 7 fluidounces of alcohol; macerate in a warm place, with frequent agitation, subsequently filtering through cotton. The mastic, sandarac and venice turpentine should be dissolved in 9 fluidounces of alcohol, and the solution also filtered through cotton. Then dissolve the shellac and aniline in the remainder of the alcohol. Now mix the three solutions, and add the glycerin.

The mixture should be dispensed in widemouth bottles, the corks being supplied with a wire which has a sponge attached to it.

#### IX.

4
2
4
2
1
nt

Mix the shellac, borax, and 14 fluidounces of water, with constant stirring, until the shellac is dissolved; then add the sugar, glycerin and nigrosin; stir until the latter is dissolved, and add enough water to make 36 fluidounces.

#### -D.

X.	
Molassesav.oz.	
Ivory blackav.oz.	$^4$
Diluted acetic acidfl.oz.	
Waterfl.oz.	12
Acacia, powdergr.	
Sulphuric acid, commercialfl.dr.	4

Mix all the ingredients, except the sulphuric acid; then add the latter gradually, with constant stirring.

#### XI.

Yellow waxav.oz.	2
Fish oilfl.oz.	2
Benzinfl.oz.	15
Tincture of green soapfl.oz.	
Lampblack sufficient to col	lor

-D.

#### Shoe Grease.

I.
Liquid petrolatum, yellowfl.oz. 14
Olive oil, commonfl.oz. 1
Ceresinav.oz. 5
Alkanningr. 10
Oil of mirbanedrops 6
Oil of citronelladrops 3

Melt the ceresin, add the petrolatum, oil and alkannin; allow to cool, and incorporate the oils of mirbane and citronella.

II.	
Yellow waxav.oz.	1
Turpentineav.oz.	1
Castor oilav.oz.	
Linseed oilfl.oz.	10
Pine targr.	120

Clean the leather; let dry, and rub the grease well in before a fire.

# III.

Resinav.oz. 2	,
Yellow waxav.oz. 3	,
Linseed oilfl.oz. 12	,
Neatsfoot oil	,
Oil of turpentinefl.oz. 4	
This has been known as Norfolk fluid.	

## Shoe Polish, Patent Leather,

_		
I.		
	Yellow wax, or ceresinav.oz.	3
	Spermacetiav.oz.	1
	Oil of turpentinefl.oz.	11
	Asphalt varnishav.oz.	1
	Borax, powdergr.	80
	Frankfort blackav.oz.	1
	Prussian bluegr.	150
	Oil of mirbanefl.dr.	

Melt the wax, add the borax, and stir until a kind of jelly has been formed. In another pan melt the spermaceti; add the varnish, previously mixed with the turpentine; stir well and add to the wax; lastly, add the colors, mix well, and incorporate the oil of mirbane.

# II.

hour.

Molassesav.oz.	
Gum arabic av.oz.	1/2
Ivory blackav.oz.	16
Mix well, and simmer together for ha	

# Shoe Varnish, Patent Leather.

I.	
Shellacav.oz.	2
Gum turpentineav.oz.	3,1
Sandaracgr.	144
Lampblack gr.	72
Oil of turpentinefl.dr.	6
Alcoholfl.oz.	19

	II.
	India rubberav.oz. ½
	Shellacav.oz. 8
Į	Camphorav.oz 1
ı	Nigrosinav.oz. 1
	Carbon disulphidefl.oz. 2
ı	Wood alcohol A 99

Wood alcohol......fl.oz. 22 Dissolve the rubber in the carbon disulphide, and add the other ingredients, previously dissolved in the alcohol.

## Shoe Dressing, Tan or Russet.

I.	
Oil of turpentinefl.oz.	10
Yellow waxav.oz.	5
Soap (ordinary bar)av.oz.	
Boiling waterfl.oz.	10

Dissolve the wax in the turpentine by the aid of the water bath, and the soap in the hot water; mix in a hot mortar, and agitate until cold.

II.	
Palm oilav.oz.	2
Common soapav.oz.	6
Oleic acidfl.oz.	
Glycerinfl.oz.	
Tannic acidgr.	

Melt the soap and palm oil together with a very gentle heat. When the soap is dissolved add the oleic acid. Dissolve the tannic in the glycerin; add to the hot mixture, and stir until cold.

111.							
Petrolat	um	 	 	 		av.oz.	12
Yellow	wax.	 • •	 	٠.		av.oz.	4
TV							

•	* •	
	Yellow waxav.oz.	2
	Fish oilav.oz.	2
	Benzinfl.oz.	
	Tincture of green soapfl.oz.	1
	Yellow ocherav.oz.	1/2
	_	D.

√.		-D.
	.av oz	1/2
Yellow waxOil of turpentineSoap shavings	fl. oz.	2
Soap shavings	gr.	280
Fish oil		
Borax, fine powder	gr.	275
Glycerin	fl. oz.	9
Yellow ocher		

Melt the wax, and add the oil of turpentine. With the fish oil incorporate the borax previously dissolved in the glycerin, add the yellow ocher, and then the wax solution; now add the soap shavings; heat the whole on a water bath until dissolved, and stir until cold.

....av.oz. 2

VI.						
Soft o	r green	soap				
Linsee	ed oil,	raw				

 Linseed oil, raw
 fl.oz. 3

 Annatto solution (in oil)
 fl.oz. 8

 Yellow wax
 av.oz. 3

 Gum turpentine
 av.oz. 8

 Water
 fl.oz. 8

Dissolve the soap in the water and add the annatto; melt the wax in the oil and turpentine, and gradually stir in the soap solution, stirring until cold.

#### VII

Yellow waxav.oz.	5
Potassium carbonategr.	264
Resin soap (common yellow bar) gr.	175
Water fl.oz.	
Oil of turpentinefl.oz.	5

Mix the first four ingredients, heat until well mixed, and when cooled to 80 degs. C. the oil may be added; now incorporate enough of an oil-soluble yellow aniline to produce the desired shade.

#### Show Globe Colors.

Colored liquids for show globes, show globe colors as they are termed, are frequently made of coal tar dyes, and very handsome colors may be produced in this way, but these, as a rule, lack permanency. However, these colors are inexpensive; so very little of the dye being required that they may be "freshened" occasionally, or they may be entirely renewed. Those who do not care to use coal tar dyes may employ the formulas mentioned below. It should be understood. however, that no show bottle colors are absolutely permanent, because they are exposed to one of the most powerful of all chemical agents, viz., light. From time to time they should be filtered or else renewed, and the bottle should be thoroughly cleansed.

Show bottle colors are liable to be subjected to considerable cold in winter time and to be frozen. This may be prevented by replacing 20 or 25 per cent of the water with alcohol or glycerin. Such replacement cannot always be made, owing to possible chemical change—for example, in purple made with potassium permanganate.

The colors most commonly employed are red, green, blue, and yellow. The formulas given below will yield these colors in several different shades. These shades may in each filter.

instance be deepened or made lighter by decreasing or increasing the amount of water.

## Show Globe Color, Amber.

Dragon's	bloodgr.	45
Sulphuric	acidfl.dr.	- 3
Distilled	watergal.	$^{2}$

Powder the dragon's blood and macerate in the acid for 20 or 30 minutes, then add the distilled water and filter.

## Show Globe Colors, Blue.

Copper sulphateav.oz.	16
Sulphuric acidfl.oz.	
Watergal.	
T.	

Copper sulphate......gr 240
Ammonia water sufficient, or fl. oz. 1 to 2
Water....gal. 2

Dissolve the copper salt in water, add ammonia water until the precipitate first formed is redissolved, and add the remainder of the water.

III. Dissolve prussian blue in water by the aid of oxalic acid, or dissolve soluble blue or indigo sulphate in water.

ïV.

Copper sulphateav.oz.	8
Alumav.oz.	8
Sulphuric acidfl.oz.	8
Distilled watergal.	2

Dissolve the alum and blue vitriol in the water, cautiously add the sulphuric acid, and filter.

#### Show Globe Colors, Crimson.

I.

Solution of iron chloride. fl.dr. 10
Water of ammonia. fl.oz. 1
Acetic acid. fl.oz. 2
Alcohol. fl.oz. 6
Distilled water. gal. 2

Add the solution of iron chloride to the water; then add the alcohol, acetic acid and water of ammonia, and filter.

II.

Iodineav.oz.	1/2
Potassium iodidefl.oz.	1/2
Hydrochloric acidfl.oz.	9
Distilled watergal.	2

Dissolve the iodine and potassium iodide in the water and add the hydrochloric acid; filter.

#### Show Globe Colors, Green.

I.	
Copper sulphate	av. oz. 12
Hydrochloric acid	
Distilled water	

Dissolve the copper sulphate in the distilled water, add the hydrochloric acid and filter.

Verdigris.....av.oz. 12 Sulphuric acid.....sufficient Distilled water.....gal. 1

Mix the verdigris with enough acid to dissolve; let stand a few minutes, add to the distilled water and filter.

#### III.

Nickelav.oz. 5	
Hydrochloric acidfl.oz. 8	
Nitrous acid	
Distilled water, enough to makegal 2	

Dissolve the nickel in the hydrochloric acid, add the water, then the nitrous acid.

IV.

Copper sulphateav.oz.	9
Ammonium chlorideav.oz.	9
Water, enough to makegal.	2

First dissolve the copper salt in the water. then add the ammonium chloride; dissolve and filter.

V .	
Copper acetate	av.oz. 1
Acetic acid	
Water, enough to make	

Add the acetic acid to the copper acetate, and triturate with the water till dissolved.

VI. Add an aqueous solution of picric acid to an aqueous solution of copper sulphate until the desired shade is produced. A beautiful grass green liquid will be the result.

#### VII.

Nickelav.oz.	1/2
Nitric acidfl.oz. 2	
Distilled watergal. 2	
Potassium bichromatesufficient	

Dissolve the nickel in the nitric acid, add the water and enough of solution of potassium bichromate to give the desired color, and filter.

#### VIII.

Copper sulphate	av.oz.	1
Water of ammonia	.fl.oz.	10
Potassium bichromate	.suffici	ent
Distilled water	9	960

tilled water, add the water of ammonia and enough of a solution of potassium bichromate to give the desired color, and filter.

#### IX.

Fluorescin	 	 	 gr.	2
Ammonia water	 	 	 .fl.ďr.	2
Water, distilled				

This makes a fluorescent grass green liquid.

X.	
Copper sulphateav.oz.	
Hydrochloric acidfl.oz.	
Subcarbonate of ironav.oz.	1
Distilled watergal.	2

Dissolve the copper sulphate in the water; dissolve the iron in the hydrochloric acid; mix the two solutions, and filter.

#### 37 T

Α1.		
Copper sulphate	av.oz.	8
Sodium chloride	av.oz.	16
Hydrochloric acid	fl.oz.	8
Water		

Mix, dissolve and filter.

#### XII.

Potassium bichromategr.	120
Copper ammonio-sulphategr.	240
Waterga	I. 2

Mix, dissolve and filter.

Instead of the copper salt may be used a solution of copper treated with ammonia water until the precipitate first formed is redissolved.

## Show Globe Colors, Orange.

I. Dissolve annatto in liquor potassa and dilute alcohol, and filter.

II. Dissolve 8 av. ounces of potassium bichromate in 2 gallons of water. The shade may be varied by adding nitric, sulphuric, or hydrochloric acid, and filter. Instead of using any of these acids, 120 grains of chromic acid may be dissolved in the water.

III. Some of the darker of the yellow show globe colors may also be employed if an orange shade is desired.

#### Show Globe Colors, Pink.

I,
Sodium salicylategr. 16
Tincture of iron chloride,
Hydrochloric acidof each, sufficient
Watergat. 2

Dissolve the sodium salicylate in the water and add (cautiously) tincture of iron in single Dissolve the copper sulphate in the dis- drops until the proper tint has been realized.

Hydrochloric acid may be used carefully (in drops) to render the tint paler. Should too much acid have been added, restore the tint by the addition of ammonia.

II.	
Cobalt oxidegr.	120
Nitric acidfl.oz.	12
Watergal.	2

Dissolve the cobalt oxide in the acid, then add the water.

## III.

Cobalt oxideav.oz.	2
Nitric acidfl.oz.	1
Hydrochloric acid fl.oz.	1
Ammonia waterfl.oz.	18
Sulphuric acidfl.oz.	1
Watergal.	2

Dissolve the cobalt oxide in the nitric and hydrochloric acids mixed, then add the remaining ingredients previously mixed; set aside for several weeks, and filter.

## Show Globe Colors, Purple.

I.	
Verdigrisgr. 60	)
Water of ammoniafl.oz. 20	)
Distilled watergal. 2	)

Mix the water and the ammonia, add the verdigris, and, when dissolved, filter.

## Π.

Salicylic acidgr. 5
Alcoholfl.oz. 1
Tincture of iron chloridefl.dr. 1
Distilled watergal. 2

Dissolve the salicylic acid in the alcohol, add the tincture of iron, then add distilled water.

#### III.

Lead acetateav.oz.	1
Cochineal, powdergr. 5	00
Distilled watergal.	1

Macerate for several days; filter, and dilute to the desired shade.

#### IV.

Potassium permanganategr. Distilled watergal.	$\frac{40}{2}$
Mix and dissolve.	

## Show Globe Colors, Purple (Brilliant.)

Copper sulphate	gr.	120
French gelatin	gr.	60
Liquor potassa	fl.oz.	32
Water	suf	ficien

Dissolve the copper salt in 2 fluidounces of water, and the gelatin in the same amount

of boiling water; mix the two solutions, add the liquor potassa, shake the mixture, let stand 10 hours, decant the clear liquid, and dilute as desired with water.

## Show Globe Colors, Red.

I. Dissolve carmine in water of ammonia or liquor potassa, and reduce with water to the desired tint.

II. Take water in which red cabbage has been boiled; add sulphuric acid to bring out the color, dilute with water to the desired tint, and filter.

#### III.

Cochineal	100
Potassium bitartrategr.	75
Sulphuric acidfl.dr.	6
Distilled watergal.	2

Boil the cochineal and potassium bitartrate in water until exhausted; allow to cool, add the sulphuric, acid and filter.

· Instead of this, the solution of cochineal (cochineal coloring) of the National Formulary may be used.

#### IV.

Cobalt carbonate	gr. 60
Hydrochloric acid,	
Ammonium carbonate	of each sufficient

Ammonium carbonate of each, sufficient Distilled water, enough to make gal. 2

Dissolve the cobalt salt in the acid and some water, add enough ammonium carbonate so that the precipitate first formed is redissolved; filter, and dilute as desired.

V. Add to the amount of water necessary to fill the show bottle, compound tincture of iodine (drop by drop) until the desired tint is obtained; then add a few scales of metallic iodine. The iodine is added in excess to prevent bleaching, which would occur very soon were it omitted.

## VI.

	Tincture of iron chloridefl.oz.	2
	Potassium or amnionium sul- phocyanide gr.	40
,	Watergal.	2

Dissolve the sulphocyanide in water, add the tincture, and filter.

#### VII.

Solution of iron chloride		
Aqua ammonia	.fl.oz.	2
Acetic acid	.fl.oz.	2
Alcohol	fl oz.	8
Alcohol	gal	9
Water	gai.	~

alum (6 grains to the gallon) and filtered; the Potassium chromate......av.oz. 8

The water should first be clarified with V.

the required color is obtained.

Dissolve and filter.

Potassium bichromate.....av.oz. 6

Sodium carbonate or bicarbonate.av.oz., 4
Distilled water....gal. 2

Dissolve the potassium bichromate in the water, add the sodium salt (dissolved); filter.

III. Add tincture of curcuma to alcohol till

 Picric acid
 av.oz.
 ½

 Water
 ...gal.
 2

II.

other ingredients mixed, and the water added,	Watergal. 2
and the whole again filtered.	Dissolve and filter.
VIII.	VI. ·
Alumgr. 100 Potassium iodideav.oz. 2½ Distilled watergal. 2  Dissolve the alum and potassium iodide in	New York chrome yellowav.oz. 1 Hydrochloric acidfl.oz. 1 Nitric acidfl.oz. 2 Watergal. 2
the distilled water, and filter.	Mix; dissolve and filter.
IX.	
Cudbearav.oz. 1 Nitric acidfl.oz. 8	Silver Plating. See "Plating with Gold, Silver," etc.
Watergal. 2	Silver, Polishing.
Mix; allow to stand for 24 hours, and filter.	See "Polishing Powders."
Show Globe Color, Violet.	The following may also be employed:
I. Cudbearav.oz. 2	Sodium thiosulphate (hyposul-
Ammonia waterfl.oz. 8	phite)
Watergal. 2	Mix; dissolve and filter.
Mix; macerate for 24 hours, and filter.	In using, apply this liquid to the oxidized
II. Cobalt nitrateav.oz. 4	silver, rub dry, and polish with one of the
Ammonium carbonategal. 2	polishing powders adapted to silver.—D.
Copper ammonio-sulphate	The following is commonly employed:
of each, sufficient Watergal. 2	Prepared chalk or whitingav.oz. 2
Dissolve the cobalt nitrate in the water	Ammonia water
saturated with ammonium carbonate, and	Water, enough to makefl.oz. 8
add of the copper ammonio-sulphate sufficient	Soap, Ox-Gall.
to produce the desired tint.	I.
Instead of the copper salt may be used a	Extract of quillaja av.oz. 1 Borax, powder
solution of copper sulphate to which is added	Borax, powderav.oz. 1 Ox-gall, freshfl.oz. 4
ammonia water until the precipitate first	Common or castile soap, powder. av. oz. 15
formed is redissolved.	Triturate together the borax, extract and
Show Globe Colors, Yellow.	ox-gall, dissolving as much of the borax as
I. Potassium bichromateav.oz. 10	possible. Then add the soap, beat the whole
Nitric acidfl.oz. 20	to a uniform consistence, and cut into cakes
Distilled watergal. 2	of the desired size.
Dissolve the potassium bichromate in the	If no extract of quillaja be at hand, soap bark (in shreds) may be exhausted by boiling
water, and add the nitric acid; filter.	bark (in shreds) may be exhausted by boiling

with water, straining, and evaporating the

liquid on a water bath. One hundred parts

of bark yield about 20 of extract.-D.

Triturate the borax with the ox-gall; then thoroughly incorporate with it the soap, previously reduced to powder, and lastly incorporate the oleic acid.

#### III.

Ox-gall, freshfl.oz.	10
Stearin soapav.oz.	9
Borax, powderav.oz.	1
Alcoholsufficie	ent

Mix the first three ingredients at a slightly elevated temperature, then add sufficient alcohol (from 1 to 2 fluidounces) to form a suitable mass; transfer the whole to a flatbottomed vessel, and when cold cut into pieces.—D.

# Soap, Shaving.

Mutton suetav.oz.	10
Cocoanut oilav.oz.	5
Caustic sodaav.oz.	2
Caustic potassagr.	170
Waterfl.oz.	71/2
Oil of carawaydrops	$25^{'}$
Oil of bergamotdrops	30
Oil of lavenderdrops	20
Oil of thyme, whitedrops	12
Oil of mirbanedrop	1

Melt the tallow and cocoanut oil; allow to cool to 50 degs. C., then add the caustic potassa and soda dissolved in the water, and warm the whole gently during one-half hour or so, stirring occasionally, until a uniform soapy mass is produced; to the latter add the volatile oils.—D.

# Soap, Shaving, Antiseptic.

To the previous soapy mixture, add 3/4 av. ounce of salol, first warming the soap to about 50 or 60 degs. C., and stir until the salol is dissolved.—D.

### Soap, Soft. (Green Soap.)

For pharmaceutical use:

Olive oilfl.oz.	25
Caustic potassaav.oz.	91/2
Waterfl.oz.	75

Dissolve 8 av.ounces of caustic potash in 50 fluidounces of water, and add 12½ fluidounces of this solution to the oil. Heat the mixture over a moderate fire, stirring until sufficiently thickened. Gradually add the remaining solution of potassa, and continue the heat, stirring occasionally until the mixture assumes a transparent, gelatinous form; dissolve the remaining 1½ av.ounces of potassa in 25 fluidounces of water; add this solution to the soap mixture, and evaporate the whole to proper consistency.

# Soap, Stearin.

Stearic acid	.av.oz.	21/4
Sodium carbonate, crystal	gr.	580
Water		
Alcohol	fl.dr.	2
Sodium chloride	fl.dr.	4

Dissolve 540 grains of the crystallized sodium carbonate in 6 fluidounces of water; transfer this solution to a water bath and gradually add the stearic acid with constant agitation; then add the alcohol; cover the vessel, and allow it to remain upon the bath for 6 hours to separate the soap; add the sodium chloride and the remainder of the sodium carbonate, dissolved in the remainder of the water; transfer the whole to a strainer, and when cold press out the remainder of the moisture.

### Soap, Whale Oil.

This may be prepared like soft soap, substituting whale oil for the olive oil.

# Stains from Fabrics, Removing.

See also the following headings: "Ox-gall Soap," "Cleansing Creams," "Cleansing Liquids," "Benzin Jelly," "Glove Cleaner," "Rust Stains, Removal of," and "Silver Nitrate Stains."

### Cleansing Pencils.

Form little rolls from ox-gall soap, half an inch thick and  $2\frac{1}{2}$  inches long, and cover with tinfoil. Instead of ox-gall soap, the following may be employed:

Borax, powderav.oz.	2
Common or castile soapav.oz.	14
Green soapav.oz.	4

Mix to a uniform mass, using a gentle heat if necessary.

#### Cleansing Liquid.

Spirit of ammoniafl.dr. 12	,
Oil of turpentinefl.dr. 10	):
Etherfl.dr. 12	3
Oil of lavender flowers fl.dr. 1	
Alcoholfl.oz. 27	
II.	
Spirit of ammoniafl.dr.	5
Ether fl.dr. 14	Ļ

Etner	100
Benzin	5
Oil of lavender flowers fl.dr.	1
Tincture of quillajafl.oz.	7
Alcoholfl.oz.	15
THE OHOU, I I I I I I I I I I I I I I I I I I I	

III.	
Oil of turpentinefl.oz.	3
Benzinfl.oz.	
Ammonia water fl.oz.	3
Alcohol	23

IV.

Benzin fl.oz.	32
Oil of turpentinedrops	8
Oil of mirbanedrops	

The last, sometimes called "Brunnersches fleckwasser," (by which term benzin only is also signified) is particularly suited for cleansing gloves.

Before applying any of the cleansers recommended to colored goods an experiment should be made, either with a sample of the goods or on some portion which will not be seen, to determine whether directions given for treating the spot will not affect the color.

The Spot is of Unknown Origin.

White Goods.—Dissolve some soap in lukewarm water and add two dessertspoonfuls of cleansing liquid (as above) and dampen the spot with a sponge soaked in this solution; finally wash out in clear water.

Colored Woolens.—Dissolve a cleansing pencil (as above) in a bottle of the solution and wash out the spot in the liquid; then rinse in clear water, and dry in the air.

Silk, Satin and Similar Delicate Fabrics.—Add to the above solution the yolks of two eggs, and spread this on the spot. Then wash in lukewarm water; rinse in cold water, and dry by a gentle heat. To press out use an iron that is warm only—not hot.

The Spot is of Dust.

White Goods.—Beat and brush.

Colored Wool, Silk, Satin, etc.—Old spots that are dried in should be painted first with yolk of egg, then with cleansing solution and allowed to dry. Scratch this off and wipe with a wet linen rag.

Milk, Soup, or Small Grease Spots Generally.

White Goods.—Wash with the warm solution of a "cleansing pencil" in water.

Colored Cotton or Woolen Goods.—Dampen with cleansing solution; remove the excess of the solution by means of blotting paper, and wash with a solution of a "cleansing pencil."

Silk, Satin, etc.—Dampen by means of a

sponge soaked in cleansing liquid I., removing any excess by means of blotting paper.

Butter, Grease, Oil, Paints, Varnish, etc.

White or Colored Woolens or Cotton Goods.

—Moisten several times with cleansing liquid
I., lay a piece of blotting paper over the spot
and press this with a hot iron. Then wash
the whole of the fabric in hot soap suds.

Silk, Satin and Delicate Fabrics.—Rub up some "white bole" or talcum; thin dough with cleansing solution II., and spread over the spot. When thoroughly dry brush off and wipe with dry bread crumbs.

When the Spots are Old.—First moisten with chloroform and then proceed as above.

The Spot is from Stearin, Wax, etc.

First remove as much as possible with a knife; then lay a damp towel under the spot and put several thicknesses of blotting paper over it and press out with a hot iron. If any stain remains after this, treat as directed under butter, grease, etc.

Resin, Tar, Axle Grease, etc.

White Goods.—Wet with good oil of turpentine, wring out, cover with blotting paper, and go over with a hot iron. Then wash in warm soap suds.

Colored Cotton or Woolen Goods.—Moisten the spot, apply butter, soap thoroughly, allow to stand for a few minutes, and then wash with oil of turpentine and hot water, alternately. If this does not help, spread over the spot the yolk of an egg previously mixed with oil of turpentine; cover with blotting paper and press with a hot iron. Then scratch off the residue and wash thoroughly. As a final resort, wash out in water slightly acidulated with hydrochloric acid.

Silk, Satin, etc.—Drench with chloroform, and, when this has evaporated, apply "white bole or talcum;" cover with blotting paper, and press with hot iron. If this does not help, mix some yolk of egg with chloroform and proceed as above, removing the residue by wiping off with bread crumbs.

Vinegar, Acid, Wines, Fruit, etc.

White Goods.—Wash out with clear water to which a little "cleansing liquid" II. has been added.

Colored Goods, Whether of Cotton, Wool

or Silk.—Moisten with cleansing liquid, allow to evaporate, and then rinse in clear water.

#### Acids.

Fresh spots may be removed by putting on a drop of cleansing liquid; old spots cannot be remedied.

Colored Fruit Stains from Peaches, Red Wine, Cherries, Strawberries, etc.

White Goods.—Dip in javelle water or solution of chlorinated soda, and immediately that the stain has disappeared wash thoroughly in clear water.

Colored Cotton or Woolen Goods.—Wash with hot soap suds, to which a smaller or larger quantity of javelle water, or solution of chlorinated soda, has been added (as the fabric is more or less delicate); rinse in water to which a little cleansing liquid has been added; finally, wash in a large quantity of clear water.

Silk, Satin, etc.—Follow directions as above, save to use very dilute solutions.

### Grass Stains.

White Goods.—Wash out with boiling water.

Colored Goods, Whether of Cotton, Wool or Silk.—Moisten the spot with a very dilute solution of tin chloride, and then wash thoroughly in a plentiful supply of clear water.

Stains from Tannin, Green Nuts, etc.

Treat with very dilute javelle water, or solution of chlorinated soda.

#### Coffee or Chocolate Stains.

Cover the spot with yolk of egg diluted with cleansing liquid; wash out in warm water, and iron (while still moist) on the wrong side of the cloth.

#### Aniline Ink Stains.

White Goods.—Wash with alcohol to which a little acetic acid has been added, and then bleach with javelle water or solution of chlorinated soda.

Colored Goods, Whether Cotton, Wool or Silk.—If the color admits of it, follow the directions for white goods. If the dye is too delicate for this wash out with strong acohol alone, as nothing else will prove of benefit.

See also "Ink Erasives."

#### Stains for Wood.

By wood stains are understood solutions of dyes, etc., used for coloring wood. They are fixed on the wood either direct or through the medium of some mordant. In many cases the color is developed in the grain only after the mordant is applied, sometimes the mordant merely changes the tone of the color.

The action of the stain is influenced not only by the mordant, but also by the natural constituents of the wood—tannin, for example. Consequently different woods sometimes give different results with the same stain.

Before applying a stain, the wood should be smoothed by sandpapering. After applying the stain, the wood should be polished so as to "bring out" the grain.

# Stain, Black or Ebony.

Solution I.

Sodium chlorateav.oz.	1
· Copper chlorideav.oz.	1
Waterfl.oz.	15

Solution II.

Aniline	e hydrochlorateav.oz. 🖇	1/2
Water	e hydrochlorate av.oz. 2 	j´ -

The dry wood is painted three times with the above solutions, applying them alternately; before each application the wood is well dried; finally, it is rubbed with linseed oil or a mixture of turpentine and wax, and polished. The color is not affected by acids or alkalies.

#### Stain, Cherry.

4
1
8

Boil until the annatto is dissolved.

II.

. 2.	
Logwood chipsav.oz	
Caustic potassaav.oz	i. ½
Waterfl.02	z. 16

Boil until the color is extracted, adding more water from time to time, to make up for the loss by evaporation.

The stain is to be "fixed" by washing the wood, after its application, with alum water.

# Stain, Mahogany.

 Madderav.oz.	2
Logwood chipsav.oz.	1/2
Waterfl.oz.	32

Mix; boil and stain. Apply to wood while hot and mordant with an aqueous solution of potassium carbonate, 60 grains to the pint. II.

Alkanetav.oz.	1/2
Aloesav.oz.	1
Dragon's bloodav.oz.	1
Alcoholfl.oz.	16

Reduce the drugs to coarse powder; mix with the alcohol, set the whole in a warm place for 3 or 4 days, agitating occasionally, and filter. Before applying, mordant with dilute nitric acid.

### Stain, Walnut.

Potassiur	m permanganateav.oz. waterfl.oz.	1/2
Distilled	waterfl.oz.	16

Apply twice in succession, and after an interval of five minutes wash with clear water.

A strong hot decoction of green walnut shells may also be applied, followed, when partially dry, with a concentrated solution of potassium bichromate.

### Stamping Powders.

These are employed for stamping embroid eries, etc. Powders of various colors are rendered adhesive by admixture with gum resins, such as resin, copal, damar or sandarac. The substances should be made into the most impalpable power by trituration and sifting.

The method employed for stamping is to perforate paper according to the pattern desired, then placing this upon the fabric, sprinkling or rubbing the powder into the perforations (carefully removing the pattern), placing a piece of unperforated paper on the cloth, and carefully passing a hot iron over the whole. The iron melts the resin and leaves the design imprinted on the material.

I. Mix equal parts powdered resin and a pigment-ultramarine or prussian blue for blue; zinc oxide or flake white for white; chrome yellow for yellow; burnt or raw umber, burnt or raw sienna, vandyke brown, etc., for brown; ivory black for black, etc.

II.

Resin, Damar resin, Copal resin, Sandarac,

Pigment.....of each, equal parts Reduce each to very fine powder, and mix

well.

# Storm Glass or Baroscope Solution. (Chemical Barometer.)

I.	
Potassium nitrate gr.	36
Ammonium chloridegr.	36
Camphorgr.	180
Absolute alcohol fl.dr.	
Acohol	6
II.	
Camphor	120
Potassium nitrate gr.	
Ammonium chloridegr.	60
Diluted alcoholfl.oz.	21/4

Mix and dissolve, and place in a glass tube about 12 inches long and 34 inch in diameter; the tube to be filled about three-fourths and tied over with a bladder.

TIT

Potassium nitrategr.	30
Ammonium chloridegr.	
Camphor gr.	
Alcoholfl.oz.	Z

Put the mixture into a bottle 18 inches in length and 3/4 inch in diameter, and cover the mouth with a piece of perforated plaster. If the weather promises to be fine the insoluble matter will settle at the bottom of the tube, while the liquid remains pellucid; but previous to a change for rain, the compound will gradually rise, the fluid remaining trans-Twenty-four hours before a storm or very high wind the substance will be partly on the surface of the liquid, apparently in the form of a leaf; the fluid in such cases will be very turbid and in a state resembling fermentation.

IV. This mixture is also used: Ammonium chloride.....gr. 60 Camphor.....gr. 60

Potassium nitrate.....gr. 60 Alcohol .....fl.oz. 21/4 Distilled water, hot.....fl.oz. -D.

# Stove Polish or Blacking.

Boiling water.....fl.oz. 16 Black lead . . . . . . . . . . . sufficient

Dissolve the soap in the water, and add enough of the black lead to form a paste.

Plumbago .....av.oz.16 Gum turpentine.....av.oz. 4 Sugar.....av.oz. 1

Knead thoroughly and keep in tin boxes.

Apply with a brush.

III. Plumbago made into a thin paste with sodium silicate or water glass. This makes an excellent stove polish and should be brushed thoroughly.

IV. Reduce graphite to an impalpable powder by grinding in a mill with water; dry; use with water first, then dry and polish. This is the base of nearly all commercial stove polishes.

V.

Bone blackav.oz. 2
Pulverized graphite 2
Copperasav.oz. 4
Watersufficient to form a creamy paste

### Tableau Lights.

See "Colored Fires."

#### Tar Stains, Removal of.

See "Stains from Fabrics, Removal of."

# Test Papers.

In preparing test papers, only the best white filter paper or letter paper should be employed. In order to remove traces of acid, which are so often present in paper, it should be macerated for 24 hours in about a 1 per cent ammonia water; then dry by suspending on lines in a room of ordinary temperature.

In making the test paper, the prepared paper above should either be drawn through the impregnating liquid, expressing the excess by means of a glass rod, or else the mixture should be applied to one side of the paper by means of a broad, soft brush. After impregnating the paper, the latter should be dried by suspending on lines.—D.

### I. Azolitmin paper:

Azolitmin gr. 15
Sodium carbonate, puregr. 8
Distilled waterfl.oz. 32
Phosphoric acidsufficient

Dissolve the azolitmin and sodium earbonate in the water, and neutralize the solution with the acid. Pass filter paper through the solution and dry, as directed above.

This paper becomes red with acids; it will indicate 1 part of sulphuric acid in 40,000, and 1 of hydrochloric acid in 50,000.

# II. Brazil wood paper:

Brazil wo	ood, raspedav.oz.	21/2
	waterfl.oz.	

Mix; macerate for 24 hours, agitating frequently; filter, and to the filtrate add enough

ammonia water, drop by drop, until it begins to acquire a blue-red color. Prepare the paper as in the preceding.

This paper will indicate 1 part of ammonia in 80,000.

#### III. Congo red paper:

Congo red (coal	tar	dye)	 gr.	15
Alcohol			 fl.oz.	28
Distilled water.				

Dissolve the dye in a mixture of the alcohol and water, and prepare the test paper as in the preceding instance.

This paper will indicate 1 part of sulphuric acid in 2,500, and 1 part of hydrochloric acid in 3,000.

A blue congo paper may be produced by adding alkali to the above solution.

### IV. Curcuma paper:

Curcuma	root, powdergr.	5
Distilled	waterfl.oz.	16
Alcohol .	sufficio	ent

Macerate the drug with 4 fluidounces of alcohol for several days, agitating frequently, and filter, adding enough alcohol through the filtrate to make 4 fluidounces. To the latter add a mixture of the water and 15½ fluidounces of alcohol. With this liquid impregnate paper as in the preceding instance.

This paper will indicate 1 part of potassium hydrate in 15,000, and 1 of ammonia in 40,000.

### V. Lead paper:

Lead ace	etate								av.oz.	3
Distilled	water								.fl.oz.	30

Dissolve, and prepare the paper as in the preceding.

### VI. Litmus paper, blue:

Litmus,	bestgr	. 730
Distilled	1 water,	

Phosphoric acid.....of each, sufficient

Macerate the litmus in 32 fluidounces of water for 12 hours; filter, and add through the filter enough water to make the filtrate measure 32 fluidounces. To the latter add phosphoric acid, drop by drop, until the liquid appears blue, with reddish cast. Then impregnate paper as in the preceding instance.

This paper will indicate 1 part of sulphuric acid in 40,000, and 1 of hydrochloric acid in 50,000.

VII. Litmus paper, red:

Litmus, best......gr. 672
Distilled water ......fl.oz. 32
Phosphoric acid......of each, sufficient

Macerate the litmus with the distilled water for 24 hours; filter, and add enough of the acid to filtrate until the fluid is red; then set aside for 24 hours, decant the clear liquid, and filter. With this liquid impregnate paper as in the preceding instance. The second filtration is necessary to remove a brownish substance which is deposited. Hydrochloric acid may be substituted for the phosphoric acid.

This paper will indicate 1 part of potassium hydrate in 20,000, and 1 of ammonia in 60,000.

VIII. Logwood paper:

Logwood, raspedgr.	585
Distilled water fl.oz.	32
Ammonia watersuffic	ient

Macerate the logwood with the water for 24 hours; filter, and to the filtrate add ammonia water, drop by drop, until the liquid assumes a dark blue-red color. With this liquid impregnate paper as in the preceding instance.

When freshly prepared, this paper will indicate 1 part of ammonia in 80,000 to 90,000.

### IX. Potassium iodide-starch paper:

Wheat starch	. áv.oz.	1
Potassium iodide		
Distilled water	fl.oz.	35 1/2

Mix the starch thoroughly with 1 fluidounce of water, gradually add the remainder of the water, in a boiling condition; heat the whole on a water bath for 30 minutes, then add the potassium iodide, and dissolve.

The test paper is prepared by painting the solution upon one surface of letter paper by means of a broad, soft brush and then drying.

# X. Starch paper:

Wheat st	arch gr	. 150
Distilled	waterfl.oz	. 32

Intimately mix the starch with 3 fluidrams of water, and then add the remainder of the water in a hot condition. With this mixture prepare test paper as in the preceding instance. A caution to be observed is that the brush must not be passed over the same spot twice

as this would loosen some of the fibers of the paper.

This paper will indicate 1 part of free iodine in 25,000.

### Tin Plating.

See "Plating with Gold, Silver," etc.

#### Twaddell's Scale.

The Twaddell is an old degree scale largely used in England, principally among the dyers' to indicate the strength of solutions of mordants, etc. It is an arbitrary standard, something similar to Baume scale. The following rule is used for converting Tw. degrees into specific gravity: Multiply Twaddell's degrees by 5, add 1000 and divide by 1000. For example: To reduce 64 degs. Tw.  $(64\times5)+1,000$  divided by 1000=1.320 sp. gr. Or another, method is to multiply by 5, cut off 3 decimal places, and add 1:  $64\times5=320$ ; a decimal point will make 0.320; then add 1=1.320.

#### Varnishes.

These, like lacquers, are resinous solutions intended as protective applications to metals, wood, etc.

### Varnish, Amber.

1.		
	Amber, coarse powderav.oz.	8
	Oil of turpentinefl.oz.	13
	Mix and dissolve.—D.	

11.		
Amber	.av.oz.	8
Oil of turpentine		
Linseed oil varnish		

Melt the amber out of contact with the air, allow to cool somewhat; add the oil, and then the varnish.—D.

#### Varnish, Anatomical.

Mastic	v.oz.	3
Sandarac		
Camphor	gr.	110
Venice turpentine	v.oz.	3/1
Alcohol	fl.oz.	28
Mix and dissolve.		

This is used for dry anatomical specimens.—H.

#### Varnish, Black.

Linseed oil varnishfl.oz.	10
Burnt umberav.oz.	2
Asphaltum, powderav.oz.	4
Oil of turpentinesufficie	

A caution to be observed is that the brush must not be passed over the same spot twice asphaltum is dissolved, then remove from the

fire and add oil of turpentine until the liquid is of proper consistency.—11.

### Varnish, Bookbinders.

I.	
Shellac	av.oz. 4
Benzoin	av. oz. $1\frac{1}{2}$
Sandarac	av.oz. 1
Mastic	
Oil of lavender	fl.dr. 2
Absolute alcohol	fl. oz. 24

Mix, macerate for some time, agitating occasionally; decant the clear liquid, and filter.—H.

#### II.

Shellac av.oz.	4
Masticav.oz.	1
Benzoin av.oz.	2
Venice turpentinegr.	150
Alcoholfl.oz.	20

Macerate a few days, agitating occasionally, and filter.—H.

### Varnish, Celluloid.

Pyroxylin (soluble gun cotton)gr.	200
Etherfl.oz.	4 1/2
Alcoholfl.oz.	6
Camphor,gr.	120

Pour the ether over the pyroxlin, add the alcohol and finally add the camphor.

This varnish may be colored by the addition of anilines. It is particularly adapted for covering paper labels.—D.

### Varnish for Chocolate Candy.

Sumatra benzoingr. 8	600
Shellac, palegr. 5	600
Vanillingr.	
Alcohol, enough to makefl.oz.	

Dissolve the first three ingredients in 14 fluidounces of alcohol; filter, and pass enough alcohol through the filter to make the filtrate measure 16 fluidounces.—D.

### Varnish, Copal.

Copal resin	 				av.oz.	8
Linseed oil	 				 .fl.oz.	6
Oil of turpentine	 				 .fl.oz.	6

Melt the copal; add the linseed oil, and when nearly cool add the oil of turpentine.

### Varnish, Dammar.

Dammar resin						.av.oz.	8
Oil of turpentine						fl. oz.	13

Melt the resin carefully over the direct flame; allow to cool, pulverize, and dissolve in the oil. This varnish may be prepared by dissolving the resin without the preliminary fusion, but the product will always be sticky.—D.

### Varnish, Furniture.

Shellacav.oz.	7
Resingr.	175
Absolute alcoholfl.oz.	20
Turpentineav.oz.	1
Talc, powderav.oz.	3/4

Warm the shellac and resin; add the absolute alcohol, and finally, the turpentine and talc. Shake vigorously for several minutes and stand in a cool place. After 8 days filter through a filter which has been previously wetted with alcohol.—D.

### Varnish, Grecian.

Balsam of fir	.av.oz.	6
Oil of turpentine	.fl.oz.	2
Alcohol	.fl.oz.	4

### Varnish, Label.

I.

1.		
	Sandaracav.oz.	41/2
	Masticav.oz.	2
	Camphor gr.	35
	Oil of lavenderfl.dr.	5
	Venice turpentinegr.	150
	Etherfl.dr.	4
	Alcohol fl.oz.	4

Macerate for several weeks, agitating frequently until dissolved, and decant or strain from impurities. The varnish dries rapidly to a colorless, smooth and glossy layer.

II.

Sandaracav.oz.	3
Masticav.oz.	3/4
Venice turpentinegr.	150
Alcoholfl.oz.	16

Macerate with repeated stirring until solution is effected, and then filter.

Paper labels are first sized with diluted mucilage, then dried, and then coated with this varnish. If the labels have been written with water-soluble inks or color, they are first coated with two coats of collodion, and then varnished.—D.

#### III.

Shellac, bleachedav.oz.	
Balsam of copaibaav.oz. Venice turpentinegr.	1/2
Alcoholfl.oz.	16

Prepare and use like the preceding.—D.

### Varnish, Linseed Oil.

I. Evaporate 100 parts of linseed oil over the naked flame, stirring constantly, until it weighs 90 parts. Allow to cool, and add 5 parts of oil of turpentine.—D.

II. Heat 100 parts of linseed oil with  $2\frac{1}{2}$  of litharge over the naked flame until there is no further effervescence. Then set aside for 14 days. The product weighs about 95 parts.—D.

III. Heat 100 parts of linseed oil with 4 parts of manganese borate over the naked flame, with constant stirring until the yellow color of the oil changes to a pale yellowishgreen. The change in color may be observed by transferring a drop of the liquid, from time to time, to a porcelain plate. The termination of the reaction is also denoted by the discontinuance of effervescence. The varnish should now be removed from the fire and cooled rapidly, if possible, by setting the dish in cold water; then set aside for 14 days. The product weighs about 93 parts.—D.

# Varnish, Map.

S	atur	ated	solu	tion	of	borax	 .fl.	oz.	12
S	hella	ac, fi	ne po	bwc	er.		 av	oz.	6
0.									

Shake together, but apply no heat.

### Jarnish for Metals.

Shellac, palegr.	500
Sandaracgr.	
Venice turpentinegr.	70
Alcohol, enough to make fl.oz.	16

Dissolve the shellac, sandarac and turpentine in 14 fluidounces of alcohol by maceration; filter and add enough alcohol to make 16 fluidounces.

This varnish may be applied to all kinds of polished metal.—D

# Varnish, Transparent.

Sandarac!av.oz.	2
Mastic	300
Venice turpentinegr.	90
Alcoholfl.oz.	12

After solution filter, and add sufficient alcohol to bring to the measure of 15 fluid-ounces.

#### Varnish Stains, Removal of.

See "Stains from Fabrics, Removal of,"

# Washing Powder. (Soap Powder.)

Washing powders, usually sold to the consumer as soap powders, may be described in a general way as mixtures of powdered soap, with about its own weight, more or less, of sodium carbonate. Some special brands are also made, which in addition contain other detergent agents, such as ammonium carbonate, sal ammoniac or borax; while still others are found, to which filling, in the form of talc, silex, etc., has been added. The soap itself may have been made by any of the processes known-cold, half-boiled, or boiled, settled, or boiled down-and the stock used may have been any fat, or mixture of fats, according to the grade of washing powder to be made. Here are some typical formulas:

I. Borax Soap Powder:

Curd (hard) soap, powderav.oz.	10
Soda ashav.oz.	
Sodium silicateav.oz.	4
Boraxav.oz.	2

Each ingredient is thoroughly dried, and all mixed together by sifting:

# II. London Soap Powder:

Yellow soapav.oz.	12
Pearl ashav.oz.	3
Palm oilav.oz.	

These ingredients are combined as well as possible without any water, and they are spread out to dry, and then ground into coarse powder. They are adapted to hard waters, as their excess of carbonated alkali neutralizes the lime in the water.

### III. Pearl Soap Powder:

Cura soap, powder	av.02. 0
Sal soda crude sodium carl	bon-
ate)	av.oz. 6
Sodium silicate	av. 0z. 4

Dry as much as possible and mix intimately.

# Window Polishing Paste.

Prepared chalkav.oz.	9
White hole av.oz.	1/2
Jeweler's rougeav.oz.	1/2
Waterfl.oz.	9
Alcohol fl.oz.	3

Make into a smooth paste and introduce into a wide-mouth bottle.

Moisten a cloth with alcohol, place upon the window glass a quantity of the paste of about the size of a bean, and rub the latter about on the glass with the cloth until dry and the powder is removed.—D.

ABC liniment	Aloes, extract of, acid	86
Abercrombie's lotion of borax 111	Extract of, fluid	86
Abernethy's pills	and myrrh, fl. extract of	86
Absinthium, (see Wormwood)	Tincture of, compound	64
Abstracts 11	Crocated	70
Acacia extract		161
Sachet	Aloin and strychnine, elixir of	32
Acacine	Strychnine, and belladonna, elixir of	32
ACE mixture		343
Acetanilid, elixir of	Alstonia constricta, fl. extract of	86
Acetic cantharidal vesicant 207	Alterative compound.	86
Acid phosphates, solution of129, 366	Elixir	57
Salicylic, elixirs containing		168
Stains, removing		
		135
		140
		118
Acne, remedies for	Alum bougies	16
Aconite chloroform		104
Collodion	Aluminated copper	24
Leaf, tincture of		129
Liniment	Aluminium acetate cotton	25
Ointment		129
Aconitine ointment		388
Acorn water, Rademacher's 159		321
Acoustic oils	Amarum, elixir	41
Actina	Amber oil, liniment of, compound 1	109
Adhæsol	Varnish	166
Adhesive and strengthening plaster 121	Ambergris 2	280
Adjuvant elixir		284
Ague cures		284
Alabaster, cement for 385	Tincture of	285
Albadermine	Ambrosia syrup	
Albolene		167
Albuminate of iron and sodium, syrup. 144	Ammonamide	
Mercury, solution of		374
Alcohol, deodorized		109
Dilution table	Solution of, anisated	129
Homeopathic		137
Alcoolatures		116
Alcooles	Ammoniated copper	$^{110}_{24}$
Alcoolats		$130^{-4}$
Aletris cordial	Ammonium benzoate, solution of I Bromide, elixir of 33; chloride, elixir of .	
Elixir of	and licorice elixir compound	33
Alizarin ink	~ .	139
		35
Alkaline ointment	and morphine valerianates, elixir	35
Ointment, camphorated	Valerianate, elixirs containing33-	
Allcock's porous plaster		130 208
Allen's (Mrs.) hair restorer	_	
Alloys of low melting point 374		208
Almond cold cream 310		245
Cream		217
Emulsion of		209
Compound		209
Meal 322	1	206
Milk	1	208
Compound		466
Mixture 79	I amount of the second of the	209
Oil, emulsion of80	1	$120^{\circ}$
Paste		309
Spirit of	Angelica, spirit of, compound 1	136
Bitter 137	Angier's petroleum emulsion S	378

A . T 11 11/1 C-:-:- 1977	Aughten heleem 9444
Angleworm oil, 114. Spirit 137	Arabian balsam
Angostura bitters	Aralia, syrup of, comp
Aniline ink 417	Arbor vitæ, elixir of
Stains, removing	Tincture of
Anisated solution of ammonia 129	Armenian cement 384
Anise, elixir of	Pills
Aniseed cordial	Arnica jelly
Annatto, extract of	Liniment
Solution of	Salve
Anodyne liniment	Arnicated cotton
Oil	Aromatic cascara
Turp. solut., Rademacher's 135	Confection
Antasthmatic remedies 164	Elixir
Anti-canker pills, Thompsonian 120	Ointment 118
Anti-catarrhal pills, Andrews' 120	Species
Anti-chill pills	Spirit
Anti-constipation pills, Carson's 120	Tea 135
Pills, Fothergill's	Vinegars
Antidiphtherikon	
Anti-dyspeptics	Arsenauro
Antifebrin, elixir of	Arsenic, elixirs containing35, 36
Anti-galactagogue	and gold bromide, sol. of
Anti-grippe pills	Paste 388
Antikamnia	Pills, Hebra's
Anti-kink hair pomade: 341	Arsenical caustic, Ratier's
Antikol	Powder, Come's
Antikrinin	Asafetida, syrup of
Anti-lacteant	
Antimonial ointment	Aseptic acid
Powder, Tyson's	A septinic acid
Antimony, butter of	Asthma cigarettes, Plant's
Chloride, solution of	Cure, Himrod's
Antinervin	in Canaries
Antineuralgic pills	Remedies
Antipyonin244A	Remedy, Upham's
Antipyrin	Astringent ointment, Thompsonian 116
Antipyrin, elixir of	Athenstædt's comp. tinct. iron 154
Antirheumatic cotton	Athlophoros
Antirheumatin	Atropine, discs of
Antiseptics	2.0
Antiseptic solution, Lister's 130	Ambergier's syrup
Seiler's	Augenstein
Tablets, Wilson's	Augentrost, Kneipp's
Antiseptin	August flower
Ants, spirit of	Australian fever bark, fl. ext., of 86
Aphrodisiae Elixir 193; pills 193	Avena sativa, tincture of
Apis mellifica, tíncture of	Axle grease
Apomorphine, hydrochlor., syrup of 139	Grease stains, removal of374, 462
Injection of	Ayer's cathartic pills
Appetite, loss of. in dogs and cats 271	Cherry pectoral
	2:0
Loss of, in horses	210
in swine	2.2011
Aqua, (see Water)	Recamier balm
Anethi	Cream 210
Castoreum	Powder
Cerasorum nigrorum 160	Soap 210
Crystallina 160	Sarsaparilla
Glandium	Vita nuova
Hungarica	Azolitmin paper
Nicotianæ	Paparation and the same
Nucum vomicarum	RABBITT'S 1776 washing powder 243
Strychni seminis	Data de la constante de la con
Aquarium cement	Balm, catarrh

...

Balm, Cream	Baume's scale	375
Ely's	Baunscheidt oil	114
Gilead, factitious	Bay rum	306
Spirit of	Bayberry ointment	116
Wizard 191	Plaster	121
Balms for skin	Syrup, Thompsonian	140
Balsam, blackberry	Beach's diaphoretic powder	122
Carminative	Bear's grease109,	340
Copaiba, factitious	Oil	337
Cough	Beauchamp's pills	211
de Maltha 13	Becker's eye salve	211
Fioravanti	Beckwith's hog cholera remedy	211
Friar's	Bedbug exterminators	376
Guaiac	Beecham's pills	211
Hair344, 345	Beef extract, elixirs containing	36
Honey	Extract of malt with	97
Linseed or flax seed, cough 179	Wine and iron	161
Lung 177	with cinchona	161
Magen	with coca	161
Malta	and iron, ext. of malt with	97
Mecca, factitious         12           Metz's         13	Bees, tincture of	151
	Begg's ague pills	211
Muskat	Bejean' gout cure.	211
	Belladonna chloroform	20
70	Collodion	20
	Infused oil of	114
	Syrup of	$\frac{140}{378}$
	Bengal lights	14
~ 0 .	Benzin, deodorized	378
Soap	Gelatinized	378
Turpentine	Jelly Benzoated cotton	26
Universal	Suet	139
Balsamic fumigation	Benzoic acid	281
Balsamum nucistæ	Benzoin, fl. ext. of	87
Bamboo brier, fl. extract of, comp 86	Fl. ext. of., comp.	86
Bandages, plaster of Paris	Tincture of	285
Bandoline 341	Benzoinol	211
Bar mucilage	Benzothymol	211
Barber's itch, remedies for 166	Berberine, elixirs containing	37
Bareel's Indian liniment	Berberis aquifolium fl. ext. of	87
Barber's nerve and bone liniment244A	Bergamot, spirit of	284
Barley, decoction of	Bibron's rattlesnake poison rem	205
Water	Bicycle liniment	191
Barometer, chemical	Oil, illuminating	378
Paper375	Oil, lubricating	378
Baroscope solution	Paint, black	378
Barrels, to determine capacity 440	Tire cement	
Barr's dental anæsthetic	Big G	
Barry's tricopherous	Biliousness, remedy for	173
Barton's lotion	Billroth's transfusion fluid	158
Bate's salve	Birch essence	363
Bateman's pectoral drops	Birch syrup	
Bath powder	Bird elixir	C 10/10/
Oils 362	Medicines	
0.0.1	Seed, mixed	
200	Bishop's gran. eff. cit. magnesia	
Perfumes.         362           Tablets         362	Bismuth, elixirs containing 37–	41
Sulphur	and pepsin, effervescent	127
Battery, filling for dry	Bismuthol	
Fluid	Bitter almond, spirit of	
Storage	Elixir	41
Battley's sedative	Species	
Baume de fioravanti	Stomach drops	158

Bitter Tea	135	Bloodroot, Vinegar of	153
Tonic tea, Boecker's			
		Bloom of Ninon	
Wine	66	Roses	
Bitters	166	Youth	-228
Ague	164	Blue-black ink	421
Angostura		Cohosh, tincture of	
Boerhaver's	212	Tincture of, comp	
Boonekamp's		Flag, elixir of	
Harter's wild cherry		Tincture of	
Hop	225	and wahoo, elixir of	42
Hostetter's		Green ink	
Swedish	64	Ink	
	14		
Thompsonian		Print paper	000
Wine	162	Bluing of copper	380
Bitterless buckthorn bark	- 88	Liquid	-380
Fl. ext. of buckthorn	88	Blutreinigungsthee, Kneipp's	108
Cascara sagrada	90	Boecker's bitter tonic tea	
Fl. ext. of cascara sagrada	90	Boerhaver's bitters	
	87		
Bittersweet, extract of		Boil remedy	
Black caustic, Velpeau's	18	Boiler compounds	
Cohosh, elixir of	41	Cover cement	-383
Elixir of, compound	42	Boiling points of solutions	-380
Fl. ext. of, comp	87	Boonekamp's bitters	211
Tincture of, comp	151	Bookbinder's glue	
Eye, remedy for	205	Mucilage	
How olivir of	41	Varnish	
Haw, elixir of			
Elixir of, compound	42	Borated cocaine cotton	20.0
Ink419,	420	Cotton	26
Liniment	109	Borax, glycerite of	104
Pepper ointment	118	Honey of	-105
Powder	122	Honey of rose and	105
Stain	463	Lotions of	
Varnish		Soap powder	468
	12	Solution, comp	1:11
Blackberry balsam			
Cordial	180	Boraxine	
Elixir41,	180	Bordeaux mixture	
Compound	41	Boric acid ointment	117
Root bark, fl. ext., comp	87	and tannic acids, glycerite of	-104
Syrup of, aromatic87,	140	Borocitrate of magnesium, solut. of	-133
Blackboard slating	378	Boroglycerin cold cream	
Blackheads, remedies for		Borol	
		Development 1	$0.44$ $\lambda$
Blacking, Day & Martin's		Borolyptol	010
Harness		Borosol	
Leather	439	Borosalyl	
Shoe	379	Bor-salicylate	-212
Stove	464	Boschee's German syrup	
Black oil		Bottle capping mixture	
Blancard's pills		Bottles, to cement labels to	
Blatta insect powder			001
		Greasy, to clean	
Bleaching of feathers		Bougies14,	101
Oils, 379. Sponges	379	Alum	16
Blessed thistle (see Carduus bened.).		Cacao butter	
Blistering liquid	207	Chloral hydrate	16
Blisters, horse	246	Ferric chloride	17
on horses	252	Gelatin	15
Plood bitters	167	Gum14,	17
Cleanser	168	Iodoform	16
Dried, defibrinated	14	Potassium iodide	16
Purifiers	168	Silver nitrate	
Purifying drops	170	Tannin	
Powder, Schultze's	238	Bouquets	282
Tea136,	168	Bowel, inflammation in cattle	263
Kneipp's	108	Bradycrotine	
Swedish	135	Branding ink	
Bloodroot, tincture of, comp	151		
		1	

Brass, lacquer for	438	Burnett's Disinfecting liquid	392
Polish		Burns, applications for	171
Brassicon		Butter of antimony	
Brazil wood paper		Comp., gilt-edge	221
Breast tea	136	Stains, removing	462
Hamburg	227	Butyl chloral hydrate, elixir of	44
Breath perfume		Syrup of	140
Bretfeld water	304	Butyromel	344 A
Brick walls, efflorescence on	38	Byrolin	344A
Brilliant stars	$\frac{400}{234}$	CACAO hyster benefic	1.4
Brilliantine, 349. Pinaud's		CACAO butter bougies	$\frac{14}{162}$
British oil	116 110		316
Brodie's liniment	43	Milk Tincture of	151
Elixir of six	218	Cachous	361
Gold and arsenic, sol. of		Cactus grandiflora, fl. ext. of	88
Bromidia		Tincture of	151
Bromine inhalation, Netolitsky's	106	Or cactine pills	213
	111	Caffeine, citrated, effervescent	125
Solution of		Elixir of	44
Water	159	with potass. brom., efferv	127
Bromo-chloralum	241	Cajeput liniment, compound	110
Bromo-seltzer	212	Mixture, compound	112
Bromophtharin	212	Tincture of, compound	152
Bronchial troches, Brown's	213	Calamine, cerate of	18
Bronchiline	213	Ointment, Rademacher's	117
Bronze paints		Calamus, confection of	23
Powder		Extract of	89
Broom, decoction of	30	Sugared	23
Brown cerate	18	Tincture of	152
Mixture, improved	178	Calcium chloride sol., Rademacher's	131
Ointment	18	Compounds, elixirs containing44.	45
Brown-Sequard's epilepsy mixt; Pills	206	Hypophos. em. cod liver oil with	$\frac{82}{99}$
Brown's bronchial troches		Ext. of malt with	
Male fern vermifuge		Syrup ofLactophos. cod liver oil with	82
Bruises, remedy for		Phos., em. of cod liver oil with	83
Brunnersches fleck wasser	462	Syrup of	
Brust elixir	63	and sod. hypophosph., syrup of	140
Tropfen	63	and sod. phosph., em. cod liver oil with	83
Bryony, fl. ext. of	87	Calder's saponaceous dentifrice	
Buchu, elixirs containing	44	Calendula cerate	
Fl. exts. containing	88	Fl. ext. of, non-alcoholic	
Helmbold's	223	and glycerin lotion	313
Infusion of	105	Jelly	318
Tincture of	151	Toilet cream	314
Buchus	189		
Bucklen's arnica salve	213	Calisaya (see also Cinchona).	
Buckthorn bark, bitterless	88	Elixir of	45
Syrup of	140	Essencia de	219
Berries, syrup of	140	La rilla	213
Decoction of, comp	30	Tonic	371
Elixir of	44	Calolactose	$\frac{214}{259}$
Extract of	88	Calves, doses for	214
F1 exts. containing	88 44	Campho-phenique	214
and senna, elixir of	377	Camphoid	11
Bugine	377	Chloral	17
Bumsted's gleet cure.		Cold cream.	310
Bunion cures		Cream	17
Burdock bitters		Ice	320
Comp. sarsaparilla, etc		Julep, Thompsonian	17
Root hair oil		Liniment	110
Seed, tincture of	151	Liniment, comp	
Burnett's cocoaine			112

Camphor, monobromated, emuls. of	80	Cascara, aromatic	90
Phenol	11	Cordial	214
Pills, comp.	120	Sagrada, bitterless	90
Powder, comp		Elixir containing45,	$\frac{46}{89}$
Salicylate	17	Extract of	91
Tincture of, comp 109,		Fl. ext. containing90,	97
Toilet cream		Malt withSyrup of	141
Chloroform	20	Tincture of	152
Dover's powder		Cascarine, tasteless	220
Emulsion	80	Casein cement	
Naphthol		Ointment, Unna's	117
Oil		Cassie sachet	294
Spirit of ether	137	Castillon's elixir	46
Soap	138	Castorbena	206
'Turpentine liniment	111	Castoria	235
Can, to determine capacity		Castor grease	109
Canada turpentine, factitious		Oil, emulsion of	80
Canary bird food	277	Hair pomade	339
Medicines		Mixture	112
Varnish	467	Substitute	206
Canker of the ear in dogs	271	Tincture of	152
Cantharides, extract of, acetic	89	Ammoniated	152
Liniment	110	Water, Rademacher's	160
Vinegar of	159	Castor oil substitute	206
Cantharidin collodion	20	Castroilina	206
Capping mixture for bottles	380	Cat's fat	109
1	110	Medicines	$\frac{271}{171}$
	155 111	Catarrh balm	
Lotion, Griffith's		Cure, royal	
	159	Remedies.	
Carbolated catgut	17		
		Remedy, Sage's	
Collodion	20 26	Saul's	237
CollodionCotton	30	Saul's	$\frac{237}{171}$
CollodionCottonSalicylic collodion	$\frac{20}{26}$	Saul's. Snuff	$\frac{237}{171}$
CollodionCotton	$\frac{20}{26}$	Saul's	237 171 268 171 367
Collodion Cotton Salicylic collodion Silk	$\begin{array}{c} 20 \\ 26 \\ 21 \\ 128 \end{array}$	Saul's Snuff	237 171 268 171 367 92
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33 Camphorated.	20 26 21 128 29 11 11	Saul's Snuff. 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut	237 171 268 171 367 92 17
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of.	20 26 21 128 29 11 11 104	Saul's Snuff. 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut Carbolated.	237 171 268 171 367 92 17
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's.	20 26 21 128 29 11 11 104 246	Saul's Snuff 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut Carbolated. Chromated	237 171 268 171 367 92 17 17
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's.	20 26 21 128 29 11 11 104 246 120	Saul's Snuff. 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut Carbolated Chromated Juniper.	237 171 268 171 367 92 17 17 18 18
Collodion. Cotton Salicylic collodion Silk Tannin cotton. Carbolic acid, No. 33. Camphorated Glycerite of. Composition, Hager's. Paste, Lister's. Salve.	20 26 21 128 29 11 11 104 246 120 194	Saul's Snuff	237 171 268 171 367 92 17 17 18 18
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball.	20 26 21 128 29 11 11 104 246 120 194 214	Saul's Snuff. 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Carbolated. Chromated Juniper. Mercuric chloride Cathartics	237 171 268 171 367 92 17 18 18 18
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33 Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor.	20 26 21 128 29 11 11 104 246 120 194 214 11	Saul's Snuff 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut Carbolated Chromated Juniper. Mercuric chloride Cathartics Cathartic elixir, compound	237 171 268 171 367 92 17 18 18 173 46
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper.	20 26 21 128 29 11 11 104 246 120 194 214 11 445	Saul's Snuff	237 171 268 171 367 92 17 18 18 18 173 46 247
Collodion. Cotton Salicylic collodion Silk Tannin cotton. Carbolic acid, No. 33. Camphorated Glycerite of. Composition, Hager's Paste, Lister's. Salve. Smoke ball. Carbolized camphor Paper Solution of iodine	20 26 21 128 29 11 104 246 120 194 214 11 445 173	Saul's Snuff. 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut Carbolated Chromated Juniper. Mercuric chloride Cathartics Cathartic elixir, compound for horses Pills, little	237 171 268 171 367 92 17 18 18 173 46 247 175
Collodion. Cotton Salicylic collodion Silk Tannin cotton. Carbolic acid, No. 33. Camphorated Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball Carbolized camphor Paper Solution of iodine Sponge	20 26 21 128 29 11 104 246 120 194 214 11 445 173 138	Saul's Snuff. 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut Carbolated Chromated Juniper. Mercuric chloride Cathartics Cathartic elixir, compound for horses. Pills, little Tea	237 171 268 171 367 92 17 18 18 173 46 247 175
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water	20 26 21 128 29 11 104 246 120 194 214 11 445 173 138 159	Saul's Snuff	237 171 268 171 367 92 17 18 18 173 46 247 175 185
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp.	20 26 21 128 29 11 104 246 120 194 214 11 445 173 138 159 89	Saul's Snuff	237 171 268 171 367 92 17 18 18 173 46 247 175
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp.	20 26 21 128 29 11 104 246 120 194 214 11 445 173 138 159	Saul's Snuff	237 171 268 171 367 92 177 18 18 18 173 46 247 175 185 259
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp.	20 26 21 128 29 11 104 246 120 194 214 11 445 173 138 159 89 137	Saul's Snuff	237 171 268 171 367 92 177 18 18 18 173 46 247 175 259 259 18 18
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of.	20 26 21 128 29 11 104 246 120 194 214 11 445 173 138 89 137 89 152	Saul's Snuff. 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut Carbolated Chromated Juniper. Mercuric chloride Cathartics Cathartics Cathartic elixir, compound for horses. Pills, little. Tea Catholicon Cattle, doses for Medicines Caustic, arsenical, Ratier's	237 171 268 171 367 92 177 18 18 18 173 46 247 175 259 259 18 18 18
Collodion. Cotton Salicylic collodion Silk Tannin cotton. Carbolic acid, No. 33. Camphorated Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor Paper Solution of iodine Sponge Water Cardamon, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of Mariæ-tinct., Rademacher's	20 26 21 128 29 11 10 42 246 120 194 214 173 138 89 159 89 152 125	Saul's Snuff	237 171 268 171 367 92 17 18 18 18 173 46 247 175 185 259 259 18 18 18 18
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of. Mariæ-tinct., Rademacher's. Carlsbad salt, artificial, efferv. Carminative balsam	20 26 21 128 29 11 11 104 246 120 119 445 173 138 139 89 137 89 152 125 137 205	Saul's Snuff	237 1711 268 1711 367 92 177 188 188 173 466 247 175 185 259 259 18 18 18 18 18 18 18 18 18 18 18 18 18
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of. Mariæ-tinct., Rademacher's. Carlsbad salt, artificial, efferv. Carmelite spirit. Carminative balsam. Oil	20 26 21 128 29 11 11 104 246 120 194 214 11 445 173 138 89 137 89 152 125 205 114	Saul's Snuff. 128, in swine Catarrhine Catawba syrup Catechu, fl. ext., comp. Catgut Carbolated Chromated Juniper. Mercuric chloride Cathartics Cathartics Cathartic elixir, compound for horses. Pills, little Tea Catholicon Cattle, doses for Medicines Caustic, arsenical, Ratier's Black, Velpeau's Vienna Cedarwood, spirit of Celandine, tinct. of, Rademacher's Celerina.	237 1711 268 1711 367 92 177 18 18 18 18 173 46 247 175 185 259 259 18 18 18 18 11 175 25 18 18 18 18 18 18 18 18 18 18 18 18 18
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of. Mariæ-tinct., Rademacher's. Carlsbad salt, artificial, efferv. Carmelite spirit. Carminative balsam Oil	20 26 21 128 29 11 11 104 246 1120 194 214 445 1173 138 159 152 125 137 205 114 136	Saul's Snuff	237 1711 268 1711 367 92 177 18 18 18 173 46 247 175 259 259 18 18 18 284 15 24 15 24 17 25 25 25 25 25 25 26 26 26 27 27 27 27 27 27 27 27 27 27 27 27 27
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of. Mariæ-tinct., Rademacher's. Carlsbad salt, artificial, efferv. Carmelite spirit. Carminative balsam Oil Species. Tincture	20 26 21 128 29 11 11 104 246 1120 194 214 117 3138 159 137 205 137 205 137 215 136 136 136 137 215 137 215 137 215 136 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 215 137 137 137 137 137 137 137 137 137 137	Saul's Snuff	237 1711 268 171 367 92 17 17 18 18 18 247 175 185 259 284 184 152 214 214 22 23
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of. Mariæ-tinct., Rademacher's. Carlsbad salt, artificial, efferv. Carminative balsam Oil Species. Tincture. Water	20 26 21 128 29 11 11 104 246 120 194 41 11 445 137 89 137 89 152 137 205 114 136 136 137	Saul's Snuff	237 171 268 171 367 17 18 18 173 46 247 175 185 259 259 18 18 18 284 152 214 192 232 192
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of. Mariæ-tinct., Rademacher's. Carlsbad salt, artificial, efferv. Carminative balsam Oil Species. Tincture Water Carmine ink	20 26 21 128 29 11 104 1246 120 194 214 11 445 173 138 159 137 89 152 125 14 136 152 160 423	Saul's Snuff	237 1711 268 171 367 17 18 18 18 173 46 247 175 259 259 18 18 18 28 4 15 22 4 4 19 22 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of. Mariæ-tinct., Rademacher's. Carlsbad salt, artificial, efferv. Carmelite spirit. Carminative balsam Oil. Species. Tincture Water Carmine ink. Carriage-top dressing.	20 26 21 128 29 11 104 246 120 194 214 445 173 189 137 205 125 137 205 160 152 160 382	Saul's Snuff	237 171 268 171 367 92 17 18 18 18 173 46 46 247 175 185 259 28 18 18 28 44 192 21 41 192 23 24 46 46 46 47 47 48 48 48 48 48 48 48 48 48 48 48 48 48
Collodion. Cotton Salicylic collodion. Silk Tannin cotton. Carbolic acid, No. 33. Camphorated. Glycerite of. Composition, Hager's. Paste, Lister's. Salve. Smoke ball. Carbolized camphor. Paper. Solution of iodine. Sponge. Water Cardamom, fl. ext. of, comp. Spirit of, comp. Carduus benedictus, extract of. Mariæ-tinct., Rademacher's. Carlsbad salt, artificial, efferv. Carminative balsam Oil Species. Tincture Water Carmine ink	20 26 21 11 11 104 246 120 194 445 173 138 89 137 89 152 137 205 114 413 60 423 423 132 132	Saul's Snuff	237 171 268 171 367 92 177 18 18 18 173 46 247 175 185 259 18 18 24 152 232 192 232 192 46 192 233 192 238 238 248 248 248 248 248 248 248 248 248 24

Cements3		Chloroform, Belladonna	20
Cement, dental		Camphorated	20
Hoof		Elixir of	47
Marine		Compound	47
for meerschaum		Glycerite of	105
Centaur liniment		Henbane	20
Cephalic snuff		Liniment, comp	110
Cerasorcin, elixir		Mitigated	
Cerate, brown		Oil	
Calamine		Paregoric	47
Calendula		Tincture of, comp.	152
Copaiba		Chlorol, chlorolin	-215
Green		Chloryl	215
Nutmeg		Chocolate candy varnish	
Rose		Stains, removing	
Soap		Syrup	367
Turner's		Chologogue	164
Zinc carbonate		Cholera, chickens	276
Ceratum æruginis	19	Hog	
Ceratum, (see Cerates).	115	Mixtures	112
Ceresin paper Cereus grandiflorus (see Cactus gr.)	445	Pills	120
Chambarlain's maliaf	04.4	Remedies	176
Chamberlain's relief	214	Christadoro's hair dye	215
Chamomile, infused oil of		Christensen's diarrhœa mixtures	112
Water		Chroatol	
Chapman's alkaline powder		Chromated catgut	18
Internal disinfectant	215	Chromograph	414
Charcoal poultice, Thompsonian		Chrysarobin collodion	20
Powder, comp		Churchill's caustic sol. of iodine	132
Tooth powder		Churchill's syrup.	$\frac{215}{166}$
Chelidonium, (see Celandine).	400	Cigarettes, asthma	402
	464	Cigars, flavor for	41
Chemical barometer		Cimicifuga, elixir of	42
Food for plants		Cinchona, beef, wine and iron with	161
Cherries, elixirs containing	46	and coce wine of	162
Syrup of		and coca, wine of	
Cherry stain		Fl. ext. comp.	91
Syrup		Hair tonic.	344
Water		Syrup of	141
Chevalier's life for the hair	$\frac{100}{215}$	Wine of	161
Chicken cholera		Compound	162
Chicken lice from coops, to remove		Cinchonidine, elixirs containing	50
Medicines		Cincho-quinine	215
Chilblain cures		Cinnamon, spirit of	284
Children's cough cure		Tincture of, comp	152
Child's cough mixture		Citric acid, saccharated	134
Chill cure or tonic		Solution of	366
Chinoidin, tincture of		Civet	280
Chirata, elixir of		Tincture of	285
Chloral, camphor	17	Clark's blood mixture	215
and ammon. chloride, elixir of	47	Classification of odors	280
Hydrate bougies	16	Cleaning of gloves	411
Elixir of	46	Cleansing creams	387
Syrup of		Greasy bottles	381
Chloralum21	5, 241	Liquids	461
Chlorides, elixirs containing	47	Marble	439
Chlorine fumigation	393	Busts	440
Water	160	Pencils	461
Chlor-lactated elixir of pepsin		Clove pink extract	285
Pepsin powder		Sachet	294
Chlorobrom		Cloves, spirit of	
Chlorodyne		Coaline headache powders	215
Cough cure		Solution of	131
Chloroform of aconite	20	Coal tar, prepared	131
		Cobb's pills.	215

Coca, beef, wine and iron with	161	Cologne waters	302
and cinchona, wine of	162	Colored fires	399
Elixirs containing50,	51	Coloring of horn and ivory black	
Emuls. of cod liver oil with	82	for syrups	363
Wine of	$\frac{163}{26}$	Colors for Easter eggs	397
Cocaine cotton	31	Colts doses for	$-400 \\ 245$
Discs of		Colts, doses for	
Morphine cotton	27	Infusion of	
Cocculus indicus, tinct., homeo		Wine of	162
Cochineal ink.		Comedones, remedies for	326
		Come's arsenical powder	121
Syrup of	152	Complexion powder	323
Cocoa milk		Tablets	324
Cocoaine	213	Composition powder	122
Cocoanut oil hair dressing	338	Compound, lithia tablets	216
Hair pomades		Concentrations	21
Codliver oil, aromatized		Condition powders	248
Combinations with malt ext	97	Darby's	249
Emulsions containing80-	83	Condy's fluid	215
with iodine		Confection, aromatic	23
with iron		Calamus	23
with iron and iodine	115	Copaiva	23
with malt extract		Figs	$\begin{array}{c} 23 \\ 23 \end{array}$
Wine of		Hollyhock	$\frac{23}{23}$
Coddington's asthma powder	215	Opium	$\frac{23}{24}$
Codeine, elixir of	51	PepperPrunes	24
Syrup of	51	Rhubarb, compound	$\frac{\sim}{24}$
Coe's dyspepsia cure		Sulphur	$\frac{\tilde{24}}{24}$
Coffee stains, removing		Tamarind	24
Syrup		Congo red paper	465
Colcothar		Congress water	371
Colchicum seed, extract of	92	Conium, infused oil of	115
Tincture of, comp	153	Ointment	117
Cold, remedies for	177	Tincture of	153
Cold creams309	-312	Conklin's salve	216
Perfumes for	312	Conserves	24
Cold in the head, remedies for	171	Constipation in birds	278
Colic in cattle		in cattle	
Oil	114	in dogs and cats	
Remedies, for horses		in horses	249
in sheep		in sheep	
in swine		Convallaria, tincture of	$\frac{153}{120}$
Colladion possits	$\frac{215}{20}$	Cook's pills	19
Collodion, aconite	20	Confection of	23
Canthardin	20	Copal varnish	
Carbolated	20	Copeland's cosmetic water	
Salicylic	$\tilde{21}$	Lotion of borax	111
Chrysarobin	20	Copper acetate tinct., Rademacher's	153
Diachylon	21	Aluminated	24
Iodized	21	Ammoniated	24
Iodoform	21	Bluing of	380
Iodol	21	Etching	398
Iron	21	Mixture, Rademacher's	113
with lead plaster	21	Subacetate, ointiment	117
Mercuric chloride	21	Copygram	414
Photographer's	21	Copygraph	414
Salol	21	Copying pad	$\frac{414}{422}$
Thiol	21	Coral ink	
Thymol	21	Corassa compound	35
Collograph	159	Blackberry	
Tincture of	153	Cascara	214
	100		

Cordial, Celery	192	Counterfeit coin detector	388
Cough	-177	Cow powder	
Diarrhœa	180	Crab apple blossom	
Drops, Warner's		Cracked heels, ointment for	
Mother's		Crampbark, elixir of, comp	51
Neutralizing	147 147	Cram's fluid lightning	216
Restorative		Cream, almond310,	$\frac{314}{171}$
Corn eradicators		Balm	$\frac{111}{219}$
Corydalis, syrup of, comp		Ely's Calendula	314
Tincture of		Camphor	214
Coryza, remedies for	171	Cleansing	
Cosmetic cream	313	Cold	
Jelly	318	Cucumber	312
Liniment, Hebra's	329	Face313,	
Liquid		Furniture	
Lotion	317	Glycerin	314
Goddard's	330	Compound	321
Manicure	334	Honey and almond	314
Vinegar		Lanolin	310
Water		Lilacs	313
Copeland's	330	Liniment	191
Lilionese	329	Oriental	
Cosmetique	349	Pearl	326
Cosmos bouquet		Roses313,	
Cotton, absorbent		Shampoo	336
Aluminum acetate	25	Shaving	333
Antirheumatic	25	Syrup	368
ArnicatedBenzoated		Toilet	$\frac{314}{352}$
Borated		Witch hazel	312
Carbolated		Creek Indian remedy	86
Cocaine	$\frac{\tilde{2}6}{26}$	Creme Simon	216
Borated	- 0	Creolin	
Morphine		Creosote, glycerite of	104
Dyeing		and hypophos. em. cod liver oil with	82
Ferrated	29	Ointment	117
Hemostatic	29	Syrup of	
Ichthyol	27	Wine of	
Iodized	27	Compound	162
Iodoform	27	Creosol water	
Iodol	27	Crew's disinfectant	
Lister's sero-sublimate		Crocus martis	438
Medicated		Tincture of	157
Mercuric cn_oride		Croton bug exterminnators	388
Naphthalin		Chloral hydrate, elixirs containing	51
Removing stains from		Syrup of	$\frac{141}{110}$
ResorcinSalicylated		Oil, liniment	
Sol. alembroth		Crystal hair pomade	340
Styptic		Water	160
Tannin carbolated		Cubeb, syrup of	141
Thymolated		Cucumber cream	312
Zinc chloride		and elder flower cream	
Couch grass, extract of	92	Milk	316
Fl. ext. of		Ointment	312
Coudray's eau de quinine		Culture fluids	388
Cough in dogs and cats		Culver's root, tincture of	153
Elixir	142	Curacao cordial or elixir	51
in horses	250	Curare, injection of	107
Powder, Thompsonian	, 122	Curcuma paper	
Remedies	177	Curlique	
Tea, Kneipp's		Currant jelly or paste	
Whooping, remedies		Cuticura ointment	216
Coumarin sugar	128	Resolvent	216
31			

Cuticura Soap	Dickson's improved anæsthetic 217
Cutting of glass	Digestive elixir, compound 58
Wrapping paper	Digitalis, syrup of
Cydonium, (see Quince seed).	Tincture of, ethereal
Cypripedium (see Ladies' slipper).	Dill water
DAMIANA compound 109	Dilution of alcohol
DAMIANA compound	Diovirburnia 217 Discs, ophthalmic 31
	Discs, ophthalmic
	Disinfectants
Wine of	Disinfecting solid
Dandelion, decoction of	Distemper remedies
Elixirs containing	Diuretic elixir
Dandruff remedies	Wayne's
Darby's condition powder	for horses
Prophylactic fluid	Liniment 110
Darlings lotion of sodium chloride 111	Species
Darwin's liniment	Tea 136
Davis' cough mixture	Wine, Hufeland's 162
Pain killer 216	Divine stone 24
Davy's urinal cakes £16	Dobell's solution
Dean's king cactus oil244B	Dog medicine
Declat's syrup of phenic acid 216	Dogs, doses for
Decoction of barley	Dog's fat or grease 109
Broom	Donovan's solution
Buckthorn, comp	Dorsenia
Dandelion, granatum 30	Doses for calves and cattle
Guaiac, comp	for dogs
Pareira 30	for horses
Sarsaparilla, comp	Rules for 374 Douglass' mixture 278
Deer suet	
Dental anæsthetics	Dover's powder, camphorated
Cement	Dressing for carriage-tops
Surprise	Leather
Dentifrices	Shoe
Dent ine 351	Driers
Dentists' amalgam fillings 388	Drops, bitter stomach
Arsenic paste	Blood purifying
Modeling wax	Cordial, Warner's 31
Nerve destroyer	Cramp 31
Deodoried alcohol	Cramp
Benzin	Earache
Deodorizers	Gold
Depilatories	Hunn's
Deshler's salve	Jesuit 32
Dewberry root, elixirs of, comp 52	Mother 32
Dextro-quinine	Nerve
Dextro-saccharin. 217 Diachylon collodion. 21	Rheumatic
Diachylon collodion	St. Peter's
Diamond cement	Dropsy tea, Kneipp's
Dust	Duffy's malt whisky
Ink	Dulcarmara (see Bittersweet).
Diaphoretic powder, Beach's 122	Dupuytren's pomade
Dialyzed iron	Dusart's syrup
Diarrhœa in birds	Dutch ague remedy
in calves	Dutchman's laudanum 109
in cattle	Dyche's capsules of guaiacol 217
in dogs and cats	Potass. iod., pills comp
in horses	Dyeing leather
in sheep	Dyes, Easter egg 397
in swine	Fabrics
Mixtures	Hair347–349
Remedies	Dysentery remedies

Dysmenorrhœa drops, Rademacher's	32	Elixir, Euonymus	78
2.1.1	181	Flavoring, No. 29, gari	53
Tablets	182	Health	$\frac{157}{241}$
FAR medicines and oils	182	Iodo-brom. calcium comp	$\frac{241}{63}$
de Raspail	32	Kola Lactated or lactinated pepsin	67
Earache drops	1	Life	64
Oils.	182	ad longam vitam	64
	182	Mitchella, compound	56
	397	Neutralizing	73
	408	Nutrans	218
	358	Opium, McMunn's	-230
	304	Paracelsi	70
	330	Pectoral63,	70
	829   850	Three phosphates	61
	$\frac{358}{916}$	Pilocarpus	63
	$rac{216}{234}$	Proprietatis	70
	463	Pulmonic	$\frac{70}{218}$
Eczema remedies		Purgans	358
Edison's polyform		Salicylic acid	218
Edward's alterative tonic bitters	217	Salutis	
	122	Sedative	42
Salts	128	Six aperiens; six hypophos six brom.	218
Effervescing powder	122	Six iodides	219
	122	Six iodides	56
Egg cream syrup		Stomachic	66
Dyes	397	Turnera	52
	275	Vegetable	142
Making food		Viscerale Hoffmanni	66
Shampoo.		Eel oil, factitious	$\frac{115}{210}$
	$\frac{115}{397}$	Ely's cream balm	$\frac{219}{397}$
Preservation of		Embalming fluids Embrocation	
Salve	$\frac{218}{218}$	Emetic tincture	
Elæosacchara		Emmenagogues	185
Elder bark, wine of	162	Emmenagogue elixir	53
	286	Powder	122
	107	Emplastra, (see Plasters).	
	107	Emplastrum fuscum molle	18
Elecampane, fl. ext. of	92	Emulsio amygdalæ	79
	387	Oleosa	80 84
Headache cures		Salicylica	
Liniment		Emulsion cod liver oil with malt ext	$-\frac{00}{97}$
	451	Kerosene	436
Electuaries	32	Liquid petrolatum, comp	84
Electuary of theriac	23	Oil	80
	218	Phosphatic	83
Elixirs	79	Tolu balsam	85
Elixir, alterative51,	57	Enamel ink	426
1	193	Skin	326
	$\frac{164}{207}$	Eno's fruit salt	219
	251	Entozoic powder	$\frac{123}{422}$
Bird	$\frac{218}{180}$	Eosin ink	273
Blackberry. Brust.	63	Erasives, ink	
Cerasorum	46	Ergot, extract of.	92
Cinchon alkaloids	70	Fluid extract of	93
Coca	50	Purified	93
Corrigens	78	Tincture of	153
Cough		Ergotin	92
Dentifrice		Injection of	107
Dinretic, Wayne's		Syrnp of	
Eriodictyon	78	Eriodictyon, elixirs containing	78

480 · INDEX.

Erysipelas remedy	206	Extracts	3-104
Essencia de calisaya		Eye diseases, Remedies for	. 183
Espey's fragrant cream		in sheep	
Esprits. Essences.		in swine	
Essence, birch.		Salves	
Bouquet extract		FACE bleach	
Sachet	001	Cream	
Fumigating	302	Lotion	317
Ginger	263	Paint, liquid	. 326
Mead	363	Powder	
Mosquito	440	Fahnestock's vermifuge	
Moth		Farina cologne	
Mouth	358	Fashion bouquet	. 286
Pennyroyal	441	Fats104,	
Pepsin	$\frac{85}{135}$	Fayard, papier	
Rennet Saccharin	85	Feathers, bleaching of	401
Sarsaparilla	363	FebritugeFebriline	010
Tamarinds	85	Feet, fetor of, remedies for	
Esterhazy bouquet	286	Perspiration of, remedies for	-01
Etching copper	398	Fellows' syrup	
Glass398,	426	Felon oil, Kneipp's	
Iron	398	Female disorders, remedies for	. 185
Steel		Pills, Harper's	
<u>Tin</u>	398	Remedy or tonic	
Zinc	398	Ferrated cotton	
Ether, lotion of, comp	111	Ferro-iodized cod liver oil	000
Spirit of, camphorated	137	Fertilizer	001
Syrup of	142 86	Fever in cattlein horses	
Eucalyptus, elixir of	53	Figs, confection of	00
Syrup of	142	Medicated	• • •
Tincture of	153	Filler, hardwood	
Euonymin	104	Fillings, dentist's amalgam	. 388
Euonymus, elixirs containing	78	Filter paper, toughened	. 399
(see also Wahoo).	240	Finger-nail polish	
Eulyptol	219	Fire extinguishers	$\frac{402}{0.400}$
Euphorbium, oil of, comp	114	Fires, colored	403
Eureka dental anæsthetic Euthymol	$\frac{219}{219}$	Fireproofing material	000
Evaporating lotion	111	Flagg's relief	000
Evening primrose	286	Flander's diffusible tonic	0.00
Exodyne	~	Flash-light powders	
Expectorant	177	Flavoring, elixir, No. 29	. 53
Tincture, King's	154	Flavor for cigars and tobacco	402
Extract of annatto	130	Flaxseed cough syrup or balsam	
Dulcamara	87	Flea exterminators	
Pinus Canadensis	94	Fleas on dogs and cats	
Root beer		Fleury's tasteless cascarine	
Vanilla	364	Floor polish	
Extracts		Florida water	
Fluid86	-104	Flowers of Ireland extract	. 287
Handkerchiefs	281	Fluid extract of Oregon grape	. 87
Liquid	25	Fluid hydrastis	. 220
Perfume285	-293	Fluid Lightning	
Powdered		Cram's	
Saccharated	101	Fly exterminators	. 405
Extractum graminis	92	Lime	
Extraits	00	PapersPastilles	
Eye-bright, Kneipp's		Pencils	
Eye diseases in cattle	261	Powders	. 405
in dogs and cats, 273; in horses	251	Foam, soda or gum	. 366
		Foods for birds	. 277

Foot spavin in horses	258	Gelatinized benzin	378
Ford's balsam of hoarhound	220	Gelatol	220
Formic acid, spirit of	137	Gentian, elixirs containing54,	
Spirit of, comp	137	Fl. ext. comp.	
Formacoll	290	Ext. malt and iron with	
Fossilin	220	Infusion of, comp	
Fothergill's asthma mixture		German bird paste	
	179	Bitters.	
	256	Cologne	
Fountains, charging of	406	Herb tea	
Four chlorides	47	Gibert's syrup	146
Fox lungs, syrup of	142	Gicht watte	
	220	Gilding	
	286	Powder	
	287	Gile's iodide of ammonia liniment	
Sachet		Gillett's washing crystal	
Fragula, elixir of	44	Gilt-edge butter comp	
	327	Essence of	363
	220	Syrup	2 0 0
	406	Tonic.	
	220	Glacialin	221
Frey's vermituge	220	Glanders246,	253
Friar's balsam	12	Glass, cement for	
	185	Cutting of	
	366	Etching of398,	
	365	Labels to bottles, to cement	
	220	to metals, to cement	
	$\begin{array}{c c} 220 \\ 219 \end{array}$	Polish for	
Eno's	163	PowderedGleet remedies	
	142	Gliricin	
	104	Glove cleaner	
	325	Glove detergent	
	302	Glover, dry process for cleaning	
Paper 301; pastilles	300	Glover's bromine lotion	111
	301	Glucose	412
	301	Glue, bookbinder's.	411
	303	Elastic 412; liquid 411; Marine	
	302   393	Mouth	
	$\frac{355}{416}$	Pocket, stick	
	$\frac{110}{407}$	Glutiform	244R
	408		
	467	Glutol	321
	205	Cold cream	
	408	Cream, comp	
		Jelly	
	185	Lotion	
Galanga, extract of, fl	93	Toilet cream	
	252	Glycerites	
	105	Glycerite of hydrastine	$\frac{132}{221}$
	$\begin{bmatrix} 117 \\ 53 \end{bmatrix}$	Glycerole yerbine compound	
Galls, elixir of, aromatic	278	Glyco-thymoline	
Garfield tea		Glycyrrhiza powder, aromatic	
Gargle species		Syrup of	142
Gargling oil	221	Glymol	
	142	Goddard's cosmetic lotion	330
Garus, elixir de	53	Elixir ammon. valerianate	33
	384	Godfrey's cordial	
Gelatin bougies14,	15	Gold and arsenic bromides, sol. of	
Mass for bougies	15	Dust washing powder	
Nutrient	444	Tropfen	31

Gold Ink	422	Gun Powder	122
Plating	447	White	
Golden oil	191	Gurania	222
Rain	400	Gutta percha cement	
Rod, tinct., Rademacher's		Purified	
	106	Solution of	
Elixirs containing	55	Guttæ ad menstrua tormentosa	
Extract of	94	Jesuitarum	32
Fl. ext., aqueous	93	TIAADI DAG H	000
Fl. ext., colorless	94	HAARLEM oil	
Powder, comp	122	Haas' hog cholera remedy	
Tincture of, comp	$\frac{154}{162}$	Hagan's magnolia balm	
Wine of. comp		Haines' golden specific	
Gonorrhœa remedies.		Hair balsams344,	
Goose grease or oil		Hair-curling liquid	
Gouraud's oriental cream		Hair dyes,	
Gout pills, Lartique's		Christadoro's.	
Remedies	196	Walnut	
Gowland's lotion		Growers	
Grafting wax		Loss of, remedies for	
Graham's cucumber and elder fl'r cream.		Oils	
Grains de sante		Perfumes for	
Granatum, decoction of	30	Oil, walnut	
Extract of	94	Pomades, perfumes for	338
Granular effervescent salts	125	Preparations for	334
Granules, liver, little		Promoters344,	, 345
Granville's counter-irritant		Restorers344	345
Grape saline	221	Tonics	1-346
Salt175,	221	Containing oil	. 346
Grass stains, removing	463	Washes344	345
Gray's glycerina tonic comp	222	Hair's asthma remedy	. 223
Gray's specific pills	222	Hall's catarrh remedy	244B
Grease eradicators	413	Hair renewer	. 223
Grease eradicators	413 331	Hair renewerHamburg breast tea	. 223 ,244B
Grease eradicators	413 331 456	Hair renewer. Hamburg breast tea. Drops	. 223 ,244B . 227
Grease eradicators. Paints. Shoe. Stains removing	$   \begin{array}{r}     413 \\     331 \\     456 \\     462   \end{array} $	Hair renewer. Hamburg breast tea. Drops Tea, Freese's	. 223 ,244B . 227 . 220
Grease eradicators.         Paints.         Shoe.         Stains, removing.         Greases.       105, 108,	413 331 456 462 109	Hair renewer.  Hamburg breast tea.  Drops.  Tea, Freese's.  Hamlin's diarrhœa mixture.	. 223 ,244B . 227 . 220 . 112
Grease eradicators.           Paints.           Shoe.           Stains, removing.           Greases.         105, 108,           Grecian varnish.	413 331 456 462 109 467	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil	. 223 ,244B . 227 . 220 . 112 . 223
Grease eradicators. Paints. Shoe. Stains, removing. Greases. Grecian varnish. Green cerate.	413 331 456 462 109 467 19	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil Handkerchief extracts	. 223 .244B . 227 . 220 . 112 . 223
Grease eradicators.         Paints.         Shoe.         Stains, removing.         Greases.       105, 108,         Grecian varnish.         Green cerate.       11, 12, 12, 12, 12, 12, 12, 12, 12, 12,	413 331 456 462 109 467 19 423	Hair renewer. Hamburg breast tea. Drops. Tea, Freese's Hamlin's diarrheea mixture. Wizard oil Handkerchief extracts Hanson's corn salve.	. 223 ,244B . 227 . 220 . 112 . 223 . 281
Grease eradicators.         Paints.         Shoe.         Stains, removing.         Greases.       105, 108,         Grecian varnish.         Green cerate.         Ink       422,         Plaster	413 331 456 462 109 467 19 423 121	Hair renewer.  Hamburg breast tea.  Drops. Tea, Freese's.  Hamlin's diarrhœa mixture.  Wizard oil.  Handkerchief extracts.  Hanson's corn salve.  Hardwood filler.	. 223 ,244B . 227 . 220 . 112 . 223 . 281 . 223
Grease eradicators.         Paints.         Shoe.         Stains, removing.         Greases.       105, 108,         Grecian varnish.         Green cerate.         Ink.       422,         Plaster.         Salve.	413 331 456 462 109 467 19 423 121 116	Hair renewer.  Hamburg breast tea.  Drops. Tea, Freese's.  Hamlin's diarrhœa mixture.  Wizard oil.  Handkerchief extracts.  Hanson's corn salve.  Hardwood filler.  Harlan's tooth paste.	. 223 .244B . 227 . 220 . 112 . 223 . 281 . 223 . 414
Grease eradicators.         Paints.         Shoe.         Stains, removing.         Greases.       105, 108,         Grecian varnish.         Green cerate.         Ink.       422,         Plaster.         Salve.         Soap.	413 331 456 462 109 467 19 423 121 116 461	Hair renewer.  Hamburg breast tea.  Drops Tea, Freese's  Hamlin's diarrhea mixture.  Wizard oil  Handkerchief extracts  Hanson's corn salve.  Hardwood filler.  Harlan's tooth paste.  Harness blacking.	. 223 .244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 357
Grease eradicators.         Paints.         Shoe.         Stains, removing.         Greases.       105, 108,         Grecian varnish.         Green cerate.         Ink.       422,         Plaster.         Salve.         Soap.         Mitigated	413 331 456 462 109 467 19 423 121 116 461 433	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhea mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking. Polish or oil.	. 223 ,244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 357 . 414
Grease eradicators.           Paints.           Shoe.           Stains, removing.           Greases.         105, 108,           Grecian varnish.           Green cerate.           Ink.         423,           Plaster.           Salve.           Soap.           Mitigated.           Green's August flower; nervura.	413 331 456 462 109 467 19 423 121 116 461 433 222	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking Polish or oil. Harney's diarrhœa mixture.	. 223 ,244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 357 . 414 . 113
Grease eradicators. Paints. Shoe. Stains, removing. Greases. 105, 108, Grecian varnish Green cerate Ink 423, Plaster Salve. Soap Mitigated Green's August flower; nervura. Greenhow's diarrhœa mixture.	413 331 456 462 109 467 19 423 121 116 461 433 222 112	Hair renewer.  Hamburg breast tea.  Drops. Tea, Freese's.  Hamlin's diarrhœa mixture. Wizard oil.  Handkerchief extracts.  Hanson's corn salve.  Hardwood filler.  Harlan's tooth paste.  Harness blacking.  Polish or oil.  Harney's diarrhœa mixture.  Harter's wild cherry bitters.	. 223 ,244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 357 . 414 . 113
Grease eradicators. Paints. Shoe. Stains, removing. Greases	413 331 456 462 109 467 19 423 121 116 461 493 222 112 123	Hair renewer.  Hamburg breast tea.  Drops. Tea, Freese's.  Hamlin's diarrhea mixture. Wizard oil.  Handkerchief extracts.  Harlan's corn salve.  Hardwood filler.  Harlan's tooth paste.  Harness blacking. Polish or oil.  Harney's diarrhea mixture.  Hatter's wild cherry bitters.  Hayden's viburnum comp.	. 223 . 244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 357 . 414 . 113 . 223 . 223
Grease eradicators. Paints. Shoe. Stains, removing. Greases	413 331 456 462 109 467 19 423 121 116 461 433 222 112 123 111	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhea mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking. Polish or oil. Harney's diarrhea mixture. Harter's wild cherry bitters. Hayden's viburnum comp. Hay fever remedies.	. 223 .244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 357 . 414 . 113 . 223 . 223
Grease eradicators. Paints. Shoe. Stains, removing. Greases	413 331 456 462 109 467 19 423 121 116 461 433 222 112 123 111	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking Polish or oil. Harney's diarrhœa mixture. Harter's wild cherry bitters. Hayden's viburnum comp. Hay fever remedies. Headache cures, electric.	. 223 .244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 357 . 414 . 113 . 223 . 218 . 218
Grease eradicators. Paints. Shoe Stains, removing. Greases	413 331 456 462 109 467 19 423 121 116 461 433 222 112 111 222	Hair renewer.  Hamburg breast tea.  Drops. Tea, Freese's.  Hamlin's diarrhœa mixture. Wizard oil.  Handkerchief extracts. Hanson's corn salve.  Hardwood filler.  Harlan's tooth paste. Harness blacking. Polish or oil.  Harney's diarrhœa mixture.  Harter's wild cherry bitters.  Hayden's viburnum comp.  Hay fever remedies.  Headache cures, electric.  Remedies.	. 223 .244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 113 . 223 . 223 . 188 . 218 . 228
Grease eradicators. Paints. Shoe Stains, removing. Greases	413 331 456 462 109 467 19 423 121 463 222 112 111 222 55	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking. Polish or oil. Harney's diarrhœa mixture. Harter's wild cherry bitters Hayden's viburnum comp. Hay fever remedies. Headache cures, electric. Remedies Headine Heaves.	. 223 .244B . 227 . 220 . 112 . 223 . 281 . 223 . 414 . 357 . 414 . 113 . 223 . 188 . 218 . 218 . 223
Grease eradicators. Paints. Shoe Stains, removing. Greases	413 331 456 462 109 467 19 423 121 433 222 123 111 222 55 206	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking. Polish or oil. Harney's diarrhœa mixture. Harter's wild cherry bitters Hayden's viburnum comp. Hay fever remedies. Headache cures, electric. Remedies Headine Heaves.	223 2448 227 220 112 223 281 281 281 414 414 414 414 414 414 414 414 414 4
Grease eradicators. Paints. Shoe. Stains, removing. Greases. 105, 108, Grecian varnish Green cerate Ink. 422, Plaster Salve. Soap. Mitigated Green's August flower; nervura. Greenhow's diarrhœa mixture. Gregory's laxative powder. Griffith's capsicum lotion. Grimault's matico injection Grindelia, elixir of. Gross' neuralgic pills. Guaiac, balsam of.	413 331 456 462 109 467 19 423 121 116 461 433 222 112 123 121 225 206 13	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking. Polish or oil. Harnesy's diarrhœa mixture. Harter's wild cherry bitters. Hayden's viburnum comp. Hay fever remedies. Headache cures, electric. Remedies Headine Heaves Hebra's arsenic pills. Cosmetic liniment.	223 244B 2247 2200 112 223 223 221 223 414 357 414 414 414 414 414 414 223 223 223 223 223 223 223 223 223 22
Grease eradicators. Paints. Shoe Stains, removing. Greases. 105, 108, Grecian varnish. Green cerate. Ink. 422, Plaster Salve. Soap. Mitigated Green's August flower; nervura. Greenhow's diarrhœa mixture. Gregory's laxative powder. Grithth's capsicum lotion. Grimauit's matico injection Grimdelia, elixir of. Gross' neuralgic pills Guaiac, balsam of. Decoction of comp. Elixir of. Glycerite of.	413 331 456 462 109 467 19 428 121 116 461 423 112 555 206 13 30 55 105	Hair renewer.  Hamburg breast tea.  Drops Tea, Freese's  Hamlin's diarrhœa mixture.  Wizard oil  Handkerchief extracts  Hanson's corn salve.  Hardwood filler.  Harlan's tooth paste.  Harness blacking.  Polish or oil.  Hanney's diarrhœa mixture.  Harter's wild cherry bitters.  Hayden's viburnum comp.  Hay fever remedies.  Headache cures, electric.  Remedies.  Headine  Heaves.  Hebra's arsenic pills.  Cosmetic liniment.  Drachylon ointment.	223 244B 227 210 210 221 221 221 221 221 221
Grease eradicators. Paints. Shoe. Stains, removing. Greases	413 331 456 462 109 467 19 423 121 123 55 206 13 30 105 142	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking. Polish or oil Hanney's diarrhœa mixture. Harter's wild cherry bitters. Hayden's viburnum comp. Hay fever remedies. Headache cures, electric. Remedies. Headine Heaves. Hebra's arsenic pills. Cosmetic liniment. Drachylon ointment. Itch ointment.	223 244B 227 220 220 221 223 281 223 414 414 414 413 357 414 414 414 113 223 281 188 238 188 238 188 238 188 188 188 188 188 188 188 1
Grease eradicators. Paints. Shoe. Stains, removing. Greases	413 331 456 462 109 467 19 423 121 123 123 55 206 13 30 55 142 154	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhea mixture Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking. Polish or oil. Harney's diarrhea mixture. Harter's wild cherry bitters. Hayden's viburnum comp. Hay fever remedies. Headache cures, electric. Remedies. Headine Heaves. Hebra's arsenic pills. Cosmetic liniment Drachylon ointment Itch ointment. Lead ointment.	223 2248 227 220 112 223 223 223 223 223 414 414 414 414 414 414 414 223 223 223 414 415 223 223 416 416 417 417 418 418 418 418 418 418 418 418 418 418
Grease eradicators. Paints. Shoe. Stains, removing. Greases	413 331 4456 462 109 467 19 423 111 461 443 123 112 123 111 223 506 13 30 55 105 142 1142 124 125 125 126 137 147 148 148 148 148 148 148 148 148	Hair renewer. Hamburg breast tea Drops Tea, Freese's Hamlin's diarrhœa mixture Wizard oil Handkerchief extracts Hanson's corn salve Hardwood filler Harlan's tooth paste Harness blacking Polish or oil Harney's diarrhœa mixture Harter's wild cherry bitters Hayden's viburnum comp Hay fever remedies Headine Headine Heaves Hebra's arsenic pills Cosmetic liniment Drachylon ointment Itch ointment Lead ointment Oriental cosmetic water	223 2244B 227 220 112 223 221 223 223 414 414 414 414 414 414 414 414 414 41
Grease eradicators. Paints. Shoe Stains, removing. Greases	413 331 4464 4662 109 467 191 116 461 1123 555 206 13 30 55 105 142 154 433 444 111 111 111 111 111 11	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhea mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking Polish or oil. Harney's diarrhea mixture. Harter's wild cherry bitters Hayden's viburnum comp. Hay fever remedies Headache cures, electric. Remedies Headine Heaves. Hebra's arsenic pills. Cosmetic liniment Drachylon ointment Itch ointment. Lead ointment Oriental cosmetic water. Sulphur paste.	2244B 2274B 2277 2200 1122 223 223 223 223 223 223 223 223 22
Grease eradicators. Paints. Shoe Stains, removing. Greases	413 831 456 4462 109 467 19 116 461 116 483 22 112 123 111 255 206 13 300 154 413 56	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking Polish or oil. Hanney's diarrhœa mixture. Harter's wild cherry bitters Hayden's viburnum comp. Hay fever remedies Headache cures, electric. Remedies Headine Heaves. Hebra's arsenic pills. Cosmetic liniment Drachylon ointment Itch ointment. Lead ointment Oriental cosmetic water. Sulphur paste. Hectograph inks. 427	2244B 2274B 2277 2200 1122 223 223 223 223 223 223 223 223 22
Grease eradicators. Paints. Shoe. Stains, removing. Greases. 105, 108, Grecian varnish Green cerate Ink. Plaster Salve. Soap Mitigated Green's August flower; nervura. Greenhow's diarrhœa mixture. Gregory's laxative powder. Griffith's capsicum lotion. Grimault's matico injection. Grindelia, elixir of. Gross' neuralgic pills. Guaiac, balsam of. Decoction of comp. Elixir of. Glycerite of. Syrup of. Tincture of, aromatic. Guaiacol capsules, Dyche's. Guano, artificial. Guarana, elixirs containing. 55, Gum bougies. 14,	413 331 4466 462 109 467 19 423 121 1166 461 4433 222 555 206 55 142 154 217 413 566 17	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking Polish or oil. Harney's diarrhœa mixture. Harter's wild cherry bitters Hayden's viburnum comp. Hay fever remedies. Headache cures, electric. Remedies. Headaine Heaves. Hebra's arsenic pills. Cosmetic liniment Drachylon ointment Itch ointment Lead ointment Oriental cosmetic water Sulphur paste. Hectograph inks. 427 Mass.	2243 2244B 227 227 220 112 223 231 223 414 357 414 113 223 238 414 113 1188 218 218 218 218 218 218 21
Grease eradicators. Paints. Shoe. Stains, removing. Greases	413 331 456 469 467 199 467 191 1166 461 122 555 206 13 30 555 142 217 413 566 17 366	Hair renewer. Hamburg breast tea Drops Tea, Freese's Hamlin's diarrhœa mixture. Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking. Polish or oil. Harney's diarrhœa mixture. Harter's wild cherry bitters. Hayden's viburnum comp. Hay fever remedies. Headine Heaves. Headine Heaves. Hebra's arsenic pills. Cosmetic liniment. Drachylon ointment Itch ointment Lead ointment Oriental cosmetic water. Sulphur paste. Heels ointment for cracked.	$\begin{array}{c} 223\\2448\\227\\227\\227\\220\\227\\220\\223\\223\\223\\223\\223\\223\\223\\223\\223$
Grease eradicators. Paints. Shoe. Stains, removing. Greases. 105, 108, Grecian varnish Green cerate Ink. Plaster Salve. Soap Mitigated Green's August flower; nervura. Greenhow's diarrhœa mixture. Gregory's laxative powder. Griffith's capsicum lotion. Grimault's matico injection. Grindelia, elixir of. Gross' neuralgic pills. Guaiac, balsam of. Decoction of comp. Elixir of. Glycerite of. Syrup of. Tincture of, aromatic. Guaiacol capsules, Dyche's. Guano, artificial. Guarana, elixirs containing. 55, Gum bougies. 14,	413 331 4464 4662 109 467 19 121 116 461 123 223 112 123 55 206 13 30 55 142 154 461 154 461 154 461 154 461 165 165 175 175 175 175 175 175 175 17	Hair renewer. Hamburg breast tea. Drops Tea, Freese's Hamlin's diarrhœa mixture Wizard oil Handkerchief extracts Hanson's corn salve. Hardwood filler. Harlan's tooth paste. Harness blacking Polish or oil. Harney's diarrhœa mixture. Harter's wild cherry bitters Hayden's viburnum comp. Hay fever remedies. Headache cures, electric. Remedies. Headaine Heaves. Hebra's arsenic pills. Cosmetic liniment Drachylon ointment Itch ointment Lead ointment Oriental cosmetic water Sulphur paste. Hectograph inks. 427 Mass.	2248 2248 227 227 220 112 223 221 223 241 414 413 357 414 414 413 223 231 251 188 188 254 118 253 118 253 118 118 253 251 251 251 251 251 251 251 251 251 251

Heliotrope Sachet	Hot Springs prescription	198
Water 306	House ink	417
White	Hubert's malvina cream	225
Hellmund's narcotico-balsamic oint 196	Lotion	225
Helmbold's buchu	Huchard's elixir	182
Helonia tablets	Huckleberry juice	107
Helonias, elixirs containing 56	Hudson's dentifrice	357
Helonin mixture, Schlotterbeck's 238	Honey of elm	225
	Hufeland's diuretic wine	162
Hemicranine	Huile antique	337
Hemorrhoid remedies	Philocome	337
Hemostatic cotton	Hungarian liniment	110
Henbane, chloroform of 20	Hunn's drops	112
Henbane, infused oil of 116	Hunter's tooth powder	357
Henry's carbolic salve 223	Hunyadi Janos water	372
Magnesia	Husband's magnesia	225
Tri-iodides	Hustenthee, Kneipp's	108
Hensel's tonicum	Hyacinth extract	287
Herpetic wash 111	Spirit of	284
Herzstaerkungs tropfen		162
Hesperis	Hydrastine, glycerite of	
1		55
	Hydrastis, elixir of	55
	Elixirs containing	93
Hind's honey and almond cream 224	Glycerite of	238
Hips, tincture of, Rademacher's 154	Mixture, Schlotterbeck's	
Hoarhound, syrup of	Solution of, colorless	132
Tar cough syrup 177	Hydrastis (see also Golden seal.)	
and tar, honey of 177	Hydrobromic acid cough mixture	179
and wild cherry, syrup of 149	Hydrocyanic acid, Scheele's	11
Hoff's malt extract	Hydroleine	225
Hog cholera remedies	Hydromel	105
Medicines	Hygienic vinegar	306
Holloway's ointment	Hygrometer paper	375
Holman's liver pad	Hyoscyamus, (see Henbane.)	
Homeopathic alcohol	Hypodermic injections106,	107
Liniment of aconite	Hypophosphites comb's with malt ext.,	
Tincture of avena sativa	etc	99
Cocculus indicus	Elixirs containing	57
Sulphur	Elixir of six	218
Honeys	Emulsion cod liver oil with	82
Honey and almond cream	Cod liver oil with creosote and	82
Balsam of	Paraffin oil with	84
Bees, tincture of	Fellows'	219
		132
	Solution of, acid	144
Hearhound and tar	Syrups containing140, 143,	1.4.4
Honeysuckle extract	TCITTINOI	27
Hoof cement	ICHTHYOL cotton	
Dressings	Idiaton	225
Hoofs, preps. for horses	Imperial cologne	302
Hoper's female pills	Incense	416
Hop bitters	Incrusting of boilers, preventing	380
Tonic 200	Indelible inks	428
Hops, elixir of		231
Horn, to color, black	Indian liniment	
Horner's rheumatic lightning 224		210
Horse and cattle food	Indigestion in cattle	261
Doses for	Infant cough mixture	178
Liniments. 255	Influenza in horses	254
Medicines	Infusions	106
Powders	Inhalation, bromine, Netolitsky's	106
Horsechestnut bark, extract of	Injection Bron	
Horseradish, spirit of, comp	Injection Bron	107
	Inks416-	433
	Ink erasives	427
Horsford's acid phosphate	Stains, removing	463
TIUSICILEI S DILLEIS	Stams, Temoving	100

Ink stains, indelible, to remove	431	Iron, Pyrophosphate, effervescent	126
for writing on photographs425,	426	Pyrophosphate, solution of	133
	432	Syrup of	1/15
		and quining gitrate offere	100
Insect bite remedy		and quinine citrate, efferv	100
Powders		Solution of	
Powder, blatta		Wine of	
Tincture of		Iodides, syrup of	144
		Salicylate, solution of	133
Insecticides	438	and sodium albuminate syrup	144
Invisible ink	428	Superphosphate, syrup of	
	226	Syrup of hypoposphites with	
	221	Tincture of, aromatic	
	58		
Iodides, elixirs containing57,	36	Athenstædt's	
Arsenic and mercury elixir	4 4 4	Compound	
	115	Tonic bitters	167
	118	Syrup	199
Solution of, carbolized	173	Wine of, bitter	162
Caustic, Churchill's	132	Sweet	163
Lugol's		Itch, barber's remedies for	166
Compound		Remedies for	188
Lugol's		Ivorine	
	132	Ivory to color blook	416
		Ivory, to color, black	410
	154	I I DOD INTO III I	90
Iodized collodion	21	ABORANDI, elixir of	63
Cotton	27	Tincture of	154
Starch	138	Jacoby's gauge	446
Iodo-bromide of calcium, elix. comp	45	Jalap and senna, fl. ext. of	103
lodoform bougies	16	Jamaica dogwood, fl. ext. of	94
Collodion	21	Japanese matches	440
Cotton	27		166
		Jaundice bitters	
	128	in cattle	261
	145	Javelle water	438
Iodol collodion	21	Java tonic	371
Cotton	$\begin{vmatrix} 21 \\ 27 \end{vmatrix}$		371 226
Cotton		Java tonic	
Cotton  Iodophenochloral	27	Java tonic	226
Cotton	27 $226$ $118$	Java tonic	226 225 318
Cotton.  Iodophenochloral Ipecac ointment. Powder, comp.	27 226 118 123	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin.	226 225 318 378
Cotton. Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of.	27 226 118 123 159	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula	226 225 318 378 318
Cotton. Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of	27 226 118 123 159 114	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic	226 225 318 378 318 318
Cotton. Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's.	27 226 118 123 159 114 154	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant.	226 225 318 378 318 318 107
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's and ammonium citrate, solut. of	27 226 118 123 159 114 154 133	Java tonic. 200, Jayne's expectorant Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin.	226 225 318 378 318 318 107 318
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's and ammonium citrate, solut. of. Phosphate syrup.	27 226 118 123 159 114 154 133 145	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose.	226 225 318 378 318 318 107 318 318
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's and ammonium citrate, solut. of Phosphate syrup Tartrate syrup.	27 226 118 123 159 114 154 133 145 145	Java tonic. 200, Jayne's expectorant Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin.	226 225 318 378 318 318 107 318 318 318
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's and ammonium citrate, solut. of Phosphate syrup Tartrate syrup.	27 226 118 123 159 114 154 133 145	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose.	226 225 318 378 318 318 107 318 318
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's and ammonium citrate, solut. of Phosphate syrup Tartrate syrup.	27 226 118 123 159 114 154 133 145 145	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses	226 225 318 378 318 318 107 318 318 318
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for Chloride bougies	27 226 118 123 159 114 154 133 145 145 384	Java tonic. 200, Jayne's expectorant Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet.	226 225 318 378 318 318 107 318 318 318 336
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies Syrup of.	27 226 118 123 159 114 154 133 145 145 145 145 145	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet Violet	226 225 318 378 318 107 318 318 318 318 318 318 319
Cotton Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of. Irish moss, mucilage of. Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies. Syrup of. Weld's.	27 226 118 123 159 114 154 133 145 145 145 145 145 243	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet Violet Witch hazel	226 225 318 378 318 318 318 318 318 318 318 318 318
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies Syrup of Weld's. Citrate, effervescent.	27 226 118 123 159 114 154 133 145 145 145 145 243 125	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet Violet Witch hazel Jesson's dental anæsthetic.	226 225 318 378 318 318 318 318 318 318 318 318 226
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup Tartrate syrup. Cement for. Chloride bougies Syrup of Weld's. Citrate, effervescent with magnesia, efferv.	27 226 118 123 159 114 154 133 145 145 145 145 126 126	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant. Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic Jesuit drops.	226 225 318 378 318 318 107 318 336 318 318 326 32
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies. Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of.	27 226 118 123 159 114 154 133 145 145 145 145 126 126 163	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic Jesuit drops. Jeweler's rouge	226 225 318 378 318 318 318 318 318 319 318 226 32 438
Cotton Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of. Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies. Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of. Collodion.	27 226 118 123 159 114 154 133 145 145 125 126 163 21	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic. Currant. Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel. Jesson's dental anæsthetic Jesuit drops. Jeweler's rouge Jewsbury & Brown's tooth paste.	226 225 318 378 318 318 318 318 318 319 318 226 32 438 226
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's and ammonium citrate, solut. of Phosphate syrup Tartrate syrup Cement for Chloride bougies Syrup of Weld's Citrate, effervescent with magnesia, efferv Wine of Collodion Comb's with malt ext., etc.  97-	27 226 118 123 159 114 154 133 145 145 145 125 126 163 21 100	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet Violet Witch hazel Jesson's dental anæsthetic Jesuit drops Jewebury & Brown's tooth paste. Jockey club, extract	226 225 318 378 318 318 318 318 318 318 326 328 226 288
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup Tartrate syrup Cement for. Chloride bougies Syrup of Weld's Citrate, effervescent with magnesia, efferv. Wine of Collodion Comb's with malt ext., etc. 97- Dialyzed	27 226 118 123 159 114 154 133 145 145 145 125 126 163 21 100	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic. Jesuit drops Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet.	226 225 318 378 318 318 318 318 318 326 328 226 288 295
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup Tartrate syrup Cement for. Chloride bougies Syrup of Weld's Citrate, effervescent with magnesia, efferv. Wine of Collodion Comb's with malt ext., etc. 97- Dialyzed	27 226 118 123 159 114 154 133 145 145 145 125 126 163 21 100	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic. Jesuit drops Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet.	226 225 318 378 318 318 318 318 318 318 326 328 226 288
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup Tartrate syrup. Cement for. Chloride bougies Syrup of Weld's. Citrate, effervescent with magnesia, efferv. Wine of Collodion Comb's with malt ext., etc. 97- Dialyzed Elixirs containing. 58-	27 226 118 123 159 114 154 133 145 145 145 126 163 21 100 133	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic. Jesuit drops Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax.	226 225 318 378 318 318 318 318 318 326 328 226 288 295
Cotton Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies. Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of. Collodion. Comb's with malt ext., etc. 97– Dialyzed. Elixirs containing. 58– Etching of.	27 226 118 123 159 114 154 133 145 145 145 126 126 100 133 63 398	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic Jesuit drops. Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax Juices	226 225 318 378 318 318 318 318 326 328 226 288 295 111
Cotton Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies. Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of. Collodion. Comb's with malt ext., etc. 97- Dialyzed. Elixirs containing. 58- Etching of. Iodohydrargyrate, syrup of.	27 226 118 123 159 114 154 133 145 145 145 126 163 21 1133 63 398 145	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet Violet Witch hazel Jesson's dental anæsthetic Jesuit drops Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax Juices Fruit.	226 225 318 378 318 318 318 318 318 326 326 226 288 295 111 107 364
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies Syrup of Weld's. Citrate, effervescent. with magnesia, efferv. Wine of Collodion. Comb's with malt ext., etc. 97- Dialyzed Elixirs containing. 58- Etching of Iodohydrargyrate, syrup of and magnesium citr. efferv.	27 226 118 123 159 145 145 145 145 145 126 163 21 103 63 398 145 126	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet Violet Witch hazel Jesson's dental anæsthetic Jesuit drops Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet Johnson's lotion of borax Juices Fruit Juniper berry syrup.	226 225 318 378 318 318 318 318 318 326 328 226 228 226 226 111 107 364 107
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup Tartrate syrup Cement for. Chloride bougies Syrup of Weld's Citrate, effervescent with magnesia, efferv. Wine of Collodion Comb's with malt ext., etc. 97– Dialyzed Elixirs containing. 58– Etching of Iodohydrargyrate, syrup of and magnesium citr., efferv. Mixture, Rademacher's	27 226 118 123 159 159 145 145 145 145 126 163 21 100 133 63 398 145 126 113	Java tonic. 200, Jayne's expectorant Vermifuge Jelly, arnica Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet Violet Witch hazel Jesson's dental anæsthetic Jesuit drops Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax Juices Fruit. Juniper berry syrup Catgut.	226 225 318 378 318 318 318 318 318 318 326 328 226 288 295 111 107 364 107
Cotton Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of. Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Cement for. Chloride bougies. Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of. Collodion. Comb's with malt ext., etc. Dialyzed. Elixirs containing. Etching of. Iodohydrargyrate, syrup of. and magnesium citr. efferv. Mixture, Rademacher's. Oxychloride, solution of.	27 226 118 123 159 114 154 133 145 145 126 126 133 63 398 145 126 113 133 145 126 1133 133 145 145 145 126 133 145 145 145 145 145 145 145 145 145 145	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic Jesuit drops. Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax Juices Fruit. Juniper berry syrup Catgut. FI. exts., containing.	226 225 318 378 318 318 318 318 336 318 326 226 228 226 228 111 107 18 88
Cotton Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies. Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of Collodion. Comb's with malt ext., etc. 97- Dialyzed Elixirs containing. 58- Etching of. Iodohydrargyrate, syrup of. and magnesium citr., efferv. Mixture, Rademacher's. Oxychloride, solution of. Phosphate, effervescent.	27 226 118 123 159 114 154 133 145 145 126 163 398 145 126 63 398 145 126 126 126 126 126 126 126 126 126 126	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic. Currant. Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel. Jesson's dental anæsthetic Jesuit drops. Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax Juices Fruit. Juniper berry syrup Catgut. FI. exts., containing. Juice.	226 225 318 318 318 318 318 318 336 338 326 328 226 328 295 111 107 364 107 188 88 107
Cotton Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of. Collodion. Comb's with malt ext., etc. 97- Dialyzed. Elixirs containing. 58- Etching of. Iodohydrargyrate, syrup of. and magnesium citr. efferv. Mixture, Rademacher's. Oxychloride, solution of. Phosphate, effervescent.	27 226 118 123 1159 114 154 133 145 384 17 144 243 125 1163 21 100 133 398 145 126 113 133 145	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic Jesuit drops. Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax Juices Fruit. Juniper berry syrup Catgut. FI. exts., containing.	226 225 318 378 318 318 318 318 336 318 326 226 228 226 228 111 107 18 88
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of. Collodion. Comb's with malt ext., etc. 97– Dialyzed. Elixirs containing. 58– Etching of. Iodohydrargyrate, syrup of. and magnesium citr. efferv. Mixture, Rademacher's. Oxychloride, solution of. Phosphate, effervescent. Solution of. and potassium tartrate, syrup.	27 226 118 159 114 154 133 145 145 145 125 126 163 398 145 126 113 133 126 113 133 145	Java tonic. 200, Jayne's expectorant Vermifuge Jelly, arnica Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet Violet Witch hazel Jesson's dental anæsthetic Jesuit drops Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet Johnson's lotion of borax Juices Fruit. Juniper berry syrup Catgut. F1. exts., containing Juice. Pomade	226 225 318 378 318 318 318 336 318 326 228 226 228 226 111 107 164 107 121
Cotton Iodophenochloral Ipecac ointment Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of. Collodion. Comb's with malt ext., etc. 97– Dialyzed. Elixirs containing. 58– Etching of. Iodohydrargyrate, syrup of. and magnesium citr. efferv. Mixture, Rademacher's. Oxychloride, solution of. Phosphate, effervescent. Solution of. and potassium tartrate, syrup.	27 226 118 159 114 154 133 145 145 145 125 126 163 398 145 126 113 133 126 113 133 145	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic Currant Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel Jesson's dental anæsthetic Jesuit drops. Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax Juices Fruit. Juniper berry syrup Catgut. Fl. exts., containing. Juice. Pomade  K ALI lemon	226 225 318 378 318 318 318 336 318 336 338 226 288 225 111 107 18 88 107 121
Cotton Iodophenochloral Ipecac ointment. Powder, comp. Vinegar of Irish moss, mucilage of Iron acetate tinct., Rademacher's. and ammonium citrate, solut. of. Phosphate syrup. Tartrate syrup. Cement for. Chloride bougies Syrup of. Weld's. Citrate, effervescent. with magnesia, efferv. Wine of. Collodion. Comb's with malt ext., etc. 97- Dialyzed. Elixirs containing. 58- Etching of. Iodohydrargyrate, syrup of. and magnesium citr. efferv. Mixture, Rademacher's. Oxychloride, solution of. Phosphate, effervescent.	27 226 118 159 1114 133 145 145 145 145 126 21 100 133 398 145 113 398 145 113 113 113 113 113 113 113 113 113 11	Java tonic. 200, Jayne's expectorant. Vermifuge Jelly, arnica. Benzin. Calendula Cosmetic. Currant. Glycerin Rose. Roses Shampoo Toilet. Violet Witch hazel. Jesson's dental anæsthetic Jesuit drops. Jeweler's rouge Jewsbury & Brown's tooth paste. Jockey club, extract Sachet. Johnson's lotion of borax Juices Fruit. Juniper berry syrup Catgut. FI. exts., containing. Juice.	226 225 318 378 318 318 318 336 318 326 228 226 228 295 111 107 18 88 88 107 107 118

Kalmia, tincture of	Lard, Anhydrous	108
Kalodont	Balsamic	108
Kalydor	Benzoated	108
Kaputine	Lards, factitious	108
Karmeliter geist	Larks, food for	
Kaskine	Lartique's gout pills	
Keg, to determine capacity of 440	Lassar's paste	120
Kendall's spavin cure	Laubach's eclectic liniment	
Kennedy's medical discovery 226	Laudanum, Dutchman's	109
Pinus canadensis	Sydenham's	150
Kerosene emulsion	Lavender extract	289
Kern's insect annihilator	Fl. ext. comp.	9.
Kickapoo Indian oil	Sachet	290
Kid shoe dressing	Spirit of	284
Kidney remedies	Water	305
Kidney wort	Laville's anti-gout liquor	228
King's expectorant tincture		228
	LavolineLawn fertilizer	
New discovery		46
Kiss-me-quick extract	Laxative elixir	
Kissingen salt, artificial. efferv 126	Powder, Gregory's	128
Water	Species	136
Kitchell's liniment	Laxol	228
Kline's nerve restorer	Lead carbonate ointment, comp	118
Kneipp's remedies, Pastor	Chloride lotion, Tuson's	111
Knox's disinfecting powder	Ointment, comp	118
Koeller's blood purifying tea 170	Hebra's	118
Koelreuter's tinct. of rhubarb 157	Paper	465
Kænig's Hamburg breast tea	Plaster collodion	21
Hamburg drops	Subacetate, glycerite of	105
Kohler's one-night corn salve244B	and sulphur hair promoters	345
Kola coca syrup	Leather blacking	379
Elixir of	Dyeing	396
Extract of, 94; wine of	Lacquer for	439
Kousso, extract of, fl	Polish or dressing	454
Krampf tropfen	Ledoyen's disinfectant	
Kummerfeld's cosmetic water 329	Lemon grass, spirit of	284
Kutnow's Laxative Powder	Juice, artificial	107
[ A DIAPHANE 323	Kali	108
Label varnish	Peel, tincture of	155
Labels to bottles, to cement 385	Salt of artificial	128
Lacquers	Salt of, artificial	284
Lactopeptine	Syrup.	368
Elixir. 218		365
Elixir with bismuth	Lemonade seltzer	900
	Leptandra, (see Culver's root.)	386
	Letters, cement for porcelain	
Combinations of	Library paste	$\frac{446}{262}$
Liquid	Lice in cattle	
Lactophosph. calc., emul. of c. l. oil with. 82	1 /	276
Lactophosphates, elixirs containing 63		439
Lactucarium, syrup of	Licorice, elixirs of	64
Ladies' shoe dressing	Extract of	95
Slipper, powder of, comp	I.	163
Lafayette cologne	Life, elixir of	64
Laird's bloom of youth	Lightning, fluid	192
Lait virginal	Renovator	387
Lallemand's gout specific	Lights, Bengal	378
Lamellæ31	,	439
Lanolin cold cream		465
Cream	Linseed oil, varnish	468
Milk		134
Toilet	Liquid cosmetics	325
Lapis divinus	Face paint	326
Medicamentosus		411
Mirabilis	Kill'em	
Lard 108	Pearl	326

	333	Lobelia and Capsicum, Lotion of, comp	111
Liquidambar, syrup of	145	Syrup of	145
Liquor, (see Solution).		Syrup of, Thompsonian	145
	129	Tincture of, comp	154
	131	Vinegar of	159
	207	Locust blossom extract	289
Naphthalini benzinatus	434	Logan's plaster	121
Sulphocarbonatus	434	Logwood, extract of, fl	95
Natri nitrici		Extract solution	418
Pectoralis	- 63 (	Ink417,	418
Picis carbonis		Paper	466
Sedans	228	London soap powder	468
Seriparus	135	Long life, elixir of	64
Lilac extract	288	Species	136
Spirit of		Look album	80
Water		Loomis' diarrhœa mixture	113
Lilacin		Loss of hair, remedies for	
Lilies, oil of		Lotions	
Lilionese		Lotion, Barton's	189
Lily of the valley, extract	288	Cosmetic	317
Tincture of			111
White extract	280	Evaporating	317
White, extract		Face, glycerin	
Lime, fly	247	Mammillary	$\frac{194}{111}$
Juice and glycerin 320,	941	Taylor's	
Linaloe, spirit of		Toilet	317
Linctus, white	80	Lovage, extract of	229
Linden water	160	Lovage, extract of	95
Limments	190	Low's magnetic liniment	229
Liniment, iodide of ammonia	221	Lozenges, cough	
Cosmetic, Hebra's	359	Pine tree tar	199
Horse	255	Worm	202
Rheumatic	109	Lubricants	439
Stable	255	Lugol's caustic sol. of iodine	132
Linoleum, polish for	439	Iodine solution	132
Linseed cough syrup or balsam	179	Rubefacient sol. of iodine	
Mucilage	114	Lung Balsam	
Oil, bleaching of		Luperine	229
Boiled		Lupulin, elixirs, containing	65
Emulsion, Thompson's	83	Tincture of	
Lip ointment	360	Lute for retorts	386
Preparations		Lydia Pinkham's veget. comp	234
Salves, pomades	359	Lyon's kathairon; tooth powder228,	229
Liriodendron extract			228
	000	Lysol	المالم
Listerine		M'DADE'S compound	86
Listol. Listol Tablets	44B		229
Lister's anticeptic fluid	130	Prescription	
Carbolic paste	120	Succus alterans	900
	28	McDougall's disinfecting powder	229 3115
Sero-sublimate cotton  Lithia tablets, comp		McGill's orange blossom	999
Lithiated bydranges		McLean's strengthening cordial	250
Lithiated hydrangea	196	McMunn's elixir of opium	230
Lithium carbonate, effervescent	196	Mabel bouquet extract	
	126	Macassar oil	
Comp's, elixirs containing	100	Machine oil, sewing	
Litmus paper		Mackenzie's eye lotion	184
Little cathartic pills, liver granules	175	Magen balsam	13
	158	Magendie's solution of iodine	132
Liver bitters	167	Morphine	-134
Granules, little	175	Magentrost, Kneipp's	108
Invigorator	192	Magic corn salve	223
Regulators	192	Magnesia, effervescing powder with	122
Lloyd's hydrastis		Milk of	112
Leontin		Magnesium borocitrate solution	133
Lobelia and capsicum, tinct., comp	155	Citrate, effervescent	
Extract of, fl. comp	95	Citrate, solution of	133
		Citiesto, Solution Olivinia and a constant	100

			2.0
Magnesium Lights	402	Melissa spirit of	192
Sulphate, effervescent		Melissa, spirit of	
Magnolia balm	222	Menthol glycerin tooth cream	
Mahogany stain	463	Plaster	
Maiden's blush		Smelling salts	308
Maidenhair, syrup of		Snuff	
Maizo-lithium	244B	Toilet cream	
Malefizoel, Kneipp's	108	Mentholin	171
Malt, elixirs containing	65	Merchant's gargling oil	
Extract of, and combinations96-	-100	Mercuric chloride catgut	18
Extract, Hoff's	224	Collodion	21
Fluid	97	Cotton	27
Malta balsam	13	Silk	128
Maltine, and combinations	229	Iodide ointment	118
Malto syrup	369	Mercury albuminate, solution of	134
Malto-yerbine	229	and arsenic iodides, sol. of	134
Malvina cream		Chloride, solution of	134
Lotion		Compounds, elixirs containing	-65
Mammillary lotion		Iodide, syrup of	146
Ointment	194	and potassium, tartrate sol	134
Manaca fi. ext	100	Merrell's vaginal discs	$\frac{230}{385}$
and salicylates, elixir of	65 322	Metals to glass, to cement  Polishing of	440
Mandelklei	022	Varnish	468
Mandrake (see Podophyllum).  Manganese hypophosph. syrup143,	144	Metcalf's cologne.	303
Iodide, syrup of	145	Mettauer's laxative aperient	129
Phosphate, syrup of	146	Solution aloes and soda	129
Mange in dogs	-	Mexican mustang liniment	230
in horses		Mezereon, extract of	100
Manicure cosmetics		Ointment	118
Manifold writer		Micajah uterine wafers	-230
Manna and senna, syrup of	147	Mildew, to remove	440
Syrup of, comp	147	Migranin	230
Map varnish		Milk, almond	79
Maple syrup		Almond, compound	. 80
Marble busts, to clean	440	Cacao, cucumber	
Cement for		Increasing of, in cows	
Cleansing and polishing of	439 296	Magnesia	
Mariana wine of coca		Powder for cows	262
Marie Stuart	289	Roses	040
Marienbad reduction pills	230	Stains, removing	462
Marine cement	385	Toilet	316
Glue	385	Virgin's	329
Marking ink	430	Millefleur sachet	296
Marrol	230	Extract	
Marrow hair pomade	340	Miller's Arabian balsam	230
True and artificial	111	Golden oil	244C
Marshall's dentifrice	4 4 0	Mineral waters, artificial371-	
Marshmallow ointment	118	Miraculous plaster, Rademacher's	
Species	136	Mirrors, polish for	411
Tea	156	Alterantiæ composita	112
Marythistle, tincture of	152 138	Smilacis compositus	86
Matches, Japanese		Mitchella, elixirs containing	56
Matico, infusion of	106	Syrup of, comp	
Elixir of, compound	65	Mitchell's eye salve	44C
May-apple (see Podophyllum).		Mixed bird seed	277
May bells extract	289	Mixtures	
Dew water	329	Mixture, almond	79
Mead essence		Brown, improved	178
Medicamentum		Phosphatic	83
Meerschaum, cement for		Mocking bird food	
Meleckel		Modeling wax, dentist's	
Melachol	250	Molds for taking impressions	440

Moles, for removing	192	Nelaton's suppositories	231
Mollisin		Nephritic powder, Rademacher's	$123^{\circ}$
Monobrom. camphor, emulsion of	180	Neroli, spirit of	284
Morphine acetate, solution of	$\frac{134}{135}$	Nerve and bone liniment	
Bimeconate, sol. of	134	and bone liniment, Barker's	
Cocaine cotton	21	Drops	187
Hydrochlor., sol. of	135	Powder	122
Injection of	107	Thompsonian	123
Oleate of	119	Tonic	192
Solution of		Nervine celery	193
Valerianate, elixir of	65	Nervous debility, remedies for	
Morrison's pills		Nervura	106
Mortars, cement for mending Mosquito essence		Netolitsky's bromine inhalation  Neuralgia remedies	
Lotion		Neurosin	001
Oil	256	Neurosine	
Pastilles	441	Neutralizing cordial	147
Tincture		Elixir	
Moss rose extract		Powder	123
Moth essence		Newmown hay extract	
Paper	441	Sachet	
Powder		Newport cologne	
Species		Newton's metal	231
Mother's cordial		Comp. cod liver oil	
Drops		Nickel bromide, syrup of	
Salve		Plating	
Seigel's curative syrup	238	Night-blooming cereus extract	290
Tinctures	155	Fluid ext. of	88
Mother-of-pearl, cement for		Tincture of	88
Mother's friend	185	Nightingales, food for	277
Mountain mint, tincture of		Nipples, fissured, cures for	
Mousseline extract		Nitric fumigation	
Mouth essence	358	Norton's chamomile pills	
Preparations		Norwood's tincture	0.04
Washes		Nouche's lotion of tin chloride	
Moxie syrup	369	Number one (Thompsonian)	150
Mucilages	-444	Two (Thompsonian)	
Mucilage, stick or bar		Three (Thompsonian)	
Muguet Padamashar's	281	Four (Thompsonian)	14
Mugwort root, tinct., Rademacher's Mund wasser		Five (Thompsonian)	
Murdock's liquid food	231	Nursery powders	
Murray's infallible system tonic		Nut stains, removing	
Musk		Nut stains, removing	418
Extract		Nutmeg balsam	13
Root, elixirs containing	77	Cerate	13
Sachet		Spirit cf	, 284
Tincture of	285	Nutrient fluids388,	444
Root, (see Sumbul)		Gelatin	$\begin{array}{c} 444 \\ 66 \end{array}$
Muskat balsam		Tincture, Rademacher's	
Preparations for		Water, Rademacher's	
Mustang liniment			
Mustard liniment	110	OATS, tincture of	511
Mutter salbe	18	Ocean spray extract	. 290
Myrrh and aloes, fl. ext. of		Oculine	231
Lotion of, comp		Odol	231
NAPHTHALIN cotton	$\frac{28}{444}$	Odontine	. 553 . 231
Paper	$\frac{444}{231}$	Odontunder	2.14C
Naphthol camphor	231	Odors, classification of	. 280
Nectar syrup		Delicate and heavy	283

Oils	Orange, elixir of	66
Oil of bitter almond	Elixir, compound	66
Black		290
Bleaching of		423
Cloves 279		284
Cloves	The state of the s	
Linusion	1	369
For watchmakers	Wine of	163
Geranium. 279 Layender: peroli bigarade 279	Compound	-66
	Oregon grape, fl. ext	87
Neroli petale	Orgeat syrup	370
Orange flowers	Oriental cologne	302
Orange peel	Cream	326
Orris	Sachet	297
Petit grain		226
Rose geranium	Tooth paste	
		280
		284
Spike	Tincture of	285
Sulphurated	Osgood's India chologogue	231
Oil-sugars	Osmunda, syrup of, comp	146
Oiled paper 445	Ox-gall soap	460
Oils, liniment of	Oxygen aquæ	231
Liniment of, comp	Oxygen aquee	
116 119 194	Compounds231,	202
Ointments	Oxymel	
Ointment, antineuralgic	Ozonized water	232
Aromatic		
for birds	P. P.P	235
Brown	Pain cure	205
for cracked heels	Dispeller	205
Cream		
for finger-nails	Expeller190,	205
10 Iniger-hans	Killer	190
Hebra's itch	Painkiller, Davis'	216
Lead	Paine's celery compound	232
for horses	Paint, black, bicycle	378
Lip 366	Black, glossy	444
Mammillary	Blackboard	378
Mayer's 118	Bronze	381
Pile	Eradicator	445
Tartar emetic	Ctoing removing	
White	Stains, removing	462
White	White enamel	444
Oleose Co		444
O'Leary's oxygen compound	Palmer's invisible powder	232
Oleate of morphine	Lily white tablet	232
Ammonium	Lotion	
Strychnine	Pancoast's cough mixture	
Tin	Pancreas, elixir of	66
Oleosaccharates	Pancreatin comb., with malt, etc 98	
Olfactorium anticatarrhoicum 173	Tancicatin combi, with mart, etc	67
One-night corn salve	Biland containing,	
Opera cologne		168
Opthalmic discs. 31		98
	[ I allereopepsiii	232
Opiated syrup	Pancropepsin	232
Opium, confection of	'i Pan·peptic tablets	232
Extract of 100		67
Fl. ext., aqueous deodorized 100	Paper, azolitmin	465
Liniment		
Liniment ammoniated 110		
Powder, comp		
Tincture of, ammoniated 156	Carbaliand	
Anisated, benzoated, compound, cro-		
	Ceresin	445
cated	Congo red	465
and saffron, tincture of		445
Wine of, compound	Curcuma	465
Licorice with	Fly	
Opodeldoc liquid	Fumicating	301
Opoponax sachet 297 Orange blossoin 226	Hygrometer	
Orange blossom	Lead	
•	Dead	200

Paper, Litmus	
Logwood	
Moth	
Naphthalin 44	
Oiled	
Paraffined 448	Pencils, cleansing
Parchment 446	Fig
Potassium iodide-starch	Pennyroyal, essence of
Starch	Pepper, confection of
Test	Ointment
Tracing	Peppermint, syrup of
Waxed	
Wrapping, to cut	
Papier Fayard	Effervescent 197
Papine	
Paraffined paper	
Paraldehyde, elixir of	
Parchment paper	
Paregoric	
Chloroform	
Concentrated 100	Perl's antikrinin
Pareira, decoction of 30	Perry Davis' painkiller
Elixir of 67	Peru balsam, syrup of 146
F1. ext. containing 88	Balsam, tincture of
Parker's tonic	
Parmese violet extract	Pestle's, cement for mending 385
Parson's local anæsthetic	Peterman's roach food
Partridgeberry, syrup of, comp 146	
Paskola	
Pasta Mack	
Pastes	
Paste, almond	Soap
(2 1 1' T' 1 1	Doap.
	Pottit's ave salve
Carbolic, Lister's	Pettit's eye salve
Currant 107	Pettit's eye salve
Currant	Phalon's hair restorative. 2440 Pheasants, gapes in 278
Currant         107           Electric         451           Furniture         408	Pettit's eye salve. 23: Phalon's hair restorative 2440 Pheasants, gapes in 278 Phenatol
Currant         107           Electric         451           Furniture         408           Gonorrhœa         186	Pettit's eye salve. 23: Phalon's hair restorative 2446 Pheasants, gapes in 278 Phenatol Phenol camphor. 11
Currant         107           Electric         45           Furniture         49           Gonorrhea         18           Lassar's         126	Pettit's eye salve. 238 Phalon's hair restorative 2440 Pheasants, gapes in 278 Phenatol Phenol camphor 11 Sodique 238
Currant       107         Electric       451         Furniture       408         Gonorrhœa       186         Lassar's       12         Library       446	Pettit's eye salve.       238         Phalon's hair restorative       2440         Pheasants, gapes in       278         Phenatol       11         Sodique       238         Phenolid       2440
Currant       107         Electric       451         Furniture       408         Gonorrhœa       186         Lassar's       12         Library       446         Phosphorus       446	Pettit's eye salve.       238         Phalon's hair restorative       2440         Pheasants, gapes in       278         Phenatol       21         Sodique       238         Phenolid       2440         Phenosolyl       238
Currant       107         Electric       451         Furniture       408         Gonorrhœa       186         Lassar's       120         Library       444         Phosphorus       446         Polishing       448	Pettit's eye salve.       23         Phalon's hair restorative       2440         Pheasants, gapes in       278         Phenatol       11         Sodique       238         Phenolid       2440         Phenosolyl       238         Phillips milk of magnesia       2440
Currant       107         Electric       451         Furniture       408         Gonorrhœa       186         Lassar's       12         Library       446	Pettit's eye salve.       238         Phalon's hair restorative       2440         Pheasants, gapes in.       278         Phenatol       11         Sodique       238         Phenolid       2440         Phenosolyl       238         Phillips milk of magnesia       2440         Phospho-mur. quinine       233
Currant       107         Electric       451         Furniture       408         Gonorrhœa       188         Lassar's       12         Library       446         Phosphorus       444         Polishing       448         for windows       468         Razor       451	Pettit's eye salve.       238         Phalon's hair restorative       2440         Pheasants, gapes in.       278         Phenatol       11         Sodique       238         Phenolid       2440         Phenosolyl       238         Phillips milk of magnesia       2440         Phospho-mur. quinine       233
Currant       107         Electric       451         Furniture       408         Gonorrhœa       188         Lassar's       12         Library       446         Phosphorus       444         Polishing       448         for windows       468         Razor       451	Pettit's eye salve.       238         Phalon's hair restorative       2440         Pheasants, gapes in       278         Phenatol       1         Phenol camphor       11         Sodique       238         Phenolid       2440         Phenosolyl       233         Phillips milk of magnesia       2440         Phospho-mur, quinine       233         Syr, wheat phosphates       2440         Wheat phosphates       2440
Currant       107         Electric       451         Furniture       408         Gonorrhœa       186         Lassar's       12         Library       44         Phosphorus       444         Polishing       448         for windows       468         Razor       451         Roach       451	Pettit's eye salve.       238         Phalon's hair restorative       2440         Pheasants, gapes in       278         Phenatol       1         Phenol camphor       11         Sodique       238         Phenolid       2440         Phenosolyl       233         Phillips milk of magnesia       2440         Phospho-mur, quinine       233         Syr, wheat phosphates       2440         Wheat phosphates       2440
Currant       107         Electric       451         Furniture       408         Gonorrhoea       186         Lassar's       12         Library       446         Phosphorus       446         Polishing       448         for windows       468         Razor       451         Roach       451         Tooth       351–358	Pettit's eye salve.       238         Phalon's hair restorative       2440         Pheasants, gapes in       278         Phenatol       218         Phenol camphor.       11         Sodique       238         Phenolid       2440         Phenosolyl       233         Phillips milk of magnesia       2440         Phospho-mur. quinine       233         Syr. wheat phosphates       2440         Wheat phosphates       2440         Philocome       337
Currant     107       Electric     451       Furniture     408       Gonorrhea     188       Lassar's     120       Library     444       Phosphorus     444       Polishing     448       for windows     468       Razor     451       Roach     451       Tooth     351-358       Wax     120	Pettit's eye salve.       33:         Phalon's hair restorative       2446         Pheasants, gapes in       278         Phenatol       11         Sodique       238         Phenolid       2440         Phenosolyl       238         Phillips milk of magnesia       2440         Phospho-mur. quinine       238         Syr. wheat phosphates       2440         Wheat phosphates       2440         Philocome       337         Phosphate, iron and quin., ext. malt, w. 100
Currant       107         Electric       451         Furniture       408         Gonorrhea       188         Lassar's       120         Library       444         Phosphorus       446         Polishing       448         for windows       465         Razor       451         Roach       451         Tooth       351–358         Wax       120         Pasteurine       232	Pettit's eye salve. 33: Phalon's hair restorative 244C Pheasants, gapes in 278 Phenatol Phenol camphor. 11 Sodique 235 Phenolid 244C Phenosolyl 233 Phillips milk of magnesia 244C Phospho-mur. quinine 233 Syr. wheat phosphates 244C Wheat phosphates 244C Philocome 337 Phosphate, iron and quin., ext. malt, w. 100 Wild cherry 365, 367
Currant       107         Electric       451         Furniture       408         Gonorrhoea       188         Lassar's       120         Library       446         Phosphorus       444         Polishing       448         for windows       468         Razor       451         Roach       451         Tooth       351-358         Wax       120         Pasteurine       232         Pastilles, fly       406	Pettit's eye salve.         33           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenotol         218           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur. quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365, 367           Phosphates, acid         366
Currant       107         Electric       451         Furniture       408         Gonorrhœa       186         Lassar's       122         Library       446         Phosphorus       444         Polishing       448         for windows       468         Razor       451         Roach       451         Tooth       351-355         Wax       120         Pasteurine       232         Pastulles, fly       406         Fumigating       300	Pettit's eye salve.         33           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenatol         218           Phenol camphor         11           Sodique         233           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur, quinine         233           Syr, wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w. 100           Wild cherry         365, 367           Phosphates, acid         366           Elixir of quinine and         71
Currant       107         Electric       451         Furniture       408         Gonorrhœa       186         Lassar's       122         Library       446         Phosphorus       444         Polishing       448         for windows       468         Razor       451         Roach       451         Tooth       351-355         Wax       120         Pasteurine       232         Pastulles, fly       406         Fumigating       300	Pettit's eye salve.         23           Phalon's hair restorative         2446           Pheasants, gapes in.         278           Phenatol.         11           Sodique         238           Phenolid         2440           Phenosolyl.         238           Phillips milk of magnesia         2440           Phospho-mur. quinine         238           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w. 100         Wild cherry         365, 367           Phosphates, acid.         366         366           Elixir of quinine and         71         71           Powder of, comp.         123
Currant         107           Electric         451           Furniture         408           Gonorrhea         188           Lassar's         120           Library         444           Phosphorus         444           Folishing         448           for windows         468           Razor         451           Roach         451           Tooth         351-358           Wax         120           Pasteurine         232           Pastulles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         168	Pettit's eye salve.         33:           Phalon's hair restorative         2440           Pheasants, gapes in.         278           Phenatol         11           Sodique         238           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur. quinine         238           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w. 100         Wild cherry         365, 367           Phosphates, acid         366         366           Elixir of quinine and         71         71           Powder of, comp         123         23           Syrup of comp., ext. of malt with         99
Currant         107           Electric         45           Furniture         408           Gonorrhea         188           Lassar's         120           Library         446           Phosphorus         446           Polishing         448           for windows         468           Razor         451           Roach         451           Tooth         351-358           Wax         120           Pasteurine         232           Pastulles, fly         406           Funigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290	Pettit's eye salve.         238           Phalon's hair restorative         2446           Pheasants, gapes in.         278           Phenatol         11           Sodique         238           Phenolid         2440           Phenosolyl.         238           Phillips milk of magnesia         2440           Phospho-mur. quinine         238           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w. 100         Wild cherry         365, 367           Phosphates, acid.         366         Elixir of quinine and         71           Powder of, comp.         123         Syrup of comp., ext. of malt with         99           Three, elixir of         61
Currant         107           Electric         451           Furniture         408           Gonorrhœa         188           Lassar's         120           Library         446           Phosphorus         444           Polishing         448           for windows         468           Razor         451           Tooth         351-358           Wax         120           Pasteurine         232           Pastulles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297	Pettit's eye salve.         33           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenotol         218           Phenolid         2440           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur, quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365         367           Phosphates, acid         366         Elixir of quinine and         71           Powder of, comp         123         Syrup of comp., ext. of malt with         99           Three, elixir of         61         Phosphatic emulsion         83
Currant         107           Electric         451           Furniture         408           Gonorrhœa         186           Lassar's         122           Library         446           Phosphorus         444           Polishing         448           for windows         468           Razor         451           Tooth         351-355           Wax         120           Pasteurine         232           Pastulles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         284	Pettit's eye salve.         238           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenatol         218           Phenol camphor.         11           Sodique         238           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur. quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w. 100         100           Wild cherry         365, 367           Phosphates, acid         366           Elixir of quinine and         71           Powder of, comp         123           Syrup of comp., ext. of malt with         99           Three, elixir of         61           Phosphatic emulsion         83           Mixture         83
Currant         107           Electric         451           Furniture         408           Gonorrhea         188           Lassar's         120           Library         444           Phosphorus         444           Polishing         448           for windows         468           Razor         451           Roach         451           Tooth         351-358           Wax         120           Pasteurine         232           Pastulles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         298           Tincture         283	Pettit's eye salve.         23           Phalon's hair restorative         2440           Pheasants, gapes in.         278           Phenatol.         11           Sodique.         238           Phenolid.         2440           Phenosolyl.         233           Phillips milk of magnesia.         2440           Phospho-mur. quinine.         238           Syr. wheat phosphates.         2440           Wheat phosphates.         2440           Philocome.         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry.         365, 367           Phosphates, acid.         366           Elixir of quinine and         71           Powder of, comp.         123           Syrup of comp., ext. of malt with         99           Three, elixir of         61           Phosphatic emulsion         83           Mixture         83           Phospho-albumen         233
Currant         107           Electric         451           Furniture         408           Gonorrhea         188           Lassar's         120           Library         446           Phosphorus         446           Polishing         448           for windows         468           Razor         451           Roach         451           Tooth         351–358           Wax         120           Pasteurine         232           Pastulles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         284           Tincture         282           Patent leather shoe polish         456	Pettit's eye salve.         33           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenotol         218           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur. quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Whilocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365           Phosphates, acid         366           Elixir of quinine and         71           Powder of, comp         123           Syrup of comp., ext. of malt with         99           Three, elixir of         61           Phosphatic emulsion         83           Mixture         83           Phospho-albumen         233           Phosphorated emulsion         84
Currant         107           Electric         451           Furniture         408           Gonorrhea         188           Lassar's         120           Library         446           Phosphorus         446           Polishing         448           for windows         468           Razor         451           Roach         451           Tooth         351–358           Wax         120           Pasteurine         232           Pastulles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         284           Tincture         282           Patent leather shoe polish         456	Pettit's eye salve.         33           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenotol         218           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur. quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Whilocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365           Phosphates, acid         366           Elixir of quinine and         71           Powder of, comp         123           Syrup of comp., ext. of malt with         99           Three, elixir of         61           Phosphatic emulsion         83           Mixture         83           Phospho-albumen         233           Phosphorated emulsion         84
Currant         107           Electric         451           Furniture         408           Gonorrhœa         188           Lassar's         120           Library         446           Phosphorus         444           Polishing         448           for windows         468           Razor         451           Tooth         351-358           Wax         120           Pasteurine         232           Pastulles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         284           Tincture         282           Patent leather shoe polish         456           Shoe varnish         456           Patterson's toothache wax         2440	Pettit's eye salve.         238           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenatol         218           Phenol camphor         11           Sodique         238           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur, quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365         367           Phosphates, acid         366         Elixir of quinine and         71           Powder of, comp.         123         Syrup of comp., ext. of malt with         99           Three, elixir of         61         Phosphatic emulsion         83           Mixture         83           Phospho-albumen         233           Phosphorated emulsion         84           Emulsion of cod liver oil         83           Phosphoric acid, solution of, comp.         129
Currant         107           Electric         451           Furniture         408           Gonorrhea         188           Lassar's         120           Library         444           Phosphorus         446           For windows         468           Razor         451           Roach         451           Tooth         351-358           Wax         120           Pasteurine         232           Pastulles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         284           Tincture         282           Patters of 's toothache wax         244C           Peach blossom extract         291	Pettit's eye salve.         238           Phalon's hair restorative         2440           Pheasants, gapes in.         278           Phenatol.         218           Phenol camphor.         11           Sodique.         238           Phenolid.         2440           Phenosolyl.         233           Phillips milk of magnesia.         2440           Phospho-mur. quinine.         233           Syr. wheat phosphates.         2440           Philocome.         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry.         365, 367           Phosphates, acid.         366           Elixir of quinine and.         71           Powder of, comp.         123           Syrup of comp., ext. of malt with.         99           Three, elixir of.         61           Phosphatic emulsion.         83           Phosphoraled emulsion.         84           Emulsion of cod liver oil.         83           Phosphoric acid, solution of, comp.         129           Phosphorus, cod liver oil with.         115
Currant         107           Electric         451           Furniture         408           Gonorrhea         188           Lassar's         120           Library         444           Phosphorus         446           Polishing         448           for windows         468           Razor         451           Roach         451           Tooth         351-358           Wax         120           Pasteurine         232           Pasteurine         232           Pasteurine         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         283           Tincture         282           Patent leather shoe polish         456           Shoe varnish         456           Patterson's toothache wax         244C           Peach blossom extract         291           Peacock's bromides         233	Pettit's eye salve.         23           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenatol         218           Phenol camphor.         11           Sodique         238           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur, quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365, 367           Phosphates, acid         366           Elixir of quinine and         71           Powder of, comp         123           Syrup of comp., ext. of malt with         99           Three, elixir of         61           Phosphatic emulsion         83           Phospho-albumen         233           Phosphorated emulsion         84           Emulsion of cod liver oil         83           Phosphoric acid, solution of, comp         129           Phosphorus, cod liver oil with         115           Elixirs contain
Currant         107           Electric         45           Furniture         408           Gonorrhea         188           Lassar's         120           Library         446           Phosphorus         446           Polishing         448           for windows         406           Razor         451           Roach         451           Tooth         351–358           Wax         120           Pasteurine         232           Pastulles, fly         406           Funigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         284           Tincture         282           Patent leather shoe polish         456           Shoe varnish         456           Patech blossom extract         291           Peacock's bromides         233           Pearl cream         326	Pettit's eye salve.         238           Phalon's hair restorative         2440           Phenatol         278           Phenol camphor         11           Sodique         238           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur, quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365           Phosphates, acid         366           Elixir of quinine and         71           Powder of, comp         123           Syrup of comp., ext. of malt with         99           Three, elixir of         61           Phosphotatic emulsion         83           Mixture         83           Phosphorated emulsion         84           Emulsion of cod liver oil         83           Phosphorus, cod liver oil with         115           Elixirs containing         69           Emulsion         84
Currant         107           Electric         451           Furniture         408           Gonorrhoea         188           Lassar's         120           Library         446           Phosphorus         444           Polishing         448           for windows         468           Razor         451           Roach         451           Tooth         351-358           Wax         120           Pasteurine         232           Pastilles, fly         406           Fumigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         284           Tincture         282           Patent leather shoe polish         456           Shoe varnish         456           Patterson's toothache wax         244C           Peacok's bromides         233           Pearl cream         326           Liquid         326	Pettit's eye salve.         238           Phalon's hair restorative         2440           Pheasants, gapes in         278           Phenotol         278           Phenol camphor         11           Sodique         238           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur, quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365         367           Phosphates, acid         366         Elixir of quinine and         71           Powder of, comp.         123         Syrup of comp., ext. of malt with         99           Three, elixir of         61         61           Phosphatic emulsion         83         Mixture         83           Phosphorablumen         233         Phosphorated emulsion         84           Emulsion of cod liver oil         83         Phosphorus, cod liver oil with         115           Elixirs containing         69         Emulsion         84
Currant         107           Electric         45           Furniture         408           Gonorrhea         188           Lassar's         120           Library         446           Phosphorus         446           Polishing         448           for windows         406           Razor         451           Roach         451           Tooth         351–358           Wax         120           Pasteurine         232           Pastulles, fly         406           Funigating         300           Mosquito         441           Pastor Kneipp's remedies         108           Patchouly extract         290           Sachet         297           Spirit of         284           Tincture         282           Patent leather shoe polish         456           Shoe varnish         456           Patech blossom extract         291           Peacock's bromides         233           Pearl cream         326	Pettit's eye salve.         238           Phalon's hair restorative         2440           Phenatol         278           Phenol camphor         11           Sodique         238           Phenolid         2440           Phenosolyl         233           Phillips milk of magnesia         2440           Phospho-mur, quinine         233           Syr. wheat phosphates         2440           Wheat phosphates         2440           Philocome         337           Phosphate, iron and quin., ext. malt, w.         100           Wild cherry         365           Phosphates, acid         366           Elixir of quinine and         71           Powder of, comp         123           Syrup of comp., ext. of malt with         99           Three, elixir of         61           Phosphotatic emulsion         83           Mixture         83           Phosphorated emulsion         84           Emulsion of cod liver oil         83           Phosphorus, cod liver oil with         115           Elixirs containing         69           Emulsion         84

Physic, Dow's; white liquid	120	Poke Root, tincture of, comp	156
Physostigmine, discs of	31	Polish, brass	381
Phytoline	234	Floor	
Pierce's comp. ext. swartweed	234	Furniture	
Favorite prescription	234	for glass, mirrors, etc	411
Medical discovery	234	Harness	414
Pigs, medicines for	268	Leather	
Pills		for linoleumfor patent leather	$\frac{439}{456}$
Ague	164		
Anti-chill	164	Shoefor silver	
Anti-grippe	120	Stove	
Anaphrodisiac	206	for windows	
Aphrodisiac		Polishing of marble	
Cathartic, little		Metals	
Cholera	120	Paste	
Damiana, comp	193	Powders	
Female, Hooper's	224	Polyform, Edison's	
Kneipp's	108		
Nervous debility	193	Pomades, flower	-341
Potass. iodide, Dyche's comp	217	Hongroise	350
Sandalwood	186	Hungarian mustache	350
Tonic, Aiken's	120	Juniper	121
Yellow	120	Lip	
Pilocarpine hair wash	347	Putz	
Pimento, spirit of	284	Pomatum, stick	349
Water	160	Pomegranate root bark (see Granatum).	
Pimple remedies		Ponca compound	235
Pinapin	234	Poppy seed oil, bleaching of	379
Pinaud's brilliantine	234	Porcelain, cement for	
Eau de quinine	234	Letters, cement for	
Pine, elixir of white, compound	69	Pot-pourris	299
Syrup of white, comp	149 199	Potassa cum calce	18- 135-
Tree tar lozenges	370	Solution of	.88
Pineapple syrupPink root, infusion of, comp	106	Acetate, solution of	135
Powder of, comp	123	Arsenite, elizir of	35
	100	Bromide, effervescent	
Fluid ext	202	Efferv., with caffeine	
Pinkham's vegetable comp	234	Compounds, elixirs containing	70
Pinus canadenses, fl. ext., non-alcoholic.	94	Citrate, effervescent	127
Piscidia, (see Jamaica dogwood).		Iodide bougies	16
Piso's consumption cure	235	Pills, Dyche's	217
Pitcher's castoria	235	Starch paper	466
Plant food	398	Permanganate solution	135
Insect exterminators	447	Sulphide ointment	118
Plant's asthma cigarettes	235	Potsdam balsam	235
Plantation bitters	167	Poultice, charcoal, Thompsonian	
Plaster, corn	176	Poultry food	275
Green	121	Medicines	
Paris bandages	14	Powders	
Plating with gold, silver, etc	447	Powder, entozoic	
Platinum plating	235	Nerve	109
Platt's chlorides	234	Thompsonian	199
Pleuricy root, powder of comp	123	Neutralizing	275
Pleurisy root, powder of, comp Podophyllin powder, comp	123	Poultry Red.	123
Podophyllum powder, comp	123	Talcum, salicylated	184
Tincture of	156	Powder, worm	
Poison ivy, remedy for poisoning by	205	Powell's balm of anise seed	235
Oak or ivy, tincture of	156	P. P. P	235
Poisoned wheat		Pozzoni's complexion powder	235
Poke ointment	118	Pray's nail polish	44C
Root, syrup of, comp.:		Preservative fluid	448
Tincture of	156	Preston salts	307

Prickly ash berries, tincture of	156	Rademacher's tincture carduus mariæ	15
	167	Tincture celandine	
"Protoxide" of iron, elixirs containing. 49,	62	Cochineal	15
Prunes, confection of	24	Copper acetate	15
Medicated	24	Colocynth seed	15
Ptelea, elixirs containing	78	Golden rod	
Pulsatilla, tincture of	196	Iron acetate	
Pulvis (see Powders).	122	Mugwort	15
	441	Nux vomica	15
	122	Shepherd's purse	16
	122		12
Vaccarum		Radway's pills	
Pungents		Ready relief	23
	423	Renovating resolvent	23
Putz pomade		Rainworm oil, artificial	
Tablets	450	Ransom's hive syrup and tolu	23
	235		19
Pyrmont water	372	Rapid relief	3
Pyrozone	235	Raspberry, elixir of, compound	73
	i		37
QUASSIA, elixir containing	70		37
water, Rademacher S	160	Rat exterminators	
C	101	Ratier's arsenical caustic	1
	235		20
Quillaja, tincture of	366	Razor paste	40
Quina-Laroche	250	Recamier preparations	20
Quince seed mucilage	100	Red bird food	16
Comp., elixirs containing70		Red clover blossoms, syr. comp	
Hair tonic		Blossoms, syrup of	49
and iron citrate, efferv	126	Lotion	11
and iron citrate, wine of	163	Oil	19
and iron iodides, syrup of		Powder	
Powder, comp		Spruce gum syrup	179
Salts, extemp. preparation of	450	Gum, tincture of	179
Sulphate, syrup of		Wash	11
Tincture of	156	Redlinger's pills	23
Ammoniated	157	Rennet essence	133
Wine of	163	Liquid	138
Quinquinia	235	Solution of; wine	13
Quionin	44C	Resin plaster, comp	12
Quitter oil	200	Stains, removing	12
R & H. three chlorides	236	Washes	2
	109	Resorbin	23
Radam's microbe killer			2
Radcliff's golden wonder remedy		Restorative cordial	
Seven seals	236	Syrup, Thompsonian	14
Rademacher's acorn water	159	Retorts, cement or lute for	38
Calamine ointment		Rheumatic drops	10
Castor water	160	Elixir	4
Copper mixture	113	Liniment	19
Diarrhœa mixture		Oil	190
Dysmenorrhœa drops	32	Tincture	103
Extract of tobacco			263
Iron mixture		in dogs and cats	214
Miraculous plaster		in horses	100
Nephritic powder		Remedies	190
Nux vomica water		in sheep	114
Ointment of shepherd's purse		Rhubarb, confection of, compound	2
Preparations, list ofQuassia water	160	Flixir containing	7
Solution calcium chloride	131	Extracts of, comp	
	135	Fluid, arom	10

Rhubarb and potassa, syrup of, comp	147	SABADILLA, vinegar of	159
Powder, comp		Saccharated citric acid	124
Syrup of, aromatic101,		Extracts	101
Tincture of, comp	157	Sodium bicarbonate	124
Ricard's urethral bougies	236	Tartaric acid	124
Richmond's Samaritan nervine2	44C	Saccharin, elixir of Essence of, 85; solution of	73
Richter's pain expeller	236	Essence of, 85; solution of	13
Riga balsam, factitious	13	Syrup of	147
Ringbone cures	257	Sachet powders294	-299
Ringworm in cattle	264	Heliotrope	
Remedies	197	Saddle gall	203
Roach food, Peterman's		Saffron, tincture of	
Paste	451	Sage, infusion of	.100
Powder	401	Infusion, compound	100
Debineen's clivin of paraldehyde	926	Sage's catarrh remedy	201
Robinson's elixir of paraldehyde		Sal alembroth cotton	28
Roche's embrocation	100	Marinum (see Sea salt).	111
Rocky mountain liniment	201	Muscatel	
Sachet		Salicyated cold cream	311
Roob juniperi	107	Cotton	28
Sambuci		Powder of talcum	
Root beer.	365	Suet	139
Extract.		Salicylates, elixirs containing73,	
Rose cerate		Salicyl bromanilid	
Cold cream	312	Salicylic acid, elixirs containing	
Geranium extract		Emulsion of	
Hair oil	337	Honey of rose with	
Hair oil	105	Carbolated collodion	21
with salicylic acid	105	Hair tonic.	346
with tannic acid		Salol collodion	
Jelly		Salt of lemon, artificial	
Moss, extract	291	Rheum remedies	183
Musk, extract	291	Sea, artificial	-128
Sachet	298	Salts, effervescent	-128
Spirit of	284	Inexhaustible	307
Compound		Preston	307
Tea, extract		Smelling	
White, extract		Violet	
Yellow, extract	292	Salubrin	
Rose's metal	950	Salves	
Roses, elixir of	110	Salve, corn	
Spirit of	138	Eye	
Compound		Lip.	
Roseter's hair regenerator		Mother	
Rotterin	237	Thompson's	
Rouges	330	Tulip	
Roup remedy		Sambucus (see Elder).	
Rourke's iodine liniment	237	Sandal, spirit of	284
Royal catarrh cure	237	Sandalwood pills	
Germeteur		Sanguinaria (see Bloodroot).	
Rubber cement		Sanguis bovinus inspissatus	14
Shoe cement	- 1	Sanitas	
Stamp ink		Sanmetto	
Rubefacient sol. of iodine, Lugol's	132	Santonin, troches of	
Rubifoam	237	Sarah Bernhardt's face powder	
Rubini's diarrhœa mixture		Sarsaparilla	168
Rubus (see Blackberry).		Sarsaparilla essence	100
Rumex (see Yellow Dock).		Extract of	
Ruppert's face bleach	237		
Ruschenberger's diarrhœa mixture		Fluid	
Russet shoe dressing.		Tea	163
Rust stains, removal of	452	Saul's catarrh remedy	237

Saunder's bloom of Ninon		243
Savin, tincture of	Shoe blacking &	379
Tincture, comp	7 Dressings	157
Saw palmetto, comp	Grease 4	456
Saxon species	Shoulder lameness in cattle	263
Tea	in horses 2	257
Scabies in horses	Show-globe colors	160
Scalds, applications for	Siccative	394
Scalp, preparations for 33	I   Silk, dyeing S	395
Scammon's diarrhœa mixture 11		161
Scarlet ink		128
Scheele's hydrocyanic acid	l   Silver ink 4	124
Schenck's pulmonic syrup 23	Nitrate bougies	16
Schinseng elixir	8   Plating 4	147
Schlotterbeck's helonin mixture244	Polishing of 4	160
Hydrastis mixture	Simmons' liver regulator	239
Schneeberger snuff	Simple elixir	74
School ink		79
Schreyer's toothache pellets 23	S Skin cream 3	313
Schreyer's toothache pellets	Food 8	313
Scott's emulsion cod liver oil 23	Preparations 3	309
Scrofulous syrup	9 Skunk cabbage, tincture of	157
Sea breeze extract	0 Oil	116
Foam liquids 33	[5] Slating, blackboard 3	378
Salt, artificial	Sloan's condition powder	239
Sealing wax	3 Smelling salts 3	307
Searle's athlophoros	8 Smith's (Mrs.) butter color	339
Sedative cordial and elixir 4	Electric oil, tonic syrup	239
	Snuffs	29
Seiler's antiseptic solution 15		171
Seltzer, lemonade 36		171
Water 37	Soap, balsam of	13
Senega, extract of		157
Senna, elixirs containing		19
Ext., fl. aqueous		161
Fluid comp 10		333
Infusion of 10		189
and jalap, fl. ext		$\frac{160}{160}$
and manna, syrup of		168
and pink root, fl. ext	-	161
and pink root, fl. ext., comp 20		
Powder, comp		
Tincture of, comp		138
Sero-sublimate cotton		461
Serpentaria, tincture of, comp 15		356
Seven Sutherland's hair grower244	Whale oil	161
Scalp cleaner	Soapine	243
Sewing machine oil		366
61		113
Shampoo cream	_	135
Egg, jelly		367
Liquids		363
Paste33		124
Powders	and calcium phosph. em. c. l. oil with.	83
Shaving cream		111
Powder		127
Soap		74
Sheep dips		135
Laurel, tincture of		127
Medicines		$\frac{88}{388}$
Sheffield's dentifrice		386 127
Shepherds purse ointment, Rademacher's 11		161.
Tincture, Rademacher's		
Sherbet syrup	O Solid perfumes	ະອອ 135

Solutions	135	Stamping powders	. 464
Solution of acid phosphates129,		Starch cement	. 386
Aloes and soda, Mettauers	129	Glycerite of	. 105
Chloro-phosphide arsenic	239	lodide, syrup of	. 147
Citric acid	366	lodized	. 138
Iodine carbolized		Mucilage of	. 114
Iodo-brom. calcium comp	241	Paper	
Phosphoric acid, comp		Starton's mixture	
for storm glass		Stavesacre, tincture of	
Tri-iodides	223	Stearin soap	. 461
Solutol	14D	Stains, removing	. 462
Solveol		C44	0445
Somnal	107	Steedman's soothing powders	
Soothing syrup	949	Steel, etching of	. <u>ə</u> yə
Winslow's	169	Stencil ink	. 432 . 292
Soup stains, removing	242	Stephanotis extract	
Sozodont		Stick mucilage	
Cures	257	Pomades	
Foot, in horses.	258	Sticky fly paper	
Species	136	Stillingia, elixirs containing	. 76
Althæa	136	Liniment, comp	. 110
ad longam vitam		Tincture of	
	301	Stock food.	
ad gargarisma		Stoddart's peerless liquid	. 240
Guaiac, comp.		Stomach bitters	
ad infusum lignorum		Catarrh in cattle	. 263
Laxantes Schrammii		in dogs and cats	
Lignorum		in horses	. 258
Moth		Consoler, Kneipp's	. 108
Quassiæ amaræ		Drops, bitter	
Resolventes		Inflammation in cattle	. 263
	84		
Spermaceti, emulsion of	84	Stomachic elixir	. 66
	84	Stomachic elixir	. 66 . 138
Spermaceti, emulsion of	84 187 116	Stomachic elixir	. 66 . 138 . 285
Spermaceti, emulsion of	84 187 116	Stomachic elixir. Stone, medicinal. Storax, tincture of.	. 66 . 138 . 285 . 464
Spermaceti, emulsion of	84 187 116	Stomachic elixir. Stone, medicinal. Storax, tincture of. Storm-glass solution.	. 66 . 138 . 285 . 464 . 240 . 464
Spermaceti, emulsion of. Spermatorrhæa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp Spirits	84 187 116 140 284 137	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish	. 66 . 138 . 285 . 464 . 240 . 464
Spermaceti, emulsion of. Spermatorrhæa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic	84 187 116 140 284 137 137	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses.	. 66 . 138 . 285 . 464 . 240 . 464 . 464
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm.	84 187 116 140 284 137 137	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses. Stramonium, infused oil of	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite.	84 187 116 140 284 137 137 137	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses. Stramonium, infused oil of Strangles	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253
Spermaceti, emulsion of. Spermatorrhea. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits	84 187 116 140 284 137 137 137 137 366	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of Strangles Stratena.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242
Spermaceti, emulsion of. Spermatorrhea. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite Nutmeg Spiritus balsamicus	84 187 116 140 284 137 137 137 137 1366 12	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of Strangles Stratena.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum	84 187 116 140 284 137 137 137 366 12 137	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena Stratena Strawberry syrup Strengthening syrup, Thompsonian.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial	84 187 116 140 284 137 137 137 136 12 137 138	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup. Strengthening syrup, Thompsonian Strong's arnica jelly.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 240
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized.	84 187 116 140 284 137 137 137 137 137 137 138 138	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup. Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 240 . 119
Spermaceti, emulsion of. Spermatorrhea. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits	84 187 116 140 284 137 137 137 137 137 137 138 12 138 138 167	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup. Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 240 . 119
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial. Carbolized. Spring bitters. Flowers extract	84 187 116 140 284 137 137 137 137 137 138 138 167 292	Stomachic elixir Stone, medicinal Storax, tincture of. Storam, glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena Stratena Strawberry syrup Strengthening syrup, Thompsonian. Strong's arnica jelly Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 240 . 119
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial. Carbolized. Spring bitters. Flowers extract Springseen's Uterine Capsules. 24	84 187 116 140 284 137 137 137 366 12 137 138 138 167 292 4D	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup Strengthening syrup, Thompsonian Strong's arnica jelly. Strychnine, oleate of Sulphate, syrup of. Acetate, solution of Tincture of, compound	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 148 . 158
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup	84 187 116 140 284 137 137 137 366 12 137 138 138 167 292 44D 179	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound. Valerianate, elixir of.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 240 . 119 . 135 . 158
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red.	84 187 116 140 284 137 137 137 136 12 137 138 167 292 44D 179	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup. Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound. Valerianate, elixir of.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 135 . 158 . 76
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing.	84 187 116 140 284 137 137 137 136 12 137 138 167 292 44D 179 56	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses. Stramonium, infused oil of Strangles Stratena Strawberry syrup Strengthening syrup, Thompsonian Strong's arnica jelly Strychnine, oleate of Sulphate, syrup of Acetate, solution of Tincture of, compound Valerianate, elixir of Styptic cotton Powder	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 135 . 158 . 76
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial. Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing Vine, syrup of, comp.	84 187 116 140 284 137 137 137 366 12 137 138 138 167 1292 44D 179 56 146	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup Strengthening syrup, Thompsonian Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound Valerianate, elixir of. Styptic cotton. Powder Sublimated (see Mercuric chloride).	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 240 . 119 . 148 . 158 . 76 . 293 . 123
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial. Carbolized. Spring bitters. Flowers extract. Springsteen's Uterine Capsules. Squaw vine, elixirs containing Vine, syrup of, comp. Squibb's cholera mixture.	84 187 116 140 284 137 137 137 366 12 137 138 138 167 292 44D 179 56 146 113	Stomachic elixir Stone, medicinal Storax, tincture of. Storamylass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound. Valerianate, elixir of. Styptic cotton. Powder. Sublimated (see Mercuric chloride). Succinic spirit of ammonia.	. 666 . 138 . 285 . 464 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 135 . 158 . 76 . 293 . 123 . 123
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing Vine, syrup of, comp. Squibb's cholera mixture Squill, extract of.	84 187 116 140 137 137 137 137 137 137 137 137 138 138 140 179 179 179 146 113 103	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound Valerianate, elixir of. Styptic cotton Powder Sublimated (see Mercuric chloride). Succinic spirit of ammonia. Alterans.	. 666 . 138 . 285 . 464 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 135 . 158 . 76 . 293 . 123 . 123
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing Vine, syrup of, comp. Squibb's cholera mixture. Squill, extract of St. Germain tea	84 187 116 140 137 137 137 137 137 138 138 167 292 44D 179 56 146 113 103 136	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena Strawberry syrup Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound Valerianate, elixir of. Styptic cotton. Powder Sublimated (see Mercuric chloride). Succinic spirit of ammonia. Alterans. Succus (see Juices).	. 66 . 138 . 285 . 464 . 240 . 464 . 464 . 257 . 370 . 148 . 135 . 158 . 76 . 29 . 123 
Spermaceti, emulsion of. Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial. Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing. Vine, syrup of, comp. Squibb's cholera mixture. Squill, extract of. St. Germain tea St. Jacob's oil.	84 187 116 140 284 137 137 137 137 138 138 138 167 179 179 56 113 103 136 240	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup. Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of Tincture of, compound Valerianate, elixir of Styptic cotton. Powder Sublimated (see Mercuric chloride). Succinic spirit of ammonia. Alterans. Succus (see Juices). Sudorific tincture.	. 66 . 138 . 285 . 464 . 240 . 464 . 464 . 257 . 116 . 253 . 370 . 148 . 240 . 119 . 148 . 135 . 158 . 76 . 29 . 29 . 29 . 123 . 137 . 29 . 29 . 29 . 29 . 29 . 29 . 29 . 29
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial. Carbolized. Spring bitters. Flowers extract. Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing. Vine, syrup of, comp. Squibb's cholera mixture. Squill, extract of St. Germain tea St. Jacob's oil. St. Peter's drops.	84 187 116 140 284 137 137 137 137 366 12 137 138 167 292 241 179 156 146 113 103 103 103 103 103 103 103 103 103	Stomachic elixir Stone, medicinal Storax, tincture of. Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup Strengthening syrup, Thompsonian Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound. Valerianate, elixir of. Styptic cotton. Powder Sublimated (see Mercuric chloride). Succinic spirit of ammonia. Alterans. Succus (see Juices). Sudorific tincture. Suet.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 240 . 119 . 123 . 123 . 123 . 129 . 123 . 137 . 229
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing Vine, syrup of, comp. Squibb's cholera mixture Squill, extract of St. Germain tea St. Jacob's oil. St. Peter's drops. Staining of gun barrels.	84 187 116 140 284 137 137 137 138 167 292 44D 179 179 146 113 103 136 146 113 146 146 113 146 146 147 147 148 149 149 149 149 149 149 149 149 149 149	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound Valerianate, elixir of. Styptic cotton Powder Sublimated (see Mercuric chloride). Succinic spirit of ammonia. Alterans. Succus (see Juices). Sudorific tincture. Suet Benzoated	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 370 . 148 . 240 . 119 . 148 . 135 . 76 . 29 . 123 . 123 . 157 . 229 . 137 . 219 . 138 . 139 . 1
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing Vine, syrup of, comp. Squibb's cholera mixture Squill, extract of St. Germain tea St. Jacob's oil. St. Peter's drops. Staining of gun barrels. Stains from fabrics, removing.	84 187 116 140 284 137 137 137 137 137 138 138 167 292 44D 179 56 113 103 136 113 136 141 141 143 143 143 143 143 143 143 143	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena Strawberry syrup Strengthening syrup, Thompsonian. Strong's arnica jelly Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound Valerianate, elixir of. Styptic cotton. Powder Sublimated (see Mercuric chloride). Succinic spirit of ammonia. Alterans. Succus (see Juices). Sudorific tincture. Suet. Benzoated Deer.	. 666 . 138 . 285 . 464 . 240 . 464 . 240 . 257 . 116 . 253 . 242 . 240 . 119 . 148 . 240 . 129 . 123 . 157 . 229 . 157 . 138 . 139 . 139 . 139
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial. Carbolized. Spring bitters. Flowers extract. Springsteen's Uterine Capsules. Sprince gum syrup Tincture of, red. Squaw vine, elixirs containing Vine, syrup of, comp. Squibb's cholera mixture. Squill, extract of St. Germain tea St. Jacob's oil. St. Peter's drops. Staining of gun barrels. Stains from fabrics, removing Ink, to remove. 427, Rust. removal of	84 187 116 140 284 137 137 137 137 137 137 138 138 167 12 12 139 140 179 179 166 113 103 136 136 136 146 113 136 136 136 136 136 136 136 136 13	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish. Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena. Strawberry syrup. Strengthening syrup, Thompsonian. Strong's arnica jelly. Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of Tincture of, compound Valerianate, elixir of Styptic cotton Powder Sublimated (see Mercuric chloride). Succinic spirit of ammonia. Alterans. Succus (see Juices). Sudorific tincture. Suet Benzoated Deer. Salicylated	. 66 . 138 . 285 . 464 . 240 . 464 . 253 . 257 . 116 . 253 . 370 . 148 . 240 . 119 . 148 . 135 . 158 . 76 . 29 . 123 . 137 . 138 . 139 . 139 . 139 . 139
Spermaceti, emulsion of Spermatorrhœa. Spigelia (see Pink root). Spike, oil of, factitious. Spikenard, syrup of, comp. Spirits. Spirit of ants. Aromatic Balm. Carmelite. Nutmeg Spiritus balsamicus Formicarum Sponge, burnt, artificial Carbolized. Spring bitters. Flowers extract Springsteen's Uterine Capsules. 24 Spruce gum syrup Tincture of, red. Squaw vine, elixirs containing Vine, syrup of, comp. Squibb's cholera mixture Squill, extract of St. Germain tea St. Jacob's oil. St. Peter's drops. Staining of gun barrels. Stains from fabrics, removing.	84 187 116 140 284 137 137 137 137 137 137 138 138 167 12 12 139 140 179 179 166 113 103 136 136 136 146 113 136 136 136 136 136 136 136 136 13	Stomachic elixir Stone, medicinal Storax, tincture of. Storm-glass solution Stoughton's bitters. Stove blacking Polish Strained sinews in horses. Stramonium, infused oil of. Strangles Stratena Strawberry syrup Strengthening syrup, Thompsonian. Strong's arnica jelly Strychnine, oleate of. Sulphate, syrup of. Acetate, solution of. Tincture of, compound Valerianate, elixir of. Styptic cotton. Powder Sublimated (see Mercuric chloride). Succinic spirit of ammonia. Alterans. Succus (see Juices). Sudorific tincture. Suet. Benzoated Deer.	. 66 . 138 . 285 . 464 . 240 . 464 . 257 . 116 . 253 . 242 . 370 . 148 . 240 . 149 . 135 . 158 . 76 . 29 . 123 . 137 . 229 . 157 . 138 . 139 . 139 . 139 . 139 . 139 . 139

Sugars, oil	1	139	Syrup, phenic acid, Declat's	
Sulphocarbolic acid, crude		11 14	Phosphates, ext. malt with	99
Sulphur, balsam of			Red clover blossoms148,	240
Baths	٠ و	24	Compound	100
Iodide ointment		19	Rhubarb, aromatic	$\frac{101}{149}$
and lead hair promoters		345	Rumex, comp	168
Lotion of, comp		11	Sarsaparilla Scrofulous	
Ointment, alkaline			Simple	
Compound	1 1	19	Soda	
Paste, Hebra's		329	Soothing	
Powder, comp	1	23	Spikenard, comp	
Tincture of		58	Spruce gum	
Homeopathic		.58	Squaw vine. comp	146
Sulphurated oil		14	Tar hoarhound cough	177
Sulphurous fumigation	3	93	Tar and wild cherry	
Sultana cold cream		312	Teething	177
Sumbul, elixirs containing		77	Tolu	103
Extract of, fl		.03	Trifolium, comp	
Sun cholera mixture	1	13	Worm	202
Superphospate of iron, syrup of	ق	328	Syrupus capilli minoris	145
Superphospate of iron, syrup of	···· 1	48	Cerasorum	141
Suppositories, pile	1	14	Coccionellæ	141
Urethral		$\begin{vmatrix} 14 \\ 240 \end{vmatrix}$	Mannatus	
Svapnia		240	Opiatus	
Swayne's ointment		240	Roborans	
Swedish bitters		64	Spinæ cervinæ	140
Blood purifying tea	1	135	Spine continuent in the second	110
Sweet briar extract			TABLEAU lights	465
Briar sachet			Tablets, antiseptic, Wilson's	
Flag, candied		23	Bath	
Gum, syrup of		148	Complexion	324
Pea extract		$292^{\circ}$	Disinfectant	
Quinine	5	241	Dyspepsia	182
Swift's specific	$\dots 24$	14D	Lithia, comp	216
Swine fever	.269, 7	3.10	Putz	
Medicines			Tamar Indien	
Plague			Tamarind, confection of	
Sydenham's laudanum	• • • • • •	190	Tamarinds, essence of	85 184
Sympathetic ink	• • • • • • •	419	Talcum, salicylated powder of	190
Syphilis remedies	• • • • • •	198	Tan, removal of	308
Syrups	367_	371	Shoe dressing	
Syrup of absinthium		149	Tannic acid, honey of rose with	105
Syrup of absinthium		164	and boric acids, glyc. of	104
Alterative		140	Tannin bougies	
Blackberry, aromatic		87	Carbolated cotton	29
Bromides	9	233	Ink417,	418
Coloring		363	Stains, removing	463
Cough			Tapeworm in sheep	268
Elder berry			Remedies	208
Eriodectyon, aromatic			Tar, elixirs containing	
Figs		240	Fumigation	
Flaxseed cough		179	Hoarhound cough syrup	1//
Frangula		140	and hoarhound, honey of	106
Gibert's		210	Stains, removing	
Iron chloride, Weld's		244	Syrup of, comp	
Iron tonic			Tolu and wild cherry	
Juniper berry			Water	
Lactucarium		209	and wild cherry syrup	177
Linseed cough			and wild cherry syrup	53
Manna, comp			Tarrant's comp. extract	241
.Partridge berry, comp			Seltzer aperient	241
artifuge berry, comp				

Tartar emetic ointment		264
Tartaric acid, saccharated	6,	258
Syrup of	, , , , , , , , , , , , , , , , , , , ,	35
Tasteless cascarine	1	190
Chill or ague cure		$\frac{138}{241}$
Fl. ext. of buckthorn		21
Taylor's lotion		111
Tea, aromatic		29
Bitter	5   Thymolyptol	241
Blood-purifying	0 Tilden's bromo-chloralum	241
Kneipp's 10	8 Chloralum	241
Swedish	Elixir iodo-bromide calcium comp	241
Boecker's bitter tonic		241
Breast		111
Cathartic 17		398
Cough, Kneipp's		439
Diuretic		334
		151 31
Garfield		157
Marshmallow		153
Rose extract		
St. Germain		31
Sarsaparilla		155
Saxon		155
Syrup 37	0 Virgæ aureæ	153
Wood	6 Tinctures	
Worm		158
Teething powders		64
Syrup 19	and the state of t	70
Terpin hydrate, elixir of		155
Emulsion of	4	$\frac{156}{152}$
Test paper		151
Tetlow's swan down		157
Tetter in cattle		151
Ointment, Heiskell's		302
Thapsia plaster		162
Theobroma, tincture of	_ 1	433
Theriac	'-	
Thielemann's cholera mixture	8 1	
Third preparation (Thompsonian) 15		
Thomas' electric oil		
Thompsonian anti-canker pills 12		440
Astringent ointment		154
Bitters		
Camphor julep 1		
Charcoal poultice		282
Confection of hollyhock		
Cough powder	1 8	
Nerve powder		109
Remedies, list of		154
		151
Strengthening syrup		358
Garlic		153
Lobelia		364
Thompson's ointment		158
		103
Emulsion of linseed oil 8	Flavor for	402
Eye water		
Three chlorides, R. & H		119
Throat affections, remedies for 19	8 Water, Rademacher's	101

Tobias' derby condition powders 241	Tyson's antimonial powder 121
Venetian liniment	
Toenail, ingrowing 466	UDDER, inflamed 264
Toilet cream313, 314	Ulyptol
Jelly 318	Uncle Sam's nerve and bone liniment 242
Lanolin	Unguenta (see Ointments).
Lotions	Unguentum bursæ pastoris
Milk	
Powders 324	Nervinum
Preparations	Universal balsam
Vinegars	Upham's asthma remedy
Waters	Financial
Tolu balsam, emulsion of	Upper-ten extract
Fluid ext. of 103 Rock and rye 241	Uricedin
Syrup of	Urinal cakes, Davy's
Tincture of	Urinary diseases in dogs and cats 274
Water	Diseases in horses
Water	in sheep
Tongaline 242	Urine, bloody, in cattle 264
Tonics	Uva ursi, fl. exts. containing 88
Tonic for birds 278	Syrup of, comp
Aphrodisiac tablets	Syrup si, competition
Bitters	VALERIAN, extract of
Calisaya	Syrup of, comp
for dogs	Tincture of, ethereal
Java 371	Valerianates, elixirs containing77, 78
Mixture, comp 113	Vanadium ink
Tea, Boecker's bitter	Van Buskirk's sozodont
Tonka, tincture of	Van Stan's stratena 243
Tooth cream	Vanilla, extract of 364
Pastes, powders, etc351, 358	Sugar
Wash	Syrup 371
Toothache drops, wax and gum 200	Tincture of
Tracing paper	Vanillin sugar
Transfusion fluids	Varnishes
Trask's magnetic ointment244D	for patent leather
Traumaticin	Shoe 456
Trifolium, syrup of, comp 148	Varnish stains, removing462, 468
Tritica244D	Vaseline (see Petrolatum).
Triticum (see Couch grass).	Vegetable elixir
Trix244D	Velpeau's black caustic 18
Troches, bronchial	Venice turpentine, factitious
Cough	
Santonin	202
Tropic fruit laxative	Vermifuges
Trousseau's diuretic wine	320
Tulio selve	
Tulip salve	224
Turkey corn, tincture of	, rotaront, opinion or recording to
	Viburnai
	Opulus, elixir of, comp
Turpentine, balsam of	Oparas, cimir or, competition
Liniment, camphorated	Vichy salt, artificial efferv
	100
Acid	The state of the s
Syrup of	202
Venice, factitious.,	Vienna caustic
Tuson's lotion of lead chloride 111	Vin Mariana
Twaddell's scale	206
	Vinaigre de toilette 306
1 wigg's (Gen.) hair restorer	Vinegars
Twigg's (Gen.) hair restorer.         345           Typewriter ink.         432	Vinaigre de toilette       300         Vinegars       159         Vinegar, aromatic       306

	306	Water, I avender; lilac	305
Fumigating	302	Mineral, artificial371	-373
77,8	306	Pepper, tincture of	158
	$371 \mid 380 \mid$	Pyrmont	
	463	Seltzer	
Status, removing transfer	306	Tar	
	306	ToiletVichy	
Vinum liquiritiæ thebaicum		Violet	
	309	White rose	
	302	Watt's anti-rheumatic pills	
	293	Wax, emulsion of	
	252	Floor	
	425	Grafting	
Jelly	319	Hungarian mustache	
Parmese, violet	293	Paste	
	298	Sealing	452
	308	Stains, removing	
	305	Toothache	200
	293	Yellow, filtered	161
	371	Waxed paper	
8	329	Wayne's diuretic elixir	
Vitalizer		Weather-glass solution	
Vitalizing tonic	$\frac{195}{275}$	Weber's transfusion fluid	
	$\frac{275}{271}$	Wei de Meyer's catarrh cure	
in swine	211	Weinman's dental anæsthetic	~
WAFER ash, elixirs containing	78	Weld's syrup of chloride of iron West-end extract	00.0
	163	Whale oil soap	
	166	Wheat, poisoned450,	451
Elixirs containing.	78	Wheeler's elixir	2441)
	103	White drink	
	158	Hellebore, ointment of	
	371	Ink	
Hair oil or dye	549	Lily extract	
	464	Linctus	
	243	Liniment	
Wampole's Laxative Compound 24		Lotion	
±	$egin{array}{c} 243 \ 243 \ \end{array}$	Oil	191
The Paris of the P	31	Pine, elixir of, comp	$\begin{array}{c} 69 \\ 149 \end{array}$
Warner's cordial drops	243	Syrup of, comp	
	204	Pond lily extract	
	111	Extract	
	358	Water	
	111	Wash	
	354	Whiteley's nipple wash	
White	111	Whooping cough remedies	
Washing of pomades	285	Wickersheimer's preservative fluid	448
Washing powders243,	468	Widow McPhelan extract	293
Wassersuchtthee, Kneipp's	108	Wiener kalk	
Waters		Wild cherry bitters	
Water, barley	29	Elixirs containing	78
Bretfeld		Emuls. cod liver oil with	
Heliotrope		and hoarhound, syrup of	
Cologne Congress	371	Phosphate	
	330	Compound	
	$\frac{329}{329}$	and tar syrup	
	216	Flowers sachet.	
Eye.		Indigo, ointment of	
Florida		Olive extract	293
Friedrichshall	371	Wilson's antiseptic tablets	150
Geranium 305; Hunyadi janos	0~0	7.	110
Kissingen	312	Zinc ointment	

Wind wasser	160	Worms in cattle	265
Window polishing paste 4	168	in dogs and cats	275
Wines	163	in horses	259
Wine, bitter	66	in swine	271
	162	Tape, in sheep	268
	163	Wormwood, extract of	104
	243	Infused oil of	116
0000, 1.1	243	Syrup of	149
Cod in con, the conference of the contract of	162	Tincture of	158
200000000000000000000000000000000000000	163	Compound	158
12014	156	Wine of	163
	66	Wrapping paper, to cut	446
Orange, comp	135	Wright's face powder	244
	463	Writing ink	
5		Writing ink	416
	162	Wuehlhuberthee, Kneipp's	108
	159	VARROW, extract of	104
	244	Yellow dock, fl. ext., comp	104
Wintergreen, syrup of	149	Syrup of, comp	149
Wistar's balsam of wild cherry		Ointment	118
Cough lozenges24		Pills	120
	158	Rose extract.	292
	312	Yerba santa, elixir containing78,	79
	312	Extract of malt with	100
Jelly :	318	Syrup of	149
Ointment	119	Aromatic	150
Vinegar E	306		293
Wither's antizymotic solution	244	Ylang ylang extract	$\frac{298}{298}$
Wizard liniment or balm	191	Sachet	284
Oil, Hamlin's	223	Spirit of	204
Woman's friend	185	ZINC acetate pills, Rademacher's	121
Wood species 1	136	Carbonate, cerate of	18
	464	Chloride cotton	29
	136	Compounds, elixirs containing	79
	293	Etching of	398
	374	Ointment, Wilson's	119
	244	Oxide ointment, comp	119
	395	Sulphate ointment	119
	461	Zittmann's decoction30,	31
Worm lozenges; powders; syrup		Zittinann's decoction	-91
Tea 1		Zymoidin	MATA
	LUU	Zymorum,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	.771)





Mont

.

